

ANNALS OF MATHEMATICS

Vacant set of random interlacements and percolation

By ALAIN-SOL SZNITMAN



SECOND SERIES, VOL. 171, NO. 3

May, 2010

ANMAAH

Vacant set of random interlacements and percolation

By ALAIN-SOL SZNITMAN

Abstract

We introduce a model of random interlacements made of a countable collection of doubly infinite paths on \mathbb{Z}^d , $d \geq 3$. A nonnegative parameter u measures how many trajectories enter the picture. This model describes in the large N limit the microscopic structure in the bulk, which arises when considering the disconnection time of a discrete cylinder $(\mathbb{Z}/N\mathbb{Z})^{d-1} \times \mathbb{Z}$ by simple random walk, or the set of points visited by simple random walk on the discrete torus $(\mathbb{Z}/N\mathbb{Z})^d$ at times of order uN^d . In particular we study the percolative properties of the vacant set left by the interlacement at level u , which is an infinite connected translation invariant random subset of \mathbb{Z}^d . We introduce a critical value u_* such that the vacant set percolates for $u < u_*$ and does not percolate for $u > u_*$. Our main results show that u_* is finite when $d \geq 3$ and strictly positive when $d \geq 7$.

0. Introduction

This article introduces a model of random interlacements consisting of a countable collection of doubly infinite trajectories on \mathbb{Z}^d , $d \geq 3$. A certain nonnegative parameter u governs the amount of trajectories which enter the picture. The union of the supports of these trajectories defines the interlacement at level u . It is an infinite connected translation invariant random subset of \mathbb{Z}^d . Our main purpose is to study whether this random “fabric” is “rainproof” or not, i.e. whether its complement, the vacant set at level u , does not, or does contain an infinite connected component. This issue is related to the broad question “how can random-walk paths create interfaces in high dimension?”. The model we construct has a special interest because in a heuristic sense it offers a microscopic description of the “texture in the bulk” for two problems related to this broad question. One problem pertains to the percolative properties in the large N limit of the vacant set left on the discrete torus $(\mathbb{Z}/N\mathbb{Z})^d$, $d \geq 3$, by the trajectory of simple random walk with uniformly

distributed starting point, up to times that are proportional to the number of sites of the discrete torus; cf. [2]. The other problem pertains to the large N behavior of the disconnection time of a discrete cylinder $(\mathbb{Z}/N\mathbb{Z})^{d-1} \times \mathbb{Z}$, $d \geq 3$, by simple random walk; cf. [6], [7], and also [19]. In this work we establish a phase transition: for $u < u_*$, the vacant set at level u does percolate, whereas for $u > u_*$, it does not. The critical value u_* is shown to be nondegenerate (i.e. positive and finite), when $d \geq 7$, and finite for all $d \geq 3$. The results presented here have triggered some progress on the questions mentioned above; see in particular [15], [20], [21], [22], and [25].

We now describe the model. We consider the spaces W_+ and W of infinite, respectively doubly infinite, nearest neighbor paths on \mathbb{Z}^d , $d \geq 3$, that spend finite time in bounded subsets of \mathbb{Z}^d . We denote with P_x , $x \in \mathbb{Z}^d$, the law on W_+ of simple random walk starting at x . This is meaningful since the walk is transient in view of the assumption $d \geq 3$. We write $X_n, n \geq 0$, or $X_n, n \in \mathbb{Z}$, for the canonical coordinates on W_+ , or on W . We also consider the set of doubly infinite trajectories modulo time-shifts

$$(0.1) \quad W^* = W / \sim, \text{ where } w \sim w', \text{ if } w(\cdot) = w'(\cdot + k), \text{ for some } k \in \mathbb{Z}.$$

We denote with $\pi^* : W \rightarrow W^*$, the canonical projection.

The random interlacements are governed by a Poisson point process $\omega = \sum_{i \geq 0} \delta_{(w_i^*, u_i)}$ on $W^* \times \mathbb{R}_+$, with intensity measure $\nu(dw^*)du$, where ν is a certain σ -finite measure on W^* , which we now describe. For any finite subset K of \mathbb{Z}^d , we denote with e_K the equilibrium measure of K ; see (1.6) for the definition, with W_K^0 the subset of W of trajectories entering K at time 0:

$$(0.2) \quad W_K^0 = \{w \in W; w(0) \in K \text{ and } w(n) \notin K, \text{ for all } n < 0\},$$

and with $W_K^* = \pi^*(W_K^0)$ the subset of W^* of equivalence classes of trajectories entering K . We show in Theorem 1.1 that there is a unique σ -finite measure ν on W^* such that

$$(0.3) \quad 1_{W_K^*} \nu = \pi^* \circ Q_K, \text{ for any finite subset } K \text{ of } \mathbb{Z}^d,$$

where Q_K is the finite measure supported on W_K^0 such that

- (0.4) i) $Q_K(X_0 \in \cdot) = e_K(\cdot)$,
- ii) when $e_K(x) > 0$, conditionally on $X_0 = x$, $(X_n)_{n \geq 0}$, and $(X_{-n})_{n \geq 0}$ are independent with respective distributions P_x and P_x conditioned on $\{X_n \notin K, \text{ for all } n \geq 1\}$.

The motivation for such a requirement stems from Theorem 3.1 and (3.13) of [2], where the large N limit of certain suitably defined excursions to a box of size $L \ll N$, by simple random walk on $(\mathbb{Z}/N\mathbb{Z})^d$ was investigated, and from the

alternative characterization of Q_K given in (1.26); see also Remarks 1.2 (2) and 1.6 (3). Similar measures appear in [24] and [16, p. 61], following an outline in [10]. The construction we give here bypasses projective limit arguments: we instead glue together expressions for ν read in “local charts”.

We denote with Ω the canonical space where ω varies, cf. (1.16), and with \mathbb{P} the law turning ω into a Poisson point process with intensity $\nu(dw^*)du$. The law \mathbb{P} enjoys a number of remarkable properties. It is invariant under translation of trajectories by a constant vector, and under time-reversal of trajectories; cf. Proposition 1.3. Also when K is a finite subset of \mathbb{Z}^d , we introduce the random point process on $W_+ \times \mathbb{R}_+$:

$$(0.5) \quad \mu_K(\omega) = \sum_{i \geq 0} \delta_{(w_i, u_i)} 1\{w_i^* \in W_K^*\}, \text{ if } \omega = \sum_{i \geq 0} \delta_{(w_i^*, u_i)},$$

where for $w_i^* \in W_K$, w_i denotes the unique trajectory in W_+ starting at time 0, where w_i^* enters K , and following from then on w_i^* step-by-step; cf. (1.18) for the precise definition. We show in Proposition 1.3 that

$$(0.6) \quad \mu_K \text{ is a Poisson point process with intensity } P_{e_K}(dw)du,$$

where $P_{e_K} = \sum_x e_K(x)P_x$. Further the point processes μ_K , as K varies, satisfy a compatibility condition; cf. (1.21), (1.46).

It may be worth pointing out that much of the above constructs, except for the aspects related to translation invariance, can be carried out in the more general set-up of a transient random walk attached to an infinite locally finite connected graph with positive weights along its edges, in place of a simple random walk on \mathbb{Z}^d , $d \geq 3$; cf. Remark 1.4.

The *interlacement at level u* is defined as

$$(0.7) \quad \mathcal{I}^u(\omega) = \bigcup_{u_i \leq u} w_i^*(\mathbb{Z}), \text{ if } \omega = \sum_{i \geq 0} \delta_{(w_i^*, u_i)} \in \Omega,$$

where $w_i^*(\mathbb{Z})$ denotes the range of any w with $\pi^*(w) = w_i^*$. The *vacant set at level u* is then

$$(0.8) \quad \mathcal{V}^u(\omega) = \mathbb{Z}^d \setminus \mathcal{I}^u(\omega).$$

Clearly \mathcal{I}^u increases with u , whereas \mathcal{V}^u decreases with u . Also one can see that the restriction of \mathcal{I}^u to K is determined by $\mu_{K'}(dw \times [0, u])$, when $K \subset K'$ are finite subsets of \mathbb{Z}^d ; cf. (1.54). Together with (0.6) one finds that the restriction of \mathcal{I}^u to K can be visualized as the trace on K of a Poisson cloud of finite trajectories. Its intensity measure is proportional to the law of a simple random walk run up to the last visit to K , with initial distribution the harmonic measure of K viewed from infinity, i.e. e_K normalized by its total mass $\text{cap}(K)$, the capacity of K , and the proportionality factor equals $u \text{cap}(K)$; cf. Remark 1.6 (3). We also show in

Corollary 2.3 and Proposition 1.5 that

(0.9) \mathbb{P} -a.s., \mathcal{G}^u is an infinite connected subset, and

$$(0.10) \quad \mathbb{P}[\mathcal{V}^u \supseteq K] = \exp\{-u \operatorname{cap}(K)\},$$

for $u > 0$, and $K \subset \mathbb{Z}^d$, finite. Formula (0.10) characterizes the law of \mathcal{V}^u ; see Remark 2.2 (2). As a special case, cf. (1.58), (1.59), one finds that for $x, y \in \mathbb{Z}^d$,

$$(0.11) \quad \mathbb{P}[x \in \mathcal{V}^u] = \exp\left\{-\frac{u}{g(0)}\right\}, \quad \mathbb{P}[\{x, y\} \subseteq \mathcal{V}^u] = \exp\left\{-\frac{2u}{g(0) + g(y-x)}\right\},$$

where $g(y-x)$ denotes the Green function; cf. (1.5). The identities in (0.11) are in essence formulas (2.26) and (3.6) of Brummelhuis-Hilhorst [4] in their theoretical physics article on the covering of a periodic lattice by a random walk; see Remark 1.6 (5). They display the presence of long range dependence in the random set \mathcal{V}^u , with a correlation of the events $\{x \in \mathcal{V}^u\}$ and $\{y \in \mathcal{V}^u\}$ decaying as $c(u)|x-y|^{-(d-2)}$, when $|x-y|$ tends to infinity.

As mentioned above, the main object of this work is to investigate the presence or absence of an infinite connected component in \mathcal{V}^u . We establish in Theorem 2.1 the ergodicity of the (properly defined) distribution of the random set \mathcal{V}^u , from which easily follows a zero-one law for the probability of occurrence of an infinite connected component in \mathcal{V}^u . It is then straightforward to see that this probability equals one precisely when

$$(0.12) \quad \eta(u) \stackrel{\text{def}}{=} \mathbb{P}[0 \text{ belongs to an infinite connected component of } \mathcal{V}^u] > 0.$$

The function $\eta(\cdot)$ is nonincreasing and just as in the case of Bernoulli percolation, cf. [8], we can introduce the critical value:

$$(0.13) \quad u_* = \inf\{u \geq 0, \eta(u) = 0\} \in [0, \infty].$$

The main results of this article concern the nondegeneracy of u_* . We show in Theorem 3.5 that \mathcal{V}^u does not percolate for large u ; i.e.,

$$(0.14) \quad u_* < \infty, \text{ for } d \geq 3,$$

and in Theorem 4.3 that when $d \geq 7$, \mathcal{V}^u percolates for small $u > 0$; i.e.,

$$(0.15) \quad u_* > 0, \text{ when } d \geq 7.$$

Subsequent developments initiated by the present article respectively relating random interacements on \mathbb{Z}^{d+1} and on \mathbb{Z}^d to the local picture left by simple random walk on $(\mathbb{Z}/N\mathbb{Z})^d \times \mathbb{Z}$ run up to times of order N^{2d} , and random walk on $(\mathbb{Z}/N\mathbb{Z})^d$ run up to times of order N^d can be found in [20], [25]. In this light the results presented here with their proofs also have a bearing on the problems investigated

in [6], [7], [2]. In particular (0.14) offers evidence that when $d \geq 2$ the laws of T_N/N^{2d} are tight, if T_N denotes the disconnection time of $(\mathbb{Z}/N\mathbb{Z})^d \times \mathbb{Z}$ studied in [6], [7]. This signals that one should be able to remove the logarithmic terms present in the (very general) upper bound of [19] and bypass the strategy based on the domination of T_N by the cover time of $(\mathbb{Z}/N\mathbb{Z})^d \times \{0\}$, by relying instead on the emergence of a nonpercolative local picture of the vacant set left by random walk. These heuristic considerations can be made precise and lead to the above claimed tightness; see [21]. Similarly (0.15) offers evidence that the lower bound on T_N in [7], which shows the tightness of N^{2d}/T_N when $d \geq 17$, should hold as soon as $d \geq 6$ (and in fact for all $d \geq 2$, in view of the recent work [15]). Analogously in the context of [2], we see that (0.14), (0.15), and [15] give support for the typical absence for large N of a giant component in the vacant set left by simple random walk on $(\mathbb{Z}/N\mathbb{Z})^d$, $d \geq 3$, run up to time uN^d , if u is large, and for its typical presence when u is chosen small.

There are many natural questions left untouched by the present article. Is there a unique infinite component when \mathcal{V}^u percolates? (See Remark 2.2 (3).) The answer is affirmative, as shown in [22]. Is $u_* > 0$, when $3 \leq d \leq 6$, as suggested by simulations? This is indeed the case; see [15]. However it is presently unknown whether the vacant set percolates at criticality, i.e. when $u = u_*$, or what the large d behavior of u_* is. We refer to Remark 4.4 (3) for further open problems.

We will now comment on the proofs of (0.14) and (0.15). Most of the work goes into the proof of (0.14). This is due to the long range dependence in the model and the fact highlighted by (0.10) that $\mathbb{P}[\mathcal{V}^u \supseteq K]$ does not decay exponentially with $|K|$. This feature creates a very serious obstruction to the Peierls-type argument commonly met in Bernoulli percolation, see [8, p. 16], when one attempts to show that \mathcal{V}^u does not percolate for large u . We instead use a renormalization technique to prove (0.14) and consider a sequence of functions on \mathbb{R}_+ :

$$(0.16) \quad p_n(u) \text{ “=” } \mathbb{P}\text{-probability that } \mathcal{V}^u \text{ contains a path from a given block of side-length } L_n \text{ to the complement of its } L_n\text{-neighborhood,}$$

(cf. (3.8) for the precise definition), where L_n is a rapidly growing sequence of length scales, cf. (3.1), (3.2),

$$(0.17) \quad L_n \approx L_0^{(1+a)^n}, \quad n \geq 0, \quad \text{with } a = \frac{1}{100d}.$$

The key control appears in Proposition 3.1, where we prove that for $L_0 \geq c$, $u_0 > c(L_0)$, and an increasing but bounded sequence u_n depending on L_0, u_0 , cf. (3.9),

$$(0.18) \quad p_n(u_n) \xrightarrow{n \rightarrow \infty} 0.$$

This immediately implies that $\eta(u) = 0$, for $u \geq u_\infty = \sup u_n (< \infty)$, and proves (0.14). The principal difficulty in proving (0.18) resides in the derivation of a

suitable recurrence relation between $p_n(\cdot)$ and $p_{n+1}(\cdot)$, cf. (3.52), due to the long range dependence in the model. In a suitable sense we use a “sprinkling technique”, where more independent paths are thrown in, so as to dominate long range dependence. This is reflected in the fact that we evaluate $p_n(\cdot)$ at an increasing but convergent sequence u_n in the key control (3.10), (a more quantitative version of (0.18)). Incidentally, the sequence of length scales appearing in (0.17) corresponds to the choice of a small a , so as to control the combinatorial complexity involved in selecting boxes of scale L_n within a given box of scale L_{n+1} , see (3.13), but also to the choice of fast enough growth, so as to discard paths making more than a certain finite number of excursions at distance of order L_{n+1} ; see below (3.25), and (3.55), (3.65).

The proof of (0.15) in Theorem 4.3 employs a similar albeit simpler renormalization strategy. One can instead employ a Peierls-type argument to show that \mathcal{V}^u percolates for small $u > 0$, when d is sufficiently large, very much in the spirit of Section 2 of [2], or Section 1 of [7]. It is based on an exponential bound on $\mathbb{P}[\mathcal{F}^u \supseteq A]$, for A finite subset of \mathbb{Z}^2 (where \mathbb{Z}^2 is viewed as a subset of \mathbb{Z}^d); cf. (2.37) in Theorem 2.4. This estimate mirrors the exponential controls derived in Theorem 2.1 of [2] and Theorem 1.2 of [2]. This strategy leads to a proof of (0.15) when:

$$(0.19) \quad 7\left(\frac{2}{d} + \left(1 - \frac{2}{d}\right) q(d-2)\right) < 1,$$

with $q(v)$ the return probability to the origin of simple random walk on \mathbb{Z}^v . In practice this means $d \geq 18$; cf. Remark 2.5 (3). The technique we use works instead as soon as $d \geq 7$. It also shows, just as the Peierls-type argument does when (0.19) holds, the existence of an infinite connected cluster in $\mathcal{V}^u \cap \mathbb{Z}^2$, for small $u > 0$.

Let us now describe the organization of the article.

In Section 1 we construct the model of random interlacements. The main task lies in the construction of the σ -finite measure ν entering the intensity of the Poisson point process we are after. This is done in Theorem 1.1. Basic properties of the model appear in Proposition 1.3, whereas Proposition 1.5 shows (0.10), (0.11).

Section 2 shows the ergodicity of the law of \mathcal{V}^u , and the zero-one law for the probability that \mathcal{V}^u percolates in Theorem 2.1. We also prove (0.9) in Corollary 2.3. In Theorem 2.4 we derive exponential bounds that provide further links of the present model to [2], [7].

Section 3 is devoted to the proof of (0.14) in Theorem 3.5. The main renormalization step is contained in Proposition 3.1.

Section 4 shows (0.15) in Theorem 4.3. The principal step appears in Proposition 4.1.

Finally let us state our convention regarding constants. Throughout the text c and c' denote positive constants which solely depend on d , with values changing from place to place. The numbered constants c_0, c_1, \dots are fixed and refer to the value at their first appearance in the text. Dependence of constants on additional parameters appears in the notation. For instance $c(L_0, u_0)$ denotes a positive constant depending on d, L_0, u_0 .

1. Basic model and some first properties

The main object of this section is to introduce the basic model and present some of its properties. As explained in the introduction the basic model comes as a Poisson point process on a suitable state space. The main task resides in the construction of the intensity measure of this point process. This is done in Theorem 1.1. We then derive some of its properties in Proposition 1.3 as well as some of the properties of the vacant set left by the interlacement at level u , cf. (0.9), in Proposition 1.5. We first begin with some notation.

We write $\mathbb{N} = \{0, 1, 2, \dots\}$ for the set of natural numbers. Given a nonnegative real number a , we write $[a]$ for the integer part of a , and for real numbers b, c , we write $b \wedge c$ and $b \vee c$ for the respective minimum and maximum of b and c . We denote with $|\cdot|$ and $|\cdot|_\infty$ the Euclidean and ℓ^∞ -distances on \mathbb{Z}^d . We write $B(x, r)$ for the closed $|\cdot|_\infty$ -ball with center x in \mathbb{Z}^d and radius $r \geq 0$, and $S(x, r)$ for the corresponding $|\cdot|_\infty$ -sphere with center x and radius r (it is empty when r is not an integer). We say that x, y in \mathbb{Z}^d are neighbors, respectively $*$ -neighbors, when $|x - y| = 1$, respectively $|x - y|_\infty = 1$. The notions of connected and $*$ -connected subsets are defined accordingly, and so are the notions of nearest neighbor or $*$ -nearest neighbor paths in \mathbb{Z}^d . For A, B subsets in \mathbb{Z}^d , we denote by $A + B$ the subset of elements of the form $x + y$, with $x \in A, y \in B$ and with $d(A, B) = \inf\{|x - y|_\infty; x \in A, y \in B\}$, the $|\cdot|_\infty$ -distance from A to B . When U is a subset of \mathbb{Z}^d , we let $|U|$ stand for the cardinality of U , ∂U for the exterior boundary of U and $\partial_{\text{int}}U$ for the interior boundary of U :

$$(1.1) \quad \partial U = \{x \in U^c; \exists y \in U, |x - y| = 1\}, \quad \partial_{\text{int}}U = \{x \in U; \exists y \in U^c, |x - y| = 1\}.$$

We write $U \subset\subset \mathbb{Z}^d$ to express U as a finite subset of \mathbb{Z}^d . In what follows, unless otherwise explicitly mentioned, we tacitly assume that $d \geq 3$.

We consider W_+ and W the spaces of trajectories:

$$(1.2) \quad \begin{aligned} W_+ &= \left\{ w \in (\mathbb{Z}^d)^{\mathbb{N}}; |w(n+1) - w(n)| = 1, \text{ for all } n \geq 0, \text{ and} \right. \\ &\quad \left. \lim_n |w(n)| = \infty \right\}, \\ W &= \left\{ w \in (\mathbb{Z}^d)^{\mathbb{Z}}; |w(n+1) - w(n)| = 1, \text{ for all } n \in \mathbb{Z}, \text{ and} \right. \\ &\quad \left. \lim_{|n| \rightarrow \infty} |w(n)| = \infty \right\}. \end{aligned}$$

We denote by $X_n, n \geq 0$, and $X_n, n \in \mathbb{Z}$, the respective canonical coordinates on W_+ and W , and write $\theta_n, n \geq 0$, and $\theta_n, n \in \mathbb{Z}$, for the respective canonical shifts. We let \mathcal{W}_+ and \mathcal{W} stand for the σ -fields on W_+ and W generated by the canonical coordinates.

Given $U \subseteq \mathbb{Z}^d, w \in W_+$, we denote with $H_U(w), T_U(w), \tilde{H}_U(w)$, the entrance time in U , the exit time from U , and the hitting time of U for the trajectory w :

$$(1.3) \quad H_U(w) = \inf\{n \geq 0; X_n(w) \in U\}, \quad T_U(w) = \inf\{n \geq 0; X_n(w) \notin U\}, \\ \tilde{H}_U(w) = \inf\{n \geq 1; X_n(w) \in U\}.$$

We often drop “ w ” from the notation and write H_x, T_x, \tilde{H}_x , when $U = \{x\}$. Also when $w \in W$, we define $H_U(w)$ and $T_U(w)$ in a similar fashion replacing “ $n \geq 0$ ” with “ $n \in \mathbb{Z}$ ” in (1.3), and $\tilde{H}_U(w)$ just as in (1.3). For $K \subseteq U$ in $\mathbb{Z}^d, w \in W_+$, we consider $R_k, D_k, k \geq 1$, the successive returns to K and departures from U of the trajectory w :

$$(1.4) \quad R_1 = H_K, \quad D_1 = T_U \circ \theta_{H_K} + H_K, \quad \text{and for } k \geq 1, \\ R_{k+1} = R_1 \circ \theta_{D_k} + D_k, \quad D_{k+1} = D_1 \circ \theta_{D_k} + D_k,$$

so that $0 \leq R_1 \leq D_1 \leq \dots \leq R_k \leq D_k \leq \dots \leq \infty$.

When X is an integrable random variable and A an event, we routinely write $E[X, A]$ in place of $E[X 1_A]$ in what follows, with E referring here to the relevant expectation. Given $x \in \mathbb{Z}^d$, we write P_x for the restriction to (W_+, \mathcal{W}_+) of the canonical law of simple random walk on \mathbb{Z}^d starting at x . Recall that $d \geq 3$, and W_+ has full measure under the canonical law. When ρ is a positive measure on \mathbb{Z}^d , we write P_ρ for the measure $\sum_{x \in \mathbb{Z}^d} \rho(x) P_x$. We denote with $g(\cdot, \cdot)$ the Green function of the walk:

$$(1.5) \quad g(x, y) = \sum_{n \geq 0} P_x[X_n = y], \quad x, y \in \mathbb{Z}^d,$$

and $g(y) = g(0, y)$ so that $g(x, y) = g(y - x)$, thanks to translation invariance. Given $K \subset \subset \mathbb{Z}^d$, we write e_K for the equilibrium measure of K , $\text{cap}(K)$ for the capacity of K , so that, cf. Chapter 2, Section 2 of [11]:

$$(1.6) \quad e_K(x) = P_x[\tilde{H}_K = \infty], \quad x \in K, \\ = 0, \quad \text{if } x \notin K$$

(note that e_K is supported on $\partial_{\text{int}}K$),

$$(1.7) \quad \text{cap}(K) = e_K(\mathbb{Z}^d) \left(= \sum_{x \in \mathbb{Z}^d} e_K(x) \right),$$

and

$$(1.8) \quad P_x[H_K < \infty] = \int_K g(x, y) e_K(dy) \left(= \sum_{y \in K} g(x, y) e_K(y) \right), \text{ for } x \in \mathbb{Z}^d.$$

The following bounds on $P_x[H_K < \infty]$, $x \in \mathbb{Z}^d$, will be useful:

$$(1.9) \quad \sum_{y \in K} g(x, y) / \sup_{z \in K} \left(\sum_{y \in K} g(z, y) \right) \leq P_x[H_K < \infty] \leq \sum_{y \in K} g(x, y) / \inf_{z \in K} \left(\sum_{y \in K} g(z, y) \right).$$

They classically follow from the $L^1(P_x)$ -convergence of the bounded martingale $M_n = \sum_{y \in K} g(X_n \wedge H_K, y)$, $n \geq 0$, towards $1_{\{H_K < \infty\}} \sum_{y \in K} g(X_{H_K}, y)$.

The state space of the Poisson point process we wish to define involves the quotient space W^* of equivalence classes of trajectories in W modulo time-shift; cf. (0.2). We recall that π^* stands for the canonical projection on W^* . We endow W^* with the canonical σ -field

$$(1.10) \quad \mathcal{W}^* = \{A \subseteq W^*; (\pi^*)^{-1}(A) \in \mathcal{W}\},$$

which is the largest σ -algebra such that $(W, \mathcal{W}) \xrightarrow{\pi^*} (W^*, \mathcal{W}^*)$ is measurable. When $K \subset \subset \mathbb{Z}^d$, we consider

$$(1.11) \quad W_K = \{w \in W; X_n(w) \in K, \text{ for some } n \in \mathbb{Z}\},$$

the subset of W of trajectories entering K . We can write $W_K \in \mathcal{W}$ as a countable partition into measurable sets (see below (1.3) for the notation):

$$(1.12) \quad W_K = \bigcup_{n \in \mathbb{Z}} W_K^n, \text{ where } W_K^n = \{w \in W; H_K(w) = n\}.$$

We then introduce

$$(1.13) \quad W_K^* = \pi^*(W_K) \left(= \pi^*(W_K^0) \right),$$

as well as the map

$$(1.14) \quad s_K : W_K^* \rightarrow W, \text{ with } s_K(w^*) = w^0 \text{ the unique element of } W_K^0 \text{ with } \pi^*(w^0) = w^*.$$

Note that $s_K(W_K^*) = W_K^0$ and s_K is a section of π^* over W_K^* ; i.e., $\pi^* \circ s_K$ is the identity map on W_K^* . It is then straightforward to check that for any $K \subset \subset \mathbb{Z}^d$,

$$(1.15) \quad W_K^* \in \mathcal{W}^* \text{ and the trace of } \mathcal{W}^* \text{ on } W_K^* \text{ coincides with } s_K^{-1}(\mathcal{W}).$$

There is no natural way to globally identify $W^* \times \mathbb{Z}$ with W , but the maps s_K enable us to identify W_K^* with W_K^0 and $W_K^* \times \mathbb{Z}$ with W_K . In a slightly pedantic way $\pi^* : (W, \mathcal{W}) \rightarrow (W^*, \mathcal{W}^*)$ endowed with the transformations θ_n , $n \in \mathbb{Z}$, on

the fiber of π^* could be viewed as a “principal fiber-bundle with group \mathbb{Z} ”; cf. [18, p. 346]. The construction of the key σ -finite measure ν in Theorem 1.1 will involve checking compatibility and patching up expressions for ν “read in the local chart s_K ”, as K varies over finite subsets of \mathbb{Z}^d .

We further need to introduce several spaces of point measures routinely used in what follows. In particular we consider Ω and M the spaces of point measures on $W^* \times \mathbb{R}_+$ and $W_+ \times \mathbb{R}_+$:

$$(1.16) \quad \Omega = \left\{ \omega = \sum_{i \geq 0} \delta_{(w_i^*, u_i)}, \text{ with } (w_i^*, u_i) \in W^* \times \mathbb{R}_+, i \geq 0, \text{ and } \omega(W_K^* \times [0, u]) < \infty, \text{ for any } K \subset\subset \mathbb{Z}^d, u \geq 0 \right\};$$

$$(1.17) \quad M = \left\{ \mu = \sum_{i \in I} \delta_{(w_i, u_i)}, \text{ with } I \text{ a variable finite or infinite subset of } \mathbb{N}, (w_i, u_i) \in W_+ \times \mathbb{R}_+, \text{ for } i \in I, \text{ and } \mu(W_+ \times [0, u]) < \infty, \text{ for } u \geq 0 \right\},$$

We endow Ω with the σ -algebra \mathcal{A} generated by the evaluation maps $\omega \rightarrow \omega(D)$, where D runs over $W^* \otimes \mathcal{B}(\mathbb{R}_+)$, cf. (1.10). Likewise we endow M with the σ -algebra \mathcal{M} generated by the evaluation maps $\mu \rightarrow \mu(D)$, where D runs over $W_+ \otimes \mathcal{B}(\mathbb{R}_+)$; cf. below (1.2). Given $K \subset\subset \mathbb{Z}^d$, we then define the measurable maps $\mu_K : \Omega \rightarrow M$ and $\Theta_K : M \rightarrow M$ via:

$$(1.18) \quad \mu_K(\omega)(f) = \int_{W_K^* \times \mathbb{R}_+} f(s_K(w^*)_+, u) \omega(dw^*, du), \text{ for } \omega \in \Omega, \text{ and } f \text{ nonnegative measurable on } W_+ \times \mathbb{R}_+,$$

where for $w \in W$, $w_+ \in W_+$ denotes the restriction of w to \mathbb{N} , so that $s_K(w^*)_+$ starts at time 0 where $w^* \in W_K^*$ enters K , and follows from then on w^* step-by-step, as well as

$$(1.19) \quad \Theta_K(\mu)(f) = \int_{\{H_K < \infty\} \times \mathbb{R}_+} f(\theta_{H_K}(w), u) \mu(dw, du), \text{ for } \mu \in M, \text{ and } f \text{ as in (1.18)}.$$

In other words,

$$\Theta_K(\mu) = \sum_{i \in I} \delta_{(\theta_{H_K}(w_i), u_i)} 1_{\{H_K(w_i) < \infty\}},$$

when $\mu = \sum_{i \in I} \delta_{(w_i, u_i)} \in M$. Given $K \subset\subset \mathbb{Z}^d$, $u \geq 0$, we will also consider the measurable function on Ω with values in the set of finite point measures on (W_+, \mathcal{W}_+) :

$$(1.20) \quad \mu_{K,u}(\omega)(dw) = \mu_K(\omega)(dw \times [0, u]), \text{ for } \omega \in \Omega.$$

We record for later use the straightforward identities valid for $K \subset K' \subset \subset \mathbb{Z}^d$:

$$(1.21) \quad \begin{aligned} \text{i)} \quad & \Theta_K \circ \mu_{K'} = \mu_K, \\ \text{ii)} \quad & \Theta_K \circ \Theta_{K'} = \Theta_K. \end{aligned}$$

We are now going to construct the σ -finite measure ν on (W^*, \mathfrak{W}^*) which enters the intensity of the Poisson point process we wish to define. For $K \subset \subset \mathbb{Z}^d$, we write \mathcal{T}_K for the countable set of finite nearest-neighbor trajectories starting and ending in the support of e_K :

$$(1.22) \quad \begin{aligned} \mathcal{T}_K = \{ \tau = (\tau(n))_{0 \leq n \leq N_\tau}; \quad & N_\tau \geq 0, |\tau(n+1) - \tau(n)| = 1, \text{ for } 0 \leq n < N_\tau, \\ & \text{and } \tau(0), \tau(N_\tau) \in \text{Supp } e_K \}. \end{aligned}$$

If $x \in \text{Supp } e_K$, we also denote with P_x^K the probability on W_+ governing the walk conditioned not to hit K :

$$(1.23) \quad P_x^K[\cdot] = P_x[\cdot | \tilde{H}_K = \infty].$$

We are now ready to state

THEOREM 1.1. *For $K \subset \subset \mathbb{Z}^d$, denote with Q_K the finite measure on W , supported on W_K^0 , such that for any $A, B \in \mathfrak{W}_+$, $x \in \mathbb{Z}^d$:*

$$(1.24) \quad Q_K[(X_{-n})_{n \geq 0} \in A, X_0 = x, (X_n)_{n \geq 0} \in B] = P_x^K[A] e_K(x) P_x[B].$$

There is a unique σ -finite measure ν on (W^, \mathfrak{W}^*) such that:*

$$(1.25) \quad 1_{W_K^*} \nu = \pi^* \circ Q_K, \text{ for any } K \subset \subset \mathbb{Z}^d.$$

Further, letting $L_K(w) = \sup\{n \geq 0; X_n(w) \in K\}$, $w \in W_K^0$, stand for the time of the last visit to K of w , one sees that the law under Q_K of $(X_n)_{0 \leq n \leq L_K}$ is supported on \mathcal{T}_K , and for $A, B \in \mathfrak{W}_+$, $\tau \in \mathcal{T}_K$,

$$(1.26) \quad \begin{aligned} & Q_K[(X_{-n})_{n \geq 0} \in A, (X_n)_{0 \leq n \leq L_K} = \tau, (X_{n+L_K})_{n \geq 0} \in B] \\ & = P_{\tau(0)}^K[A] e_K(\tau(0)) P_{\tau(0)}[X_n = \tau(n), 0 \leq n \leq N_\tau] e_K(\tau(N_\tau)) P_{\tau(N_\tau)}^K[B]. \end{aligned}$$

$$(1.27) \quad \begin{aligned} & \nu \text{ is invariant under the time reversal involution on } W^*, w^* \rightarrow \tilde{w}^*, \\ & \text{where } \tilde{w}^* = \pi^*(\tilde{w}), \text{ with } \pi^*(w) = w^* \text{ and } \tilde{w}(n) = w(-n), \text{ for } n \in \mathbb{Z}. \end{aligned}$$

$$(1.28) \quad \begin{aligned} & \nu \text{ is invariant under the translations on } W^*: w^* \rightarrow w^* + x, x \in \mathbb{Z}^d, \\ & \text{where } w^* + x = \pi^*(w + x), \text{ with } \pi^*(w) = w^*. \end{aligned}$$

Proof. We begin with the proof of the existence and uniqueness of ν . Since $W^* = \bigcup_{m \geq 0} W_{K_m}^*$, where $K_m \uparrow \mathbb{Z}^d$, with K_m finite, for $m \geq 0$, the uniqueness of

ν satisfying (1.25) is immediate. As for the existence of ν , denote for $K \subset\subset \mathbb{Z}^d$ with ν_K the finite measure supported on $W_K^* = \pi^*(W_K^0)$ on the right-hand side of (1.25):

$$(1.29) \quad \nu_K = \pi^* \circ Q_K .$$

The existence of ν will follow once we show that for $K \subset K' \subset\subset \mathbb{Z}^d$,

$$1_{W_K^*} \nu_{K'} = \nu_K .$$

This in turn will follow once we prove that:

$$(1.30) \quad (s_K \circ s_{K'}^{-1}) \circ (1_{s_{K'}(W_K^*)} Q_{K'}) = Q_K ,$$

where $s_{K'}^{-1}$ denotes the restriction of π^* to $W_{K'}^0$. Indeed it suffices simply to take the image of both sides under π^* . We now write $s_{K'}(W_K^*)$ as the at most countable partition into measurable sets:

$$(1.31) \quad s_{K'}(W_K^*) = \bigcup_{\sigma \in \Sigma} W_{K',\sigma}^0 ,$$

where Σ denotes the set of finite nearest-neighbor trajectories $\sigma = (\sigma(n))_{0 \leq n \leq N_\sigma}$, with $\sigma(0) \in K'$, $\sigma(n) \notin K$ for $n < N_\sigma$, and $\sigma(N_\sigma) \in K$, and

$$(1.32) \quad W_{K',\sigma}^0 = \{w \in W_{K'}^0; X_n(w) = \sigma(n), \text{ for } 0 \leq n \leq N_\sigma\} .$$

One then has the identity:

$$(1.33) \quad s_K \circ s_{K'}^{-1}(w) = w(\cdot + N_\sigma) = \theta_{N_\sigma}(w), \text{ for } w \in W_{K',\sigma}^0 .$$

As a result, denoting with Q the left-hand side of (1.30), we find that

$$(1.34) \quad Q = \sum_{\sigma \in \Sigma} \theta_{N_\sigma} \circ (1_{W_{K',\sigma}^0} Q_{K'}) .$$

Thus given an arbitrary collection $A_i, i \in \mathbb{Z}$, of subsets of \mathbb{Z}^d , we see that

$$(1.35) \quad \begin{aligned} Q[X_i \in A_i, i \in \mathbb{Z}] &= \sum_{\sigma \in \Sigma} Q_{K'}[X_{i+N_\sigma} \in A_i, i \in \mathbb{Z}, X_n = \sigma(n), 0 \leq n \leq N_\sigma] \\ &= \sum_{\sigma \in \Sigma} Q_{K'}[X_i \in A_{i-N_\sigma}, i \in \mathbb{Z}, X_n = \sigma(n), 0 \leq n \leq N_\sigma] \end{aligned}$$

$$(1.6),(1.24) \quad \begin{aligned} &= \sum_{\sigma \in \Sigma} \sum_{x \in \text{Supp}(e_{K'})} P_x^{K'}[X_m \in A_{-m-N_\sigma}, m \geq 0] P_x[\tilde{H}_{K'} = \infty] \\ &\quad P_x[X_n \in A_{n-N_\sigma}, n \geq 0, X_n = \sigma(n), 0 \leq n \leq N_\sigma] \end{aligned}$$

$$(1.23), \text{Markov} \quad \begin{aligned} &= \sum_{\sigma \in \Sigma} \sum_{x \in \text{Supp}(e_{K'})} P_x[X_m \in A_{-m-N_\sigma}, m \geq 0, \tilde{H}_{K'} = \infty] \\ &\quad P_x[X_n = \sigma(n) \in A_{n-N_\sigma}, 0 \leq n \leq N_\sigma] P_{\sigma(N_\sigma)}[X_n \in A_n, n \geq 0] . \end{aligned}$$

It follows from the reversibility of the walk that for $y \in K$:

$$\begin{aligned}
 (1.36) \quad & \sum_{\sigma: \sigma(N_\sigma)=y} \sum_{x \in \text{Supp}(e_{K'})} P_x[X_m \in A_{-m-N_\sigma}, m \geq 0, \tilde{H}_{K'} = \infty] \\
 & P_x[X_n = \sigma(n) \in A_{n-N_\sigma}, 0 \leq n \leq N_\sigma] \\
 & = \sum_{x \in \text{Supp}(e_{K'})} \sum_{\substack{\sigma: \sigma(N_\sigma)=y \\ \sigma(0)=x}} P_x[X_m \in A_{-m-N_\sigma}, m \geq 0, \tilde{H}_{K'} = \infty] \\
 & P_y[X_n = \sigma(N_\sigma - n) \in A_{-n}, 0 \leq n \leq N_\sigma] \\
 & \stackrel{\text{Markov}}{=} \sum_{x \in \text{Supp}(e_{K'})} \sum_{\substack{\sigma: \sigma(N_\sigma)=y \\ \sigma(0)=x}} P_y[X_n = \sigma(N_\sigma - n) \in A_{-n}, 0 \leq n \leq N_\sigma, \\
 & \qquad \qquad \qquad X_n \in A_{-n}, n \geq N_\sigma, \tilde{H}_{K'} \circ \theta_{N_\sigma} = \infty] \\
 & = \sum_{x \in \text{Supp}(e_{K'})} P_y[\tilde{H}_K = \infty, \text{the last visit to } K' \text{ occurs at } x, \text{ and} \\
 & \qquad \qquad \qquad X_n \in A_{-n}, \text{ for } n \geq 0] \\
 & = P_y[\tilde{H}_K = \infty, X_n \in A_{-n}, n \geq 0].
 \end{aligned}$$

Inserting this identity in the last line of (1.35) we find that:

$$\begin{aligned}
 (1.37) \quad & Q[X_i \in A_i, i \in \mathbb{Z}] = \sum_{y \in K} P_y[\tilde{H}_K = \infty, X_n \in A_{-n}, n \geq 0] P_y[X_n \in A_n, n \geq 0] \\
 & = \sum_{y \in \text{Supp}(e_K)} P_y^K[X_n \in A_{-n}, n \geq 0] e_K(y) P_y[X_n \in A_n, n \geq 0] \\
 & \stackrel{(1.24)}{=} Q_K[X_n \in A_n, n \in \mathbb{Z}].
 \end{aligned}$$

This proves that (1.30) holds and thus concludes the proof of the existence of ν satisfying (1.25), which is automatically σ -finite.

We now turn to the proof of (1.26). We consider $\tau(n), 0 \leq n \leq N$, some finite sequence in \mathbb{Z}^d . Observe that $Q_K[(X_n)_{0 \leq n \leq L_K} = \tau]$ vanishes unless τ is nearest neighbor and $\tau(N) \in K$. Moreover when this is the case it follows from the use of the Markov property at time N that:

$$\begin{aligned}
 (1.38) \quad & Q_K[(X_n)_{0 \leq n \leq L_K} = \tau] = Q_K[X_n = \tau(n), 0 \leq n \leq N, \tilde{H}_K \circ \theta_N = \infty] \\
 & \stackrel{(1.24), (1.6)}{\stackrel{\text{Markov}}{=}} e_K(\tau(0)) P_{\tau(0)}[X_n = \tau(n), 0 \leq n \leq N] e_K(\tau(N)).
 \end{aligned}$$

This shows that the law of $(X_n)_{0 \leq n \leq L_K}$ under Q_K is supported by \mathcal{T}_K . Also repeating the argument which yielded (1.38), we see that for $A, B \in \mathcal{W}_+, \tau \in \mathcal{T}_K$

the left-hand side of (1.26) equals (writing N in place of N_τ for simplicity):

$$\begin{aligned} & Q_K[(X_{-n})_{n \geq 0} \in A, X_n = \tau(n), 0 \leq n \leq N, \theta_N^{-1}(\{\tilde{H}_K = \infty\} \cap \{X_n \in B, n \geq 0\})] \\ & \stackrel{(1.24)}{=} P_{\tau(0)}^K[A] e_K(\tau(0)) P_{\tau(0)}[X_n = \tau(n), 0 \leq n \leq N, \\ & \qquad \qquad \qquad \theta_N^{-1}(\{\tilde{H}_K = \infty\} \cap \{X_n \in B, n \geq 0\})] \\ & \stackrel{\text{Markov}}{\stackrel{(1.6),(1.23)}{=}} P_{\tau(0)}^K[A] e_K(\tau(0)) P_{\tau(0)}[X_n = \tau(n), 0 \leq n \leq N] e_K(\tau(N)) P_{\tau(N)}^K[B], \end{aligned}$$

and this proves (1.26).

To prove (1.27), observe that for $K \subset \subset \mathbb{Z}^d$, $w^* \rightarrow \check{w}^*$ leaves W_K^* invariant and $X_n(s_K(\check{w}^*)) = X_{L_K-n}(s_K(w^*))$, for $n \in \mathbb{Z}$, $w^* \in W_K^*$. Denoting $\check{\gamma}$ the image under $w^* \rightarrow \check{w}^*$ of a measure γ on W^* , we find for $C \in \mathcal{W}$

$$\begin{aligned} (1.39) \quad s_K \circ (1_{W_K^*} \check{\nu})(C) &= s_K \circ ((1_{W_K^*} \nu)^\check{)}(C) = s_K \circ (1_{W_K^*} \nu)((X_{L_K-}) \in C) \\ & \stackrel{(1.25)}{=} Q_K((X_{L_K-}) \in C). \end{aligned}$$

Hence with (1.26), $A, B \in \mathcal{W}_+$, $\tau \in \mathcal{T}_K$, and C denotes the event in the probability in the first line of (1.26); writing N in place of N_τ , for simplicity, we find that

$$\begin{aligned} (1.40) \quad s_K \circ (1_{W_K^*} \check{\nu})(C) &= Q_K[(X_{-n})_{n \geq 0} \in B, \\ & (X_n)_{0 \leq n \leq L_K} = \tau(N - \cdot), (X_{n+L_K})_{n \geq 0} \in A] \stackrel{(1.26)}{=} P_{\tau(N)}^K[B] e_K(\tau(N)) \\ & P_{\tau(N)}[(X_n)_{0 \leq n \leq N} = \tau(N - n)] e_K(\tau(0)) P_{\tau(0)}^K[A] \stackrel{\text{reversibility}}{=} P_{\tau(0)}^K[A] e_K(\tau(0)) \\ & P_{\tau(0)}[X_n = \tau(n), 0 \leq n \leq N] e_K(\tau(N)) P_{\tau(N)}^K[B] = Q_K(C) \stackrel{(1.25)}{=} \\ & s_K \circ (1_{W_K^*} \nu)(C). \end{aligned}$$

It now readily follows that $s_K \circ (1_{W_K^*} \check{\nu}) = s_K \circ (1_{W_K^*} \nu)$ and hence $1_{W_K^*} \check{\nu} = 1_{W_K^*} \nu$ for any $K \subset \subset \mathbb{Z}^d$, whence $\check{\nu} = \nu$. This proves (1.27).

Finally for the proof of (1.28), we note that for $x \in \mathbb{Z}^d$, $K \subset \subset \mathbb{Z}^d$, $w^* \rightarrow w^* + x$ maps W_K^* one-to-one onto W_{K+x}^* , and $s_K(w^* + x) = s_{K-x}(w^*) + x$, for $w^* \in W_{K-x}^*$. Denoting by γ^x the image under $w^* \rightarrow w^* + x$ of a measure γ on W^* , we see that for $C \in \mathcal{W}$, we have

$$\begin{aligned} (1.41) \quad s_K \circ (1_{W_K^*} \nu^x)(C) &= s_K \circ ((1_{W_{K-x}^*} \nu)^x)(C) = s_{K-x} \circ (1_{W_{K-x}^*} \nu)((X_n) + x \in C) \\ & \stackrel{(1.25)}{=} Q_{K-x}((X_n) + x \in C). \end{aligned}$$

Hence with $A, B \in \mathcal{W}_+, y \in \mathbb{Z}^d$ and C denoting the event in the left-hand side of (1.24), where x is replaced by y , we find that:

$$s_K \circ (1_{W^*} \nu^x)(C) = Q_{K-x}[(X_{-n})_{n \geq 0} \in A - x, X_0 = y - x, (X_n)_{n \geq 0} \in B - x] \\ \stackrel{(1.24)}{=} P_{y-x}^{K-x}[A - x] e_{K-x}(y - x) P_{y-x}[B - x] = Q_K[C] \stackrel{(1.25)}{=} s_K \circ (1_{W_K^*} \nu)(C),$$

using (1.24) and translation invariance in the third equality. This readily implies that $\nu^x = \nu$ and concludes the proof of Theorem 1.1. \square

Remark 1.2. 1) Let us say a few words on why the quotient space W^* is better suited for our purpose than W . One can of course use the sections s_K , with K growing along an increasing sequence of finite sets exhausting \mathbb{Z}^d to construct “by patching” a σ -finite measure on (W, \mathcal{W}) projecting down to ν under π^* . However there is no measure on (W, \mathcal{W}) invariant under translation of trajectories by constant vectors projecting down to ν . Indeed if such a measure ρ existed then for any $K \subset \subset \mathbb{Z}^d$ we would have

$$\text{cap}(K) = \nu(W_K^*) = \rho(W_K) \geq \rho(X_0 \in K) = \rho(X_0 = 0) |K|,$$

using translation invariance in the last equality. However capacity grows more slowly than volume when K is of the form $B(0, L)$, with L tending to infinity; cf. below (3.24). This would imply that $\rho(X_0 = 0) = 0$ and hence $\rho = 0$, due to translation invariance, thus leading to a contradiction. More obstructions can be brought to light, which make measures on (W, \mathcal{W}) projecting down to ν definitely less natural than ν .

2) The expression in the right-hand member of (1.38) when $K = B(0, L)$ coincides up to a normalization factor with the expression (3.13) of [2] governing the limit law of certain properly recentered excursions of a simple random walk on $(\mathbb{Z}/N\mathbb{Z})^d$ to a box of side-length $2L$; see Theorem 3.1 of [2]. This limiting result played a key role in the control of fluctuations of certain averages; cf. (4.43) and Proposition 4.2. of [2]. \square

We will now endow the space (Ω, \mathcal{A}) , cf. (1.16), with a probability measure and thereby complete the construction of the basic model of interlacements. To this end we note that the infinite measure $\nu(dw^*)du$ on $W^* \times \mathbb{R}_+$ gives finite mass to the sets $W_K^* \times [0, u]$, for $K \subset \subset \mathbb{Z}^d$ and $u \geq 0$. We can thus construct on (Ω, \mathcal{A}) the law \mathbb{P} of a Poisson point measure with intensity $\nu(dw^*)du$. We denote by $\mathbb{E}[\cdot]$ the corresponding expectation. The law \mathbb{P} is for instance characterized by the fact that, cf. [14, p. 129],

$$(1.42) \quad \mathbb{E} \left[\exp \left\{ - \int_{W^* \times \mathbb{R}_+} f \omega(dw^*, du) \right\} \right] = \exp \left\{ - \int_{W^* \times \mathbb{R}_+} (1 - e^{-f}) \nu(dw^*)du \right\},$$

for any nonnegative $\mathcal{W}^* \otimes \mathcal{B}(\mathbb{R}_+)$ -measurable function f .

In a similar fashion we can also realize on (M, \mathcal{M}) , cf. (1.17), the law of the Poisson point measure on $W_+ \times \mathbb{R}_+$ with intensity $P_{e_K}(dw)du$, when $K \subset\subset \mathbb{Z}^d$. We call it \mathbb{P}_K and write $\mathbb{E}_K[\cdot]$ for the corresponding expectation. It is characterized by the fact that:

$$(1.43) \quad \mathbb{E}_K \left[\exp \left\{ - \int_{W_+ \times \mathbb{R}_+} f \mu(dw, du) \right\} \right] = \exp \left\{ - \int_{W_+ \times \mathbb{R}_+} (1 - e^{-f}) P_{e_K}(dw)du \right\},$$

for any nonnegative $W_+ \otimes \mathcal{B}(\mathbb{R}_+)$ -measurable function f .

We will now collect some straightforward properties of the laws \mathbb{P} and \mathbb{P}_K . Given $\omega = \sum_{i \geq 0} \delta_{(w_i^*, u_i)}$, we write

$$(1.44) \quad \check{\omega} = \sum_{i \geq 0} \delta_{(\check{w}_i^*, u_i)} \in \Omega,$$

$$\tau_x \omega = \sum_{i \geq 0} \delta_{(w_i^* - x, u_i)} \in \Omega, \text{ for } x \in \mathbb{Z}^d.$$

We also recall the notation from (1.18), (1.19).

PROPOSITION 1.3 ($K \subset K' \subset\subset \mathbb{Z}^d$).

- (1.45) \mathbb{P}_K is the law of μ_K under \mathbb{P} .
- (1.46) $\Theta_K \circ \mathbb{P}_{K'} = \mathbb{P}_K$.
- (1.47) \mathbb{P} is invariant under $\omega \rightarrow \check{\omega}$ (time-reversal invariance).
- (1.48) \mathbb{P} is invariant under τ_x for any $x \in \mathbb{Z}^d$ (translation invariance).

Proof. We begin with (1.45), and note that μ_K due to (1.18) is distributed as a Poisson point process on $W_+ \times \mathbb{R}_+$ with intensity measure $\gamma(dw du)$ such that for f as in (1.18)

$$(1.49) \quad \int_{W_+ \times \mathbb{R}_+} f \gamma(dw, du) = \int_{W_K^* \times \mathbb{R}_+} f(s_K(w^*)_+, u) \nu(dw^*)du$$

$$\stackrel{(1.25), (1.24)}{=} \int_{W_+ \times \mathbb{R}_+} f(w, u) P_{e_K}(dw)du.$$

This shows that the law of μ_K coincides with \mathbb{P}_K . Then (1.46) immediately follows from (1.21) i) and (1.45), whereas (1.47), (1.48) respectively follow from (1.27), (1.28). □

Remark 1.4. The constructions we have made here in the case of a simple random walk on \mathbb{Z}^d , $d \geq 3$, can be straightforwardly generalized to the case of an infinite locally finite connected graph $\Gamma = (G, \mathcal{E})$ with vertex set G and (undirected) edge set \mathcal{E} , endowed with positive weights

$$(1.50) \quad \lambda(e) > 0, e \in \mathcal{E},$$

so that the corresponding nearest-neighbor walk on G with transition probability

$$(1.51) \quad p_{x,y} = \frac{\lambda(\{x, y\})}{\sum_{z:\{x,z\} \in \mathcal{E}} \lambda(\{x, z\})}, \text{ if } \{x, y\} \in \mathcal{E}, \\ = 0, \text{ otherwise,}$$

is transient. This walk is reversible with respect to the measure

$$(1.52) \quad \lambda_x = \sum_{y:\{x,y\} \in \mathcal{E}} \lambda(\{x, y\}), \quad x \in G.$$

In this set-up some of our definitions need to be modified. For instance if P_x stands for the law of the walk starting from $x \in G$, one divides the right-hand side of (1.5) by λ_y , and multiplies the right-hand side of (1.6) by λ_x ; cf. [23].

The results we stated in Theorem 1.1 and Proposition 1.3, except for (1.28), (1.48), which explicitly refer to the additive structure of \mathbb{Z}^d , can easily be extended to this set-up. We refrain from doing this here since the main results of this article will pertain to percolation properties of the vacant set, which we introduce below, and rely on the structure of \mathbb{Z}^d . \square

We can now define for $\omega \in \Omega$, the *interlacement at level u* , as the subset of \mathbb{Z}^d :

$$(1.53) \quad \mathcal{I}^u(\omega) = \bigcup_{u_i \leq u} \text{range}(w_i^*), \text{ if } \omega = \sum_{i \geq 0} \delta_{(w_i^*, u_i)} \in \Omega, \quad u \geq 0, \\ \stackrel{(1.20)}{=} \bigcup_{K \subset \subset \mathbb{Z}^d} \bigcup_{w \in \text{Supp } \mu_{K,u}(\omega)} w(\mathbb{N}),$$

where for $w^* \in W^*$, $\text{range}(w^*) = w(\mathbb{Z})$, for any $w \in W$ with $\pi^*(w) = w^*$. Note that in view of (1.18), (1.20), the following identity holds:

$$(1.54) \quad \mathcal{I}^u(\omega) \cap K = \bigcup_{w \in \text{Supp } \mu_{K',u}(\omega)} w(\mathbb{N}) \cap K, \text{ for any } K \subset K' \subset \subset \mathbb{Z}^d.$$

The *vacant set at level u* is then defined as

$$(1.55) \quad \mathcal{V}^u(\omega) = \mathbb{Z}^d \setminus \mathcal{I}^u(\omega), \quad \omega \in \Omega, \quad u \geq 0.$$

Obviously with (1.53), (1.55), $\mathcal{I}^u(\omega)$ increases with u , whereas $\mathcal{V}^u(\omega)$ decreases with u . In the next proposition we collect some simple properties of these random subsets. Given $K, \tilde{K} \subset \subset \mathbb{Z}^d$, we say that \tilde{K} separates K from infinity when any nearest neighbor path starting in K and tending to infinity enters \tilde{K} .

PROPOSITION 1.5 ($u \geq 0, K, \tilde{K} \subset \subset \mathbb{Z}^d$).

$$(1.56) \quad \mathcal{I}^u(\omega) \cap K \neq \emptyset \iff \mu_{K,u}(\omega) \neq 0, \\ \text{for } \omega \in \Omega, \text{ and } \mathcal{I}^u, \mathcal{V}^u \text{ depend measurably on } \omega.$$

$$(1.57) \quad \mathbb{P}[K \subseteq \mathcal{V}^u] = \exp\{-u \operatorname{cap}(K)\}.$$

$$(1.58) \quad \mathbb{P}[x \in \mathcal{V}^u] = \exp\left\{-\frac{u}{g(0)}\right\}, \text{ for } x \in \mathbb{Z}^d.$$

$$(1.59) \quad \mathbb{P}[\{x, y\} \subseteq \mathcal{V}^u] = \exp\left\{-\frac{2u}{g(0) + g(y-x)}\right\}, \text{ for } x, y \in \mathbb{Z}^d.$$

If \tilde{K} separates K from infinity then the following inclusion holds:

$$(1.60) \quad \{\mathcal{V}^u \supseteq \tilde{K}\} \subseteq \{\mathcal{V}^u \supseteq K\}, \text{ (screening effect).}$$

Proof. The claim (1.56) immediately follows from (1.54) when $K' = K$, and (1.18), (1.20). The measurability of the sets $\mathcal{G}^u, \mathcal{V}^u$ (understood as the measurability of the maps $1\{x \in \mathcal{G}^u\}$ and $1\{x \in \mathcal{V}^u\}$ for all $x \in \mathbb{Z}^d$) is a direct consequence of the above statement. With (1.56), we thus see that

$$(1.61) \quad \mathbb{P}[\mathcal{V}^u \supseteq K] = \mathbb{P}[\mu_{K,u} = 0] \stackrel{(1.20),(1.43)}{=} \exp\{-u P_{e_K}(W_+)\} \stackrel{(1.7)}{=} \exp\{-u \operatorname{cap}(K)\},$$

and this proves (1.57). As a result of (1.6) or (1.8) one finds that

$$(1.62) \quad \operatorname{cap}(\{x\}) = g(0)^{-1}, \text{ for } x \in \mathbb{Z}^d,$$

and (1.58) follows from (1.57). As for (1.59), we can assume without loss of generality that $x \neq y$, and note that for suitable $\rho_x, \rho_y > 0$, one has

$$(1.63) \quad e_{\{x,y\}} = \rho_x \delta_x + \rho_y \delta_y, \quad \operatorname{cap}(\{x, y\}) = \rho_x + \rho_y,$$

so that with (1.8) one finds:

$$g(z, x) \rho_x + g(z, y) \rho_y = 1, \text{ for } z = x, y.$$

Solving this system of equations we see that $\rho_x = \rho_y = (g(0) + g(y-x))^{-1}$, and hence

$$(1.64) \quad \operatorname{cap}(\{x, y\}) = \frac{2}{g(0) + g(y-x)}, \text{ for } x, y \in \mathbb{Z}^d.$$

The claim (1.59) now follows from (1.57).

Finally note that when \tilde{K} separates K from infinity, $w^* \in W_K^* \implies w^* \in W_{\tilde{K}}^*$, and with (1.56) we see that $\mathcal{G}^u(\omega) \cap K \neq \emptyset \implies \mathcal{G}^u(\omega) \cap \tilde{K} \neq \emptyset$, whence (1.60). \square

Remark 1.6. 1) Using estimates on the capacity of a large cube, cf. for instance (2.4) in Lemma 2.2 of [3] and [17, p. 341], one sees that for $u \geq 0$,

$$(1.65) \quad \mathbb{P}[\mathcal{V}^u \supseteq B(0, L)] = \exp\{-cu L^{d-2}(1 + o(1))\}, \text{ as } L \rightarrow \infty.$$

In particular there is no general exponential decay with $|A|$ of $\mathbb{P}[\mathcal{V}^u \supseteq A]$. This feature is drastically different from what happens for Bernoulli site percolation; see [8]. This creates very serious difficulties when trying to prove that for large u , \mathcal{V}^u does not percolate; see Section 3. Also (1.65) can be compared with (4.58), (4.62)

of Benjamini-Sznitman [2], in the case of the vacant set left by simple random walk on $(\mathbb{Z}/N\mathbb{Z})^d$ up to time $[uN^d]$.

Incidentally, in spite of the fact that \mathcal{V}^u displays a tendency to contain bigger boxes than Bernoulli site percolation, no matter how small $u > 0$, the law Q_u of $1_{\{x \in \mathcal{V}^u\}}$, $x \in \mathbb{Z}^d$, on $\{0, 1\}^{\mathbb{Z}^d}$, does not stochastically dominate Bernoulli site percolation with parameter close to 1. Indeed the complement \mathcal{F}^u of \mathcal{V}^u always percolates.

2) As a direct consequence of (1.57), and the inequality $\text{cap}(K \cup K') \leq \text{cap}(K) + \text{cap}(K')$, we see that

$$(1.66) \quad \mathbb{P}[K \cup K' \subseteq \mathcal{V}^u] \geq \mathbb{P}[K \subseteq \mathcal{V}^u] \mathbb{P}[K' \subseteq \mathcal{V}^u], \text{ for } K, K' \subset \subset \mathbb{Z}^d, u \geq 0,$$

i.e. the events $\{K \subseteq \mathcal{V}^u\}$, $\{K' \subseteq \mathcal{V}^u\}$ are positively correlated. However we do not know whether the FKG Inequality holds under the law Q_u mentioned in 1).

3) As a direct consequence of (1.54) and (1.26), for $K \subset \subset \mathbb{Z}^d$ we can visualize $\mathcal{F}^u \cap K$ as the trace left on K by a Poisson point process of finite trajectories belonging to the space \mathcal{T}_K of (1.22). More precisely for any $u \geq 0$,

$$(1.67) \quad \mathcal{F}^u \cap K \text{ has the same distribution under } \mathbb{P} \text{ as the trace on } K \text{ of a Poisson point process of trajectories on } \mathcal{T}_K \text{ with intensity measure } \rho_K^u(\tau) = u e_K \tau(0) P_{\tau(0)}[X_n = \tau(n), 0 \leq n \leq N_\tau] e_K(\tau(N_\tau)), \text{ for } \tau \in \mathcal{T}_K.$$

This has a very similar flavor to some of the results in Sections 3 and 4 of [2].

4) With standard estimates on the behavior of $g(\cdot)$ at infinity, cf. [11, p. 31], one sees that for any $u \geq 0$,

$$(1.68) \quad \text{cov}_{\mathbb{P}}(1_{\{x \in \mathcal{V}^u\}}, 1_{\{y \in \mathcal{V}^u\}}) \sim \frac{2u}{g(0)^2} g(y-x) e^{-\frac{2u}{g(0)}} \\ \sim \frac{cu}{|y-x|^{d-2}} e^{-cu}, \text{ as } |y-x| \rightarrow \infty,$$

where $\text{cov}_{\mathbb{P}}$ denotes the covariance under \mathbb{P} . This displays the presence of long range correlations in the random set \mathcal{V}^u .

5) Formulas (1.58), (1.59) are in essence (2.26) and (3.6) in Brummelhuis-Hilhorst [4], concerning the large N behavior of the probability that one or two given points in $(\mathbb{Z}/N\mathbb{Z})^d$ are not visited by simple random walk up to time $t = [uN^d]$. The prefactors present in formulas (2.26), (3.6) of [4] stem from the fact that the walk under consideration starts at the origin and not with the uniform distribution as in [2]. For a similar interpretation of (1.58) see also Aldous-Fill [1, Ch. 3, Prop. 20, and Chap. 13, Prop. 8]. One can also compare (1.57) with Propositions 20 and 37 in Chapter 3 of [1]. □

2. A zero-one law and an exponential bound

In this section we exploit the translation invariance of the basic model in a more substantial way. We prove that the probability that \mathcal{V}^u , the vacant set at level u , contains an infinite connected components, is either zero or one. This zero-one law comes as a consequence of the ergodicity of the law of \mathcal{V}^u , cf. Theorem 2.1. We also show in Corollary 2.3 that with probability one \mathcal{F}^u is connected. In Theorem 2.4 we prove an exponential bound on the probability that \mathcal{F}^u contains a given subset of an m -dimensional discrete subspace of \mathbb{Z}^d , with $m \leq d - 3$. This result has a similar flavor to Theorem 2.1 of [2], or Theorem 1.2 of [7], but has a more algebraic proof due to the nature of our basic model. Combined with a Peierls-type argument, cf. Remark 2.5, this can be used to show that when d is large enough, \mathcal{V}^u percolates when u is chosen sufficiently small. In Section 4 we will present a more powerful method proving such a result as soon as $d \geq 7$. We begin with some notation.

We denote with Q_u , the law on $\{0, 1\}^{\mathbb{Z}^d}$ of $(1\{x \in \mathcal{V}^u\})_{x \in \mathbb{Z}^d}$, for $u \geq 0$. We write $Y_x, x \in \mathbb{Z}^d$, for the canonical coordinates on $\{0, 1\}^{\mathbb{Z}^d}$, \mathcal{Y} for the canonical σ -algebra, and $t_x, x \in \mathbb{Z}^d$, for the canonical shift. We also consider for $u \geq 0$ the event

$$(2.1) \text{ Perc}(u) = \{\omega \in \Omega; \mathcal{V}^u(\omega) \text{ contains an infinite connected component}\},$$

as well as

$$(2.2) \quad \eta(u) = \mathbb{P}[0 \text{ belongs to an infinite connected component of } \mathcal{V}^u].$$

The first main result of this section is:

THEOREM 2.1 ($d \geq 3$).

$$(2.3) \quad \text{For any } u \geq 0, (t_x)_{x \in \mathbb{Z}^d} \text{ is a measure-preserving flow on } (\{0, 1\}^{\mathbb{Z}^d}, \mathcal{Y}, Q_u) \text{ which is ergodic,}$$

$$(2.4) \quad \text{For any } u \geq 0, \mathbb{P}[\text{Perc}(u)] = 0 \text{ or } 1.$$

Proof. Beginning with the proof of (2.3), we denote by $\psi_u : \Omega \rightarrow \{0, 1\}^{\mathbb{Z}^d}$, the map $\psi_u(\omega) = (1\{x \in \mathcal{V}^u(\omega)\})_{x \in \mathbb{Z}^d}$, so that $Q_u = \psi_u \circ \mathbb{P}$. Note that with (1.44), (1.53), (1.55), one has

$$(2.5) \quad t_x \circ \psi_u = \psi_u \circ \tau_x, \text{ for } x \in \mathbb{Z}^d.$$

Since \mathbb{P} is invariant under (τ_x) , cf. (1.48), it follows that Q_u is invariant under (t_x) . To prove the ergodicity of (t_x) , we argue as follows. Consider $u \geq 0$, and note that the claim will follow once we show that for any $K \subset \subset \mathbb{Z}^d$, and any $[0, 1]$ -valued

$\sigma(Y_z, z \in K)$ -measurable function f on $\{0, 1\}^{\mathbb{Z}^d}$, one has

$$(2.6) \quad \lim_{|x| \rightarrow \infty} E^{Q_u} [f \circ t_x] = E^{Q_u} [f]^2.$$

Indeed the indicator function of any $A \in \mathcal{G}$ invariant under $(t_x)_{x \in \mathbb{Z}^d}$ can be approximated in $L^1(Q_u)$ by functions f as above. With (2.6) one classically deduces that necessarily $Q_u(A) = Q_u(A)^2$, whence $Q_u(A) \in \{0, 1\}$. In view of (1.54), with $K = K'$, and (2.5), the claim (2.6) will follow once we show that for any $K \subset \subset \mathbb{Z}^d$:

$$(2.7) \quad \lim_{|x| \rightarrow \infty} \mathbb{E}[F(\mu_{K,u}) F(\mu_{K,u}) \circ \tau_x] = \mathbb{E}[F(\mu_{K,u})]^2,$$

for any $[0, 1]$ -valued measurable function F on the set of finite point-measures on W_+ , endowed with its canonical σ -field. With (1.20), (1.44), we can find G (depending on x), with properties similar to F , such that the expectation on the left-hand side of (2.7) equals $\mathbb{E}[F(\mu_{K,u}) G(\mu_{K+x,u})]$.

From now on we assume $|x|$ large enough so that $K \cap (K + x) = \emptyset$. To control the above expectation we are going to express both $\mu_{K,u}$ and $\mu_{K+x,u}$ in terms of $\mu_{K \cup (K+x),u}$, with the help of (1.21) i), and extract the desired asymptotic independence. We will recurrently use this type of decomposition in what follows. Namely with $V = K \cup (K + x)$ we write:

$$(2.8) \quad \begin{aligned} \mu_{V,u} &= \mu_{1,1} + \mu_{1,2} + \mu_{2,1} + \mu_{2,2}, \text{ where} \\ \mu_{1,1}(dw) &= 1\{X_0 \in K, H_{x+K} = \infty\} \mu_{V,u}(dw), \\ \mu_{1,2}(dw) &= 1\{X_0 \in K, H_{x+K} < \infty\} \mu_{V,u}(dw), \end{aligned}$$

and similar formulas for $\mu_{2,2}$ and $\mu_{2,1}$ with the role of K and $K + x$ exchanged. It follows from (1.20), (1.45) that

$$(2.9) \quad \mu_{i,j}, 1 \leq i, j \leq 2, \text{ are independent Poisson point processes on } W_+,$$

and their respective intensity measures are:

$$(2.10) \quad \begin{aligned} \gamma_{1,1} &= u 1\{X_0 \in K, H_{K+x} = \infty\} P_{e_V}, \quad \gamma_{1,2} = u 1\{X_0 \in K, H_{K+x} < \infty\} P_{e_V}, \\ \gamma_{2,1} &= u 1\{X_0 \in K+x, H_K < \infty\} P_{e_V}, \quad \gamma_{2,2} = u 1\{X_0 \in K+x, H_K = \infty\} P_{e_V}. \end{aligned}$$

As a consequence of (1.20), (1.21) i), we see that

$$(2.11) \quad \begin{aligned} \mu_{K,u} &= \mu_{1,1} + \mu_{1,2} + \bar{\mu}_{2,1}^K, \\ \mu_{K+x,u} &= \mu_{2,2} + \mu_{2,1} + \bar{\mu}_{1,2}^{K+x}, \end{aligned}$$

where given $U \subset \subset \mathbb{Z}^d$, and $\mu(dw) = \sum_{0 \leq i \leq N} \delta_{w_i}$ a finite point measure on W_+ , $\bar{\mu}^U(dw) = \sum_{0 \leq i \leq N} \delta_{\theta_{H_U}(w_i)} 1\{H_U(w_i) < \infty\}$, and we have used in (2.11) the fact that $\bar{\mu}_{2,2}^K = 0$, and $\bar{\mu}_{1,1}^{K+x} = 0$. Therefore introducing auxiliary independent

Poisson point processes $\mu'_{1,2}, \mu'_{2,1}$, independent of $\mu_{i,j}, 1 \leq i, j \leq 2$, with the same distribution as $\mu_{1,2}, \mu_{2,1}$ respectively, we find that

$$(2.12) \quad \mu'_{K,u} \stackrel{\text{def}}{=} \mu_{1,1} + \mu_{1,2} + \overline{\mu'}_{2,1}^K, \quad \mu'_{K+x,u} \stackrel{\text{def}}{=} \mu_{2,2} + \mu_{2,1} + \overline{\mu'}_{1,2}^{K+x},$$

are independent point processes respectively distributed as $\mu_{K,u}$ and $\mu_{K+x,u}$. With the same notation as in (1.68) we find that

$$(2.13) \quad \begin{aligned} & \left| \text{cov}_{\mathbb{P}}(F(\mu_{K,u}), G(\mu_{K+x,u})) \right| = \left| \mathbb{E}[F(\mu_{K,u})G(\mu_{K+x,u}) - F(\mu'_{K,u})G(\mu'_{K+x,u})] \right| \\ & \stackrel{(2.11),(2.12)}{\leq} \mathbb{P}[\mu_{1,2} \text{ or } \mu_{2,1} \text{ or } \mu'_{1,2} \text{ or } \mu'_{2,1} \text{ is different from } 0] \\ & \stackrel{(2.9),(2.10)}{\leq} 2(1 - \exp\{-\gamma_{1,2}(W_+)\}) + 2(1 - \exp\{-\gamma_{2,1}(W_+)\}) \\ & \leq 2u(P_{e_V}[X_0 \in K, H_{K+x} < \infty] + P_{e_V}[X_0 \in K+x, H_K < \infty]), \end{aligned}$$

where in the last step we have used the inequality $1 - e^{-v} \leq v$, for $v \geq 0$, in addition to (2.10). Observe now that

$$(2.14) \quad \begin{aligned} P_{e_V}[X_0 \in K, H_{K+x} < \infty] &= \sum_{z \in K} e_V(z) P_z[H_{K+x} < \infty] \\ &\stackrel{(1.8)}{=} \sum_{z \in K, y \in K+x} e_V(z) g(z, y) e_{K+x}(y) \\ &\stackrel{(1.6)}{\leq} \sum_{z \in K, y \in K+x} e_K(z) g(z, y) e_{K+x}(y) \leq c \frac{\text{cap}(K)^2}{d(K, K+x)^{d-2}}, \end{aligned}$$

with the notation introduced above (1.1), as well as standard bounds on the Green function, cf. [11, p. 31], and translation invariance. A similar bound holds for $P_{e_V}[X_0 \in K+x, H_K < \infty]$, and with (2.13) we see that for $u \geq 0, K \subset\subset \mathbb{Z}^d, x \in \mathbb{Z}^d, F, G$ -measurable functions on the set of finite point measures on W_+ with values in $[0, 1]$,

$$(2.15) \quad \left| \text{cov}_{\mathbb{P}}(F(\mu_{K,u}), G(\mu_{K+x,u})) \right| \leq c u \frac{\text{cap}(K)^2}{d(K, K+x)^{d-2}}.$$

This implies (2.7) and thus concludes the proof of (2.3). As for (2.4), note that $\text{Perc}(u) = \psi_u^{-1}(A)$, where $A \in \mathcal{Q}$ stands for the invariant event consisting of configurations in $\{0, 1\}^{\mathbb{Z}^d}$ such that there is an infinite connected component in the subset of \mathbb{Z}^d where the configuration takes the value 1. It now follows from (2.3) that $Q_u(A) = \mathbb{P}[\text{Perc}(u)]$ is either 0 or 1. This proves (2.4). \square

Remark 2.2. 1) Note that (2.15) has a similar flavor to (1.68), which mirrors the long range dependence built into the basic model. Taming this effect will bring some serious difficulties in Section 3.

2) One can characterize Q_u as the unique probability on $(\{0, 1\}^{\mathbb{Z}^d}, \mathfrak{y})$ such that

$$(2.16) \quad Q_u(Y_z = 1, \text{ for } z \in K) = \exp\{-u \text{cap}(K)\}, \text{ for any } K \subset\subset \mathbb{Z}^d.$$

Indeed the collection of events which appear in (2.16) is stable under finite intersection and generates \mathfrak{y} . In a slightly more constructive fashion, we see with a classical inclusion exclusion argument that for any disjoint finite subsets K, K' of \mathbb{Z}^d , one has

$$(2.17) \quad Q_u[Y_z = 1, \text{ for } z \in K, Y_z = 0, \text{ for } z \in K'] \\ = E^{Q_u} \left[\prod_{z \in K} Y_z \prod_{z \in K'} (1 - Y_z) \right] = \sum_{A \subseteq K'} (-1)^{|A|} \exp\{-u \text{cap}(K \cup A)\}.$$

3) The present work does not address the question of whether there is a unique infinite connected component in \mathfrak{V}^u when it percolates and u is positive. The answer to this question is affirmative, as proved in [22]. The classical results of Burton-Keane [5], see also [9, pp. 326, 332], implying such a uniqueness do not apply because, as one easily sees, Q_u fails to fulfill the so-called finite energy condition:

$$0 < Q_u(Y_x = 1 | Y_z, z \neq x) < 1, \text{ } Q_u\text{-a.s.}, \text{ for all } x \in \mathbb{Z}^d.$$

Loosely speaking the problem stems from the fact that the set of sites w , where Y_w takes the value 0, has no bounded component, and on some configurations turning a value 0 into a value 1, say at the origin, can lead to a forbidden configuration (see also (1.67)). In Corollary 2.3 we are able to adapt the argument of Burton-Keane in the case of \mathfrak{F}^u , and prove that with probability one, \mathfrak{F}^u is connected. In the case of \mathfrak{V}^u the construction of so-called trifurcations is more delicate, and can be found in [22].

4) Denote with $\mathbb{E}_d = \{\{x, y\}; x, y \text{ in } \mathbb{Z}^d \text{ with } |x - y| = 1\}$, the collection of nearest neighbor edges on \mathbb{Z}^d . Given $\omega \in \Omega$ and $u \geq 0$, one can consider the subset $\tilde{\mathfrak{F}}^u(\omega)$ of \mathbb{E}_d consisting of the edges which are traversed by at least one of the trajectories at level u entering ω :

$$(2.18) \quad \tilde{\mathfrak{F}}^u(\omega) = \{e \in \mathbb{E}_d; \text{ for some } i \geq 0, \text{ with } u_i \leq u \text{ and } n \in \mathbb{Z}, \\ e = \{w_i(n), w_i(n + 1)\}\}, \text{ if } \omega = \sum_{i \geq 0} \delta_{(w_i^*, u_i)} \in \Omega,$$

and w_i is any element of W with $\pi^*(w_i) = w_i^*$.

Connected components of \mathbb{Z}^d induced by $\tilde{\mathfrak{F}}^u(\omega)$ are either singletons in $\mathfrak{F}^u(\omega)^c$ or infinite components partitioning $\mathfrak{F}^u(\omega)$. Denoting with $\tilde{\psi}_u: \Omega \rightarrow \{0, 1\}^{\mathbb{E}_d}$ the map $\tilde{\psi}_u(\omega) = (1\{e \in \tilde{\mathfrak{F}}^u(\omega)\})_{e \in \mathbb{E}_d}$, one can consider the image \tilde{Q}_u on $(\{0, 1\}^{\mathbb{E}_d}, \tilde{\mathfrak{y}})$ of \mathbb{P} under $\tilde{\psi}_u$, where $\tilde{\mathfrak{y}}$ stands for the canonical σ -algebra on $\{0, 1\}^{\mathbb{E}_d}$. With \tilde{t}_x ,

$x \in \mathbb{Z}^d$, the canonical shift on $\{0, 1\}^{\mathbb{E}^d}$, one finds exactly as in (2.5) that $\tilde{t}_x \circ \tilde{\psi}_u = \tilde{\psi}_u \circ \tau_x$, for $x \in \mathbb{Z}^d$. The same proof as in (2.3), see in particular (2.7), now yields

$$(2.19) \quad \text{For any } u \geq 0, (\tilde{t}_x)_{x \in \mathbb{Z}^d} \text{ is a measure-preserving flow on } (\{0, 1\}^{\mathbb{E}^d}, \tilde{\mathcal{Y}}, \tilde{Q}_u) \text{ which is ergodic. } \square$$

The first statement below is an immediate consequence of Theorem 2.1 and (2.2).

COROLLARY 2.3 ($d \geq 3$). *For $u \geq 0$, one has the equivalences*

$$(2.20) \quad \begin{aligned} \text{i)} \quad & \mathbb{P}[\text{Perc}(u)] = 1 \iff \eta(u) > 0, \\ \text{ii)} \quad & \mathbb{P}[\text{Perc}(u)] = 0 \iff \eta(u) = 0. \end{aligned}$$

$$(2.21) \quad \text{For } u > 0, \mathbb{P}\text{-a.s.}, \mathcal{I}^u \text{ is an infinite connected subset of } \mathbb{Z}^d .$$

Proof. We begin with (2.20). One simply needs to observe that

$$\eta(u) \leq \mathbb{P}[\text{Perc}(u)] \leq \sum_{x \in \mathbb{Z}^d} \mathbb{P} [x \text{ belongs to an infinite connected component of } \mathcal{V}^u] ,$$

and in view of (1.48) all summands on the right-hand side equal $\eta(u)$. The claim (2.20) now follows from the zero-one law (2.4).

We now turn to the proof of (2.21), which is an adaptation of the argument of Burton-Keane [5]. The consideration of $\tilde{\mathcal{I}}^u$, cf. Remark 2.2 (4), will be helpful; see in particular the observation below (2.18). With the ergodicity property (2.19), it follows that the total number N_u of infinite connected components determined by $\tilde{\mathcal{I}}^u$ is \mathbb{P} -a.s. equal to a positive, possibly infinite, constant. With the observation below (2.18) our claim (2.21) will follow once we show that this constant equals 1. The first step, see also [13], is to argue that

$$(2.22) \quad \text{for } 2 \leq k < \infty, \mathbb{P}[N_u = k] = 0 .$$

Assume instead that for some $2 \leq k < \infty$, $\mathbb{P}[N_u = k] = 1$. Then we can find $K = B(0, L)$ such that $\mathbb{P}[A] > 0$, where A denotes the event $\{N_u = k \text{ and } K \text{ intersects two distinct infinite components determined by } \tilde{\mathcal{I}}^u(\omega)\}$. Note that under \mathbb{P}

$$(2.23) \quad \omega_K^1 = 1_{W_K^* \times \mathbb{R}_+} \omega \text{ and } \omega_K^0 = 1_{(W_K^*)^c \times \mathbb{R}_+} \omega \text{ are two independent Poisson point processes with respective intensity measures } 1_{W_K^* \times \mathbb{R}_+} dv du \text{ and } 1_{(W_K^*)^c \times \mathbb{R}_+} dv du .$$

For each $z \in S(0, L)$, the ‘‘surface of K ’’, we now pick a nearest neighbor loop in K starting and ending at z , and passing through 0. We then define a map φ from W_K^* into itself such that for $w^* \in W_K^*$, $\varphi(w^*)$ is the trajectory (modulo time-shift) obtained by ‘‘inserting in w^* ’’ just after the entrance in K , the loop attached to the entrance point of w^* in K . The map φ is in fact injective and one checks with (1.25), (1.26) that the image measure $\varphi \circ (1_{W_K^*} \nu)$ is absolutely continuous with

respect to $1_{W_K^* \nu}$. We extend φ to W^* , by letting φ be the identity map on $(W_K^*)^c$. It now follows from the above observations that the measurable map Φ from Ω into itself defined by:

$$\Phi(\omega) = \sum_{u_i \leq u} \delta_{(\varphi(w_i^*), u_i)} + \sum_{u_i > u} \delta_{(w_i^*, u_i)}, \text{ for } \omega = \sum_{i \geq 1} \delta_{(w_i^*, u_i)},$$

is such that

$$(2.24) \quad \Phi \circ \mathbb{P} \text{ is absolutely continuous with respect to } \mathbb{P}.$$

By construction $\Phi(\omega)$ links together all infinite connected components of $\tilde{\mathcal{F}}^u(\omega)$, which intersect K , and hence $\Phi(A) \subseteq \{N_u < k\}$, where A appears below (2.22). We thus find that

$$(2.25) \quad \Phi \circ (1_A \mathbb{P})[N_u < k] = \mathbb{P}[A \cap \Phi^{-1}(N_u < k)] = \mathbb{P}[A] > 0,$$

and due to (2.24) we see that $\mathbb{P}[N_u < k] > 0$, a contradiction. This proves (2.22). The claim (2.21) will now follow once we show that

$$(2.26) \quad \mathbb{P}[N_u = \infty] = 0.$$

The heart of the matter, cf. [8, p. 199], or [9, p. 297], is to show that with positive \mathbb{P} -probability there is a trifurcation in $\tilde{\mathcal{F}}^u$, i.e. a site $x \in \mathcal{J}^u(\omega)$ with exactly three $\tilde{\mathcal{F}}^u(\omega)$ -neighbors and the removal of x splits the infinite connected component of x determined by $\tilde{\mathcal{F}}^u(\omega)$ in exactly three infinite components.

Assume by contradiction that $\mathbb{P}[N_u = \infty] = 1$, then for arbitrarily large $L > 0$, one has with $K = B(0, L)$,

$$(2.27) \quad \mathbb{P}[K \text{ intersects more than } 4|B(0, 100)| \text{ infinite connected components of } \tilde{\mathcal{F}}^u(\omega)] > 0.$$

We fix L large enough, such that (2.27) holds and for any three couples of points $(z_1, z_2), (z_3, z_4), (z_5, z_6)$ on the $|\cdot|_\infty$ -sphere $S(0, L)$, for which no point in a given pair may be within $|\cdot|_\infty$ -distance 100 from any other pair (but points within a pair may be arbitrarily close or even coincide), we can construct τ_1, τ_2, τ_3 finite nearest neighbor trajectories in K with respective starting points z_1, z_3, z_5 and end points z_2, z_4, z_6 , so that any two trajectories only meet in 0, each trajectory visits 0 only once, and this occurs by crossing an edge touching 0 and immediately crossing the same edge in the reverse direction. With (2.23) and (2.27) we see that

$$\mathbb{P} \otimes (1_{W_K^* \nu})^{\otimes m} [K \text{ intersects more than } 4|B(0, 100)| \text{ infinite connected components determined by } \tilde{\mathcal{F}}^u(\omega_K^0 + \sum_{i=1}^m \delta_{(w_i^*, u)})] > 0, \text{ for some } m > 4|B(0, 100)|,$$

where ω and $w_i^*, 1 \leq i \leq m$, are the respective Ω - and W_K^* -valued coordinates on the product space, by notation from (2.18) and (2.23). From the above event we

can select three trajectories within the w_1^*, \dots, w_m^* with supports lying in distinct infinite connected components and corresponding pairs of entrance and last exit points of K with mutual $|\cdot|_\infty$ -distance bigger than 100. As a result we see that

$$(2.28) \quad \mathbb{P} \otimes (1_{W_K^* \nu})^{\otimes m} [C_m] > 0, \text{ for some } m \geq 3,$$

where C_m stands for the event

$\{\tilde{\mathcal{F}}^u(\omega_K^0 + \sum_{i=1}^m \delta_{(w_i^*, u)})$ has at least three infinite connected components meeting K respectively containing $w_{i_1}^*(\mathbb{Z}), w_{i_2}^*(\mathbb{Z}), w_{i_3}^*(\mathbb{Z})$ for some distinct i_1, i_2, i_3 in $\{1, \dots, m\}$, and the three corresponding pairs of entrance and last exit points of K have mutual $|\cdot|_\infty$ -distance bigger than 100 $\}$.

Observe now that without loss of generality we can assume $m = 3$ in (2.28).

We denote with γ the map from $(W_K^*)^3$ into itself such that $\gamma(w_1^*, w_2^*, w_3^*) = (\bar{w}_1^*, \bar{w}_2^*, \bar{w}_3^*)$, where γ simply coincides with the identity if the three pairs of entrance and last exit points for K for w_1^*, w_2^*, w_3^* do not fulfill the condition appearing below (2.27), and otherwise such that $\bar{w}_1^*, \bar{w}_2^*, \bar{w}_3^*$ are obtained from w_1^*, w_2^*, w_3^* by replacing the respective portions of trajectory between first entrance in K and last exit from K by τ_1, τ_2, τ_3 . With (1.25), (1.26) one checks that

$$(2.29) \quad \gamma \circ (1_{W_K^* \nu})^{\otimes 3} \text{ is absolutely continuous with respect to } (1_{W_K^* \nu})^{\otimes 3}.$$

Note that in the event C_3 , 0 is a trifurcation point for $\tilde{\mathcal{F}}^u(\omega_K^0 + \sum_{i=1}^3 \delta_{(\bar{w}_i^*, u)})$, where the notation is the same as in the above paragraph. With a calculation similar to (2.25) we see that

$$\mathbb{P} \otimes (1_{W_K^* \nu})^{\otimes 3} [0 \text{ is a trifurcation point for } \tilde{\mathcal{F}}^u(\omega_K^0 + \sum_{i=1}^3 \delta_{(w_i^*, u)})] > 0.$$

With (2.23) this readily implies that

$$(2.30) \quad \mathbb{P}[0 \text{ is a trifurcation point for } \tilde{\mathcal{F}}^u(\omega)] > 0.$$

The proof of (2.26) now runs just as in [8, p. 200–202]. This concludes the proof of (2.21). □

Just as in the case of Bernoulli percolation, cf. [8, p. 13], we can introduce the critical value

$$(2.31) \quad u_* = \inf\{u \geq 0, \eta(u) = 0\} \in [0, \infty].$$

Is this critical value nondegenerate? We will see in Section 3 that $u_* < \infty$, cf. Theorem 3.5, and in fullrefsecfo that $u_* > 0$, as soon as $d \geq 7$, cf. Theorem 4.3.

We are now going to discuss the exponential bound mentioned at the beginning of this section. For $1 \leq m \leq d$, we write \mathcal{L}_m for the collection of m -dimensional affine subspaces of \mathbb{Z}^d generated by m distinct vectors of the canonical

basis $(e_i)_{1 \leq i \leq d}$ of \mathbb{R}^d :

(2.32)

$$\mathcal{L}_m = \left\{ F \subseteq \mathbb{Z}^d; \text{ for some } I \subseteq \{1, \dots, d\} \text{ with } |I| = m \text{ and some } y \in \mathbb{Z}^d, \right. \\ \left. F = y + \sum_{i \in I} \mathbb{Z} e_i \right\},$$

and introduce

(2.33) $\mathcal{A}_m =$ the collection of finite subsets A with $A \subseteq F$ for some $F \in \mathcal{L}_m$.

We denote with $q(v)$ the return probability to the origin of simple random walk in \mathbb{Z}^v , i.e. with (we hope) obvious notation:

(2.34)
$$q(v) = P_0^{\mathbb{Z}^v} [\tilde{H}_0 < \infty], \text{ for } v \geq 1.$$

The promised exponential estimate comes in the following:

THEOREM 2.4 ($d \geq 4, 1 \leq m \leq d - 3$). Assume that $\lambda > 0$ satisfies

(2.35)
$$\chi(\lambda) \stackrel{\text{def}}{=} e^\lambda \left(\frac{m}{d} + \left(1 - \frac{m}{d} \right) q(d - m) \right) < 1;$$

then for $u \geq 0, A \in \mathcal{A}_m$ and $A \subseteq K \subset \subset \mathbb{Z}^d$, with the notation

$$f_A(w) = \sum_{n \geq 0} 1_{\{X_n(w) \in A\}},$$

for $w \in W_+$, one has

(2.36)
$$\mathbb{E}[\exp\{\lambda \langle \mu_{K,u}, f_A \rangle\}] \leq \exp \left\{ u \operatorname{cap}(A) \frac{e^\lambda - 1}{1 - \chi(\lambda)} \right\},$$

and the left-hand side does not depend on K as above.

Moreover there exists $u_1(d, m, \lambda) > 0$, such that:

(2.37)
$$\mathbb{P}[\mathcal{G}^u \supseteq A] \leq \exp\{-\lambda |A|\}, \text{ for all } A \in \mathcal{A}_m \text{ and } u \leq u_1.$$

Proof. Consider $A \in \mathcal{A}_m, F \in \mathcal{L}_m$ containing A ; then for $A \subseteq K \subseteq K' \subset \subset \mathbb{Z}^d$, we find that

$$\langle \mu_{K,u}, f_A \rangle \stackrel{(1.21)i)}{=} \langle \mu_{K',u}, f_A \circ \theta_{H_K} 1_{\{H_K < \infty\}} \rangle = \langle \mu_{K',u}, f_A \rangle.$$

So the left-hand side of (2.36) does not depend on $K \subset \subset \mathbb{Z}^d$ containing A . In particular picking $K = A$, we find that it equals

(2.38)
$$\mathbb{E}[\exp\{\lambda \langle \mu_{A,u}, f_A \rangle\}] \stackrel{(1.20),(1.43)}{=} \exp\{u E_{e_A}[e^{\lambda f_A} - 1]\}.$$

Introducing the function

(2.39)
$$\phi(x) = E_x[e^{\lambda f_A}], \text{ for } x \in \mathbb{Z}^d,$$

and writing $R_F \stackrel{\text{def}}{=} T_F + H_F \circ \theta_{T_F}$, the return time to F , see (1.3) for notation, we find:

$$\begin{aligned} e^{\lambda f_A} &\leq e^{\lambda T_F} (1_{\{R_F = \infty\}} + 1_{\{R_F < \infty\}} e^{\lambda f_A} \circ \theta_{R_F}) \\ &= e^{\lambda T_F} (1 + 1_{\{R_F < \infty\}} (e^{\lambda f_A} \circ \theta_{R_F} - 1)). \end{aligned}$$

With the strong Markov property at times R_F and then T_F , we thus obtain:

$$\begin{aligned} (2.40) \quad \phi(x) &\leq E_x[e^{\lambda T_F}] + E_x[e^{\lambda T_F} P_{X_{T_F}}[H_F < \infty]](\|\phi\|_\infty - 1) \\ &\stackrel{(2.34)}{=} E_x[e^{\lambda T_F}](1 + q(d - m)(\|\phi\|_\infty - 1)), \end{aligned}$$

considering in the last step the motion of the walk in the components “transversal” to F . Note that when $z \notin F$, $T_F = 0$, P_z -a.s., whereas when $z \in F$, T_F has geometric distribution with success probability $1 - \frac{m}{d}$. Hence with λ satisfying (2.35) we find that:

$$(2.41) \quad E_z[\exp\{\lambda T_F\}] = \sum_{k \geq 1} \left(1 - \frac{m}{d}\right) \left(\frac{m}{d}\right)^{k-1} e^{\lambda k} = e^{\lambda} \left(1 - \frac{m}{d}\right) \left(1 - e^{\lambda} \frac{m}{d}\right)^{-1} \stackrel{\text{def}}{=} \alpha.$$

With a routine approximation argument of f_A by a finite sum, to exclude the possibility that $\|\phi\|_\infty$ is infinite, and (2.40) we see that:

$$\|\phi\|_\infty \leq \frac{\alpha(1 - q(d - m))}{1 - q(d - m)\alpha},$$

and hence

$$(2.42) \quad \|\phi\|_\infty - 1 \leq \frac{\alpha - 1}{1 - q(d - m)\alpha} = \frac{e^\lambda - 1}{1 - \chi(\lambda)}.$$

Coming back to (2.38), and using (1.7) we find (2.36). As for (2.37), note with (1.6), (1.62) that

$$\text{cap}(A) \leq \sum_{x \in A} \text{cap}(\{x\}) = \frac{|A|}{g(0)}.$$

Further, on the event $\{\mathcal{G}^u \supseteq A\}$ we have $\langle \mu_{A,u}, f_A \rangle \geq |A|$. So choosing $\tilde{\lambda}(d, m, \lambda) > \lambda$, such that $1 - \chi(\tilde{\lambda}) = \frac{1}{2} (1 - \chi(\lambda))$, we now see that for $A \in \mathcal{A}_m$:

$$\begin{aligned} (2.43) \quad \mathbb{P}[\mathcal{G}^u \supseteq A] &\stackrel{(2.36)}{\leq} \exp\left\{-\tilde{\lambda}|A| + \frac{|A|}{g(0)} u \frac{e^{\tilde{\lambda}} - 1}{1 - \chi(\tilde{\lambda})}\right\} \\ &\leq \exp\{-\lambda|A|\}, \end{aligned}$$

if $u \leq u_1(d, m, \lambda)$. This proves (2.37). □

Remark 2.5. 1) The proof of Theorem 2.4 is very similar to the proofs of Theorem 2.1 of [2] and Theorem 1.2 of [7]; however it has a somewhat more algebraic character due to the nature of the basic model we work with.

2) There is no bound of type (2.37) valid uniformly for \mathcal{A}_d , the collection of subsets of \mathbb{Z}^d . The argument is in essence the same as in Remark 2.4 (2) of [2]. One can for instance consider $A_L = B(0, L)$ and note that for large L , when the random walk starts in A_L , conditionally on not leaving A_{2L} up to time $c L^d \log L$ (with c a large enough constant), it covers A_L with probability at least $\frac{1}{2}$; cf. (2.33) of [2]. From this it follows that for large L ,

$$\begin{aligned} \mathbb{P}[\mathcal{G}^u \supseteq A_L] &\geq \mathbb{P}[\mu_{A_L, u} \neq 0] \frac{1}{2} \inf_{x \in A_L} P_x[T_{A_{2L}} > c L^d \log L] \\ &\geq c(1 - \exp\{-u \operatorname{cap}(A_L)\}) \exp\{-c L^{d-2} \log L\}. \end{aligned}$$

As a result no matter how small $u > 0$, one finds that

$$(2.44) \quad \lim_{L \rightarrow \infty} |A_L|^{-1} \log \mathbb{P}[\mathcal{G}^u \supseteq A_L] = 0.$$

3) One can combine (2.37) with a Peierls-type argument by considering the collection of $*$ -nearest neighbor circuits separating 0 from infinity in some $F \in \mathcal{L}_2$ containing 0; cf. Corollary 2.5 of [2] or Corollary 1.5 of [7]. One sees that when d satisfies

$$(2.45) \quad 7\left(\frac{2}{d} + \left(1 - \frac{2}{d}\right) q(d-2)\right) < 1,$$

then for small $u > 0$, \mathcal{V}^u percolates; i.e.,

$$(2.46) \quad \mathbb{P}[\operatorname{Perc}(u)] = 1, \text{ for small } u > 0.$$

The factor 7 in (2.45) simply stems from the fact that there are at most eight 7^{n-1} $*$ -nearest neighbor circuits with n steps in \mathbb{Z}^2 that start at the origin. It is known that $q(v) \sim (2v)^{-1}$, as $v \rightarrow \infty$, cf. (5.4) of [12], and hence (2.45) holds for large d . Clearly (2.45) forces $d > 14$, and with the help of tables of values for $q(\cdot)$, one can see that in effect (2.45) holds exactly when $d \geq 18$; cf. Remark 2.1 of [7]. In section 4 we will show that (2.46) holds when $d \geq 7$. \square

3. Absence of percolation for large u

The principal object of this section is to show in Theorem 3.5 that when $d \geq 3$, for large enough u , \mathbb{P} -almost surely all connected components of \mathcal{V}^u are finite. We know from Remark 1.6 (1) or (1.57) that in general $\mathbb{P}[\mathcal{V}^u \supseteq A]$ does not decay exponentially with $|A|$. This creates an obstruction to the classical Peierls-type argument, which is used in the context of Bernoulli percolation. It substantially complicates the matter. The strategy of the proof we present here is instead based on a renormalization argument. We establish in Proposition 3.1 key estimates on the probability of existence of certain crossings at scale L_n in \mathcal{V}^{u_n} , cf. (3.7), (3.8), on an increasing sequence of length scales L_n and an increasing but bounded sequence of values u_n . The proof of Proposition 3.1 uses a recurrence propagating

certain controls, cf. (3.10), from one scale to the next along a sequence of level-values as in (3.9). Once Proposition 3.1 is established it is a simple matter to deduce Theorem 3.5. We will now introduce some notation.

We consider the positive number a and an integer L_0 :

$$(3.1) \quad a = \frac{1}{100d}, \quad L_0 > 1.$$

We then define an increasing sequence of length scales via

$$(3.2) \quad L_{n+1} = \ell_n L_n, \text{ where } \ell_n = 100[L_n^a](\geq L_n^a), \text{ for } n \geq 0,$$

so that $L_n, n \geq 0$, quickly grows to infinity:

$$(3.3) \quad L_n \geq L_0^{(1+a)^n}, \text{ for } n \geq 0.$$

We organize \mathbb{Z}^d in a hierarchical way with L_0 corresponding to the bottom scale and $L_1 < L_2 < \dots$ representing coarser and coarser scales. For this purpose, given $n \geq 0$, we consider the set of labels at level n :

$$(3.4) \quad I_n = \{n\} \times \mathbb{Z}^d.$$

To each label at level $n, m = (n, i) \in I_n$, we associate the boxes:

$$(3.5) \quad C_m = (iL_n + [0, L_n)^d) \cap \mathbb{Z}^d, \\ \tilde{C}_m = \bigcup_{m' \in I_n: d(C_{m'}, C_m) \leq 1} C_{m'},$$

where we refer to the notation above (1.1). It is straightforward to see that $C_m, m \in I_n$, is a partition of \mathbb{Z}^d into boxes of side-length $L_n - 1$, and \tilde{C}_m simply stands for the union of C_m and its “*-neighboring” boxes of level n . Also when $m \in I_{n+1}$, then C_m is the disjoint union of the ℓ_n^d boxes $C_{\bar{m}}$ at level n it contains. We denote with \tilde{S}_m the interior boundary of \tilde{C}_m :

$$(3.6) \quad \tilde{S}_m = \partial_{\text{int}} \tilde{C}_m, \text{ for } m \in I_n, n \geq 0.$$

In what follows we investigate the probability of the existence of certain vacant crossings defined for $u \geq 0, n \geq 0, m \in I_n$, via:

$$(3.7) \quad A_m^u = \{\omega \in \Omega; \text{ there is a nearest neighbor path in } \mathcal{V}^u(\omega) \cap \tilde{C}_m \text{ from } C_m \text{ to } \tilde{S}_m\}.$$

It follows from translation invariance, cf. Theorem 2.1 or (1.48), that

$$(3.8) \quad p_n(u) = \mathbb{P}[A_m^u], \quad u \geq 0, n \geq 0, \text{ with } m \in I_n,$$

is well-defined, i.e. does not depend on which $m \in I_n$ enters the right-hand side. Clearly the functions $p_n(\cdot)$ are nonincreasing on \mathbb{R}_+ . Our main task consists in

the derivation of recurrence relations on the functions $p_n(\cdot)$. The key control is provided by the following

PROPOSITION 3.1 ($d \geq 3$). *There exist positive constants c_1, c_2 , cf. (3.34), (3.52), such that defining for $u_0 > 0$ and $r \geq 1$ integer*

$$(3.9) \quad u_n = u_0 \prod_{0 \leq n' < n} (1 + c_1 \ell_{n'}^{-(d-2)})^{r+1}, \text{ for } n \geq 0,$$

one has for $L_0 \geq c, u_0 \geq c(L_0), r \geq c(L_0, u_0)$,

$$(3.10) \quad c_2 \ell_n^{2(d-1)} p_n(u_n) \leq L_n^{-1}, \text{ for all } n \geq 0.$$

Proof. In the course of the proof of Proposition 3.1, we will use the expression “for large L_0 ”, in place of “for $L_0 \geq c$ ”, with c a positive constant as explained at the end of the introduction. We first consider $n \geq 0, m \in I_{n+1}$, as well as $0 < u' < u$. We are first going to bound $p_{n+1}(u)$ in terms of $p_n(u')$, when $\frac{u'}{u}$ is sufficiently away from 1, cf. (3.45), (3.52).

We write \mathcal{H}_1 for the collection of labels at level n of boxes contained in C_m touching $\partial_{\text{int}} C_m$:

$$(3.11) \quad \mathcal{H}_1 = \{\bar{m} \in I_n; C_{\bar{m}} \subseteq C_m \text{ and } C_{\bar{m}} \cap \partial_{\text{int}} C_m \neq \emptyset\},$$

as well as

$$(3.12) \quad \mathcal{H}_2 = \left\{ \bar{m} \in I_n; C_{\bar{m}} \cap \left\{ z \in \mathbb{Z}^d : d(z, C_m) = \frac{L_{n+1}}{2} \right\} \neq \emptyset \right\},$$

for the collection of labels of n -level boxes containing some point at $|\cdot|_\infty$ -distance $\frac{L_{n+1}}{2}$ from C_m (with similar notation as above (1.1)).

Observe that any nearest neighbor path in \mathcal{V}^u originating in C_m and ending in \tilde{S}_m must go through some $C_{\bar{m}_1}, \bar{m}_1 \in \mathcal{H}_1$, reach $\tilde{S}_{\bar{m}_1}$, and then go through some $C_{\bar{m}_2}, \bar{m}_2 \in \mathcal{H}_2$, and reach $\tilde{S}_{\bar{m}_2}$. Therefore we see that

$$(3.13) \quad p_{n+1}(u) \leq \sum_{\bar{m}_1 \in \mathcal{H}_1, \bar{m}_2 \in \mathcal{H}_2} \mathbb{P}[A_{\bar{m}_1}^u \cap A_{\bar{m}_2}^u] \leq c \ell_n^{2(d-1)} \sup_{\bar{m}_1 \in \mathcal{H}_1, \bar{m}_2 \in \mathcal{H}_2} \mathbb{P}[A_{\bar{m}_1}^u \cap A_{\bar{m}_2}^u],$$

using a rough counting argument to bound $|\mathcal{H}_1|$ and $|\mathcal{H}_2|$ in the last step. We will now focus our attention on the probability which appears in the last member of (3.13). We write $V = \tilde{C}_{\bar{m}_1} \cup \tilde{C}_{\bar{m}_2}$ for given $\bar{m}_1 \in \mathcal{H}_1, \bar{m}_2 \in \mathcal{H}_2$, and just as in (2.8) introduce the decomposition

$$(3.14) \quad \mu_{V,u} = \mu_{1,1} + \mu_{1,2} + \mu_{2,1} + \mu_{2,2},$$

where $\tilde{C}_{\bar{m}_1}, \tilde{C}_{\bar{m}_2}$ respectively play the role of K and $K + x$ in (2.8). In particular $\mu_{i,j}, 1 \leq i, j \leq 2$, are independent Poisson point processes on W_+ with intensity measures $\gamma_{i,j}, 1 \leq i, j \leq 2$, as in (2.10). The following notation will be convenient. When Λ is a random point process on W_+ defined on Ω , i.e. a measurable map




Figure 1. A schematic illustration of the event A_m^u . The path drawn lies in \mathcal{V}^u .

from Ω into the space of pure point measures on W_+ , we denote with $A_{\bar{m}}(\Lambda)$, for $\bar{m} \in I_n$, the event:

$$(3.15) \quad A_{\bar{m}}(\Lambda) = \left\{ \omega \in \Omega; \text{there is a nearest neighbor path in } \tilde{C}_{\bar{m}} \setminus \left(\bigcup_{w \in \text{Supp}(\Lambda(\omega))} w(\mathbb{N}) \right) \text{ from } C_{\bar{m}} \text{ to } \tilde{S}_{\bar{m}} \right\}.$$

For instance with (1.54) we see that for any $\bar{m} \in I_n$:

$$(3.16) \quad A_{\bar{m}}^u = A_{\bar{m}}(\mu_{K,u}), \text{ for any } K \supseteq \tilde{C}_{\bar{m}}.$$

We can apply this identity to $\bar{m} = \bar{m}_i, i = 1, 2$, with $K = V$. Noting that $w \in \text{Supp} \mu_{2,2}$ implies $w(\mathbb{N}) \cap \tilde{C}_{\bar{m}_1} = \emptyset$, we find that

$$\begin{aligned} A_{\bar{m}_1}^u \cap A_{\bar{m}_2}^u &= A_{\bar{m}_1}(\mu_{V,u}) \cap A_{\bar{m}_2}(\mu_{V,u}) \\ &= A_{\bar{m}_1}(\mu_{1,1} + \mu_{1,2} + \mu_{2,1}) \cap A_{\bar{m}_2}(\mu_{V,u}) \\ &\subseteq A_{\bar{m}_1}(\mu_{1,1} + \mu_{1,2} + \mu_{2,1}) \cap A_{\bar{m}_2}(\mu_{2,2}). \end{aligned}$$

With the help of the independence properties mentioned above we find that

$$(3.17) \quad \begin{aligned} \mathbb{P}[A_{\bar{m}_1}^u \cap A_{\bar{m}_2}^u] &\leq \mathbb{P}[A_{\bar{m}_1}(\mu_{1,1} + \mu_{1,2} + \mu_{2,1})] \mathbb{P}[A_{\bar{m}_2}(\mu_{2,2})] \\ &= p_n(u) \mathbb{P}[A_{\bar{m}_2}(\mu_{2,2})]. \end{aligned}$$

Our next task is to bound $\mathbb{P}[A_{\bar{m}_2}(\mu_{2,2})]$ from above. For this purpose we decompose the $\mu_{i,j}$, $1 \leq i, j \leq 2$, in (3.14) into

$$(3.18) \quad \mu_{i,j} = \mu'_{i,j} + \mu^*_{i,j},$$

where the $\mu'_{i,j}, \mu^*_{i,j}$, $1 \leq i, j \leq 2$, are independent Poisson point processes on W_+ , with $\mu'_{i,j}$ defined as $\mu_{i,j}$, with $u' (< u)$ replacing u in (3.14), and $\mu^*_{i,j}$ defined analogously as in (3.14), but with the role of $\mu_{V,u}(dw)$ replaced by $\mu_V(dw \times (u', u))$, cf. (1.18), (1.20), (1.45), which is also a Poisson point process on W_+ . We write $\gamma'_{i,j}$ and $\gamma^*_{i,j}$, $1 \leq i, j \leq 2$, for the intensity measures of these point processes, and note that

$$(3.19) \quad \gamma^*_{2,2}(dw) = (u - u') 1\{X_0 \in \tilde{C}_{\bar{m}_2}, H_{\tilde{C}_{\bar{m}_1}} = \infty\} P_{e_V}(dw).$$

Our aim is to bound from above $\mathbb{P}[A_{\bar{m}_2}(\mu_{2,2})] = \mathbb{P}[A_{\bar{m}_2}(\mu'_{2,2} + \mu^*_{2,2})]$ in terms of quantities involving $p_n(u') = \mathbb{P}[A_{\bar{m}_2}(\mu'_{2,2} + \mu'_{2,1} + \mu'_{1,2})]$. The rough idea is to try to dominate the influence on $\tilde{C}_{\bar{m}_2}$ of $\mu'_{2,1} + \mu'_{1,2}$ by that of $\mu^*_{2,2}$. This is a kind of “sprinkling technique” where the discrepancy between u and u' in the form of $\mu^*_{2,2}$ is used to dominate the long range interaction reflected by $\mu'_{2,1} + \mu'_{1,2}$.

With this in mind we introduce an integer $r \geq 1$, and further decompose $\mu'_{2,1}$, $\mu'_{1,2}$ and $\mu^*_{2,2}$ into:

$$(3.20) \quad \begin{aligned} \mu'_{2,1} &= \sum_{1 \leq \ell \leq r} \rho^{\ell}_{2,1} + \bar{\rho}_{2,1}, \quad \mu'_{1,2} = \sum_{1 \leq \ell \leq r} \rho^{\ell}_{1,2} + \bar{\rho}_{1,2}, \\ \mu^*_{2,2} &= \sum_{1 \leq \ell \leq r} \rho^{\ell}_{2,2} + \bar{\rho}_{2,2}, \end{aligned}$$

where denoting with $R_k, D_k, k \geq 1$, the successive returns to $\tilde{C}_{\bar{m}_2}$ and departures from $U = \{z \in \mathbb{Z}^d; d(z, \tilde{C}_{\bar{m}_2}) \leq \frac{1}{10} L_{n+1}\}$, cf. (1.4) and the notation above (1.1), we have set for $1 \leq i \neq j \leq 2, \ell \geq 1$,

$$(3.21) \quad \begin{aligned} \rho^{\ell}_{i,j} &= 1\{R_{\ell} < D_{\ell} < R_{\ell+1} = \infty\} \mu'_{i,j}, \quad \bar{\rho}_{i,j} = 1\{R_{r+1} < \infty\} \mu'_{i,j} \\ \rho^{\ell}_{2,2} &= 1\{R_{\ell} < D_{\ell} < R_{\ell+1} = \infty\} \mu^*_{2,2}, \quad \bar{\rho}_{2,2} = 1\{R_{r+1} < \infty\} \mu^*_{2,2}, \end{aligned}$$

(note that $\{R_1 < D_1 < \infty\}$ has full measure under each of $\mu'_{2,1}, \mu'_{1,2}$ and $\mu^*_{2,2}$).

We then see that with the above definitions and the independence property mentioned below (3.18),

$$(3.22) \quad \mu'_{2,2}, \rho^{\ell}_{i,j}, 1 \leq \ell \leq r, \bar{\rho}_{i,j}, 1 \leq i, j \leq 2, \text{ with } i \text{ or } j \neq 1,$$

are independent Poisson point processes on W_+ . Letting $\bar{\xi}_{2,1}$ and $\bar{\xi}_{1,2}$ stand for the respective intensity measures on W_+ of $\bar{\rho}_{2,1}$ and $\bar{\rho}_{1,2}$, we have

$$\begin{aligned}
 (3.23) \quad \bar{\xi}_{2,1}(W_+) &= u' P_{e_V} [X_0 \in \tilde{C}_{\bar{m}_2}, H_{\tilde{C}_{\bar{m}_1}} < \infty, R_{r+1} < \infty] \\
 &\stackrel{(1.6),(1.7)}{\leq} u' \text{cap}(\tilde{C}_{\bar{m}_2}) \sup_{x \in \tilde{C}_{\bar{m}_2}} P_x [R_{r+1} < \infty] \\
 &\leq u' \text{cap}(\tilde{C}_{\bar{m}_2}) \left(\sup_{x \in U^c} P_x [H_{\tilde{C}_{\bar{m}_2}} < \infty] \right)^r,
 \end{aligned}$$

where we used the strong Markov property at times D_r, D_{r-1}, \dots, D_1 , in the last step. With the right-hand inequality of (1.9) as well as [11, p. 31], we thus find that:

$$\sup_{x \in U^c} P_x [H_{\tilde{C}_{\bar{m}_2}} < \infty] \leq c L_{n+1}^{-(d-2)} \frac{L_n^d}{L_n^2} \stackrel{(3.2)}{=} c \ell_n^{-(d-2)},$$

and hence

$$(3.24) \quad \bar{\xi}_{2,1}(W_+) \leq u' \text{cap}(\tilde{C}_{\bar{m}_2}) (c \ell_n^{-(d-2)})^r \leq u' c^r L_n^{(d-2)-a(d-2)r}.$$

In the last step we used (3.2) as well as the right-hand inequality of the standard capacity estimate:

$$c L^{(d-2)} \leq \text{cap}(B(0, L)) \leq c' L^{(d-2)}, \text{ for } L \geq 1,$$

(which follows from (1.8), (1.9) with $K = B(0, L)$, letting x tend to infinity in (1.8) and using (1.9) to bound $c |x|^{-(d-2)} \text{cap}(B(0, L)) \sim P_x [H_{B(0,L)} < \infty]$, for $|x| \rightarrow \infty$). In a similar way we find that

$$\begin{aligned}
 (3.25) \quad \bar{\xi}_{1,2}(W_+) &= u' P_{e_V} [X_0 \in \tilde{C}_{\bar{m}_1}, H_{\tilde{C}_{\bar{m}_2}} < \infty, R_{r+1} < \infty] \\
 &\leq u' c^r L_n^{(d-2)-a(d-2)r}.
 \end{aligned}$$

We will now seek to show that the trace left on $\tilde{C}_{\bar{m}_2}$ by paths in the supports of $\mu'_{2,1} - \bar{\rho}_{2,1} = \sum_{1 \leq \ell \leq r} \rho_{2,1}^\ell$ and $\mu'_{1,2} - \bar{\rho}_{1,2} = \sum_{1 \leq \ell \leq r} \rho_{1,2}^\ell$ is dominated by the corresponding trace of paths in the support of $\mu^*_{2,2}$. The point processes $\bar{\rho}_{2,1}$ and $\bar{\rho}_{1,2}$ are then viewed as correction terms to be controlled with the help of (3.24), (3.25).

With this perspective we consider the space W_f of finite nearest neighbor paths on \mathbb{Z}^d , and for $\ell \geq 1$, the measurable map ϕ^ℓ from $\{D_\ell < R_{\ell+1} = \infty\} \subseteq W_+$ into the product space $W_f^{\times \ell}$ defined by:

$$(3.26) \quad \phi^\ell(w) = (w(R_k + \cdot)_{0 \leq \cdot \leq D_k - R_k})_{1 \leq k \leq \ell} \in W_f^{\times \ell} \text{ for } w \in \{D_\ell < R_{\ell+1} = \infty\}.$$

In other words $\phi^\ell(w)$ for w in the above event keeps track of the ℓ portions of the trajectory w corresponding to times going from the successive returns to $\tilde{C}_{\bar{m}_2}$ up to departure from U . We can view the various $\rho_{i,j}^\ell$, i or $j \neq 1$, with $\ell \geq 1$ fixed as point processes on $\{D_\ell < R_{\ell+1} = \infty\} (\subseteq W_+)$. We then denote with $\tilde{\rho}_{i,j}^\ell$ their respective images under ϕ^ℓ , which are Poisson point processes on $W_f^{\times \ell}$. We write

$\tilde{\xi}_{i,j}^\ell$ for their corresponding intensity measures. As a result of (3.22), we see that

$$(3.27) \quad \mu'_{2,2}, \tilde{\rho}_{i,j}^\ell, 1 \leq \ell \leq r, \bar{\rho}_{i,j}, 1 \leq i, j \leq 2, i \text{ or } j \neq 1$$

are independent Poisson point processes. We will see that when $u' < u$ are sufficiently far apart, cf. (3.34), $\tilde{\rho}_{2,2}^\ell$ has an intensity measure on $W_f^{\times \ell}$ which is bigger than the intensity measure of $\tilde{\rho}_{2,1}^\ell + \tilde{\rho}_{1,2}^\ell$, for $1 \leq \ell \leq r$. The following lemma will be helpful; we refer to (3.11), (3.12) and below (3.20) for the notation.

LEMMA 3.2. *For large L_0 , for all $n \geq 0, m \in I_{n+1}, \bar{m}_1 \in \mathcal{H}_1, \bar{m}_2 \in \mathcal{H}_2, x \in \partial U, y \in \partial_{\text{int}} \tilde{C}_{\bar{m}_2}$, we have:*

$$(3.28) \quad P_x[H\tilde{C}_{\bar{m}_1} < R_1 < \infty, X_{R_1} = y] \leq c \ell_n^{-(d-2)} P_x[H\tilde{C}_{\bar{m}_1} > R_1, X_{R_1} = y],$$

$$(3.29) \quad P_x[H\tilde{C}_{\bar{m}_1} < \infty, R_1 = \infty] \leq c \ell_n^{-(d-2)} P_x[R_1 = \infty = H\tilde{C}_{\bar{m}_1}].$$

Proof. We begin with the proof of (3.28). Recalling the notation introduced below (1.4). For $z \in \partial U, y \in \partial_{\text{int}} \tilde{C}_{\bar{m}_2}$ we have

$$(3.30) \quad \begin{aligned} &P_z[H\tilde{C}_{\bar{m}_1} < R_1 < \infty, X_{R_1} = y] \\ &\stackrel{\text{strong Markov}}{=} E_z[H\tilde{C}_{\bar{m}_1} < R_1, P_{X_{H\tilde{C}_{\bar{m}_1}}} [R_1 < \infty, X_{R_1} = y]] \\ &= E_z[H\tilde{C}_{\bar{m}_1} < R_1, E_{X_{H\tilde{C}_{\bar{m}_1}}} [H_{\partial U} < \infty, P_{X_{H_{\partial U}}} [R_1 < \infty, X_{R_1} = y]]], \end{aligned}$$

where in the last step we used for $z' \in \tilde{C}_{\bar{m}_1}$ the $P_{z'}$ -almost sure identity $R_1 = H_{\partial U} + R_1 \circ \theta_{H_{\partial U}}$, and the strong Markov property at time $H_{\partial U}$. As a result we see that:

$$(3.31) \quad \begin{aligned} &\sup_{z \in \partial U} P_z[H\tilde{C}_{\bar{m}_1} < R_1 < \infty, X_{R_1} = y] \leq \sup_{z \in \partial U} P_z[H\tilde{C}_{\bar{m}_1} < \infty] \\ &\quad \times \sup_{z \in \partial U} P_z[R_1 < \infty, X_{R_1} = y] \stackrel{(1.9)}{\leq} c \ell_n^{-(d-2)} \sup_{z \in \partial U} P_z[R_1 < \infty, X_{R_1} = y], \end{aligned}$$

with a similar bound as above (3.24) in the last step. Note that

$$P_z[R_1 < \infty, X_{R_1} = y] = P_z[H\tilde{C}_{\bar{m}_2} < \infty, X_{H\tilde{C}_{\bar{m}_2}} = y], z \in \tilde{C}_{\bar{m}_2}^c,$$

is a positive harmonic function, and using Harnack's inequality, cf. Theorem 1.7.2 of [11], together with a standard covering argument, we see that:

$$(3.32) \quad \sup_{z \in \partial U} P_z[R_1 < \infty, X_{R_1} = y] \leq c \inf_{z \in \partial U} P_z[R_1 < \infty, X_{R_1} = y].$$

Therefore coming back to (3.31) we see that

$$\begin{aligned} \sup_{z \in \partial U} P_z[H_{\tilde{C}_{\bar{m}_1}} < R_1 < \infty, X_{R_1} = y] &\leq c' \ell_n^{-(d-2)} \inf_{z \in \partial U} P_z[R_1 < \infty, X_{R_1} = y] \\ &= c' \ell_n^{-(d-2)} \inf_{z \in \partial U} \left(P_z[H_{\tilde{C}_{\bar{m}_1}} < R_1 < \infty, X_{R_1} = y] \right. \\ &\quad \left. + P_z[R_1 < \infty, X_{R_1} = y, H_{\tilde{C}_{\bar{m}_1}} > R_1] \right). \end{aligned}$$

For large L_0 , we have $c' \ell_n^{-(d-2)} \leq \frac{1}{2}$, for all $n \geq 0$, with c' as in the last line of (3.32), and thus we see that for $x \in \partial U$,

$$(3.33) \quad P_x[H_{\tilde{C}_{\bar{m}_1}} < R_1 < \infty, X_{R_1} = y] \leq 2c' \ell_n^{-(d-2)} P_x[H_{\tilde{C}_{\bar{m}_1}} > R_1, X_{R_1} = y].$$

This proves (3.28). We now turn to the proof of (3.29) which is more elementary. Indeed one has

$$\inf_{x \in \partial U} P_x[R_1 = \infty, H_{\tilde{C}_{\bar{m}_1}} = \infty] \geq c,$$

as follows from the invariance principle used to let the walk move at a distance from $V = \tilde{C}_{\bar{m}_1} \cup \tilde{C}_{\bar{m}_2}$, which is a multiple of L_{n+1} , as well as (1.9) and standard bounds on the Green function. On the other hand the left-hand side of (3.29) with a similar inequality as above (3.24) is bounded by $c \ell_n^{-(d-2)}$. Our claim follows. \square

The main control on the intensity measure $\tilde{\xi}_{1,2}^\ell + \tilde{\xi}_{2,1}^\ell$ of $\tilde{\rho}_{1,2}^\ell + \tilde{\rho}_{2,1}^\ell$ in terms of the intensity measure $\tilde{\xi}_{2,2}^\ell$ of $\tilde{\rho}_{2,2}^\ell$ is provided by the next

LEMMA 3.3. *For large L_0 , one has*

$$(3.34) \quad \tilde{\xi}_{1,2}^\ell + \tilde{\xi}_{2,1}^\ell \leq \frac{u'}{u - u'} \left[\left(1 + \frac{c_1}{\ell_n^{d-2}} \right)^{\ell+1} - 1 \right] \tilde{\xi}_{2,2}^\ell, \text{ for } \ell \geq 1.$$

Proof. The measure $\tilde{\xi}_{2,1}^\ell$ on $W_f^{\times \ell}$ is the image under ϕ^ℓ of the intensity measure $\xi_{2,1}^\ell$ on $\{D_\ell < R_{\ell+1} = \infty\} (\subseteq W_+)$ of $\rho_{2,1}^\ell$, which in view of (3.21) equals

$$(3.35) \quad \xi_{2,1}^\ell(dw) = u' P_{e_V} [dw, X_0 \in \tilde{C}_{\bar{m}_2}, H_{\tilde{C}_{\bar{m}_1}} < \infty, D_\ell < R_{\ell+1} = \infty].$$

As a result we hopefully find that with obvious notation

$$\begin{aligned} (3.36) \quad \tilde{\xi}_{2,1}^\ell(dw_1, \dots, dw_\ell) &= u' P_{e_V} [X_0 \in \tilde{C}_{\bar{m}_2}, H_{\tilde{C}_{\bar{m}_1}} < \infty, D_\ell < R_{\ell+1} = \infty, \\ &\quad (X_{R_k+})_{0 \leq \cdot \leq D_k - R_k} \in dw_k, 1 \leq k \leq \ell] \\ &= u' P_{e_V} [X_0 \in \tilde{C}_{\bar{m}_2}, \bigcup_{k=1}^\ell \{H_{\tilde{C}_{\bar{m}_1}} \circ \theta_{D_k} + D_k < R_{k+1}\}, D_\ell < R_{\ell+1} = \infty, \\ &\quad (X_{R_k+})_{0 \leq \cdot \leq D_k - R_k} \in dw_k, 1 \leq k \leq \ell] \end{aligned}$$

$$= u' \sum_{\phi \neq B \subseteq \{1, \dots, \ell\}} P_{e_V} [X_0 \in \tilde{C}_{\bar{m}_2}, H_{\tilde{C}_{\bar{m}_1}} \circ \theta_{D_k} + D_k < R_{k+1} \text{ exactly when} \\ k \in B, \text{ for } 1 \leq k \leq \ell, D_\ell < R_{\ell+1} = \infty, \\ (X_{R_k+})_{0 \leq \cdot \leq D_k - R_k} \in dw_k, 1 \leq k \leq \ell].$$

The generic term of the above sum evaluated on $(w_1, \dots, w_\ell) \in W_f^{\times \ell}$ equals:

$$(3.37) \\ u' P_{e_V} [X_0 \in \tilde{C}_{\bar{m}_2}, H_{\tilde{C}_{\bar{m}_1}} \circ \theta_{D_k} + D_k < R_{k+1}, \text{ exactly when } k \in B, 1 \leq k \leq \ell, \\ D_\ell < R_{\ell+1} = \infty, (X_{R_k+})_{0 \leq \cdot \leq D_k - R_k} = w_k(\cdot), 1 \leq k \leq \ell] \\ = u' E_{e_V} [X_0 \in \tilde{C}_{\bar{m}_2}, H_{\tilde{C}_{\bar{m}_1}} \circ \theta_{D_k} + D_k < R_{k+1}, \text{ exactly when} \\ k \in B \cap \{1, \dots, \ell - 1\} \text{ for } 1 \leq k \leq \ell - 1, D_\ell < \infty, \\ (X_{R_k+})_{0 \leq \cdot \leq D_k - R_k} = w_k(\cdot), 1 \leq k \leq \ell, \\ E_{X_{D_\ell}} [R_1 = \infty, 1\{\ell \notin B\} 1\{H_{\tilde{C}_{\bar{m}_1}} = \infty\} + 1\{\ell \in B\} 1\{H_{\tilde{C}_{\bar{m}_1}} < \infty\}]],$$

by the strong Markov property at time D_ℓ in the last step.

If we denote with w_k^s and w_k^e the starting point and the end point of w_k , for $1 \leq k \leq \ell$, the above expression vanishes unless $w_k^s \in \tilde{C}_{\bar{m}_2}$ and $w_k^e \in \partial U$ for each $k \in \{1, \dots, \ell\}$. If these conditions are fulfilled, using the strong Markov property repeatedly at times $R_\ell, D_{\ell-1}, R_{\ell-1} \dots D_1$, we see that the last line of (3.37) equals:

$$u' P_{e_V} [X_0 \in \tilde{C}_{\bar{m}_2}, (X_\cdot)_{0 \leq \cdot \leq D_1} = w_1(\cdot)] E_{w_1^e} [1\{1 \notin B\} 1\{H_{\tilde{C}_{\bar{m}_1}} > R_1\} \\ + 1\{1 \in B\} 1\{H_{\tilde{C}_{\bar{m}_1}} < R_1\}, R_1 < \infty, X_{R_1} = w_2^s] P_{w_2^e} [(X_\cdot)_{0 \leq \cdot \leq D_1} = w_2(\cdot)] \dots \\ E_{w_\ell^e} [1\{\ell \notin B\} 1\{H_{\tilde{C}_{\bar{m}_1}} = \infty\} + 1\{\ell \in B\} 1\{H_{\tilde{C}_{\bar{m}_1}} < \infty\}, R_1 = \infty].$$

We can use Lemma 3.2 for all terms in the above expression where $k \in B$, and repeatedly apply the Markov property to come back to an expression similar to (3.37). In this fashion we see that the above expression is at most:

$$(3.38) \\ (c \ell_n^{-(d-2)})^{|B|} u' P_{e_V} [X_0 \in \tilde{C}_{\bar{m}_1}, H_{\tilde{C}_{\bar{m}_1}} \circ \theta_{D_k} + D_k \geq R_{k+1}, \text{ for } 1 \leq k \leq \ell, \\ D_\ell < R_{\ell+1} = \infty, (X_{R_k+})_{0 \leq \cdot \leq D_k - R_k} = w_k(\cdot), 1 \leq k \leq \ell].$$

Summing over the various nonempty subsets B of $\{1, \dots, \ell\}$, we see by (3.36) that

$$(3.39) \\ \tilde{\xi}_{2,1}^\ell(dw_1, \dots, dw_\ell) \leq u' \sum_{\phi \neq B \subseteq \{1, \dots, \ell\}} (c \ell_n^{-(d-2)})^{|B|} P_{e_V} [X_0 \in \tilde{C}_{\bar{m}_2}, H_{\tilde{C}_{\bar{m}_1}} = \infty, \\ D_\ell < R_{\ell+1} = \infty, (X_{R_k+})_{0 \leq \cdot \leq D_k - R_k} \in dw_k, 1 \leq k \leq \ell] \\ = \frac{u'}{u - u'} [(1 + c \ell_n^{-(d-2)})^\ell - 1] \tilde{\xi}_{2,2}^\ell(dw_1, \dots, dw_\ell),$$

where we recall that $\tilde{\xi}_{2,2}^\ell$ stands for the intensity measure of $\tilde{\rho}_{2,2}^\ell$.

We can proceed in a similar fashion to bound $\tilde{\xi}_{1,2}^\ell(dw_1, \dots, dw_\ell)$. The only difference stems from the fact that under $\xi_{1,2}^\ell(dw)$ paths start in $\tilde{C}_{\bar{m}_1}$ and B , cf. last line of (3.36), can also be the empty set. In an analogous fashion to (3.38) we then obtain the following bound:

$$(3.40) \quad \begin{aligned} & u' P_{e_V} [X_0 \in \tilde{C}_{\bar{m}_1}, H_{\tilde{C}_{\bar{m}_1}} \circ \theta_{D_k} + D_k < R_{k+1}, \text{ exactly when } k \in B, \text{ for } 1 \leq k \leq \ell, \\ & \quad D_\ell < R_{\ell+1} = \infty, (X_{R_k+})_{0 \leq \cdot \leq D_k - R_k} \in w_k(\cdot), 1 \leq k \leq \ell] \\ & \leq (c \ell_n^{-(d-2)})^{|B|} u' P_\rho [X_0 \in \tilde{C}_{\bar{m}_2}, H_{\tilde{C}_{\bar{m}_1}} \circ \theta_{D_k} + D_k \geq R_{k+1}, \text{ for } 1 \leq k \leq \ell, \\ & \quad D_\ell < R_{\ell+1} = \infty, (X_{R_k+})_{0 \leq \cdot \leq D_k - R_k} = w_k(\cdot), 1 \leq k \leq \ell], \end{aligned}$$

with ρ the measure

$$(3.41) \quad \begin{aligned} \rho(y) &= \sum_{x \in \tilde{C}_{\bar{m}_1}} e_V(x) P_x [H_{\tilde{C}_{\bar{m}_2}} < \infty, X_{\tilde{C}_{\bar{m}_2}} = y], \text{ for } y \in \partial_{\text{int}} \tilde{C}_{\bar{m}_2}, \\ &= 0, \text{ otherwise.} \end{aligned}$$

We now see that for $y \in \partial_{\text{int}} \tilde{C}_{\bar{m}_2}$ with a similar calculation as in (1.36)

$$(3.42) \quad \begin{aligned} \rho(y) &\stackrel{(1.6)}{=} \sum_{x \in \tilde{C}_{\bar{m}_1}, n \geq 0} P_x [\tilde{H}_V = \infty] P_x [\tilde{H}_{\tilde{C}_{\bar{m}_2}} = n, X_n = y] \\ &\stackrel{\text{reversibility}}{=} \sum_{x \in \tilde{C}_{\bar{m}_1}, n \geq 0} P_x [\tilde{H}_V = \infty] P_y [\tilde{H}_{\tilde{C}_{\bar{m}_2}} > n, X_n = x] \\ &\stackrel{\text{Markov}}{=} \sum_{x \in \tilde{C}_{\bar{m}_1}, n \geq 0} P_y [\tilde{H}_{\tilde{C}_{\bar{m}_2}} > n, X_n = x, \tilde{H}_V \circ \theta_n = \infty] \\ &= P_y [\tilde{H}_{\tilde{C}_{\bar{m}_2}} = \infty, H_{\tilde{C}_{\bar{m}_1}} < \infty], \end{aligned}$$

summing over the time n and location x of the last visit of the path to $\tilde{C}_{\bar{m}_1}$ in the last step. Using the strong Markov property at time T_U , (recall that $\tilde{C}_{\bar{m}_1} \subseteq U^c$), we thus find

$$(3.43) \quad \begin{aligned} \rho(y) &= E_y [\tilde{H}_{\tilde{C}_{\bar{m}_2}} > T_U, P_{X_{T_U}} [H_{\tilde{C}_{\bar{m}_1}} < \infty, H_{\tilde{C}_{\bar{m}_2}} = \infty]] \\ &\stackrel{(3.29)}{\leq} c \ell_n^{-(d-2)} E_y [\tilde{H}_{\tilde{C}_{\bar{m}_2}} > T_U, P_{X_{T_U}} [H_{\tilde{C}_{\bar{m}_1}} = \infty = H_{\tilde{C}_{\bar{m}_2}}]] \\ &= c \ell_n^{-(d-2)} E_y [\tilde{H}_V > T_U, P_{X_{T_U}} [H_V = \infty]] = c \ell_n^{-(d-2)} e_V(y), \end{aligned}$$

for $y \in \partial_{\text{int}} \tilde{C}_{\bar{m}_2}$, by the strong Markov property and (1.6) in the last step. Therefore summing (3.40) over $B \subseteq \{1, \dots, \ell\}$, we find that

$$(3.44) \quad \tilde{\xi}_{1,2}^\ell(dw_1, \dots, dw_\ell) \leq \frac{u'}{u-u'} \frac{c}{\ell_n^{d-2}} \left(1 + \frac{c}{\ell_n^{d-2}}\right)^\ell \tilde{\xi}_{2,2}^\ell(dw_1, \dots, dw_\ell).$$

Summing (3.39) and (3.44) we obtain the claim (3.34). □

We now suppose L_0 large enough so that Lemma 3.2 holds, and also that

$$(3.45) \quad u = \left(1 + \frac{c_1}{\ell_n^{d-2}}\right)^{r+1} u' \left(\text{hence } \frac{u'}{u-u'} \left[\left(1 + \frac{c_1}{\ell_n^{d-2}}\right)^{r+1} - 1\right] = 1\right).$$

We will now derive the promised upper bound on $\mathbb{P}[A_{\bar{m}_2}(\mu_{2,2})]$ in terms of $p_n(u') = \mathbb{P}[A_{\bar{m}_2}^{u'}]$. Observe that the restriction of the interlacement at level u' to $\tilde{\mathcal{C}}_{\bar{m}_2}$ satisfies:

$$(3.46) \quad \begin{aligned} \mathcal{I}^{u'} \cap \tilde{\mathcal{C}}_{\bar{m}_2} &\stackrel{(1.54)}{=} \bigcup_{w \in \text{Supp}(\bar{\nu}_{\cdot, u'})} w(\mathbb{N}) \cap \tilde{\mathcal{C}}_{\bar{m}_2} = \bigcup_{w \in \text{Supp}(\mu'_{2,2} + \mu'_{2,1} + \mu'_{1,2})} w(\mathbb{N}) \cap \tilde{\mathcal{C}}_{\bar{m}_2} \\ &= \mathcal{I}' \cup \tilde{\mathcal{I}} \cup \bar{\mathcal{I}}, \end{aligned}$$

where we have set

$$(3.47) \quad \begin{aligned} \mathcal{I}' &= \bigcup_{w \in \text{Supp}(\mu'_{2,2})} w(\mathbb{N}) \cap \tilde{\mathcal{C}}_{\bar{m}_2}, \\ \tilde{\mathcal{I}} &= \bigcup_{1 \leq \ell \leq r} \bigcup_{(w_1, \dots, w_\ell) \in \text{Supp} \tilde{\rho}_{1,2}^\ell + \tilde{\rho}_{2,1}^\ell} (\text{range } w_1 \cup \dots \cup \text{range } w_\ell) \cap \tilde{\mathcal{C}}_{\bar{m}_2}, \\ \bar{\mathcal{I}} &= \bigcup_{w \in \text{Supp} \bar{\rho}_{1,2} + \bar{\rho}_{2,1}} w(\mathbb{N}) \cap \tilde{\mathcal{C}}_{\bar{m}_2}, \end{aligned}$$

and we used (3.20) together with the fact that for any $\ell \geq 1$, $w \in \text{Supp} \rho_{1,2}^\ell \cup \text{Supp} \rho_{2,1}^\ell$, with $\phi^\ell(w) = (w_1, \dots, w_\ell)$ due to (3.26):

$$w(\mathbb{N}) \cap \tilde{\mathcal{C}}_{\bar{m}_2} = (\text{range } w_1 \cup \dots \cup \text{range } w_\ell) \cap \tilde{\mathcal{C}}_{\bar{m}_2}.$$

If we now define \mathcal{I}^* by replacing $\tilde{\rho}_{1,2}^\ell + \tilde{\rho}_{2,1}^\ell$ in the second line of (3.47) by $\tilde{\rho}_{2,2}^\ell$, we see from (3.27) that

$$(3.48) \quad \text{the random sets } \mathcal{I}', \tilde{\mathcal{I}}, \bar{\mathcal{I}}, \mathcal{I}^* \text{ are independent under } \mathbb{P}.$$

We also see from (3.34), (3.27) and the choice (3.45) that for each $1 \leq \ell \leq r$, the Poisson point process $\tilde{\rho}_{1,2}^\ell + \tilde{\rho}_{2,1}^\ell$ is stochastically dominated by $\tilde{\rho}_{2,2}^\ell$ so that

$$(3.49) \quad \tilde{\mathcal{I}} \text{ is stochastically dominated by } \mathcal{I}^*.$$

With (3.48), (3.49) we thus find in view of (3.15) that

$$\begin{aligned}
 (3.50) \quad & \mathbb{P}\left[A_{\bar{m}_2}\left(\mu'_{2,2} + \sum_{1 \leq \ell \leq r} \rho_{2,2}^\ell\right)\right] \\
 &= \mathbb{P}\left[\text{there is a crossing in } \tilde{C}_{\bar{m}_2} \setminus (\mathcal{F}' \cup \mathcal{F}^*) \text{ from } C_{\bar{m}_2} \text{ to } \tilde{S}_{\bar{m}_2}\right] \\
 &\leq \mathbb{P}\left[\text{there is a crossing in } \tilde{C}_{\bar{m}_2} \setminus (\mathcal{F}' \cup \tilde{\mathcal{F}}) \text{ from } C_{\bar{m}_2} \text{ to } \tilde{S}_{\bar{m}_2}\right] \\
 &= \mathbb{P}\left[A_{\bar{m}_2}\left(\mu'_{2,2} + \sum_{1 \leq \ell \leq r} \rho_{2,1}^\ell + \rho_{1,2}^\ell\right)\right],
 \end{aligned}$$

so that

$$\begin{aligned}
 (3.51) \quad & \mathbb{P}[A_{\bar{m}_2}(\mu_{2,2})] \stackrel{(3.18)}{=} \mathbb{P}[A_{\bar{m}_2}(\mu'_{2,2} + \mu_{2,2}^*)] \stackrel{(3.20)}{\leq} \mathbb{P}\left[A_{\bar{m}_2}\left(\mu'_{2,2} + \sum_{1 \leq \ell \leq r} \rho_{2,2}^\ell\right)\right] \\
 &\stackrel{(3.50)}{\leq} \mathbb{P}\left[A_{\bar{m}_2}\left(\mu'_{2,2} + \sum_{1 \leq \ell \leq r} \rho_{2,1}^\ell + \rho_{1,2}^\ell\right)\right] \\
 &\stackrel{(3.20)}{\leq} \mathbb{P}[A_{\bar{m}_2}(\mu'_{2,2} + \mu'_{2,1} + \mu'_{1,2}), \bar{\rho}_{2,1} = 0 = \bar{\rho}_{1,2}] \\
 &\quad + \mathbb{P}[\bar{\rho}_{2,1} \text{ or } \bar{\rho}_{1,2} \neq 0] \\
 &= \mathbb{P}[A_{\bar{m}_2}(\mu_{V,u'}), \bar{\rho}_{2,1} = 0 = \bar{\rho}_{1,2}] + \mathbb{P}[\bar{\rho}_{2,1} \text{ or } \bar{\rho}_{1,2} \neq 0] \\
 &\leq p_n(u') + 1 - e^{-\bar{\xi}_{2,1}(W^+)} + 1 - e^{-\bar{\xi}_{1,2}(W^+)} \\
 &\stackrel{(3.24),(3.25)}{\leq} p_n(u') + 2u'c^r L_n^{(d-2)-a(d-2)r}.
 \end{aligned}$$

This is the promised upper bound on $\mathbb{P}[A_{\bar{m}_2}(\mu_{2,2})]$. We can now come back to (3.13), (3.17) and obtain that when L_0 is large for $n \geq 0, r \geq 1, 0 < u' < u$ satisfying (3.45) one has

$$(3.52) \quad p_{n+1}(u) \leq c_2 \ell_n^{2(d-1)} p_n(u) (p_n(u') + u' c_3^r L_n^{(d-2)(1-ar)}).$$

Given $u_0 > 0, r \geq 1$, we thus define the increasing sequence

$$(3.53) \quad u_{n+1} = \left(1 + \frac{c_1}{\ell_n^{(d-2)}}\right)^{r+1} u_n, \text{ for } n \geq 0,$$

as well as the sequence

$$(3.54) \quad a_n = c_2 \ell_n^{2(d-1)} p_n(u_n), \text{ } n \geq 0.$$

We will now prove a lemma that uses inequality (3.52) to set-up an induction scheme ensuring that a_n is at most L_n^{-1} for all $n \geq 0$. Note that (3.52) deteriorates when u' becomes large. This is compensated by picking r sufficiently big and checking (3.56) ii) at each step. In the end, to be able to initiate the induction, we will need to pick L_0 large, then $u_0 \geq c(L_0)$ to check (3.56) i) for $n = 0$, and finally

$r \geq c(L_0, u_0)$, see (3.65), thus influencing the whole sequence u_n , so as to ensure that (3.56) ii) holds for $n = 0$.

LEMMA 3.4. *If $L_0 \geq c$, then for r such that*

$$(3.55) \quad (d - 2)ar \geq 4d ,$$

and any $u_0 > 0$, when for some $n \geq 0$,

$$(3.56) \quad \text{i) } a_n \leq L_n^{-1}, \text{ and ii) } u_n \leq L_n^{(d-2)a\frac{r}{2}},$$

then (3.56) holds as well with $n + 1$ in place of n .

Proof. Since $p_n(\cdot)$ is a nonincreasing function, we see from (3.52) that for $u_0 > 0, r \geq 1$, one has for $n \geq 0$,

$$a_{n+1} \leq a_n \left(\left(\frac{\ell_{n+1}}{\ell_n} \right)^{2(d-1)} a_n + c_2 \ell_{n+1}^{2(d-1)} u_n c_3^r L_n^{(d-2)(1-ar)} \right),$$

and since one also has

$$(3.57) \quad \frac{\ell_{n+1}}{\ell_n} \stackrel{(3.2)}{=} \frac{[L_{n+1}^a]}{[L_n^a]} \leq c \left(\frac{L_{n+1}}{L_n} \right)^a \stackrel{(3.2)}{=} c \ell_n^a \leq c L_n^{a^2}, \text{ for } n \geq 0,$$

we find:

$$(3.58) \quad a_{n+1} \leq c_4 a_n (L_n^{2(d-1)a^2} a_n + L_n^{2(d-1)a(1+a)} u_n c_3^r L_n^{(d-2)(1-ar)}), \text{ for } n \geq 0.$$

We will now seek to propagate (3.56) i) from n to $n + 1$. For this purpose it suffices to show that the following two inequalities hold:

$$(3.59) \quad c_4 L_n^{2(d-1)a^2} a_n \leq \frac{1}{2} \frac{L_n}{L_{n+1}},$$

and

$$(3.60) \quad c_4 u_n c_3^r L_n^{2(d-1)a(1+a)+(d-2)(1-ar)} \leq \frac{1}{2} \frac{L_n}{L_{n+1}}.$$

To check (3.59) observe that:

$$(3.61) \quad c_4 L_n^{2(d-1)a^2} a_n \stackrel{(3.56)i)}{\leq} c_4 L_n^{2(d-1)a^2-1} \stackrel{(3.1)}{\leq} c_4 L_n^{a-1} \leq \frac{1}{200} L_n^{-a} \stackrel{(3.2)}{\leq} \frac{1}{2} \frac{L_n}{L_{n+1}},$$

with $L_0 \geq c$ and (3.1) in the next to last inequality.

We now turn to (3.60) and observe that when r satisfies (3.55) then

$$(3.62) \quad u_n c_3^r L_n^{(d-2)(1-ar)} \stackrel{(3.56)ii)}{\leq} c_3^r L_n^{(d-2)(1-a\frac{r}{2})} \stackrel{(3.55)}{\leq} \left(c_3 L_n^{-(d-2)\frac{a}{4}} \right)^r L_n^{-2} \leq L_n^{-2}, \text{ if } L_0 \geq c.$$

As a result we see that the left-hand side of (3.60) is smaller than

$$c_4 L_n^{2(d-1)a(1+a)-2} \stackrel{(3.1)}{\leq} c_4 L_n^{-1} \leq \frac{1}{200} L_n^{-a} \leq \frac{1}{2} \frac{L_n}{L_{n+1}}, \text{ if } L_0 \geq c .$$

Recalling (3.61), we see that for large L_0 we can propagate (3.56) i) from n to $n + 1$. We now turn to (3.56) ii). We have with (3.53):

(3.63)

$$\begin{aligned} u_{n+1} &= \left(1 + \frac{c_1}{\ell_n^{d-2}}\right)^{r+1} u_n \stackrel{(3.56) \text{ (ii)}}{\leq} \left(1 + \frac{c_1}{\ell_n^{d-2}}\right)^{r+1} L_n^{(d-2)a\frac{r}{2}} \\ &= L_{n+1}^{(d-2)a\frac{r}{2}} \ell_n^{-(d-2)a\frac{r}{2}} \left(1 + \frac{c_1}{\ell_n^{d-2}}\right)^{r+1} \\ &\stackrel{r \geq 1}{\leq} L_{n+1}^{(d-2)a\frac{r}{2}} \left[\left(1 + \frac{c_1}{\ell_n^{d-2}}\right)^2 \ell_n^{-(d-2)\frac{a}{2}}\right]^r \\ &\stackrel{(3.2)}{\leq} L_{n+1}^{(d-2)a\frac{r}{2}} \left[(1 + c_1)^2 (100[L_0^a])^{-(d-2)\frac{a}{2}}\right]^r \leq L_{n+1}^{(d-2)a\frac{r}{2}}, \text{ if } L_0 \geq c . \end{aligned}$$

Hence for $L_0 \geq c$, we can propagate (3.56) ii) from n to $n + 1$ as well, and this concludes the proof of Lemma 3.4. □

We now choose L_0 large so that for any $u_0 > 0$ and $r \geq 1$ satisfying (3.55), when (3.56) holds for $n = 0$, then it holds for all $n \geq 0$. If we now choose $u_0 \geq c(L_0)$, we see that for any $m \in I_0$, (recall I_0 is the set of labels at level 0),

$$\begin{aligned} (3.64) \quad a_0 &\stackrel{(3.54)}{=} c_2 \ell_0^{2(d-1)} p_0(u_0) = c_2 \ell_0^{2(d-1)} \mathbb{P}[A_m^{u_0}] \\ &\leq c_2 \ell_0^{2(d-1)} \mathbb{P}[\mathcal{V}^{u_0} \cap \tilde{S}_m \neq \emptyset] \stackrel{(1.58)}{\leq} c_2 \ell_0^{2(d-1)} |\tilde{S}_m| e^{-u_0/g(0)} \\ &\leq c L_0^{2(d-1)a+d-1} e^{-u_0/g(0)} \leq L_0^{-1} , \end{aligned}$$

using $u_0 \geq c(L_0)$ in the last step. Similarly given $L_0 \geq c$ and $u_0 \geq c(L_0)$ as above, we can pick $r \geq c(L_0, u_0)$ such that:

$$(3.65) \quad u_0 \leq L_0^{(d-2)a\frac{r}{2}} .$$

With such choices, as noted above, it follows that $a_n \leq L_n^{-1}$, for all $n \geq 0$, and this completes the proof of Proposition 3.1. □

This now brings us to the main result of this section. We recall the definition of the critical value u_* in (2.31).

THEOREM 3.5 ($d \geq 3$). *For large u the vacant set \mathcal{V}^u does not percolate, i.e.*

$$(3.66) \quad u_* < \infty ,$$

and for $u > u_*$, $\mathbb{P}[\text{Perc}(u)] = 0$.

Proof. With Corollary 2.3 we only need to prove (3.66). We choose L_0, u_0, r as in Proposition 3.1, so that with u_n as in (3.9) we find:

$$(3.67) \quad c_2 \ell_n^{2(d-1)} \mathbb{P}[A_m^{u_n}] \leq L_n^{-1}, \text{ for any } n \geq 0 \text{ and } m \in I_n.$$

With (3.2) we know that $L_n \geq L_0^{(1+a)^n}$, and hence $\sum_n \ell_n^{-(d-2)} < \infty$, and we thus see that

$$(3.68) \quad u_\infty = u_0 \prod_{n \geq 0} \left(1 + \frac{c_1}{\ell_n^{d-2}}\right)^{r+1} = u_0 \left(\prod_{n \geq 0} \left(1 + \frac{c_1}{\ell_n^{d-2}}\right)\right)^{r+1} < \infty.$$

Consequently for any $n \geq 0$, and $m \in I_n$ such that $0 \in C_m$, we find as a consequence of (2.2) and (3.7) that

$$(3.69) \quad \eta(u_\infty) \leq \mathbb{P}[A_m^{u_\infty}] \leq c L_n^{-1}.$$

Letting n tend to infinity we see that $\eta(u_\infty) = 0$, and (3.66) follows. □

Remark 3.6. Once we know that \mathcal{V}^u does not percolate for large u , it is natural to wonder how large the vacant cluster at the origin can be. An exponential tail bound on the number of sites of the vacant cluster at the origin of subcritical Bernoulli percolation is known to hold; cf. [8, pp. 132 and 350]. Such an estimate cannot be true in the case of \mathcal{V}^u due to (1.65). The exact nature of the tail of this random variable is an interesting problem. □

4. Percolation for small u

The main objective of this section is to show that when $d \geq 7$, the vacant set \mathcal{V}^u percolates for small $u > 0$, or equivalently that $u_* > 0$; see Theorem 4.3. In spite of the fact that \mathcal{V}^u tends to contain bigger boxes than what is the case for Bernoulli percolation, cf. (1.65), it does not stochastically dominate Bernoulli percolation in the highly percolative regime as noted in Remark 1.6 (1). This fact precludes a strategy based on a direct comparison argument. We develop here a similar but simpler renormalization procedure as in the previous section. It yields a sharper result than the strategy based on the combination of the exponential bound (2.37) and a Peierls-type argument, as outlined in Remark 3.5 (3). Such a proof only works for $d \geq 18$.

We begin with some notation. We recall the definitions of $a > 0$ and $L_n, n \geq 0$, in (3.1), (3.2). Throughout we identify \mathbb{Z}^2 with the subset of points $z = (z_1, \dots, z_d)$ in \mathbb{Z}^d , such that $z_3 = z_4 = \dots = z_d = 0$. For $n \geq 0$, we define the set J_n of labels of level n just as in (3.4), but with d replaced by 2. For $n \geq 0$ and $m \in J_n$ we attach the boxes in $\mathbb{Z}^2, D_m \subseteq \tilde{D}_m$, with a similar definition as in (3.5), but with d replaced by 2, and I_n by J_n . We write \tilde{V}_m for the relative interior boundary in \mathbb{Z}^2 of \tilde{D}_m , i.e. the set of points of \tilde{D}_m neighboring $\mathbb{Z}^2 \setminus \tilde{D}_m$. In this section, parallel

to (3.7), a crucial role is played by the “occupied crossing” events. Namely for $u \geq 0, n \geq 0, m \in J_n$, we define:

$$(4.1) \quad B_m^u = \{\omega \in \Omega; \text{there is a } * \text{-nearest neighbor path in } \mathcal{J}^u(\omega) \cap \tilde{D}_m \text{ from } D_m \text{ to } \tilde{V}_m\}.$$

As a consequence of translation invariance, cf. Theorem 2.1 or (1.48),

$$(4.2) \quad q_n(u) = \mathbb{P}[B_m^u], \quad u \geq 0, n \geq 0, \text{ with } m \in J_n,$$

is well defined. It is also a nondecreasing function of u . Our main task consists in showing that when u is chosen small $q_n(u)$ tends sufficiently rapidly to 0. Our key control stems from

PROPOSITION 4.1 ($d \geq 7$). *There exists a positive constant c_5 , cf. (4.13), such that for $L_0 \geq c$, and $u \leq c(L_0)$ one has*

$$(4.3) \quad c_5 \ell_n^2 q_n(u) \leq L_n^{-\frac{1}{2}}, \text{ for all } n \geq 0.$$

Proof. In analogy with (3.11), (3.12), we define for $n \geq 0$ and $m \in J_{n+1}$ the collection of labels of boxes at level n “at the boundary of D_m ”:

$$(4.4) \quad \mathcal{K}_1 = \{\bar{m} \in J_n; D_{\bar{m}} \subseteq D_m \text{ and some point of } D_{\bar{m}} \text{ neighbors } \mathbb{Z}^2 \setminus D_m\}.$$

We also consider the collection of labels of boxes at level n containing some point at $|\cdot|_\infty$ -distance $\frac{L_{n+1}}{2}$ from D_m :

$$(4.5) \quad \mathcal{K}_2 = \left\{ \bar{m} \in J_n; D_{\bar{m}} \cap \left\{ z \in \mathbb{Z}^2; d(z, D_m) = \frac{L_{n+1}}{2} \right\} \neq \emptyset \right\}.$$

The argument leading to (3.13) applies here as well and shows that

$$(4.6) \quad q_{n+1}(u) \leq c \ell_n^2 \sup_{\bar{m}_1 \in \mathcal{K}_1, \bar{m}_2 \in \mathcal{K}_2} \mathbb{P}[B_{\bar{m}_1}^u \cap B_{\bar{m}_2}^u], \text{ for } u \geq 0.$$

From now on we assume that

$$(4.7) \quad u \leq 1.$$

For $\bar{m}_1 \in \mathcal{K}_1, \bar{m}_2 \in \mathcal{K}_2$, we define $V = \tilde{D}_{\bar{m}_1} \cup \tilde{D}_{\bar{m}_2}$ and write

$$(4.8) \quad \mu_{V,u} = \delta_{1,1} + \delta_{1,2} + \delta_{2,1} + \delta_{2,2},$$

with a similar definition as in (3.14) or (2.8), so that the $\delta_{i,j}(dw), 1 \leq i, j \leq 2$, are independent Poisson point processes on W_+ with respective intensity measures $\zeta_{i,j}, 1 \leq i, j \leq 2$, given by analogous formulas as in (2.10), except that K and $K+x$ are now respectively replaced by $\tilde{D}_{\bar{m}_1}$ and $\tilde{D}_{\bar{m}_2}$. With notation similar to (3.15),

when $\Lambda(dw)$ is a random point process on W_+ , we write:

$$(4.9) \quad B_{\bar{m}}(\Lambda) = \left\{ \omega \in \Omega; D_{\bar{m}} \text{ and } \tilde{V}_{\bar{m}} \text{ are connected by a } * \text{-nearest-neighbor path} \right. \\ \left. \text{in } \tilde{D}_{\bar{m}} \cap \left(\bigcup_{w \in \text{Supp}(\Lambda(\omega))} w(\mathbb{N}) \right) \right\}, \bar{m} \in J_n.$$

For instance we now see with (1.54) that

$$(4.10) \quad B_{\bar{m}}^u = B_{\bar{m}}(\mu_{K,u}) \text{ for any } K \supseteq \tilde{D}_{\bar{m}}, \bar{m} \in J_n.$$

Specializing to $K = V$, and noting that

$$\tilde{D}_{\bar{m}_i} \cap \left(\bigcup_{w \in \text{Supp}(\delta_{j,j})} w(\mathbb{N}) \right) = \emptyset, \text{ when } \{i, j\} = \{1, 2\},$$

we obtain the identities:

$$(4.11) \quad B_{\bar{m}_1}^u = B_{\bar{m}_1}(\mu_{V,u}) = B_{\bar{m}_1}(\delta_{1,1} + \delta_{1,2} + \delta_{2,1}), \\ B_{\bar{m}_2}^u = B_{\bar{m}_2}(\mu_{V,u}) = B_{\bar{m}_2}(\delta_{2,2} + \delta_{2,1} + \delta_{1,2}).$$

Due to the independence of the $\delta_{i,j}$, $1 \leq i, j \leq 2$, it follows that for $\bar{m}_i \in \mathcal{H}_i$, $i = 1, 2$, we have:

$$(4.12) \quad \mathbb{P}[B_{\bar{m}_1}^u \cap B_{\bar{m}_2}^u] = \mathbb{P}[B_{\bar{m}_1}(\delta_{1,1} + \delta_{1,2} + \delta_{2,1}) \cap B_{\bar{m}_2}(\delta_{2,2} + \delta_{2,1} + \delta_{1,2})] \\ \leq \mathbb{P}[B_{\bar{m}_1}(\delta_{1,1}) \cap B_{\bar{m}_2}(\delta_{2,2}), \delta_{1,2} = \delta_{2,1} = 0] + \mathbb{P}[\delta_{1,2} \text{ or } \delta_{2,1} \neq 0] \\ \leq \mathbb{P}[B_{\bar{m}_1}(\delta_{1,1})] \mathbb{P}[B_{\bar{m}_2}(\delta_{2,2})] + \mathbb{P}[\delta_{1,2} \neq 0] + \mathbb{P}[\delta_{2,1} \neq 0] \\ (4.2), (4.11) \quad \leq q_n(u)^2 + 1 - e^{-\xi_{1,2}(W_+)} + 1 - e^{-\xi_{2,1}(W_+)} \\ \leq q_n(u)^2 + u(P_{e_V}[X_0 \in \tilde{D}_{\bar{m}_1}, H_{\tilde{D}_{\bar{m}_2}} < \infty] + P_{e_V}[X_0 \in \tilde{D}_{\bar{m}_2}, H_{\tilde{D}_{\bar{m}_1}} < \infty]) \\ (4.7) \quad \leq q_n(u)^2 + c L_n^2 \frac{L_n^2}{L_{n+1}^{d-2}},$$

where in the last step we used (1.6), (1.8) together with standard bounds on the Green function, cf. [11, p. 31]. With (4.6), (4.12), we thus see that

$$(4.13) \quad q_{n+1}(u) \leq c_5 \ell_n^2 (q_n^2(u) + L_n^4 L_{n+1}^{-(d-2)}).$$

We thus define

$$(4.14) \quad b_n = c_5 \ell_n^2 q_n(u), \text{ for } n \geq 0,$$

and see that

$$b_{n+1} \leq \left(\frac{\ell_{n+1}}{\ell_n} \right)^2 b_n^2 + c(\ell_{n+1} \ell_n)^2 L_n^4 L_{n+1}^{-(d-2)}, \text{ for } n \geq 0.$$

With (3.2) we know that $(\ell_n \ell_{n+1})^2 \leq c L_n^{2a} L_{n+1}^{2a} \leq c L_n^{4a+2a^2}$, and with (3.57) we know that $\frac{\ell_{n+1}}{\ell_n} \leq c L_n^{a^2}$. As a result we obtain:

$$(4.15) \quad b_{n+1} \leq c(L_n^{2a^2} b_n^2 + L_n^{-(d-2)(1+a)+4+4a+2a^2}) \stackrel{d \geq 7}{\leq} c_6 L_n^{2a^2} (b_n^2 + L_n^{-1}).$$

We will now use the following induction lemma.

LEMMA 4.2 ($d \geq 7$). *If $L_0 \geq c$, then for any $u \leq 1$, when for some $n \geq 0$,*

$$(4.16) \quad b_n \leq L_n^{-\frac{1}{2}},$$

then (4.16) holds as well with $n + 1$ in place of n .

Proof. With (4.15) we see that

$$(4.17) \quad b_{n+1} \leq 2c_6 L_n^{2a^2-1} \stackrel{(3.2)}{\leq} c L_{n+1}^{-\frac{1}{2}} L_n^{\frac{1}{2}(1+a)+2a^2-1} \stackrel{(3.1)}{\leq} c L_{n+1}^{-\frac{1}{2}} L_0^{-\frac{1}{4}} \leq L_{n+1}^{-\frac{1}{2}},$$

when $L_0 \geq c$. This proves our claim. □

We now choose $L_0 \geq c$, so that for any $u \leq 1$, when $b_0 \leq L_0^{-\frac{1}{2}}$ holds then $b_n \leq L_n^{-\frac{1}{2}}$ for all $n \geq 0$. Further picking $u \leq c(L_0)(\leq 1)$, we see that for $m \in J_0$,

$$(4.18) \quad b_0 \stackrel{(4.14)}{=} c_5 \ell_0^2 q_0(u) \leq c_5 \ell_0^2 \mathbb{P}[\mathcal{F}^u \cap \tilde{D}_m \neq \phi] \leq c L_0^{2(1+a)} \mathbb{P}[0 \in \mathcal{F}^u] \\ \stackrel{(1.58)}{=} c L_0^{2(1+a)} (1 - e^{-u/g(0)}) \leq L_0^{-\frac{1}{2}}.$$

With this choice of L_0 and u we thus find that $b_n \leq L_n^{-\frac{1}{2}}$ for all $n \geq 0$, and this concludes the proof of Proposition 4.1. □

THEOREM 4.3. ($d \geq 7$) *For small $u > 0$ the vacant set \mathcal{V}^u does percolate, i.e.*

$$(4.19) \quad u_* > 0,$$

and for $u < u_$, $\mathbb{P}[\text{Perc}(u)] = 1$.*

Proof. We only need to prove (4.19) thanks to Corollary 2.3. We choose L_0 large and $u \leq c(L_0)$ so that (4.3) holds for all $n \geq 0$. Then for $n_0 \geq 0$ and $M = L_{n_0} - 1$, we can write:

(4.20)

$$\begin{aligned}
 1 - \eta(u) &\leq \mathbb{P}[0 \text{ does not belong to an infinite connected component of } \mathcal{V}^u \cap \mathbb{Z}^2] \\
 &\leq \mathbb{P}[\mathcal{F}^u \cap B(0, M) \cap \mathbb{Z}^2 \neq \emptyset] + \mathbb{P}[\mathcal{F}^u \cap (\mathbb{Z}^2 \setminus B(0, M)) \text{ contains a} \\
 &\quad \text{*-nearest neighbor circuit surrounding } 0] \stackrel{(1.58)}{\leq} c M^2 (1 - e^{-u/g(0)}) \\
 &\quad + \sum_{n \geq n_0} \mathbb{P}[\mathcal{F}^u \cap (\mathbb{Z}^2 \setminus B(0, M)) \text{ contains a *-nearest neighbor circuit} \\
 &\quad \text{containing } 0 \text{ and passing through a point in } [L_n, L_{n+1} - 1] e_1] \\
 &\leq c L_{n_0}^2 u + \sum_{n \geq n_0} \sum_m \mathbb{P}[B_m^u],
 \end{aligned}$$

where m runs over the collection of labels at level n of boxes D_m intersecting the segment $[L_n, L_{n+1} - 1] e_1$, $((e_1, \dots, e_d)$ stands for the canonical basis of \mathbb{R}^d , and recall we identified \mathbb{Z}^2 with $\mathbb{Z}e_1 + \mathbb{Z}e_2$). With (3.2) this collection has cardinality at most $\ell_n \leq c L_n^a$, and we thus find that

$$(4.21) \quad 1 - \eta(u) \leq c L_{n_0}^2 u + \sum_{n \geq n_0} c L_n^a L_n^{-\frac{1}{2}} \stackrel{(3.1)}{\leq} c(L_{n_0}^2 u + \sum_{n \geq n_0} L_n^{-\frac{1}{4}}).$$

Choosing n_0 large and then $u \leq c(L_0, n_0)$, we find that $1 - \eta(u) < 1$, and this proves (4.19). □

Remark 4.4. 1) Combining Theorems 3.5 and 4.3 we find that when $d \geq 7$,

$$(4.22) \quad 0 < u_* < \infty,$$

i.e. u_* is a nondegenerate critical value. This has been extended to all $d \geq 3$ in [15].

2) As a matter of fact the above proof combined with an ergodicity argument, cf. (2.6), shows that when $d \geq 7$, for small $u > 0$, \mathcal{V}^u percolates in $\mathbb{Z}^2 \subset \mathbb{Z}^d$. This feature remains true for all $d \geq 3$, see Theorem 3.4 of [15].

3) Some other very natural questions remain open. When $u > u_*$, how large is the vacant cluster at the origin? When $u < u_*$, how large can a vacant cluster at the origin be, if it does not meet the infinite cluster, (which is known to be unique thanks to [22])? Is there percolation at criticality (i.e. is $\eta(u_*) > 0$)? What is the asymptotic behavior of u_* for large d ? These are just a few examples of many unresolved issues concerning percolative properties of the vacant set left by the random interlacements model described in this work. □

Acknowledgements. We wish to thank Yves Le Jan for useful references.

References

[1] D. ALDOUS and J. FILL, *Reversible Markov Chains and Random Walks on Graphs*, Book in preparation. arXiv <http://www.stat.berkeley.edu/~aldous/RWG/book.html>

- [2] I. BENJAMINI and A.-S. SZNITMAN, Giant component and vacant set for random walk on a discrete torus, *J. Eur. Math. Soc. (JEMS)* **10** (2008), 133–172. MR 2008i:60076 Zbl 1141.60057
- [3] E. BOLTHAUSEN and J.-D. DEUSCHEL, Critical large deviations for Gaussian fields in the phase transition regime. I, *Ann. Probab.* **21** (1993), 1876–1920. MR 95c:60023 Zbl 0801.60018
- [4] M. J. A. M. BRUMMELHUIS and H. J. HILHORST, Covering of a finite lattice by a random walk, *Phys. A* **176** (1991), 387–408. MR 92m:82058
- [5] R. M. BURTON and M. KEANE, Density and uniqueness in percolation, *Comm. Math. Phys.* **121** (1989), 501–505. MR 90g:60090 Zbl 0662.60113
- [6] A. DEMBO and A.-S. SZNITMAN, On the disconnection of a discrete cylinder by a random walk, *Probab. Theory Related Fields* **136** (2006), 321–340. MR 2007i:60053 Zbl 1105.60029
- [7] ———, A lower bound on the disconnection time of a discrete cylinder, in *In and Out of Equilibrium. 2*, *Progr. Probab.* **60**, Birkhäuser, Basel, 2008, pp. 211–227. MR 2477383 Zbl 05363764
- [8] G. GRIMMETT, *Percolation*, second ed., *Grundle Math. Wissen.* **321**, Springer-Verlag, New York, 1999. MR 2001a:60114 Zbl 0926.60004
- [9] O. HÄGGSTRÖM and J. JONASSON, Uniqueness and non-uniqueness in percolation theory, *Probab. Surv.* **3** (2006), 289–344. MR 2007m:60297
- [10] G. A. HUNT, Markoff chains and Martin boundaries, *Illinois J. Math.* **4** (1960), 313–340. MR 23 #A691 Zbl 0094.32103
- [11] G. F. LAWLER, *Intersections of Random Walks, Probability and its Applications*, Birkhäuser, Boston, MA, 1991. MR 92f:60122 Zbl 0925.60078
- [12] E. W. MONTROLL, Random walks in multidimensional spaces, especially on periodic lattices, *J. Soc. Indust. Appl. Math.* **4** (1956), 241–260. MR 19,470d Zbl 0080.35101
- [13] C. M. NEWMAN and L. S. SCHULMAN, Infinite clusters in percolation models, *J. Statist. Phys.* **26** (1981), 613–628. MR 83e:82038 Zbl 0509.60095
- [14] S. I. RESNICK, *Extreme Values, Regular Variation, and Point Processes, Applied Probability. A Series of the Applied Probability Trust* **4**, Springer-Verlag, New York, 1987. MR 89b:60241 Zbl 0633.60001
- [15] V. SIDORAVICIUS and A.-S. SZNITMAN, Percolation for the vacant set of random interacements, *Comm. Pure Appl. Math.* **62** (2009), 831–858. MR 2512613 Zbl 1168.60036
- [16] M. L. SILVERSTEIN, *Symmetric Markov Processes, Lecture Notes in Math.* **426**, Springer-Verlag, New York, 1974. MR 52 #6891 Zbl 0296.60038
- [17] F. SPITZER, *Principles of Random Walks*, second ed., *Graduate Texts Math.* **34**, Springer-Verlag, New York, 1976. MR 52 #9383
- [18] ———, *Differential Geometry*, vol. 2, Publish or Perish, Berkeley, CA, 1979.
- [19] A.-S. SZNITMAN, How universal are asymptotics of disconnection times in discrete cylinders?, *Ann. Probab.* **36** (2008), 1–53. MR 2008m:60143 Zbl 1134.60061
- [20] ———, Random walks on discrete cylinders and random interacements, *Probab. Theory Related Fields* **145** (2009), 143–174. MR 2520124 Zbl 05601403
- [21] ———, Upper bound on the disconnection time of discrete cylinders and random interacements, *Ann. Probab.* **37** (2009), 1715–1746.
- [22] A. TEIXEIRA, On the uniqueness of the infinite cluster of the vacant set of random interacements, *Ann. Appl. Probab.* **19** (2009), 454–466. MR 2498684 Zbl 1158.60046
- [23] A. TELCS, *The Art of Random Walks, Lecture Notes in Math.* **1885**, Springer-Verlag, New York, 2006. MR 2007d:60001 Zbl 1104.60003

- [24] M. WEIL, Quasi-processus, in *Séminaire de Probabilités, IV (Univ. Strasbourg, 1968/69), Lecture Notes in Math.* **124**, Springer-Verlag, New York, 1970, pp. 216–239. MR 42 #1212 Zbl 0211.21203
- [25] D. WINDISCH, Random walk on a discrete torus and random interlacements, *Electron. Commun. Probab.* **13** (2008), 140–150. MR 2008k:60249 Zbl pre05636396

(Received April 30, 2007)

(Revised June 8, 2008)

E-mail address: alain-sol.sznitman@math.ethz.ch

ETH, DEPARTEMENT MATHEMATIK, HG G 36.2, RÄMISTRASSE 101, 8092 ZÜRICH,
SWITZERLAND

<http://www.math.ethz.ch/u/sznitman>

