

ANNALS OF MATHEMATICS

Teichmüller curves, triangle groups, and Lyapunov exponents

By IRENE I. BOUW and MARTIN MÖLLER



SECOND SERIES, VOL. 172, NO. 1

July, 2010

ANMAAH

Teichmüller curves, triangle groups, and Lyapunov exponents

By IRENE I. BOUW and MARTIN MÖLLER

Abstract

We construct a Teichmüller curve uniformized by a Fuchsian triangle group commensurable to $\Delta(m, n, \infty)$ for every $m, n \leq \infty$. In most cases, for example when $m \neq n$ and m or n is odd, the uniformizing group is equal to the triangle group $\Delta(m, n, \infty)$. Our construction includes the Teichmüller curves constructed by Veech and Ward as special cases. The construction essentially relies on properties of hypergeometric differential operators. For small m , we find billiard tables that generate these Teichmüller curves. We interpret some of the so-called Lyapunov exponents of the Kontsevich-Zorich cocycle as normalized degrees of a natural line bundle on a Teichmüller curve. We determine the Lyapunov exponents for the Teichmüller curves we construct.

1. Introduction

Let C be a smooth curve defined over \mathbb{C} . The curve C is a *Teichmüller curve* if there exists a generically injective, holomorphic map from C to the moduli space M_g of curves of genus g which is geodesic for the Teichmüller metric. Consider a pair (X, ω_X) , where X is a Riemann surface of genus g and ω_X is a holomorphic 1-form on X . If the projective affine group Γ of (X, ω_X) is a lattice in $\mathrm{PSL}_2(\mathbb{R})$, then $C := \mathbb{H}/\Gamma$ is a Teichmüller curve. Such a pair (X, ω_X) is called a *Veech surface*. Moreover, the curve X is a fiber of the family of curves \mathcal{X} corresponding to the map $C \rightarrow M_g$. We refer to Section 2 for precise definitions and more details.

Teichmüller curves arise naturally in the study of dynamics of billiard paths on a polygon in \mathbb{R}^2 . Veech ([35]) constructed a first class of Teichmüller curves $C = C_n$ starting from a triangle. The corresponding projective affine group is commensurable to the triangle group $\Delta(2, n, \infty)$. Ward ([38]) also found triangles which generate Teichmüller curves, with projective affine group $\Delta(3, n, \infty)$. Several authors tried to find other triangles which generate Teichmüller curves, but only sporadic examples were found. Many types of triangles were disproven to yield Veech surfaces ([37], [16], [32]).

In this paper we show that for all m, n there is a Teichmüller curve $C(m, n, \infty)$ whose projective affine group is commensurable to $\Delta(m, n, \infty)$. It is in fact equal to $\Delta(m, n, \infty)$ in most cases, for example if $m \neq n$ and one of m and n odd. (Since Teichmüller curves are never complete ([35]), triangle groups $\Delta(m, n, k)$ with $k \neq \infty$ do not occur.) We use a different construction from previous authors; we construct the family \mathcal{X} of curves defined by C rather than the individual Veech surface (which is a fiber of \mathcal{X}). However, starting from our description, we compute an algebraic equation for the corresponding Veech surface. The family \mathcal{X} is given as the quotient of an abelian cover $\mathcal{Y} \rightarrow \mathbb{P}^1$ by a finite group.

Under the simplifying assumption that $m < n < \infty$ and n is odd, we relate the Veech surface corresponding to the Teichmüller curve $C(m, n, \infty)$ to a rational polygon. This polygon has $(m + 3)/2$ edges if m is odd and $(m + 4)/2$ edges if m is even. This polygon does not have self-crossings if and only if $m \leq 5$. Therefore, for $m \leq 5$ we obtain the Veech surface by unfolding a polygon.

From our construction we obtain new information even for the Teichmüller curves found by Veech and Ward. Namely, we determine the complete decomposition of the relative de Rham cohomology $R^1 f_* \mathbb{C}_{\mathcal{X}}$ and the Lyapunov exponents (see below).

There exist Teichmüller curves whose projective affine group is not a triangle group. McMullen ([24]) constructed a series of such examples in genus $g = 2$. It would be interesting to try to extend our method to Fuchsian groups other than triangle groups. This would probably be much more involved due to the appearance of so-called *accessory parameters*.

We now give a more detailed description of our results. Suppose that $m \geq 4$ and $m < n \leq \infty$ or that $m \geq 2$ and $3 \leq n < \infty$. We consider a family of N -cyclic covers

$$\mathcal{Y}_t : \quad y^N = x^{a_1}(x-1)^{a_2}(x-t)^{a_3}$$

of the projective line branched at four points. Note that \mathcal{Y} defines a family over $C = \mathbb{P}_t^1 - \{0, 1, \infty\}$. It is easy to compute the differential equation corresponding to the eigenspaces $\mathbb{L}(i)$ of the action of \mathbb{Z}/N on the relative de Rham cohomology of \mathcal{Y} (see §4). These eigenspaces are local systems of rank 2, and the corresponding differential equation is hypergeometric. Cohen and Wolfart ([6]) showed that we may choose N and a_i in terms of n and m such that the projective monodromy group of at least one of the eigenspaces $\mathbb{L}(i)$ is the triangle group $\Delta(m, n, \infty)$.

First consider the case that m and n are finite and relatively prime. Here we show that the particular choice of N and the a_i implies that, after replacing C by a finite unramified cover, the automorphism group of \mathcal{Y} contains a subgroup isomorphic to $\mathbb{Z}/N \rtimes H$, where $H \simeq \mathbb{Z}/2 \times \mathbb{Z}/2$. If n is infinite the group H has order 2. This case corresponds to half of Veech's series of Teichmüller curves (see §5). If m and n are not relatively prime we replace \mathcal{Y} by a suitable G_0 -Galois cover of the projective line, where G_0 is some subgroup of $\mathbb{Z}/N \times \mathbb{Z}/N$. The description of \mathcal{Y} in this case is just as explicit (§6).

THEOREMS 5.2 AND 6.2. *The quotient family $\mathcal{X} := \mathcal{Y}/H$ is the pullback to C of the universal family over the moduli space of curves. The curve C is an unramified cover of a Teichmüller curve.*

The proof of this result relies on a Hodge-theoretical characterization of Teichmüller curves ([29]). Another key ingredient of the proof is the characterization of the vanishing of the Kodaira-Spencer map in terms of invariants of the hypergeometric differential equation corresponding to $\mathbb{L}(i_0)$ (Proposition 3.2). Here i_0 is chosen such that the projective monodromy group of $\mathbb{L}(i_0)$ is the triangle group $\Delta(m, n, \infty)$. The statement on the Kodaira-Spencer map translates to the following geometric property of \mathcal{X} . A fiber \mathcal{X}_c of \mathcal{X} is singular if and only if the monodromy around c of the local system induced by $\mathbb{L}(i_0)$ is infinite (Proposition 4.2). This is one of the central observations of the paper. This is already apparent in our treatment of the relatively straightforward case of Veech's families of Teichmüller curves in Section 5.

THEOREM 6.12. *Suppose $m \neq n$. Then the projective affine group of \mathcal{X}*

- (a) *is the (m, n, ∞) -triangle group if at least one of m and n is odd,*
- (b) *contains a subgroup of index 2 in the (m, n, ∞) -triangle group otherwise.*

If $m = n$ is odd, then the projective affine group of the translation surface (X, ω_X) either equals the (m, m, ∞) -triangle group or contains it as subgroup of index 2.

We determine the projective affine group of our Teichmüller curves directly from the construction of the family \mathcal{X} and do not need to consider the corresponding Veech surfaces, as done by Veech and Ward. For example, we determine the number of zeros of the generating differential of a Veech surface corresponding to $C(m, n, \infty)$ in terms of n and m by algebraic methods (Theorem 6.13).

In Section 8 we change perspective, and discuss the question of realizing our Teichmüller curves via unfolding of rational polygons (or: billiard tables). This section may be read independently of the rest of the paper. For $m \leq 5$ we construct a billiard table $T(m, n, \infty)$ and show that it defines a Teichmüller curve, via unfolding. For $m = 2, 3$ this gives the triangles considered by Veech ([35]) and Ward ([38]). For $m = 4, 5$ we find new billiard tables which are rational 4-gons. We interpret the Veech surfaces corresponding to these billiard tables as fiber of the family $\mathcal{X} \rightarrow C$ of curves. A key ingredient here is a theorem of Ward ([38, Th. C']) which relates a cyclic cover of the projective line to a polygon, via the Schwarz-Christoffel map. We then use that certain fibers of \mathcal{X} are a cyclic cover of the projective line (Theorem 6.14).

For $m \geq 6$ the same procedure still produces rational polygons $T(m, n, \infty)$, but they have self-crossings and therefore do not define billiard tables. In principle, one could still describe the translation surface corresponding to $T(m, n, \infty)$, but these would be hard to visualize.

Our last main result concerns Lyapunov exponents. Let V be a flat normed vector bundle on a manifold with flow. The *Lyapunov exponents* measure the rate of growth of the length of vectors in V under parallel transport along the flow. We refer to Section 9 for precise definitions and a motivation of the concept. We express the Lyapunov exponents for an arbitrary Teichmüller curves in terms of the degree of certain local systems.

Let $f : \mathcal{X} \rightarrow C$ be the universal family over an unramified cover of an arbitrary Teichmüller curve. The relative de Rham cohomology $R^1 f_* \mathbb{C}_{\mathcal{X}}$ has r local subsystems $\mathbb{L}(i)$ of rank 2. The associated vector bundles carry a Hodge filtration (see Theorem 2.1). The $(1, 0)$ -parts of the Hodge filtration are line bundles $\mathcal{L}(i)$ and the ratios

$$\lambda_i := 2 \deg(\mathcal{L}(i)) / (2g(C) - 2 + s) \quad \text{and} \quad s = \text{card}(\bar{C} \setminus C)$$

are unchanged if we pass to an unramified cover of C .

THEOREM 9.2: *The ratios λ_i are r of g nonnegative Lyapunov exponents of the Kontsevich-Zorich cocycle over the Teichmüller geodesic flow on the canonical lift of a Teichmüller curve to the 1-form bundle over the moduli space.*

A sketch of the relation between the degree of $f_* \omega_{\mathcal{X}/C}$ and the sum of all Lyapunov exponents has already appeared in [18].

Now suppose that C is an unramified cover of $C(m, n, \infty)$ (Theorems 5.2 and 6.2), and let $f : \mathcal{X} \rightarrow C$ be the corresponding family of curves. In Corollaries 5.3 (Veech's series), 5.6 and 6.9 we give an explicit expression for all Lyapunov exponents of C . For Veech's series of Teichmüller curves and for a series of square-tiled coverings the Lyapunov exponents were calculated independently by Kontsevich and Zorich (unpublished). In these cases they form an arithmetic progression. Example 6.10 shows that this does not hold in general.

It is well-known that the largest Lyapunov exponent $\lambda_1 = 1$ occurs with multiplicity one. We interpret $1 - \lambda_i$ as the number of zeros of the Kodaira-Spencer map of $\mathbb{L}(i)$, counted with multiplicity (see §2), up to a factor. For the Teichmüller curves constructed in Theorems 5.2 and 6.2, we determine the position of the zeros of the Kodaira-Spencer map. These zeros are related to elliptic fixed points of the projective affine group Γ (Propositions 3.2 and 4.4). For an arbitrary Teichmüller curve it is an interesting question to determine the position of the zeros of the Kodaira-Spencer map. Precise information on the zeros of the Kodaira-Spencer map might shed new light on the defects $1 - \lambda_i$ of the Lyapunov exponents.

The starting point of this paper was a discussion with Pascal Hubert and Anton Zorich on Lyapunov exponents. The second named author thanks them heartily. Both authors acknowledge support from the DFG-Schwerpunkt 'Komplexe Mannigfaltigkeiten'. We thank Frits Beukers for suggesting the proof of Proposition 8.4, Silke Notheis for cartographic support, and Pat Hooper for pointing out a mistake in an earlier version and for sending us the manuscript [13].

2. Teichmüller curves

A *Teichmüller curve* is a generically injective, holomorphic map $C \rightarrow M_g$ from a smooth algebraic curve C to the moduli space of curves of genus g which is geodesic for the Teichmüller metric. A Teichmüller curve arises as quotient $C = \mathbb{H}/\Gamma$, where $\mathbb{H} \rightarrow T_g$ is a complex Teichmüller geodesic in Teichmüller space T_g . Here Γ is the subgroup in the Teichmüller modular group fixing \mathbb{H} as a subset of T_g (setwise, not pointwise) and C is the normalization of the image $\mathbb{H} \rightarrow T_g \rightarrow M_g$.

Veech showed that a Teichmüller curve C is never complete ([35, Prop. 2.4]). Let \bar{C} be a smooth completion of C and $S := \bar{C} \setminus C$. In the sequel, rather than consider Teichmüller curves themselves, it will be convenient to consider finite unramified covers of C that satisfy two conditions: the corresponding subgroup of Γ is torsion free and the moduli map factors through a fine moduli space of curves (e.g. with level structure $M_g^{[n]}$). We nevertheless stick to the notation C for the base curve and let $f : \mathcal{X} \rightarrow C$ be the pullback of the universal family over $M_g^{[n]}$ to C . We will use $\bar{f} : \bar{\mathcal{X}} \rightarrow \bar{C}$ for the family of stable curves extending f . See also [29, §1.3].

Teichmüller curves, or more generally geodesic discs in Teichmüller space, are generated by a pair (X, q) of a Riemann surface and a quadratic differential $q \in \Gamma(X, (\Omega_X^1)^{\otimes 2})$. These pairs are called *translation surfaces*. If a pair (X, q) generates a Teichmüller curve, the pair is called a *Veech surface*. Any smooth fiber of f together with the suitable quadratic differential is a Veech surface. Theorem 2.1 below characterizes Teichmüller curves where $q = \omega^2$ is the square of a holomorphic 1-form $\omega \in \Gamma(X, \Omega_X^1)$. The examples that we construct will have this property, too. Hence:

From now on the notion ‘Teichmüller curve’ includes ‘generated by a 1-form’.

For a pair (X, ω) we let $\text{Aff}^+(X, \omega)$ be the group of orientation preserving diffeomorphisms of X that are affine with respect to the charts provided by integrating ω . Associating with an element of $\text{Aff}^+(X, \omega)$ its matrix part gives a well-defined map to $\text{SL}_2(\mathbb{R})$. The image of this map in $\text{SL}(X, \omega)$ is called the *affine group* of (X, ω) . The matrix part of an element of $\text{Aff}^+(X, \omega)$ is also called its *derivative*. The stabilizer group Γ of $\mathbb{H} \hookrightarrow T_g$ coincides, up to conjugation, with the affine group $\text{SL}(X, \omega)$ ([24]). Throughout this article, we denote the trace field by $K = \mathbb{Q}(\text{tr}(\gamma, \gamma \in \Gamma))$ and let $r := [K : \mathbb{Q}]$. We call the image of $\text{SL}(X, \omega)$ in $\text{PSL}_2(\mathbb{R})$ the *projective affine group* and denote it by $\text{PSL}(X, \omega)$.

We refer to [17] and [16] for a systematic description of Teichmüller curves in terms of billiards.

We recall from [29, Ths. 2.6 and 5.5] a description of the variation of Hodge structures (VHS) over a Teichmüller curve, and a characterization of Teichmüller curves in these terms.

Let \mathbb{L} be a rank 2 irreducible \mathbb{C} -local system on an affine curve C . Suppose that the Deligne extension \mathcal{E} of $\mathbb{L} \otimes_{\mathbb{C}} \mathbb{C}$ ([9, Prop. II.5.2]) to \bar{C} carries a Hodge

filtration of weight one $\mathcal{L} := \mathcal{E}^{(1,0)} \subset \mathcal{L}$. We denote by ∇ the corresponding logarithmic connection on \mathcal{E} . The *Kodaira-Spencer map* (also: Higgs field, or: second fundamental form) with respect to S is the composition map

$$(1) \quad \Theta : \mathcal{L} \rightarrow \mathcal{E} \xrightarrow{\nabla} \mathcal{E} \otimes \Omega_{\bar{C}}^1(\log S) \rightarrow (\mathcal{E}/\mathcal{L}) \otimes \Omega_{\bar{C}}^1(\log S).$$

A VHS of rank 2 and weight one whose Kodaira-Spencer map with respect to some S vanishes nowhere on \bar{C} is called *maximal Higgs* in [36]. The corresponding vector bundle \mathcal{E} is called *indigenous bundle*. See [5] or [27] for appearances of such bundles with more emphasis on characteristic $p > 0$.

THEOREM 2.1. (a) *Let $f : \mathcal{X} \rightarrow C$ be the universal family over a finite unramified cover of a Teichmüller curve. Then we have a decomposition of the VHS of f as*

$$(2) \quad R^1 f_* \mathbb{Q} = \mathbb{W} \oplus \mathbb{M} \quad \text{and} \quad \mathbb{W} \otimes_{\mathbb{Q}} \mathbb{C} = \bigoplus_{i=1}^r \mathbb{L}_i.$$

In this decomposition the \mathbb{L}_i are Galois conjugate, irreducible, pairwise nonisomorphic, \mathbb{C} -local systems of rank 2. The \mathbb{L}_i are in fact defined over some field $F \subset \mathbb{R}$ that is Galois over \mathbb{Q} and contains the trace field K . Moreover, \mathbb{L}_1 is maximal Higgs.

(b) *Conversely, suppose $f : \mathcal{X} \rightarrow C$ is a family of smooth curves such that $R^1 f_* \mathbb{C}$ contains a local system of rank 2 which is maximal Higgs with respect to the set $S = \bar{C} \setminus C$. Then f is the universal family over a finite unramified cover of a Teichmüller curve.*

Note that ‘maximal Higgs’ depends on S . We will encounter cases where \mathbb{L} extends over some points of S and becomes maximal Higgs with respect to a smaller set $S_u \subset S$, but it is not maximal Higgs with respect to S . See also Proposition 4.2 and Remark 4.3.

3. Local exponents of differential equations and zeros of the Kodaira-Spencer map

In this section we provide a dictionary between local systems plus a section on the one side and differential equations on the other side. In particular, we translate local properties of a differential operator into vanishing of the Kodaira-Spencer map. In Sections 5 and 6 we essentially start with a hypergeometric differential equation whose local properties are well-known. Via Proposition 3.2 the vanishing of the Kodaira-Spencer map of the corresponding local system is completely determined. This knowledge is then exploited in a criterion (Proposition 4.2) for a family of curves $f : \mathcal{X} \rightarrow C$ to be the universal family over a Teichmüller curve.

Let \mathbb{L} be an irreducible \mathbb{C} -local system of rank 2 on an affine curve C , not necessarily a Teichmüller curve. Let $C \hookrightarrow \bar{C}$ be the corresponding complete curve,

and let \mathbb{E} be the Deligne extension of \mathbb{L} (see §2). We suppose that \mathbb{L} carries a polarized VHS of weight one and choose a section s of $(\mathbb{L} \otimes_{\mathbb{C}} \mathbb{O}_C)^{(1,0)}$. Let t be a coordinate on C . We denote by $D := \nabla(\partial/\partial t)$. Since \mathbb{L} is irreducible, the sections s and Ds are linearly independent. Hence s satisfies a differential equation $Ls = 0$, where

$$L = D^2 + p(t)D + q(t),$$

for some meromorphic functions p, q on \bar{C} . Note that we may interpret L as a second order differential operator $L : \mathbb{O}_C \rightarrow \mathbb{O}_C$, by interpreting D as derivation with respect to t .

Conversely, the set of solutions of a second order differential operator $L : \mathbb{O}_C \rightarrow \mathbb{O}_C$ forms a local system $\text{Sol} \subset \mathbb{O}_C$. If L is obtained from \mathbb{L} then $\text{Sol} \cong \mathbb{L}^\vee$ ([9, §1.4]). The canonical map

$$\varphi : \text{Sol} \otimes_{\mathbb{C}} \mathbb{O}_C \rightarrow \mathbb{O}_C, \quad f \otimes g \mapsto fg$$

hence defines a section $s = s_\varphi$ of $\mathbb{L} \otimes_{\mathbb{C}} \mathbb{O}_C$.

A point $c \in \bar{C}$ is a *singular point* of L if p or q has a pole at c . In what follows, we always assume that L has regular singularities. Let t be a local parameter at $c \in \bar{C}$. Recall that L has a *regular singularity* at c if $(t - c)p$ and $(t - c)^2q$ are holomorphic at c , by Fuchs' theorem. Note that there is a difference between the notions 'singularity of the Deligne extension of the local system \mathbb{L} ' and 'singularities of the differential operator L '. We refer to [15, §11], for a definition of the notion regular singularity of a flat vector bundle. (The essential difference between the two notions is that the basis of [15, (11.2.1)], need not be a cyclic basis ([15, §11.4]).) Unless stated explicitly, we only use the notion of singularity of the differential operator.

The *local exponents* γ_0, γ_1 of L at c are the roots of the characteristic equation

$$t(t - 1) + tp_{-1} + q_{-2} = 0,$$

where $p = \sum_{i=-1}^{\infty} p_i(t - c)^i$ and $q = \sum_{i=-2}^{\infty} q_i(t - c)^i$. The table recording singularities and the local exponents is usually called *Riemann scheme*. See, e.g., [42, §2.5] for more details.

Note that L and the local exponents not only depend on \mathbb{L} but also on the section chosen. Replacing s by αs shifts the local exponents at c by the order of the function α at c . The exponentials $e^{2\pi i t_1}$ and $e^{2\pi i t_2}$ of the local exponents are the eigenvalues of the local monodromy matrix of L at c . The following criterion is well-known (e.g. [42, I.2.6]).

LEMMA 3.1. *All local monodromy matrices of Sol are unipotent if and only if both local exponents are integers for all $c \in \bar{C}$.*

In the classical case that $\bar{C} \cong \mathbb{P}^1$ the differential operator L is determined by the local exponents exactly if the number of singularities is three; this is the case of hypergeometric differential equations. We will exploit this fact in the next sections.

If the number of singularities is larger than three, L is no longer determined by the local exponents and the position of the singularities, but also depends on the *accessory parameters* ([42, §I.3.2]).

In the rest of this section we suppose that all local monodromy matrices of L are unipotent. We define $S_u = \bar{C} - C$ as the set of points where the monodromy is nontrivial. Let $S \subset \bar{C}$ be a set containing the singularities of the Deligne extension of \mathbb{L} . The reader should think of S being the set of singular fibers of a family of curves over \bar{C} . In particular $S \supset S_u$.

The following proposition expresses the order of vanishing of the Kodaira-Spencer map (1) at $c \in \bar{C}$ in terms of the local exponents at c . If $c \in C$ we suppose that the section s is chosen such that the local exponents are $(0, n_c)$ with $n_c \geq 0$. This is always possible, multiplying s with a power of a local parameter if necessary.

PROPOSITION 3.2. (a) *Let $c \in C$. Then $n_c \geq 1$.*

(b) *Suppose that $c \notin S$. The order of vanishing of Θ at b is $n_c - 1$.*

(c) *Suppose that $c \in S \setminus S_u$. The order of vanishing of Θ at b is n_c .*

(d) *If $c \in S_u$ then Θ does not vanish at c .*

Proof. Suppose that $c \in C$. Our assumptions imply that the local exponents $(0, n_c)$ at c are nonnegative integers. Since \mathbb{L} is a local system on C , it has two linearly independent algebraic section in a neighborhood of c . This implies that $n_c \geq 1$ ([42, §I.2.5]). This proves (a).

If $c \notin S$ the differential operator L has solutions s_1, s_2 with leading terms 1 and t^{n_c} , respectively ([42, I.2.5]). We want to determine the vanishing order of $D(s)$ in $\mathcal{E}/(s \otimes_{\mathbb{C}} \mathbb{O}_C)$. By the above correspondence between the local system and the differential equation we may as well calculate the vanishing order of $D(\varphi)$ in $(\text{Sol}^\vee \otimes_{\mathbb{C}} \mathbb{O}_C)/(\varphi \otimes_{\mathbb{C}} \mathbb{O}_C)$. A basis of $\text{Sol}^\vee \otimes_{\mathbb{C}} \mathbb{O}_C$ around c is

$$s_i^\vee : s_1 \otimes g_1 + s_2 \otimes g_2 \mapsto s_i g_i \quad (i = 1, 2).$$

By definition of the dual connection and the flatness of s_i one calculates that $D(\varphi)$ is the class of

$$s_1 \otimes g_1 + s_2 \otimes g_2 \mapsto g_1 s_1' + g_2 s_2'$$

in $(\text{Sol}^\vee \otimes_{\mathbb{C}} \mathbb{O}_C)/(\varphi \otimes_{\mathbb{C}} \mathbb{O}_C)$. Since both φ and s_1 do not vanish at c , we conclude that the order of vanishing of $D(\varphi)$ at c is $n_c - 1$. This proves (b).

In the case that $c \in S$ we should consider the contraction against $t \partial / \partial t$. This increases the order of vanishing of Θ by one. This proves (c).

We now treat the case that $c \in \bar{C} \setminus C$. Consider the residue map $\text{Res}_c(\nabla) \in \text{End}(\mathcal{E}_c)$. Suppose that the Kodaira-Spencer map vanishes at c . This implies that $\text{Res}_c(\nabla)$ is a diagonal matrix in a basis consisting of an element from \mathcal{L}_c and an element from its orthogonal complement. But $\text{Res}_c(\nabla)$ is nilpotent ([8, Prop. II.5.4 (iv)]), hence zero. This implies that two linearly independent sections of \mathbb{L} extend to c . This contradicts the hypothesis on the monodromy around c . This proves (d). \square

The ratios $\lambda(\mathbb{L}, S) := 2 \deg(\mathcal{L}) / \Omega_{\mathbb{C}}^1(\log S)$ will be of central interest in the sequel. The factor 2 is motivated by Section 9, where we interpret the $\lambda(\mathbb{L}, S)$ as Lyapunov exponents. Therefore we call the $\lambda(\mathbb{L}, S)$ from now on *Lyapunov exponents*. We will suppress S if it is clear from the context.

Remark 3.3. We will only be interested in local \mathbb{C} -systems \mathbb{L} that arise as local subsystems of $R^1 f_* \mathbb{C}$ for a family of curves $f : \mathcal{X} \rightarrow C$. In this case a Hodge filtration exists on \mathbb{L} and is unique ([8, Prop. 1.13]). Therefore we only have to keep track of the local system, but not of the VHS.

The following lemma is noted for future reference. The proof is straightforward.

LEMMA 3.4. *The ratio $\lambda(\mathbb{L}, S)$ does not change by taking unramified coverings.*

4. Cyclic covers of the projective line branched at four points

Let $N > 1$ be an integer, and suppose given a 4-tuple of integers (a_1, \dots, a_4) with $0 < a_\mu < N$ and $\sum_{\mu=1}^4 a_\mu = (k+1)N$, for some integer k . We denote by \mathbb{P}^1 the projective line with coordinate t , and put $\mathbb{P}^* = \mathbb{P}^1 - \{0, 1, \infty\}$. Let $\mathcal{P} \simeq \mathbb{P}^1 \times \mathbb{P}^* \rightarrow \mathbb{P}^*$ be the trivial fibration with fiber coordinate x . Let $x_1 = 0$, $x_2 = 1$, $x_3 = t$, $x_4 = \infty$ be sections of $\mathcal{P} \rightarrow \mathbb{P}^*$. We fix an injective character $\chi : \mathbb{Z}/N \rightarrow \mathbb{C}^*$. Let $g : \mathcal{X} \rightarrow \mathbb{P}^*$ be the N -cyclic cover of type (x_μ, a_μ) ([3, Def. 2.1]). This means that \mathcal{X} is the family of projective curves with affine model

$$(3) \quad \mathcal{X}_t : \quad z^N = x^{a_1}(x-1)^{a_2}(x-t)^{a_3}.$$

We suppose, furthermore, that $\gcd(a_1, a_2, a_3, a_4, N) = 1$. This implies that the family is connected. The genus of \mathcal{X}_t is $N + 1 - (\sum_{\mu=1}^4 \gcd(a_\mu, N))/2$.

In this section, we collect some well-known facts on such cyclic covers. We write

$$\sigma_\mu(i) = \langle i a_\mu / N \rangle = a_\mu(i) / N,$$

where $\langle \cdot \rangle$ denotes the fractional part. Let $k(i) + 1 = \sum_{\mu=1}^4 \sigma_\mu(i)$. We fix an injective character $\chi : \mathbb{Z}/N \rightarrow \mathbb{C}^*$ such that $h \in \text{Gal}(\mathcal{X}/\mathcal{P}) \cong \mathbb{Z}/N$ acts as $h \cdot z = \chi(h)z$.

LEMMA 4.1. *For $0 < i < N$, we let $s(i)$ be the number of a_μ unequal to $0 \bmod N / \gcd(i, N)$. Put $\mathbb{L}(i) = H_{\text{dR}}^1(\mathcal{X}/\mathbb{P}^*)$. Then*

- (a) $\dim_{\mathbb{C}} \mathbb{L}(i) = s(i) - 2$.
- (b) $\text{rank } g_*(\Omega_{\mathcal{X}/\mathbb{P}^*}^1)_{\chi^i} = s(i) - 2 - k(i)$, $\text{rank}(R^1 g_* \mathbb{C}_{\mathcal{X}})_{\chi^i} = k(i)$.
- (c) *If $k(i) = 1$, then*

$$\omega_i := \frac{z^i dx}{x^{1+[i\sigma_1]}(x-1)^{1+[i\sigma_2]}(x-t)^{1+[i\sigma_3]}}$$

is a nonvanishing section of $g_*(\Omega_{\mathcal{X}/\mathbb{P}^*}^1)_{\chi^i}$. It is a solution of the hypergeometric differential operator

$$L(i) := \nabla \left(\frac{\partial}{\partial t} \right)^2 + \frac{(A(i) + B(i) + 1)t - C(i)}{t(t-1)} \nabla \left(\frac{\partial}{\partial t} \right) + \frac{A(i)B(i)}{t(t-1)},$$

where

$$\begin{aligned} A(i) &= 1 - \sigma_3(i), \quad B(i) = 2 - (\sigma_1(i) + \sigma_2(i) + \sigma_3(i)), \\ C(i) &= 2 - (\sigma_1(i) + \sigma_3(i)). \end{aligned}$$

Proof. The second statement of (b) is proved in [2, Lemma 4.3]. The first statement follows from Serre duality and [2, Lemma 4.5]. Part (a) follows immediately from (b). The statement that ω_i is holomorphic and nonvanishing is a straightforward verification. The statement that $L(i)\omega_i = 0$ in $H_{\text{dr}}^1(\mathcal{X}/\mathbb{P}^*)_{\chi^i}$ is proved for example in [4, Lemma 1.1.4]. \square

The differential operator $L(i)$ corresponds to the local system

$$\mathbb{L}(i) = H_{\text{dr}}^1(\mathcal{X}/\mathbb{P}^*)_{\chi^i}$$

together with the choice of a section ω_i via the correspondence described at the beginning of Section 3. It has singularities precisely at 0, 1, and ∞ . Its local exponents are summarized in the Riemann scheme

$$(4) \quad \left[\begin{array}{ccc} t=0 & t=1 & t=\infty \\ 0 & 0 & A(i) \\ \gamma_0 := 1 - C(i) & \gamma_1 := C(i) - A(i) - B(i) & B(i) \end{array} \right].$$

A (Fuchsian) (m, n, p) -triangle group for $m, n, p \in \mathbb{N} \cup \{\infty\}$ satisfying $1/m + 1/n + 1/p < 1$ is a Fuchsian group in $\text{PSL}_2(\mathbb{R})$ generated by matrices M_1, M_2, M_3 satisfying $M_1 M_2 M_3 = 1$ and

$$\text{tr}(M_1) = \pm 2 \cos(\pi/m), \quad \text{tr}(M_2) = \pm 2 \cos(\pi/n), \quad \text{tr}(M_3) = \pm 2 \cos(\pi/p).$$

A triangle group is determined, up to conjugation in $\text{PSL}_2(\mathbb{R})$, by the triple (m, n, p) . It is well-known that the projective monodromy groups of the hypergeometric differential operators $L(i)$ are triangle groups under suitable conditions on $A(i), B(i), C(i)$. These conditions are met in the cases we consider in Section 5 and 6.

We are interested in determining the order of vanishing of the Kodaira-Spencer map. Note that if $k(i) = 0$ or $k(i) = 2$ then the Hodge filtration on the corresponding eigenspace is trivial and hence the Kodaira-Spencer map is zero.

Let $\bar{\pi} : \bar{C} \rightarrow \mathbb{P}^1$ be a finite cover, unbranched outside $\{0, 1, \infty\}$, such that the monodromy of the pullback of \mathcal{X} via $\bar{\pi}$ is unipotent for all $c \in \bar{C}$.

Let $S_u = S_u(i) \subset \pi^{-1}(0, 1, \infty)$ be the set of points such that $\mathbb{L}(i)$ has nontrivial local monodromy. Our assumption implies that the monodromy at $c \in S_u$ is infinite. In what follows, the set S_u will be nonempty. It is therefore no restriction to suppose that $\pi^{-1}(\infty)$ is contained in S_u . In terms of the invariants a_μ this means that

$a_3(i) + a_4(i) \equiv 0 \pmod{N}$. It follows that $A(i) = B(i)$. Let b_0 (resp. b_1) be the common denominator of the local exponents $\gamma_0(i)$ (resp. $\gamma_1(i)$) for $1 \leq i < N$. Write $|\gamma_0(i)| = n_0(i)/b_0$ and $|\gamma_1(i)| = n_1(i)/b_1$. Note that $\pi^{-1}(t = \mu) \subset S_u(i)$ if and only if $\gamma_\mu(i) = 0$. Therefore $S_u(i) = \pi^{-1}(\{0, 1, \infty\})$ if and only if $\gamma_0(i) = \gamma_1(i) = 0$. It easily follows that the set S_u is in fact independent of i .

The following proposition is the basic criterion we use for constructing Teichmüller curves.

PROPOSITION 4.2. *Consider a family of curves $\mathcal{X} \rightarrow \mathbb{P}_t^*$ as in (3). Let $0 < i_0 < N$ be an integer such that*

$$(5) \quad \gamma_\mu(i_0) = 1/b_\mu \quad \text{for all } \mu \in \{0, 1\} \text{ with } \gamma_\mu(i_0) \neq 0.$$

There is a finite cover $\pi : \bar{C} \rightarrow \mathbb{P}_t^1$ branched of order exactly b_μ at $t = \mu \in \{0, 1\}$ for all μ such that $\gamma_\mu(i_0) \neq 0$. Moreover, we require that the local monodromy of the pullback of $\mathbb{L}(i_0)$ to \bar{C} is unipotent, for all $c \in \bar{C}$. Write \mathcal{X}_C for the pullback of \mathcal{X} to C .

Choose a subgroup H of $\text{Aut}(\mathcal{X}_C)$ and define $\mathcal{X} := \mathcal{X}_C/H$. Suppose that

- \mathcal{X} extends to a smooth family over $\tilde{C} := \bar{C} \setminus S_u$,
- there is a local system \mathbb{L} isomorphic to $\mathbb{L}(i_0)$ which descends to \mathcal{X} .

Then the moduli map $\tilde{C} \rightarrow M_g$ is an unramified cover of a Teichmüller curve.

This criterion will be applied to subgroups H that intersect trivially with $\text{Gal}(\mathcal{X}_C/\mathcal{P}_C)$.

Proof. If $\gamma_\mu(i_0) \neq 0$ the monodromy of g at $t = \mu$ becomes trivial after pullback by a cover which is branched at $t = \mu$ of order b if and only if $b_\mu | b$. Hence if the cover $\bar{\pi}$ is sufficiently branched at points over S_u , the local monodromy of the pullback of $\mathbb{L}(i_0)$ to C is unipotent by Lemma 3.1.

The local exponents of the pullback of $\mathbb{L}(i_0)$ to \bar{C} are the original ones multiplied by the ramification index. Hence, for all $c \in \tilde{C}$, the local exponents are $(0, 1)$. By definition, the same holds for the local exponents of the bundle \mathbb{L} . The hypothesis on the singular fibers of \mathcal{X} implies that $c \in \tilde{C}$ is not a singularity of the flat bundle \mathbb{L} ([15, §14]). Therefore we may apply Proposition 3.2 (b) and (d) to \mathbb{L} with $S = S_u$. We conclude that the Kodaira-Spencer map of \mathbb{L} vanishes nowhere. The proposition therefore follows from Theorem 1.1. \square

Remark 4.3. The structure of the stable model $g_{\bar{C}}$ of the family $g_C : \mathcal{X}_C \rightarrow C$ is given in the next subsection. It implies that all fibers of preimages of $\{0, 1, \infty\}$ are singular. Hence applying Proposition 4.2 to $g_{\bar{C}}$ with $H = \{1\}$, we find that $g_{\bar{C}}$ defines a Teichmüller curve if and only if $S_u = \pi^{-1}(\{0, 1, \infty\})$. This happens for example for the families

$$y^2 = x(x-1)(x-t) \quad \text{and} \quad y^4 = x(x-1)(x-t).$$

Here $\bar{C} = \mathbb{P}_t^1$, and the uniformizing group is the triangle group $\Delta(\infty, \infty, \infty)$. Clearly, this is a very special situation.

PROPOSITION 4.4. *Let $0 < i < N$ be an integer with $k(i) = 1$. Denote by $\mathcal{L}(i)$ the $(1, 0)$ -part of the local system $\mathbb{L}(i)$ over C . Then*

$$\deg \mathcal{L}(i) = \frac{\deg(\pi)}{2} \left(1 - \frac{n_0(i)}{b_0} - \frac{n_1(i)}{b_1} \right)$$

with the convention that $1/b_\mu = 0$ if $n_\mu = 0$. In particular, the Lyapunov exponent

$$\lambda(\mathbb{L}(i), S_u) = \left(1 - \frac{n_0(i)}{b_0} - \frac{n_1(i)}{b_1} \right) / \left(1 - \frac{1}{b_0} - \frac{1}{b_1} \right)$$

is independent of the choice of $\bar{\pi}$.

Proof. We only treat the case that both $n_0(i)$ and $n_1(i)$ are nonzero, leaving the few modifications in the other cases to the reader. One checks that

$$\deg \Omega_{\bar{C}}^1(\log S_u) = \deg(\bar{\pi}) \left(1 - \frac{1}{b_0} - \frac{1}{b_1} \right)$$

is independent of the ramification order of g over $t = \infty$. It follows from the definition (1) of the Kodaira-Spencer map Θ that $2 \deg \mathcal{L}_{\chi^i} - \deg \Omega_{\bar{C}}^1(\log S_u)$ is the number of zeros of Θ , counted with multiplicity. Therefore the proposition follows from Proposition 3.2. \square

4.1. Degenerations of cyclic covers. We now describe the stable model of the degenerate fibers of \mathcal{X} . For simplicity, we only describe the fiber \mathcal{X}_0 above $t = 0$. The other degenerate fibers may be described similarly, by permuting $\{0, 1, t, \infty\}$. A general reference for this is [39, §4.3]. However, since we consider the easy situation of cyclic covers of the projective line branched at four points, we may simplify the presentation.

As before, we let $\mathcal{P} \rightarrow \mathbb{P}^*$ be the trivial fibration with fiber coordinate x . We consider the sections $x_1 = 0, x_2 = 1, x_3 = t, x_4 = \infty$ of $\mathcal{P} \rightarrow \mathbb{P}^*$ as marking on \mathcal{P} . We may extend \mathcal{P} to a family of stably marked curves over $\mathbb{P}(= \mathbb{P}_t^1)$, which we still denote by \mathcal{P} . The fiber P_0 of \mathcal{P} at $t = 0$ consists of two irreducible components which we denote by P_0^1 and P_0^2 . We assume that x_1 and x_3 (resp. x_2 and x_4) specialize to the smooth part of P_0^1 (resp. P_0^2). We denote the intersection point of P_0^1 and P_0^2 by ξ . It is well-known that the family of curves $f : \mathcal{X} \rightarrow \mathcal{P}$ over \mathbb{P}^* extends to a family of *admissible covers* over \mathbb{P}_t^1 . See for example [12] or [40]. For a short overview we refer to [41, §2.1].

The definition of the type of a cover ([3, Def. 2.1]) immediately implies that the restriction of the admissible cover $f_0 : \mathcal{X}_0 \rightarrow \mathcal{P}_0$ to P_0^1 (resp. P_0^2) has type $(x_1, x_3, \xi; a_1, a_3, a_2 + a_4)$ (resp. $(x_2, x_4, \xi; a_2, a_4, a_1 + a_3)$). (In our situation, admissibility amounts to $(a_1 + a_3) + (a_2 + a_4) \equiv 0 \pmod{N}$.) Let Z_0^j be a connected component of the restriction of \mathcal{X}_0 to P_0^j . Choosing suitable coordinates, Z_0^1

(resp. Z_0^2) is a connected component of the smooth projective curve defined by the equation $z^N = x^{a_1}(x-1)^{a_3}$ (resp. $z^N = x^{a_2}(x-1)^{a_4}$).

Denote by $H^j = \text{Gal}(Z_0^j, P_0^j) \subset H \simeq \mathbb{Z}/N$ the subgroups obtained by restricting the Galois action. Then \mathcal{X}_0 is obtained by suitably identifying the points in the fiber above ξ of $\text{Ind}_{H^1}^H Z_0^1$ and $\text{Ind}_{H^2}^H Z_0^2$.

Proposition 4.5 follows from the explicit description of the components of \mathcal{X}_0 . Put $\beta_1 = \gcd(a_1, a_3, N)$ and $\beta_2 = \gcd(a_2, a_4, N)$.

PROPOSITION 4.5. (a) *The degree of $Z_0^1 \rightarrow P_0^1$ (resp. $Z_0^2 \rightarrow P_0^2$) is N/β_1 (resp. N/β_2).*

(b) *The genus of Z_0^1 (resp. Z_0^2) is*

$$(N - \gcd(a_1, N) - \gcd(a_3, N) - \gcd(a_1 + a_3, N))/2\beta_1$$

$$\text{(resp. } (N - \gcd(a_2, N) - \gcd(a_4, N) - \gcd(a_1 + a_3, N))/2\beta_2).$$

(c) *The number of singular points of \mathcal{X}_0 is $\gcd(a_1 + a_3, N)$.*

5. Veech's n -gons revisited

In this section we realize the (n, ∞, ∞) -triangle groups as the affine groups of a Teichmüller curve. This result is due to Veech, but our method is different. An advantage of our method is that we obtain the Lyapunov exponents in Corollary 5.3 with almost no extra effort. The reader may take this section as a guideline to the more involved next section. In this section the family of cyclic covers which we consider has only one elliptic fixed point. A $(\mathbb{Z}/2\mathbb{Z})$ -quotient of this family is shown to be a Teichmüller curve. In the next section there are two elliptic fixed points and we need a $(\mathbb{Z}/2\mathbb{Z})^2$ -quotient. Moreover, in the next section, common divisors of m and n make a fiber product construction necessary, that does not show up here.

Let $n = 2k \geq 4$ be an even integer and fix a primitive n -th root of unity ζ_n . We specialize the results of Section 4 to the family $g : \mathcal{X} \rightarrow \mathbb{P}^*$ of curves of genus $n-1$ given by the equation

$$\mathcal{X}_t : z^n = x(x-1)^{n-1}(x-t),$$

i.e., we consider the case that $N = n$, $a_1 = a_3 = 1$, and $a_2 = a_4 = n-1$. Let

$$\varphi(x, y) = (x, \zeta_n y)$$

be a generator of $\text{Gal}(\mathcal{X}/\mathcal{P})$. The geometric fibers of g admit an involution covering $x \mapsto t/x$. We choose this involution to be

$$\sigma(x, y) = \begin{cases} \left(\frac{t}{x}, \frac{t^{2/n}(x-1)(x-t)}{xy} \right) & \text{if } k \text{ is even,} \\ \left(\frac{t}{x}, \zeta_n \frac{t^{2/n}(x-1)(x-t)}{xy} \right) & \text{if } k \text{ is odd.} \end{cases}$$

LEMMA 5.1. *The exponents a_i are chosen such that*

- (a) *condition (5) is satisfied for $i = (n + 1)/2$,*
- (b) *the projective monodromy group of the local systems*

$$\mathbb{L}((n - 1)/2) \text{ and } \mathbb{L}((n + 2)/2)$$

is the triangle group $\Delta(n, \infty, \infty)$.

Proof. Part (a) follows by direct verification. Part (b) is proved in [6]. \square

Let $\pi : C \rightarrow \mathbb{P}^*$ be defined by $s = t^{n/2}$. Then σ extends to an automorphism of the family of curves $g_C : \mathcal{X}_C \rightarrow C$. As before, we let $\bar{\pi} : \bar{C} \rightarrow \mathbb{P}^1$ be the extension of π to a smooth completion. Moreover, the local monodromy matrices of the pullback of the local systems $\mathbb{L}(i)$ to \bar{C} are unipotent.

Let $f : \mathcal{X} = \mathcal{X}/\langle \sigma \rangle \rightarrow C$ and let $\bar{f} : \bar{\mathcal{X}} \rightarrow \bar{C}$ be the stable model of f . Our goal is to show that the fibers \mathcal{X}_c of \bar{f} are smooth for all $c \in \tilde{C} := \bar{\pi}^{-1}(\mathbb{P}^1 \setminus \{1, \infty\})$. This allows us to apply the criterion (Proposition 4.2) for \tilde{C} to be the cover of a Teichmüller curve.

THEOREM 5.2. *Let $g = (n - 2)/2$. The natural map $m : \tilde{C} \rightarrow M_g$ induced by \bar{f} exhibits \tilde{C} as the unramified cover of a Teichmüller curve.*

Proof. We first determine the degeneration of g_C at $c \in \bar{C}$ with $\bar{\pi}(c) \in \{0, 1, \infty\}$. Our assumption on the local monodromy matrices implies that the fiber \mathcal{X}_c is a semi-stable curve, and we may apply Proposition 4.5. For $\bar{\pi}(c) \in \{1, \infty\}$ the fiber \mathcal{X}_c consists of two irreducible components, which have genus 0. The local monodromy matrices of $\mathbb{L}(i)$ at c are unipotent and of infinite order for all i , as can be read off from the local exponents.

Similarly, the local monodromy at c with $\bar{\pi}(c) = 0$ is finite. The definition of C implies therefore that it is trivial. The set $S_u \subset \bar{C}$ (notation of §4) consists exactly of $\bar{\pi}^{-1}\{1, \infty\}$.

One checks that σ acts on the holomorphic 1-forms ω_i (Lemma 4.1(c)) as follows:

$$(6) \quad \sigma^* \omega_i = (-1)^i d(i) \omega_{n-i} \quad \text{for } i \neq n/2, \quad \sigma^* \omega_{n/2} = -\omega_{n/2},$$

where $d(i) = t^{2i/n-1}$ if k is odd and $d(i) = t^{2i/n-1} \zeta_n^i$ if k is even. This implies that the generic fiber of \mathcal{X} has genus $n/2 - 1$.

We claim that \mathcal{X}_c is smooth for all $c \in \bar{C} - S_u$. We only need to consider $c \in \bar{C}$ such that $\pi(c) = 0$. Proposition 4.5 implies that the degenerate fiber \mathcal{X}_c consists of two components of genus $n/2 - 1$. Note that σ acts as the permutation $(0 \infty)(1 t)$ on the branch points of $\mathcal{X} \rightarrow \mathcal{P}$. Hence σ interchanges the two components of \mathcal{X}_c . We conclude that the quotient \mathcal{X}_c of \mathcal{X}_c by ρ is a smooth curve of genus $n/2 - 1$.

Consider the local system $\mathbb{M} = \mathbb{L}((n-2)/2) \oplus \mathbb{L}((n+2)/2)$ in $R^1(g_C)_* \mathbb{C}$ on C . It is invariant under σ . The part of \mathbb{M} on which σ acts trivially is a local subsystem $\mathbb{L} \subset \mathbb{M}$. This \mathbb{L} is necessarily of rank 2, since $\omega_{(n-2)/2} + d((n-2)/2) \omega_{(n+2)/2}$ is

σ -invariant (resp. anti-invariant), if k is odd (resp. even). Furthermore, $\omega_{(n-2)/2} - d((n-2)/2)\omega_{(n+2)/2}$ is σ -anti-invariant (resp. invariant) for k odd (resp. even). This also implies that the compositions

$$\mathbb{L} \rightarrow \mathbb{L}((n-2)/2) \oplus \mathbb{L}((n+2)/2) \rightarrow \mathbb{L}((n-2)/2)$$

and

$$\mathbb{L} \rightarrow \mathbb{L}((n-2)/2) \oplus \mathbb{L}((n+2)/2) \rightarrow \mathbb{L}((n+2)/2)$$

are nontrivial. Since the monodromy group Γ of both $\mathbb{L}((n-2)/2)$ and $\mathbb{L}((n+2)/2)$ contains two noncommuting parabolic elements, we conclude that $\mathbb{L}((n-2)/2)$ is an irreducible local system, and hence that

$$\mathbb{L} \cong \mathbb{L}((n-2)/2) \oplus \mathbb{L}((n+2)/2).$$

From Proposition 4.2 and Lemma 5.1(a) we conclude that \mathcal{X} is the universal family over an unramified cover of a Teichmüller curve as claimed. \square

Corollary 5.3 follows from Proposition 4.4.

COROLLARY 5.3. *The VHS of the family $f : \mathcal{X} \rightarrow C$ decomposes as*

$$R^1 f_* \mathbb{C} \cong \bigoplus_{j=1}^{(n-2)/2} \mathbb{L}_j,$$

where \mathbb{L}_j is a rank 2 local system isomorphic to $\mathbb{L}((n-2j)/2)$. Moreover,

$$\lambda(\mathbb{L}_j) = \frac{k-j}{k-1}.$$

Anton Zorich has communicated to the authors that he (with Maxim Kontsevich) independently calculated these Lyapunov exponents.

Remark 5.4. The trace field of $\Delta(n, \infty, \infty)$ is $K = \mathbb{Q}(\zeta_n + \zeta_n^{-1})$, hence $r = [K : \mathbb{Q}] \leq \phi(n/2)$. Corollary 5.3 allows to decompose the VHS of \mathcal{X} completely into rank 2 pieces. This is much finer than Theorem 2.1 that predicts only r pieces of rank 2 plus some rest.

Each fiber \mathcal{X}_t admits an extra isomorphism, namely

$$\tau(x, y) = \left(\frac{x-t}{x-1}, y \frac{t-1}{(x-1)^2} \right).$$

It extends to an automorphism of the family $g_{\tilde{C}} : \mathcal{X}_{\tilde{C}} \rightarrow \tilde{C}$. One checks that τ and σ commute. Hence τ descends to an automorphism of \mathcal{X} , which we also denote by τ . Let $p : \mathcal{U} = (\bar{\mathcal{X}}|_{\tilde{C}})/\langle \tau \rangle \rightarrow \tilde{C}$ be the quotient family. One calculates that

$$\tau^* \omega_i = (-1)^{i+1} \omega_i.$$

From this we deduce that the fibers of \mathcal{X} are Veech surfaces that cover nontrivially Veech surfaces of smaller genus, the fibers of the fibers of p .

THEOREM 5.5. (a) *The moduli map $\tilde{C} \rightarrow M_{g(\mathfrak{U})}$ of the family of curves $p : \mathfrak{U} \rightarrow \tilde{C}$ is an unramified covering of Teichmüller curve. Its VHS decomposes as*

$$R^1 p_* \mathbb{C} \cong \bigoplus_{j=0}^{t(n)} \mathbb{L}(1+2j),$$

where $\mathbb{L}(j)$ is the local system appearing in the VHS of f and $t(n) = (n-6)/4$ if k is odd (resp. $t(n) = (n-4)/4$ if k is even).

(b) *The genus of \mathfrak{U} is $t(n) + 1$ and*

$$\lambda(\mathbb{L}(1+2j)) = \frac{k - (1+2j)}{k-1}.$$

Proof. Both for k odd and k even the generating holomorphic 1-form in $\mathbb{L}(1)$ is τ -invariant. Hence this local system descends to \mathfrak{U} . The property of being a Teichmüller curve now follows from Proposition 4.2. The remaining statements are easily deduced from Corollary 5.3. \square

Let U be a fiber of \mathfrak{U} . We denote by $\omega_X \in \Gamma(X, \Omega_X^1)$ (resp. $\omega_U \in \Gamma(U, \Omega_U^1)$) the differential that pulls back to $\omega_{(n-2)/2} \pm d(i)\omega_{(n+2)/2}$ on \mathcal{X}_c , where the sign depends on the parity of n and refer to it as the *generating differential* of the Teichmüller curve.

COROLLARY 5.6. *The Teichmüller curve \mathcal{X} is the one generated by the regular n -gon studied in [35].*

Proof. Let c be a point of \bar{C} with $\pi(c) = 0$. The fiber \mathcal{X}_c consists of two components isomorphic to

$$\mathcal{X}_0 : y^n = x(x-1)$$

which are interchanged by σ . The generating differential ω_X specializes to the differential

$$\omega_0 = y^{(n-2)/2} dx/x(x-1)$$

on \mathcal{X}_0 . There is an obvious isomorphism between the curve $w^n - 1 = z^2$ and \mathcal{X}_0 such that ω_0 pulls back to the differential dw/z considered by Veech ([35, Th. 1.1]). \square

Actually the family \mathcal{X} is isomorphic (after some base change) to

$$y^2 = p_t(z) = \prod_{i=1}^n (x - \zeta_n^i - t\zeta_n^{-1}).$$

This was shown by Lochak ([20]; see also [26]).

The following proposition is shown in [35, Th. 1.1]. We give an alternative proof in our setting.

PROPOSITION 5.7. *The projective affine group of a fiber of \mathcal{X} together with the generating differential contains the (n, ∞, ∞) -triangle group. The same holds for the fibers of \mathfrak{U} .*

Proof. We first consider \mathcal{X} . We have to show that the moduli map $C \rightarrow M_g$ given by \mathcal{X} factors through $\pi : C \rightarrow \mathbb{P}^*$. That is, we have to show that two generic fibers \mathcal{X}_p and $\mathcal{X}_{\tilde{p}}$ with $p, \tilde{p} \in C$ such that $\pi(p) = \pi(\tilde{p})$ are isomorphic. Equivalently, we have to show that for p, \tilde{p} as above there is an isomorphism $i_0 : \mathcal{X}_p \rightarrow \mathcal{X}_{\tilde{p}}$ which is σ -equivariant. It suffices to show the existence of i_0 after any base change $\pi : C' \rightarrow \mathbb{P}^*$ such that σ is defined on $\mathcal{Y}_{C'}$. We may suppose that $\bar{\pi} : \bar{C}' \cong \mathbb{P}_s^1 \rightarrow \mathbb{P}_t^1$ is given by $t = s^{n/2}$. The hypothesis $\pi(p) = \pi(\tilde{p})$ implies that $p = \zeta_n^{2e} \tilde{p}$, for some p . It follows that the canonical isomorphism $i : \mathcal{X}_p \rightarrow \mathcal{X}_{\tilde{p}}$, given by $(x, y) \mapsto (x, y)$, satisfies

$$\sigma \circ i = \varphi^{2e} \circ i \circ \sigma.$$

Hence $i_0 = \varphi^e \circ i$ is the isomorphism we were looking for.

The proof for the family \mathcal{U} is similar. □

We record for completeness:

COROLLARY 5.8. *All (n, ∞, ∞) -triangle groups for $n \geq 4$ arise as projective affine groups.*

Remark 5.9. For n odd the same construction works with N and a_i chosen as above. The local exponents of $(\mathbb{L}(i), \omega_i)$ at $t = 0$ are then $1 - 2i/n$. The local system $\mathbb{L}(i_0)$ becomes maximal Higgs for $i_0 = (n+1)/2$, after a base change $\pi : C \rightarrow \mathbb{P}^*$ whose extension to $\bar{C} \rightarrow \mathbb{P}^1$ is branched of order n at 0. The quotient family $f : \mathcal{X} = \mathcal{Z}/\langle \sigma \rangle \rightarrow C$ may be constructed in the same way as above. Its moduli map yields as above a Teichmüller curve $\tilde{C} \rightarrow M_g$ where $g = (n-1)/2$. The corresponding translation surfaces are again the ones studied in [35]. Veech also determines that the affine group is not $\Delta(n, \infty, \infty)$ but the bigger group $\Delta(2, n, \infty)$, containing $\Delta(n, \infty, \infty)$ with index 2. We obtain the same family of curves also as a special case of the construction in Section 6, by putting $m = 2$. For this family we calculate, using Proposition 4.4, that

$$\lambda(\mathbb{L}(i)) = \frac{2i}{n-1}, \quad i = 1, \dots, (n-1)/2.$$

6. Realization of $\Delta(m, n, \infty)$ as projective affine group

Let $m, n > 1$ be integers with $mn \geq 6$. Let

$$\sigma_1 = \frac{nm + m - n}{2mn}, \quad \sigma_2 = \frac{nm - m + n}{2mn}, \quad \sigma_3 = \frac{nm + m + n}{2mn}, \quad \sigma_4 = \frac{nm - m - n}{2mn},$$

and let N be the least common denominator of these fractions. Let $a_i = N\sigma_i$ and consider the family of curves $g : \mathcal{Z} \rightarrow \mathbb{P}^*$ given by

$$\mathcal{Z}_t : \quad z^N = x^{a_1}(x-1)^{a_2}(x-t)^{a_3}.$$

The family g cyclically covers the constant family $\mathcal{P} \cong \mathbb{P}_x^1 \times \mathbb{P}^* \rightarrow \mathbb{P}^* (= \mathbb{P}_t^1 - \{0, 1, \infty\})$.

The plan of this section is as follows. We construct a cover $\mathcal{Y} \rightarrow \mathcal{X}$ such that the involutions

$$(7) \quad \begin{aligned} \sigma(x) &= (t(x-1)/(x-t)), \\ \tau(x) &= (t/x) \end{aligned}$$

of $\mathcal{P} \rightarrow \mathbb{P}^*$ lift to involutions of the family $\mathcal{Y}_C \rightarrow C$ obtained from $\mathcal{Y} \rightarrow \mathbb{P}^*$ by a suitable unramified base change $\pi : C \rightarrow \mathbb{P}^*$. We denote these lifts again by σ and τ . If m and n are relatively prime then in fact \mathcal{Y} equals \mathcal{X} .

Remark 6.1. The exponents a_i are chosen such that the local system \mathbb{L}_χ has as projective monodromy group the triangle group $\Delta(m, n, \infty)$; see again, e.g., [6]. We modify the lifts τ and σ by appropriate powers of a generator of $\text{Aut}(\mathcal{X}/\mathcal{P})$ such that the group $H = \langle \tau, \sigma \rangle$ is still isomorphic to $(\mathbb{Z}/2)^2$ and such that σ, τ , and $\sigma\tau =: \rho$ have ‘as many fixed points as possible’.

We consider the quotient family $f : \mathcal{X} = \mathcal{Y}/H \rightarrow C$. Its stable model $\bar{f} : \bar{\mathcal{X}} \rightarrow \bar{C}$ has smooth fibers over $\bar{C} = \bar{\pi}^{-1}(\mathbb{P}^1 \setminus \{\infty\})$, where $\bar{\pi} : \bar{C} \rightarrow \mathbb{P}^1$ extends π .

Together with an analysis of the action of H on differentials we can apply Proposition 4.2 to produce Teichmüller curves.

THEOREM 6.2. *Via the natural map $m : \bar{C} \rightarrow M_g$ induced from \bar{f} , the curve \bar{C} is an unramified cover of a Teichmüller curve. The genus g is given in Corollary 6.6.*

As corollaries to this result we calculate the precise VHS of f and the projective affine group of the translation surfaces corresponding to f . In Section 6.1 we show that for $m = 3$ we rediscover Ward’s Teichmüller curves ([38]).

Remark 6.3. The notation in the proof of Theorem 6.2 is rather complicated, due to the necessary case distinction. We advise the reader to restrict to the case that m and n are odd and relatively prime on a first reading. This considerably simplifies the notation, but all main features of the proof are already visible. In this case $\mathcal{Y} = \mathcal{X}$, $m = m'$, $n = n'$, $\gamma = \beta = 1$, and $N = \hat{N}$.

We start with some more notation. We write Z (resp. P, X, Y) for the geometric generic fiber of \mathcal{X} (resp. $\mathcal{P}, \mathcal{X}, \mathcal{Y}$). We choose a primitive N -th root of unity $\zeta_N \in \mathbb{C}$ and define the automorphism $\varphi_1 \in \text{Aut}(\mathcal{X}/\mathcal{P})$ by

$$\varphi_1(x, z) = (x, \zeta_N z).$$

We need to determine the least common denominator N of the $\sigma_i, i = 1, \dots, 4$, precisely. Let $m = 2^\mu m', n = 2^\nu n'$ with m', n' odd. We may suppose that $\mu \geq \nu$. Define

$$\gamma_1 = \gcd(2mn, mn + m - n), \quad \gamma_2 = \gcd(2mn, mn + m + n), \quad \gamma = \gcd(m, n),$$

and write $\gamma = 2^\nu \gamma'$. We determine $N = 2mn / \gcd(\gamma_1, \gamma_2)$. For this we need to distinguish four cases.

- Case O $\mu = \nu = 0$, $N = 2mn/\gamma$, $\hat{N} = N/\gamma = 2^\delta m'n'/\gamma'^2$;
- Case OE $\mu > \nu = 0$, $N = 2mn/\gamma$, $\hat{N} = N/\gamma = 2^\delta m'n'/\gamma'^2$;
- Case DE $\mu > \nu > 0$, $N = 2mn/\gamma$, $\hat{N} = 2N/\gamma = 2^\delta m'n'/\gamma'^2$;
- Case S $\mu = \nu \neq 0$, $N = mn/\gamma$, $\hat{N} = N/\gamma = mn/\gamma^2$.

It is useful to keep in mind that $\gamma = \gcd(\gamma_1, \gamma_2)$, except in case S where $2\gamma = \gcd(\gamma_1, \gamma_2)$. Let $\delta := 0$ in case S, and $\delta := \min\{\mu - \nu + 2, \mu + 1\}$, otherwise.

Our first goal is to determine the maximal intermediate covering of $Z \rightarrow P$ to which τ lifts. This motivates the definition of \hat{N} . Let $0 < \bar{\alpha} < \hat{N}$ be the integer satisfying

$$\bar{\alpha} \equiv 1 \pmod{m'/\gamma'}, \quad \bar{\alpha} \equiv -1 \pmod{n'/\gamma'}, \text{ and}$$

$$\bar{\alpha} \equiv \begin{cases} 1 & \pmod{2^\delta} \text{ cases O, OE, S,} \\ \frac{n' + 2^{\mu-\nu}m'}{n' - 2^{\mu-\nu}m'} \equiv 1 + 2^{\delta-1} & \pmod{2^\delta} \text{ case DE.} \end{cases}$$

For convenience, we lift $\bar{\alpha}$ to an element α in $\mathbb{Z}/N\mathbb{Z}$ such that $\alpha^2 = 1$. Note that $\bar{\alpha}$ may be one.

Recall that for a rational number σ , we write $\sigma(i) := \langle i\sigma \rangle$ (the fractional part). Similarly, for an integer a we write $a(i) = a(i; \nu) = \nu \langle ia/\nu \rangle$, where ν is mostly clear from the context. For each integer $0 < i < N$ which is prime to N , we write

$$z(i) = \frac{z^i}{x^{[i\sigma_1]}(x-1)^{[i\sigma_2]}(x-t)^{[i\sigma_3]}},$$

hence

$$z(i)^N = x^{a_1(i)}(x-1)^{a_2(i)}(x-t)^{a_3(i)}.$$

LEMMA 6.4. (a) *In the cases O, OE and DE the covering $Z \rightarrow P$ has ramification order $\gamma N/\gamma_1$ (resp. $\gamma N/\gamma_2$) in points of Z over $x = 0, 1$ (resp. $x = t, \infty$). In case S the ramification orders are $\gamma N/2\gamma_1$ (resp. $\gamma N/2\gamma_2$). Therefore*

$$g(Z) = \begin{cases} 1 + N - (\gamma_1 + \gamma_2)/2\gamma & \text{case S,} \\ 1 + N - (\gamma_1 + \gamma_2)/\gamma & \text{other cases.} \end{cases}$$

- (b) *The automorphism σ of P lifts to an automorphism σ of Z of order 2.*
- (c) *The automorphism τ of P lifts to an automorphism τ of order 2 of $\hat{Z} := Z/\langle \varphi_1^{\hat{N}} \rangle$. Moreover, we may choose the lifts such that σ, τ commute as elements of $\text{Aut}(\hat{Z})$.*
- (d) *We may choose the lifts σ, τ such that, moreover, τ has $4m/\gamma$ fixed points (resp. $2m/\gamma$ in case S) and such that $\rho := \sigma\tau$ has $4n/\gamma$ fixed points on \hat{Z} (resp. $2n/\gamma$ in case S).*
- (e) *With σ and τ chosen as in (d) the automorphism σ has no (2 in case S) fixed points both on Z and on \hat{Z} .*

Proof. The statements in (a) are immediate from the definitions. For (b) and (c) we choose once and for all elements $t^{1/n}, (t-1)^{1/m} \in \overline{\mathbb{C}(t)}$. Define

$$(8) \quad c = (t-1)^{\sigma_2+\sigma_3}, \quad d = t^{\sigma_1+\sigma_3}.$$

Then

$$\sigma(z) = cd \frac{x(x-1)}{z(x-t)} = cd \frac{z(N-1)}{(x-t)^2}$$

defines a lift of σ to Z , since $\sigma_1 + \sigma_2 = \sigma_3 + \sigma_4 = 1$. Moreover, this lift has order 2. We denote it again by σ . The quotient curve \hat{Z} is defined by the equation

$$\bar{z}^{\hat{N}} = x^{\bar{a}_1} (x-1)^{\bar{a}_2} (x-t)^{\bar{a}_3},$$

where \bar{a}_i denotes $a_i \bmod \hat{N}$. Put $\bar{\sigma}_i = \bar{a}_i / \hat{N}$. One computes that α satisfies:

$$(9) \quad (\bar{a}_1(\alpha), \bar{a}_2(\alpha), \bar{a}_3(\alpha), \bar{a}_4(\alpha)) = (\bar{a}_4, \bar{a}_3, \bar{a}_2, \bar{a}_1).$$

This implies that

$$\tau(\bar{z}) = (-1)^{\bar{a}_2+\bar{a}_3} t^{\bar{\sigma}_1+\bar{\sigma}_3} \frac{\bar{z}(\alpha)}{x^2}$$

defines a lift of τ to \hat{Z} which has order 2. Note that $\bar{a}_2 + \bar{a}_3$ is odd in case S and even in all other cases. Moreover,

$$\bar{\sigma}_1 + \bar{\sigma}_3 \equiv \begin{cases} \gamma/n \bmod \hat{N} & \text{in cases O, OE and S,} \\ \gamma/2n \bmod \hat{N} & \text{in case DE.} \end{cases}$$

It is easy to check that τ commutes with the image of σ on \hat{Z} . This proves (b). Furthermore, one checks that σ is an involution and that

$$(10) \quad \tau\varphi_1\tau = \varphi_1^\alpha \in \text{Aut}(\hat{Z}) \quad \text{and} \quad \sigma\varphi_1\sigma = \varphi_1^{-1} \in \text{Aut}(Z).$$

This proves (c).

We start with the proof of (d). Let $x_1 = \sqrt{t}$ be one of the fixed points of τ on P and let R be a point in the fiber of $\hat{Z} \rightarrow P$ over x_1 . We may describe the whole fiber by $R_a := \varphi_1^a R$ for $a = 0, \dots, \hat{N} - 1$. Suppose that $\tau R = R_{a_0}$, hence $\tau R_a = R_{a_0+\alpha a}$. Since τ is an involution, a_0 satisfies necessarily $a_0 \equiv 0 \bmod m'/\gamma'$ and $2a_0 \equiv 0 \bmod 2^\delta$. Furthermore, R_a is a fixed point of τ if and only if

$$(11) \quad a_0 \equiv 2a \bmod n'/\gamma' \quad \text{and} \quad a_0 \equiv 2^{\mu-\nu+1} 2^{\delta-1} a \bmod 2^\delta \quad (\text{case DE only}).$$

Hence if τ has a fixed point in this fiber, it has precisely $2m/\gamma$ fixed points in this fiber (m/γ in case S). Since τ and σ commute, σ bijectively maps fixed points of τ over x_1 to fixed points of τ over $x_2 = -\sqrt{t}$. Hence, if τ has a fixed point, then the number of fixed points is as stated in (d).

Similarly, let $x_3 = 1 + \sqrt{1+t}$ be one of the fixed points of ρ on P and let S be a point in the fiber over x_3 . Write $S_b = \varphi_1^b S$ for the whole fiber. Write $\rho S = S_{b_0}$. As above we deduce that $b_0 \equiv 0 \bmod m'/\gamma'$. In case DE, we have moreover that $b_0 \equiv 0 \bmod 2$. The point S_b is fixed by ρ if and only if

$$(12) \quad b_0 \equiv 2b \pmod{m'/\gamma'} \quad \text{and} \quad b_0 \equiv \begin{cases} 2b \pmod{2^\delta} & \text{cases O, OE, S,} \\ (2 + 2^{\delta-2})b \pmod{2^\delta} & \text{case DE.} \end{cases}$$

Analogous to the argument for τ , one checks that if ρ has a fixed point, then it has as many fixed points as claimed in (d).

Note that we may replace σ by $\varphi_1^i \sigma$ and τ by $\varphi_1^j \tau$ such that they still commute, and without changing the orders of these elements, if the following conditions are satisfied:

$$(13) \quad j(1 + \alpha) \equiv 0 \pmod{\hat{N}}, \quad (i - j)(1 - \alpha) \equiv 0 \pmod{\hat{N}}.$$

The only obstruction for τ and ρ to have fixed points consists in a condition modulo 2^δ . We check in each case that we can modify τ and ρ respecting (13) such that this obstruction vanishes. This proves (d).

For (e) we check, with the same argument as above, that σ has 0 or 4 (resp. 0 or 2 in case S) fixed points. Checking case by case one finds that $\hat{Z} \rightarrow P$ is totally ramified over $\{0, 1, t, \infty\}$. Hence $g(\hat{Z}) = \hat{N} - 1$. The Riemann-Hurwitz formula implies that σ does not have fixed points on \hat{Z} , hence also not on Z in cases O, OE, and DE. The number of fixed points of σ in case S may be checked directly by counting fixed points of τ on \hat{Z} . \square

Let Z^τ be the conjugate of Z under τ . Define Y as the normalization of $Z \times_{\hat{Z}} Z^\tau$. As remarked above, the definition of \hat{N} implies that $\hat{Z} \rightarrow P$ is the largest subcover of $Z \rightarrow P$ such that τ lifts to \hat{Z} . In other words, $Y \rightarrow \hat{P} := P / \langle \sigma, \tau \rangle$ is the Galois closure of $Z \rightarrow \hat{P}$. This implies that Y is connected. The particular choice of \hat{N} is used precisely to guarantee that the Veech surfaces constructed in Theorem 6.2 are connected.

By construction, σ lifts to Z acting on both Z and Z^τ and τ lifts to Z by exchanging the two factors of the fiber product. These two involutions commute and $\rho := \sigma\tau$ also has order 2. We have defined the following coverings. The labels indicate the Galois group of the morphism with the notation introduced in the following lemma.

$$\begin{array}{ccccc}
 & & Y & & \\
 & \swarrow \langle \psi_2 \rangle & & \searrow \langle \psi_1^{\hat{N}} \psi_2^{-1} \rangle & \\
 Z & & & & Z^\tau \\
 & \searrow \langle \varphi_1^{\hat{N}} \rangle & & \swarrow \langle \varphi_2^{\hat{N}} \rangle & \\
 & & \hat{Z} & & \\
 & & \downarrow \langle \varphi_1 \bmod \hat{N} \rangle = \langle \varphi_2 \bmod \hat{N} \rangle & & \\
 & & P & & \\
 & & \downarrow \langle \sigma, \tau \rangle & & \\
 & & \hat{P} & &
 \end{array}$$

LEMMA 6.5. (a) *We may choose a generator φ_2 of $\text{Aut}(Z^\tau/P)$ such that the Galois group, G_0 , of Y/P is*

$$G_0 \cong \{(\varphi_1^i, \varphi_2^j), i, j, \in \mathbb{Z}/N\mathbb{Z}, i \equiv j \pmod{\hat{N}}\} \subset \langle \varphi_1 \rangle \times \langle \varphi_2 \rangle \cong (\mathbb{Z}/N\mathbb{Z})^2.$$

We fix generators $\psi_1 = (\varphi_1, \varphi_2)$ and $\psi_2 = (0, \varphi_2^{\hat{N}})$ of G_0 . The Galois group, G , of the covering Y/\hat{P} is generated by $\psi_1, \psi_2, \sigma, \tau$, satisfying

$$\psi_1^N = \psi_2^\beta = \sigma^2 = \tau^2 = 1, \quad [\psi_1, \psi_2] = [\sigma, \tau] = 1,$$

$$\sigma \psi_i \sigma = \psi_i^{-1} \quad (i = 1, 2), \quad \tau \psi_1 \tau = \psi_1^\alpha, \quad \text{and} \quad \tau \psi_2 \tau = \psi_1^{\alpha N} \psi_2^{-\alpha} (= (\varphi_1^{\alpha N}, 0)).$$

- (b) *The genus of Y is $g(Y) = 1 + N\beta - 2\beta$, where $\beta = \gamma/2$ in case DE and $\beta = \gamma$ in the other cases.*
- (c) *The number of fixed points of τ on Y is $4m$ in case O and OE (resp. $2m$ in case DE and S).*
- (d) *The number of fixed points of ρ on Y is $4n$ in case O and OE (resp. $2n$ in case DE and S).*
- (e) *The involution σ has no fixed points on Y .*

Proof. The presentation in (a) follows from the above construction. To prove (b), we remark that Z^τ is given by the equation

$$\tilde{z}^N = x^{a_4}(x-1)^{a_3}(x-t)^{a_2},$$

compare to (9). Recall that $\hat{Z} \rightarrow P$ is totally ramified over $\{0, 1, t, \infty\}$. Hence at each of the γ_1/γ points (resp. $\gamma_1/2\gamma$ in case S) over 0 and 1 in Z the map $Z \rightarrow \hat{Z}$ is branched of order γ^2/γ_1 (resp. $2\gamma^2/\gamma_1$ in case S and $\gamma^2/2\gamma_1$ in case DE). The other covering $Z^\tau \rightarrow \hat{Z}$ is branched at the corresponding γ_1/γ (resp. $\gamma_1/2\gamma$ in case S) points of order γ^2/γ_2 (resp. $2\gamma^2/\gamma_2$ in case S and $\gamma^2/2\gamma_2$ in case DE). Over t and ∞ instead of 0 and 1 the roles of γ_1 and γ_2 are interchanged.

It follows from Abhyankar's lemma that $Y \rightarrow \hat{Z}$ is ramified in all cases at each point over 0, 1, t , ∞ of order β . Hence these fibers of $Y \rightarrow P$ consist of β points in each case.

For (c), (d), and (e) note that $Z \rightarrow P$ is unramified over the fixed points of τ , σ , and ρ . Hence Y is indeed the fiber product in neighborhoods of these points. Since τ interchanges the two factors, exactly β of the β^2 preimages in Y of a fixed point of τ on Z will be fixed by the lift of τ to Y . This completes the proof of (c).

For (d) note that $\text{id} \times \sigma : Z \times_{\hat{Z}} Z^\tau \rightarrow Z \times_{\hat{Z}} Z^\tau$ is an isomorphism and we may now argue as in (c).

If σ has a fixed point on Y it has a fixed point on Z . This implies (e) for cases O, OE and DE. In case S we argue as in the proof of Lemma 6.4, and conclude that σ has zero or two fixed points in Y above each fixed point in \hat{Z} . We deduce the claim from the Riemann-Hurwitz formula applied to $Y \rightarrow Y/H$. \square

COROLLARY 6.6. *The genus of $X = Y/H$ is*

$$g(X) = \begin{cases} (mn - m - n - \gamma)/2 + 1 & \text{in cases O and OE,} \\ (mn - m - n - \gamma)/4 + 1 & \text{in case DE,} \\ (mn - m - n - 2\gamma)/4 + 1 & \text{in case S.} \end{cases}$$

Notation 6.7. Until now we have been working on the geometric generic fiber of $g : \mathcal{Y} \rightarrow \mathbb{P}^*$, etc. Let $\pi : C \rightarrow \mathbb{P}^*$ be the unramified cover obtained by adjoining the elements c and d defined in (8) to $\mathbb{C}(t)$. Then $H = \langle \sigma, \tau \rangle$ is a subgroup of $\text{Aut}(\mathcal{Y}_C)$. Passing to a further unramified cover, if necessary, we may suppose that the VHS of the pullback family $h_C : \mathcal{Y}_C \rightarrow C$ is unipotent. We write $\bar{\pi} : \bar{C} \rightarrow \mathbb{P}_t^1$ for the corresponding (branched) cover of complete curves. Then h_C extends to a family $h_C : \bar{\mathcal{Y}}_C \rightarrow \bar{C}$ of stable curves over this base curve.

The following lemma describes the action of H on the degenerate fibers of h_C .

LEMMA 6.8. *Let $c \in \bar{C}$ be a point with $\pi(c) \in \{0, 1\}$. The quotient $\mathcal{X}_c := (\mathcal{Y}_C)_c/H$ is smooth and*

$$g(\mathcal{X}_c) = \begin{cases} (mn - m - n - \gamma)/2 + 1 & \text{cases O and OE,} \\ (mn - m - n - \gamma)/4 + 1 & \text{case DE,} \\ (mn - m - n - 2\gamma)/4 + 1 & \text{case S.} \end{cases}$$

Proof. Choose $c \in \pi^{-1}(0)$. The case that $c \in \pi^{-1}(1)$ is similar, and left to the reader. By Proposition 4.5 the fiber $(\mathcal{X}_C)_c$ consists of two irreducible components which we call Z_0^1 and Z_0^2 ; we make the convention that the fixed points $x = 0, t$ of φ_1 on \mathcal{X}_C specialize to Z_0^1 . Choosing suitable coordinates, the curve Z_0^1 is given by

$$(14) \quad z_0^N = x_0^{a_1}(x_0 - 1)^{a_3}.$$

The components Z_0^1 and Z_0^2 intersect in $2m/\gamma$ points (resp. m/γ in case S). We write P_0^j for the quotient of Z_0^j by $\langle \varphi_1 \rangle \cong \mathbb{Z}/N$.

We claim that the fiber $(\mathcal{Y}_C)_c$ consists of 2 irreducible components Y_0^1, Y_0^2 , as well. Let \mathcal{N} be the normalization of the fiber product $(\mathcal{X}_C)_c \times_{(\mathcal{X}_C)_c} (\mathcal{X}_C)_c^\tau$. By Abhyankar's lemma, $\mathcal{N} \rightarrow (\mathcal{X}_C)_c$ is étale at the preimages of the intersection point of the two components of $(\mathcal{P}_C)_c$. Hence \mathcal{N} consists of two curves: the fiber products over $Z_0^j/\langle \varphi_1^{\hat{N}} \rangle$ of Z_0^j with its τ -conjugate, for $j = 1, 2$. These two curves intersect transversally in $2m\beta/\gamma$ (resp. $m\beta/\gamma$ in case S) points. This implies that \mathcal{N} is a stable curve and indeed the fiber $(\mathcal{Y}_C)_c$.

One computes that $g(Y_0^j) = 1 + mn\beta/\gamma - m\beta/\gamma - \beta$ in cases O, OE, and DE, and $g(Y_0^j) = mn - m/2 + 1 - \gamma$ in case S. Since ρ acts on the points $\{0, 1, t, \infty\}$ as the permutation $(0t)(1\infty)$ we conclude that ρ fixes the components Y_0^j while σ and τ interchange them. Clearly, for a coordinate x_0 as in (14) we have that $\rho(x_0) = 1 - x_0$, i.e. ρ fixes the points $1/2$. This is a specialization of one of the

two fixed points $1 \pm \sqrt{1-t} \in P$. Since, by Lemma 6.5, the automorphism ρ fixes $2n$ (n in case S) points in Y above each of these points of P it follows that ρ fixes $2n$ (resp. n) points of Y_0^j with $x_0 = 1/2$. It remains to compute the number r_∞ of fixed points of ρ over $x_0 = \infty$.

Suppose we are not in case S. Then by the Riemann-Hurwitz formula

$$g(\mathcal{X}_c) = g(Y_0^j / \langle \rho \rangle) = (mn - m - n - \gamma)\beta/2\gamma + 1 - r_\infty/4.$$

Applying the Riemann-Hurwitz formula to the quotient map $Z_0^j \rightarrow Z_0^j / \langle \rho \rangle$, we conclude that $r_\infty \equiv 0 \pmod{4}$. Represent the fiber of Z_0^j over ∞ as $\varphi_1^b R$, for $b = 1, \dots, 2m/\gamma$. As in the proof of Lemma 6.4, we conclude that r_∞ equals zero or two. It follows that $r_\infty = 0$.

In case S we have

$$g(\mathcal{X}_c) = (mn - m - n - 2\gamma)/4 + 1 - r_\infty/4.$$

and we conclude as above that $r_\infty = 0$. Genus comparison shows that the fiber $(\mathcal{X}_C)_c$ is smooth. \square

Proof. We now prove Theorem 6.2. We have shown in Lemma 6.8 that $\bar{\mathcal{X}}_c$ is smooth for $c \notin S_u = \pi^{-1}(\infty)$. We have to show that the VHS of $f : \mathcal{X} \rightarrow C$ contains a local subsystem of rank 2 which is maximal Higgs.

We decompose the VHS of g into the characters

$$\chi(i, j) : \begin{cases} G_0 & \rightarrow \mathbb{C} \\ \psi_1 & \mapsto \zeta_N^i \\ \psi_2 & \mapsto (\zeta_N^{\hat{N}})^j. \end{cases}$$

We let $\mathbb{L}(i, j) \subset R^1 h_* \mathbb{C}$ be the local system on which G acts via $\chi(i, j)$. Local systems with $j = 0$ arise as pullbacks from \mathcal{X} . By Lemma 4.1 the local systems $\mathbb{L}(i, 0)$ are of rank 2 if i does not divide N . Using the presentation of G one checks that $\sigma^* \mathbb{L}(i, j) = \mathbb{L}(-i, -j)$ and $\tau^* \mathbb{L}(i, j) = \mathbb{L}(-\alpha i, \alpha(i - j))$.

The local exponents of $(\mathbb{L}(1, 0), \omega_1)$ at $t = 0$ (resp. $t = 1$) are $(0, 1/n)$ (resp. $(0, 1/m)$). Therefore, the definition of $\bar{\pi} : \bar{C} \rightarrow \mathbb{P}_t^1$ (Notation 6.7) implies that condition (5) is satisfied for $\mathbb{L}(1, 0)$.

Consider the local system

$$\mathbb{M} := \mathbb{L}(1, 0) \oplus \mathbb{L}(-1, 0) \oplus \mathbb{L}(-\alpha, \alpha) \oplus \mathbb{L}(\alpha, -\alpha)$$

on \mathcal{X}_C . Since H permutes the four factors of \mathbb{M} transitively, we conclude that for each character ξ of H there is a rank 2 local subsystem of \mathbb{M} on which H acts via ξ . Moreover the projection of the subsystem $\mathbb{L} := \mathbb{M}^H$ to each summand is nontrivial. Since the four summands of \mathbb{M} are irreducible by construction, this implies that

$$\mathbb{L} \cong \mathbb{L}(1, 0) \cong \mathbb{L}(-1, 0) \cong \mathbb{L}(-\alpha, \alpha) \cong \mathbb{L}(\alpha, -\alpha).$$

Hence \mathbb{L} descends to \mathcal{X} and is maximal Higgs with respect to S_u . Proposition 4.2 implies that the extension of f to $\pi^{-1}(\mathbb{P}^1 \setminus \{\infty\})$ is the pullback of the universal family of curves to an unramified cover of a Teichmüller curve. \square

The proof of Theorem 6.2 contains more information on the VHS of f and on the Lyapunov exponents $\lambda(\mathbb{L}_i)$. We work out the details in the most transparent case that m and n are odd integers which are relatively prime. The interested reader can easily work out the Lyapunov exponents in the remaining cases, too. In this case the curves \mathcal{X} and \mathcal{Y} coincide (Remark 6.3) and the local system $\mathbb{L}(i, j)$ is $\mathbb{L}(i)$ in the notation of Lemma 4.1.

We deduce from the arguments of the proof of Theorem 6.2 that, for each i not divisible by m or n , there is an H -invariant local system \mathbb{L}_i with

$$\mathbb{L}_i \cong \mathbb{L}(i) \cong \mathbb{L}(\alpha i) \cong \mathbb{L}(-\alpha i) \cong \mathbb{L}(-i).$$

Since those i fall into $(m-1)(n-1)/2$ orbits under $\langle \pm 1, \pm \alpha \rangle$, we have the complete description of the VHS of h . Let $c_j(i) = \sigma_j(i) + \sigma_3(i) - 1$.

COROLLARY 6.9. *Let m and n be odd integers which are relatively prime.*

(a) *The VHS of f splits as*

$$R^1 f_* \mathbb{C} \cong \bigoplus_{j \in J} \mathbb{L}(j),$$

where $\mathbb{L}(j)$ is an irreducible rank 2 local system and j runs through a set of representatives of

$$J = \{0 < i < N, m \nmid i, n \nmid i\} / \sim, \quad \text{where } i \sim -i \sim \alpha i \sim -\alpha i.$$

(b) *The Lyapunov exponents are*

$$\lambda(\mathbb{L}(i)) = \frac{mn - e_1(i)m - e_2(i)m}{mn - m - n},$$

where $e_1(i) = n|c_1(i)|$ and $e_2(i) = m|c_2(i)|$.

Proof. This follows directly from Proposition 4.4. \square

Example 6.10. We calculate the Lyapunov exponents explicitly for $m = 3$ and $n = 5$. Then $N = 2nm = 30$ and hence $\alpha = 19$. We need to calculate the $\lambda(\mathbb{L}(i))$ only up to the relation ‘ \sim ’ and hence expect at most four different values. One checks:

$$\lambda(\mathbb{L}(i)) = \begin{cases} 7/7 & \text{if } i \sim 1, \\ 4/7 & \text{if } i \sim 2, \\ 2/7 & \text{if } i \sim 4, \\ 1/7 & \text{if } i \sim 7. \end{cases}$$

In particular, we see that in general the $\lambda(\mathbb{L}(i))$ do not form an arithmetic progression as one might have guessed from studying Veech’s n -gons.

Remark 6.11. Note that $K := \mathbb{Q}(\cos(\pi/n), \cos(\pi/m))$ is the trace field of the $\Delta(m, n, \infty)$ -triangle group. Hence $r = [K : \mathbb{Q}] \leq \phi(mn)/4 \leq (m-1)(n-1)/4$. Here again the decomposition of the VHS is finer than predicted by Theorem 2.1; cf. the remark after Corollary 5.3.

Let X be any fiber of f . We denote by $\omega_X \in \Gamma(X, \Omega_X^1)$ a generating differential, i.e., a holomorphic differential that generates $(1, 0)$ -part of the maximal Higgs local system when restricted to the fiber X . This condition determines ω_X uniquely up to scalar multiples.

THEOREM 6.12. *Suppose $m \neq n$. Then the projective affine group of the translation surface (X, ω_X)*

- (a) *is the (m, n, ∞) -triangle group if at least one of m and n is odd (i.e. in the cases O and OE);*
- (b) *contains a subgroup of index 2 in the (m, n, ∞) -triangle group in the case DE and S.*
- (c) *If $m = n$ is odd, then projective affine group of the translation surface (X, ω_X) equals the (m, m, ∞) -triangle group or equals a triangle group that contains the (m, m, ∞) -triangle group as a subgroup of index 2.*

Hooper has shown in [13] that this result is close to optimal concerning the realization of triangle groups: he shows that the triangle groups $\Delta(m, n, \infty)$ in case S and in case DE with $v = 1$ cannot occur as projective affine groups.

Proof. We first determine when the projective affine group of (X, ω_X) contains the triangle group $\Delta(m, n, \infty)$. As in the proof of Proposition 5.7, we take two fibers \mathcal{Y}_p and $\mathcal{Y}_{\tilde{p}}$ with $\pi(p) = \pi(\tilde{p})$. We need to show the existence of an isomorphism $\mathcal{Y}_p \rightarrow \mathcal{Y}_{\tilde{p}}$ which is equivariant with respect to H . By construction of σ and τ , it suffices to find $i_0 : \mathcal{X}_p \rightarrow \mathcal{X}_{\tilde{p}}$ which is equivariant with respect to σ and φ_1 , and such that the induced isomorphism $\hat{i}_0 : \hat{\mathcal{X}}_p \rightarrow \hat{\mathcal{X}}_{\tilde{p}}$ is equivariant with respect to τ .

Since $\pi(p) = \pi(\tilde{p})$, the map $(x, z) \mapsto (x, z)$ defines an isomorphism $i : \mathcal{X}_p \rightarrow \mathcal{X}_{\tilde{p}}$. We try $i_0 := \varphi_1^j \circ i$, for a suitably chosen j . Then i_0 is automatically φ_1 -equivariant. Let π_1 (resp. π_2) denote the maps from C to the intermediate cover given by $s_1^n = t$ (resp. $s_2^m = (t-1)$). By hypothesis we have $\pi_1(p) = \zeta_n^{e_1} \pi_1(\tilde{p})$ and $\pi_2(p) = \zeta_m^{e_2} \pi_2(\tilde{p})$ for some integers e_1, e_2 , where $\zeta_m := \zeta_N^{N/m}$ (resp. $\zeta_n := \zeta_N^{N/n}$) is a primitive m -th (resp. n -th) root of unity. By the definition of e_1 and e_2 we have

$$i(s_1) = \zeta_n^{e_1} s_1 = \zeta_N^{e_1 N/n} \quad \text{and} \quad i(s_2) = \zeta_m^{e_2} s_2 = \zeta_N^{e_2 N/m} s_2.$$

We conclude that

$$\tau \circ \hat{i} = \varphi_1^k \circ \hat{i} \circ \tau, \quad \text{and} \quad \sigma \circ i = \varphi_1^{e_1 N/n + e_2 N/m} \circ i \circ \sigma,$$

with

$$k \equiv \begin{cases} e_1 2m/\gamma \bmod \hat{N} & \text{in cases O and OE,} \\ e_1 m/\gamma \bmod \hat{N}/2 & \text{in case DE,} \\ e_1 m/\gamma \bmod \hat{N} & \text{in case S.} \end{cases}$$

Recall that $\varphi_1(\bar{z}) = \zeta_N^{N/\hat{N}} \bar{z}$.

The equivariance properties for $i_0 = \varphi_1^j \circ i$ together with the conjugation formulas in (10) impose the two conditions on j :

$$j(\alpha - 1) + k \equiv 0 \bmod \hat{N} \quad \text{and} \quad -2j + e_1 N/n + e_2 N/m \equiv 0 \bmod N.$$

Recall that $\hat{N} = 2^\delta m' n' / (\gamma')^2$ with $\gcd(m'/\gamma', n'/\gamma') = 1$. We start by solving these congruences modulo m'/γ' and n'/γ' . Modulo m'/γ' the first condition is trivial and the second condition is solvable. Modulo n'/γ' both conditions are identical and solvable.

It remains to solve the conditions modulo 2^δ . In case O and OE the first condition is trivial modulo 2^δ and the second condition is solvable modulo 2^δ . In case DE the first condition is solvable if and only if e_1 is even. The second condition is solvable. In case S the first condition is trivial, since \hat{N} is odd. The second condition becomes

$$-2j + e_1 m/\gamma + e_2 n/\gamma \equiv 0 \bmod 2^\nu.$$

This condition is solvable if and only if $e_1 + e_2$ is even, since m/γ and n/γ are both odd.

So far we have shown that in the cases O and OE the projective affine group contains $\Delta(m, n, \infty)$ and in the cases S and DE it contains an index 2 subgroup.

To see that the projective affine group is not larger than $\Delta(m, n, \infty)$ in the cases O and OE, we note that a larger projective affine group is again a triangle group. Singerman ([33]) shows that any inclusion of triangle groups is a composition of inclusions in a finite list. Only the cases where $m = n$ fit into his list. This proves (a) and the second part of the theorem. \square

We determine the basic geometric invariant of the Teichmüller curves constructed in Theorem 6.2.

THEOREM 6.13. *In cases S and DE the generating differential ω_X has $\gamma/2$ zeros and in the cases O and OE the generating differential ω_X has γ zeros.*

Proof. We only treat the cases O and OE. The cases S and DE are similar. We calculate the zeros of the pullback ω_Y of ω_X to the corresponding fiber Y of \mathcal{Y} . The differential ω_i on Z has zeros of order $a_1(i)\gamma/\gamma_1 - 1$ (resp. $a_2(i)\gamma/\gamma_1 - 1$) at the γ_1/γ points over 0 (resp. 1). It has zeros of order $a_3(i)\gamma/\gamma_2 - 1$ (resp. $a_4(i)\gamma/\gamma_2 - 1$) at the γ_2/γ points over t (resp. ∞). Therefore, the pullback of ω_i to Y has zeros of order $a_\mu(i) - 1$ at the γ preimages of $\mu = 0, 1, t, \infty$. The differential ω_Y is a linear combination with nonzero coefficients of ω_1 , ω_{-1} , and two differentials

that are pulled back from Z^τ . The vanishing orders of these differentials on Z^τ are obtained from those of ω_1 and ω_{-1} on Z by replacing a_1 by a_4 , a_2 by a_3 , and conversely. Since the a_μ are pairwise distinct, we conclude that ω_Y vanishes at the (in total) 4γ preimages of $\{0, 1, t, \infty\}$ of order $\min\{a_1, a_2, a_3, a_4\} - 1 = a_4 - 1$. Since ω_Y vanishes also at the $4m + 4n$ ramification points of $Y \rightarrow X$ we deduce that it vanishes there to first order and nowhere else. The 4γ zeros at the nonramification points yield the γ zeros of ω_X . \square

6.1. Comparison with Ward's results. In this section we compute an explicit equation for one particular fiber of the family $\bar{f} = \bar{f}(m, n) : \bar{\mathcal{X}} \rightarrow \bar{C}$. This fiber, \mathcal{X}_c , is chosen such that \mathcal{X}_c is a cyclic cover of a projective line. This result is used in Section 8 to realize \mathcal{X}_c via unfolding a billiard table, for small m . In this section we show moreover that for $m = 3$, the family $\bar{f} : \bar{\mathcal{X}} \rightarrow \bar{C}$ coincides with the family of curves constructed by Ward [38].

The assumptions on m and n in the following theorem are not necessary. We include them to avoid case distinctions. The reader can easily work out the corresponding statement in the general situation, as well. We use the same notation as in the rest of this section. In particular, $\bar{\pi} : \bar{C} \rightarrow \mathbb{P}_t^1$ denotes the natural projection of \bar{C} to the t -line defined in Notation 6.7. One may of course interchange the role of m and n in the theorem. In that case one should consider the fiber of \mathcal{X} in a point of \bar{C} above $t = 1$, instead.

THEOREM 6.14. *Suppose that m and n are relatively prime and n is odd. Then a fiber of $\bar{f} : \bar{\mathcal{X}} \rightarrow \bar{C}$ over a point of $\bar{\pi}^{-1}(0)$ is a $2n$ -cyclic cover of the projective line branched at $(m + 3)/2$ points if m is odd and $(m + 4)/2$ points if m is even.*

(a) *For m odd this cover is given by the equation*

$$X_0 : \quad y^{2n} = (u - 2) \prod_{k=1}^{(m-1)/2} \left(u - 2 \cos \left(\frac{2k\pi}{m} \right) \right)^2.$$

The generating differential of the Teichmüller curve is

$$\omega_0 = \frac{y \, du}{(u - 2) \prod_{k=1}^{(m-1)/2} (u - 2 \cos(2k\pi/m))}.$$

(b) *For m even this cover is given by the equation*

$$X_0 : \quad y^{2n} = (u - 2)^n \prod_{k=1}^{m/2} \left(u - 2 \cos \left(\frac{(2k-1)\pi}{m} \right) \right)^2.$$

The generating differential of the Teichmüller curve is

$$\omega_0 = \frac{y \, du}{(u - 2) \prod_{k=1}^{m/2} (u - 2 \cos((2k-1)\pi/m))}.$$

(c) *For $m = 3$ the surface (X_0, ω_0) is the translation surface found by Ward.*

Proof. Our simplifying assumptions imply that $\gamma = 1$ and $\mathcal{X} \cong \mathcal{Y}$ (Remark 6.3). Let c be a point of \bar{C} with $\bar{\pi}(c) = 0$. Then the fiber \mathcal{Y}_c of \mathcal{Y} consists of two isomorphic irreducible components, Y_0^j , given by the affine equation $y^N = x^{a_1}(x-1)^{a_3}$. Note that $Y_0^1 \rightarrow \mathbb{P}_x^1$ is branched at $x = \infty$ of order m . The fiber $X_0 := \mathcal{X}_c$ of \mathcal{X} is the quotient of Y_0^1 by ρ .

From the presentation of G (Lemma 6.5) we deduce that φ_1^k commutes with ρ if and only if k is a multiple of m . We denote by A the abelian subgroup of $\text{Aut}(Y_0^1)$ generated by ρ and φ_1^m . One computes that the quotient of Y_0^1 by $\langle \varphi_1^m \rangle$ has genus zero. We denote this quotient by \mathbb{P}_z^1 . Here z is a parameter on \mathbb{P}_z^1 such that $\mathbb{P}_z^1 \rightarrow \mathbb{P}_x^1$ is given by

$$z^m = \left(\frac{x-1}{x} \right)^n.$$

Let \mathbb{P}_u^1 be the quotient of Y_0^1 by A . The subscript u denotes a coordinate which is defined below. We obtain the following diagram of covers

$$\begin{array}{ccccc} & Y_0^1 & & & \\ \swarrow \langle \rho \rangle & & \searrow \langle \varphi_1^m \rangle & & \\ X_0 & & \mathbb{P}_z^1 & & \\ \searrow q_1 & & \swarrow q_2 & & \searrow \langle \varphi_1 \bmod m \rangle \\ & \mathbb{P}_u^1 & & & \mathbb{P}_x^1 \end{array}$$

Suppose that m is odd. After replacing y by $z^{(n+1)/2}/y$, we find that $Y_0^1 \rightarrow \mathbb{P}_z^1$ is given by

$$(15) \quad y^{2n} = \frac{(z^m - 1)^2}{z^m}.$$

Here we use that n is odd. Recall that $\rho \in \text{Aut}(\mathbb{P}_x^1)$ is given by $\rho(x) = 1 - x$. It follows from Lemma 6.4 that ρ lifts to an automorphism of order 2 of \mathbb{P}_z^1 which has one fixed point in the fiber above $x = 1/2$. Without loss of generality, we may assume that $\rho(z) = 1/z$. Therefore, $u := z + 1/z$ is an invariant of ρ and it is a parameter on \mathbb{P}_u^1 . We find an equation for X_0 by rewriting (15) in terms of y and u . Noting that

$$u - (\zeta_m^i + \zeta_m^{-i}) = \frac{(z - \zeta_m^i)(z - \zeta_m^{-i})}{z},$$

we find the equation in (a). The differential form ω_0 in (a) is a holomorphic differential form with a zero only in $u = 0$. Therefore Theorem 6.13 implies that ω_0 is a generating differential form.

Specializing to $m = 3$, we find the equation found by Ward ([38, §5]). This proves (c).

Suppose now that m is even. After replacing y by $z^{m/2}/y$, we find that $Y_0^1 \rightarrow \mathbb{P}_x^1$ is given by

$$y^{2n} = \frac{(z^m - 1)^2}{z^{m+n}}.$$

In this case the automorphism ρ of \mathbb{P}_x^1 lifts to an automorphism of \mathbb{P}_z^1 with two fixed points in the fiber above $x = 1/2$. Without loss of generality, we may suppose that $\rho(z) = \zeta_m/z$. Therefore, $u := \zeta_{2m}^{-1}z + \zeta_{2m}/z$ is an invariant of ρ which we regard as parameter on \mathbb{P}_u^1 . Here ζ_{2m} is a square root of ζ_m . One computes that

$$u - (\zeta_{2m}^{2i-1} + \zeta_{2m}^{-2i+1}) = \zeta_{2m}^{-1} \frac{(z - \zeta_m^i)(z - \zeta_m^{1-i})}{z}, \text{ and } u - 2 = \zeta_{2m}^{-1} \frac{(z - \zeta_{2m})^2}{z}.$$

After replacing y by $c(z - \zeta_{2m})/y$ for a suitable root of unity c , we find the equation for X_0 which is stated in (b). The expression for ω_0 follows as in the proof of (a). \square

7. Primitivity

A *translation covering* is a covering $q : X \rightarrow Y$ between translation surfaces (X, ω_X) and (Y, ω_Y) such that $\omega_X = q^*\omega_Y$. A translation surface (X, ω_X) is called *geometrically primitive* if it does not admit a translation covering to a surface Y with $g(Y) < g(X)$.

A Veech surface (X, ω) is called *algebraically primitive* if the degree r of the trace field extension over \mathbb{Q} equals $g(X)$. Algebraic primitivity implies geometric primitivity, but the converse does not hold ([28]). In [loc. cit., Th. 2.6] it is shown that a translation surface of genus greater than one covers a unique primitive translation surface.

Obviously the Veech examples $(p : \mathcal{U} \rightarrow \tilde{C}$ in the notation of Theorem 5.5) for $n = 2\ell$ and ℓ prime and those for $(2, n, \infty)$ (cf. Remark 5.9) are algebraically primitive. We will not give a complete case by case discussion of primitivity of the (m, n, ∞) -Teichmüller curves, but restrict to the case that m and n are odd and relatively prime. Comparing the degree of the trace field $[\mathbb{Q}(\zeta_m + \zeta_m^{-1}, \zeta_n + \zeta_n^{-1}) : \mathbb{Q}] = r \leq \phi(m)\phi(n)/4$ with the genus (Corollary 6.6), we deduce that the fibers of $\mathcal{X} \rightarrow C$ are never algebraically primitive. Nevertheless, we show that there are infinitely many geometrically primitive ones.

THEOREM 7.1. *Let m and n be distinct odd primes. Then the Veech surfaces arising from the (m, n, ∞) -Teichmüller curve $f : \mathcal{X} \rightarrow C$ of Theorem 6.2 are geometrically primitive.*

Proof. Let (X, ω_X) be such a Veech surface and suppose there is a translation covering $q : X \rightarrow Y$. Then $g(Y) \geq r$, by [28, Th. 2.6]. Theorem 6.13 implies that the generating differential has only one zero on \mathcal{X}_c . Therefore the cover q is totally ramified at this zero, and nowhere else. This contradicts the Riemann-Hurwitz formula. Namely, a degree two cover cannot be branched in exactly one point. If the degree d of q is larger than 2, we obtain a contradiction with $g(Y) \geq r$. \square

Remark 7.2. At the time of writing the authors are aware of the following series of examples of Teichmüller curves: the triangle constructions in [35] and [38] and the Weierstrass eigenform or Prym eigenform constructions in [24] and [25]. Besides them there is a finite number of sporadic examples.

COROLLARY 7.3. *The Veech surfaces arising from the case (m, n, ∞) with m and n sufficiently large distinct primes are not translation covered by any of the Veech surfaces listed in Remark 7.2.*

Proof. Recall that translation coverings between Veech surfaces preserve the affine group up to commensurability. In particular, they preserve the trace field.

Choose m and n sufficiently large such that the trace field K of the (m, n, ∞) -triangle group is none of the trace fields occurring in the sporadic examples and such that the genus of the Veech surface is larger than 5. This implies that the surface cannot be one of examples in [24] and [25]. There is only a finite list of arithmetic triangle groups ([34]). We choose $m > 3$ and $n > 5$ such that K is not one of the trace fields in this finite list. Nonarithmetic lattices have a unique maximal element ([21]) in its commensurability class and the (m, n, ∞) -triangle groups are the maximal elements in their classes. Since the $(2, n, \infty)$ - and $(3, n, \infty)$ -triangle groups are the maximal elements in the commensurability classes of the examples of [35] and [38], these examples cannot be a translation cover of the examples given by Theorem 6.2 for (m, n) chosen as above. \square

Remark 7.4. Even in the cases that the Veech surfaces with affine group $\Delta(m, n, \infty)$ are geometrically primitive, Theorem 2.6 of [28] does not exclude that there are other primitive Veech surfaces with the same affine group. Such examples are provided by Theorem 3' of [14] for $n = \infty$. By Remark 3.3 we know a rank $2r$ subvariation of Hodge structures of the family of curves generated by such a Veech surface. In particular, we know r of the Lyapunov exponents $\lambda(\mathbb{L}_i)$.

8. Billiards

In this section we approach Teichmüller curves uniformized by triangle groups in the way Veech and Ward did in [35] and [38]. We start by presenting two series of billiard tables $T(m, n, \infty)$, for $m = 4, 5$. These tables are (rational) 4-gons in the complex plane. We show that the affine group of the translation surface $X(m, n, \infty)$ attached to $T(m, n, \infty)$ is the (m, n, ∞) -triangle group, for $m = 4, 5$. This part is independent of the previous sections, and requires only elementary notions of translation surfaces (see [23] or §2). The proof we give that these billiard tables define Teichmüller curves is combinatorially more complicated than the analogous proof for the series of Teichmüller curves found by Veech and Ward. This suggests that it would have been difficult to find these billiards by a systematic search among 4-gons.

In Section 8.3 we relate these explicitly constructed billiard tables to our main realization result (Theorem 6.2). Denote by $f = f(m, n) : \mathcal{X} \rightarrow \mathcal{C}$ the family

of curves constructed in Section 6. This family defines a finite map from C to M_g , for a suitable integer $g \geq 2$. The image of this map is a Teichmüller curve whose (projective) affine group is the (m, n, ∞) -triangle group. We have shown in Theorem 6.14 that a suitable fiber \mathcal{X}_c of \mathcal{X} is a $2n$ -cyclic cover of the projective line which we described explicitly. In this situation, one may use a result of Ward to find the corresponding billiard table $T[m, n, \infty]$. We show that $T[m, n, \infty]$ may be embedded in the complex plane (i.e., without self-crossings) if and only if $m \leq 5$. For $m = 2, 3$ we find back the billiard tables found by Veech and Ward. We show that the tables we obtain for $m = 4, 5$ are the ones we have already constructed.

Consider a compact polygon $P \subset \mathbb{R}^2 \cong \mathbb{C}$ in the plane whose interior angles are rational multiples of π . The linear parts of reflections along the sides of the polygon generate a finite subgroup $G \subset O_2(\mathbb{R})$. If s is a side of P we write σ_s for the linear part of the reflection in the side s . One checks that for sides s and t of P we have $\sigma_{\sigma_s(t)} = \sigma_s \sigma_t \sigma_s$.

We define an equivalence relation on G as follows. We write $\sigma_1 \sim \sigma_2$ if the reflected polygons $\sigma_1(P)$ and $\sigma_2(P)$ differ by a translation in \mathbb{C} . Let $G_0 \subset G$ represent the equivalence classes of this relation. By gluing copies of P we obtain a compact Riemann surface

$$X = \left(\coprod_{g \in G_0} gP \right) / \approx,$$

where \approx denotes the following identification of edges: if gP is obtained from $\tilde{g}P$ by a reflection σ along a side s of $\tilde{g}P$, then s is glued to the edge $\sigma(s)$ of gP by a translation.

The holomorphic 1-form dz on P and its copies are translation invariant, hence define a 1-form ω on X . We say that the translation surface (X, ω) is *obtained by unfolding* P . The trajectories of a billiard ball on P correspond to straight lines on X . In [24] X is called the *small surface* attached to P . The translation surface has a finite number of points where the total angle exceeds 2π . These are called *singular points*. They correspond to the zeros of ω .

8.1. *The tables $T(5, n, \infty)$.* Let $n \geq 7$ be an odd integer which is not divisible by 5. We define a billiard table $T(5, n, \infty)$ as follows (Figure 1). The billiard table $T(5, n, \infty)$ is a 4-gon in the complex plane with angles $\alpha = \beta = \pi/n$ and $\gamma = \pi/2n$, as indicated in the picture. We denote by I_1, \dots, I_4 the vectors corresponding to the sides of the polygon which we regard as complex numbers. We rotate and scale the billiard table such that $I_4 = 1$ and

$$|\operatorname{Re}(I_3)| = \cos(\pi/n) + \cos(\pi/5).$$

In particular, I_4 points in the direction of the positive x -axis. This determines the table uniquely.




Figure 1. Billiard table $T(5, n, \infty)$, for $n = 9$

We now construct the translation surface obtained by unfolding the table $T(5, n, \infty)$ (Figure 2). Reflecting the billiard table $2n$ times in the (images of the) sides I_2 and I_3 yields the upper star; it consists of alternating long and short points. The second star is obtained by reflecting the first star in the side I_4 of the billiard table (this is the side marked by 15 in Figure 2). The two stars can be glued together to a translation surface $X := X(5, n, \infty)$: sides denoted by the same letters or numbers are glued by translations. Note that the tips of the ‘long points’ (resp. the ‘short points’) of the stars correspond to one point of the translation surface X ; both points of X are not singularities, since the total angle is 2π . There is one singularity; it corresponds to the angle δ . The genus of X is $g = 2(n - 1)$.

THEOREM 8.1. *Let $n \geq 7$ be odd and not divisible by 5. Then the affine group of $X(5, n, \infty)$ contains the elements*

$$R = \begin{pmatrix} \cos(\pi/n) & -\sin(\pi/n) \\ \sin(\pi/n) & \cos(\pi/n) \end{pmatrix} \quad \text{and} \quad T = \begin{pmatrix} 1 & 2 \frac{\cos(\pi/n) + \cos(\pi/5)}{\sin(\pi/n)} \\ 0 & 1 \end{pmatrix}.$$

The elements $R, T \in \text{PSL}_2(\mathbb{R})$ generate the Fuchsian triangle group $\Delta(5, n, \infty)$. In particular, $X(5, n, \infty)$ is a Veech surface.

Proof. Rotation around the center of the stars defines an affine diffeomorphism of the surface $X(5, n, \infty)$. Its derivative is R .

We rotate $X(5, n, \infty)$ as in Figure 1 and Figure 2, i.e., such that the edge I_4 resp. the one with label 15 is horizontal and to the left of the center of the star.

We now consider the horizontal foliation defined by ω . Recall that a *saddle connection* is a leaf of the foliation that begins and ends in a singularity. In a dense set of directions, the saddle connections divide X into *metric cylinders*, see for example [23, §4.1]. We claim that, in the horizontal direction, X decomposes into $g = 2(n - 1)$ metric cylinders. We distinguish two types of cylinders. Each cylinder corresponds to one shading style in Figure 2.

The cylinders of type 1, denoted by C_i , are those that are glued together from pieces from both stars. An example is the checkered cylinder. Since the second star is obtained from the first by reflection, the cylinders C_i appear in pairs, as can




Figure 2. Cylinder decomposition of $X(5, 9, \infty)$

be seen from Figure 2. There is a bijection between the cylinders of type 1 and pairs of long points. For example, the checkered cylinder corresponds to the long points 17-18 and 5-6. Here a ‘pair’ consists of an orbit of length two of long points under the reflection in the vertical axis. The two vertical long points correspond

to orbits of length one, and hence do not correspond to a cylinder of type 1. We conclude that the number of cylinders of type 1 is $n - 1$.

The cylinders of type 2, denoted by \tilde{C}_i , are those consisting of pieces of one star. An example is the black cylinder. These cylinders also come in pairs. There is a bijection between cylinders of type 2 and pairs of short points. Therefore the number of cylinders of type 2 is also $n - 1$.

The width and the height of a pair of cylinders of type 1, for an appropriate numbering, is given by

$$(16) \quad \begin{aligned} w_k &= 2|I_3| \cos \frac{(n-2k)\pi}{2n} \quad \text{and} \\ h_k &= |I_4| \left(\sin \frac{(n+1-2k)\pi}{2n} - \sin \frac{(n-1-2k)\pi}{2n} \right) \\ &= 2|I_4| \sin \frac{\pi}{2n} \cos \frac{(n-2k)\pi}{2n} \end{aligned}$$

for $k = 1, \dots, (n-1)/2$. This is seen by cutting the points of the stars into pieces, and translating these pieces so that one obtains $2(n-1)$ connected cylinders, one for each shading style. One then uses the rotation and reflection symmetries of the original star.

Similarly, the widths and heights of pairs of cylinders \tilde{C}_i , for an appropriate numbering, are given by

$$(17) \quad \begin{aligned} \tilde{w}_k &= 2|I_2| \cos \frac{(n-2k)\pi}{2n} \quad \text{and} \\ \tilde{h}_k &= |I_1| \left(\sin \frac{(n+2-2k)\pi}{2n} - \sin \frac{(n-2-2k)\pi}{2n} \right) \\ &= 2|I_1| \sin \frac{2\pi}{2n} \cos \frac{(n-2k)\pi}{2n} \end{aligned}$$

for $k = 1, \dots, (n-1)/2$.

The moduli of these cylinders are

$$m_k := h_k/w_k = |I_4| \sin \frac{\pi}{2n} / |I_3| \quad \text{and} \quad \tilde{m}_k := \tilde{h}_k/\tilde{w}_k = |I_1| \sin \frac{2\pi}{2n} / |I_2|.$$

Note that m_k and \tilde{m}_k are independent of k .

We claim that $m_k/\tilde{m}_k = |I_4||I_2| \sin(\pi/n)/|I_3||I_1| \sin(\pi/2n) = 1$, that is that the moduli of all the cylinders are identical. This is equivalent to

$$(18) \quad \frac{|I_2|}{|I_1|} = \frac{|I_3|}{|I_4|} \frac{\sin(\pi/n)}{\sin(\pi/2n)}.$$

Since we assumed that $I_4 = 1$, the right-hand side is equal to $2|\operatorname{Re}(I_3)|$.

Using the geometry of the billiard table one shows that

$$\begin{aligned} |I_2| \cos(3\pi/2n) - |I_1| \cos(5\pi/2n) &= |\operatorname{Re}(I_3)| - |I_4| = |\operatorname{Re}(I_3)| - 1, \\ |I_2| \sin(3\pi/2n) - |I_1| \sin(5\pi/2n) &= \operatorname{Im}(I_3) = \operatorname{Re}(I_3) \tan(\pi/2n). \end{aligned}$$

This implies that

$$(19) \quad \frac{|I_2|}{|I_1|} = \frac{-(\operatorname{Re}(I_3) - 1) \sin(5\pi/n) + \operatorname{Re}(I_3) \tan(\pi/2n) \cos(5\pi/n)}{-(\operatorname{Re}(I_3) - 1) \sin(3\pi/n) + \operatorname{Re}(I_3) \tan(\pi/2n) \cos(3\pi/n)}.$$

The minimal polynomial of $\operatorname{Re}(I_3)$ over $\mathbb{Q}(\cos(\pi/n))$ is

$$(20) \quad X^2 - (2 \cos(\pi/n) + 1/2)X + (\cos^2(\pi/n) + \cos(\pi/n)/2 - 1/4).$$

One deduces (18) from (19), (20), and the addition laws for sines and cosines.

From the claim (18), we deduce that T is contained in the affine group of X . Namely, fixing the horizontal lines and postcomposing local charts of the cylinders by T defines an affine diffeomorphism whose derivative is T (cf. [35, Prop. 2.4] or [24, Lemma 9.7]).

It remains to prove that R and T generate the $(5, n, \infty)$ -triangle group. One constructs the hyperbolic triangle in the extended upper half-plane with corners $i\infty, i$, and

$$z_0 = \frac{\cos(\pi/n) + \cos(\pi/5)}{\sin(\pi/n)} + i \frac{\sin(\pi/5)}{\sin(\pi/n)}$$

bounded by the vertical axes through i and z_0 and the circle around $\cot(\pi/n)$ with radius $1/\sin(\pi/n)$. The interior angles at i and z_0 are indeed π/n and $\pi/5$. By Poincaré's theorem this triangle plus its reflection along the imaginary axis is a fundamental domain for the group generated by R and T .

The last claim follows now from the standard criterion to detect Teichmüller curves; see e.g., [24, Cor. 3.3]. \square

Remark 8.2. Assuming the comparison results which will be proved in Section 8.3 below, the number of cylinders in, say, the horizontal direction is already determined by results of the previous sections.

Consider the family of translation surfaces $\operatorname{diag}(e^t, e^{-t}) \cdot (X_0, \omega_0)$, where (X_0, ω_0) is as in Theorem 6.14. This family converges for $t \rightarrow \infty$ to a singular fiber of $\bar{f} : \bar{\mathcal{X}} \rightarrow \bar{C}$ and by [22] the number of cylinders in the horizontal direction equals the number of singularities of the singular fiber X_∞ .

Since all the local systems \mathbb{L}_i as in the proof of Theorem 6.2 have nontrivial parabolic monodromy around points in $\bar{\pi}^{-1}(\infty)$, the arithmetic genus of X_∞ is zero. Since ω_0 has only one zero, X_∞ is irreducible and hence the number of singularities of the fiber X_∞ equals $g(X_\infty)$.

8.2. The tables $T(4, n, \infty)$. Let $n \geq 5$ be odd. We define a billiard table $T(4, n, \infty)$ as follows. The billiard table is again a 4-gon in the complex plane with angles $\alpha = \pi/2$ and $\beta = \gamma = \pi/n$, as indicated in Figure 3. We denote by I_1, \dots, I_4 the vectors corresponding to the sides of the polygon. We regard these vectors as complex numbers. We scale and rotate the billiard table such that $I_4 = 1$ and such that $|I_3| = 2(\cos(\pi/n) + \cos(\pi/4))$. This determines the table uniquely.

The translation surface $X := X(4, n, \infty)$, obtained by unfolding $T(4, n, \infty)$, looks similar to the one obtained from $T(5, n, \infty)$. It can be obtained by identifying




Figure 3. Billiard table $T(4, n, \infty)$, for $n = 9$




Figure 4. Billiard table $T(4, 9, \infty)$, partially unfolded

parallel sides of two stars. The first star is illustrated in Figure 4. The second star is obtained from the first by reflection in the horizontal axis. The translation surface $X(4, n, \infty)$ has one singularity which corresponds to the vertex of the billiard table with angle δ . Its genus is $g = 3(n - 1)/2$.

THEOREM 8.3. *Let $n \geq 5$ be odd. Then the affine group of $X(4, n, \infty)$ contains the elements*

$$R = \begin{pmatrix} \cos(\pi/n) & -\sin(\pi/n) \\ \sin(\pi/n) & \cos(\pi/n) \end{pmatrix} \quad \text{and} \quad T = \begin{pmatrix} 1 & 2 \frac{\cos(\pi/n) + \cos(\pi/4)}{\sin(\pi/n)} \\ 0 & 1 \end{pmatrix}.$$

The elements $R, T \in \text{PSL}_2(\mathbb{R})$ generate the Fuchsian triangle group $\Delta(4, n, \infty)$. In particular, $X(4, 9, \infty)$ is a Veech surface.

Proof. Rotation around the center of each of the stars defines an affine diffeomorphism of $X(4, n, \infty)$ whose derivative is R , as in the case $(5, n, \infty)$.

We describe the cylinders in the horizontal direction. As for $X(5, n, \infty)$, we distinguish two types of cylinders. The cylinders of type 1, denoted by C_i , are those that are glued together from pieces of both stars. They correspond to pairs of sides which connect two points. Here a pair of sides consists of two distinct sides which are interchanged by reflection in the vertical axis. There are $(n - 1)/2$ such cylinders. The widths and heights of these cylinders, in an appropriate numbering,

are given by

$$w_k = 4|I_2| \cos\left(\frac{(n-2k)\pi}{2n}\right) \text{ and } h_k = 2|I_1| \sin\left(\frac{k\pi}{n}\right), \quad k = 1, \dots, (n-1)/2.$$

There are two cylinders with the same width and height, due to the symmetry.

The cylinders of type 2, denoted by \tilde{C}_i , are those that consist of pieces of one star only. They correspond to pairs of points of the stars. Here we use the same convention for pairs as above. The number of such cylinders is also $(n-1)/2$. The widths and heights of these cylinders are

$$\begin{aligned} \tilde{w}_k &= 2|I_3| \cos\left(\frac{(n-2k+2)\pi}{2n}\right) \text{ and} \\ \tilde{h}_k &= 2|I_4| \cos\left(\frac{(n-2k+2)\pi}{2n}\right) \sin\left(\frac{\pi}{n}\right), \quad k = 1, \dots, (n-1)/2. \end{aligned}$$

The moduli of the cylinders are

$$m_k := h_k/w_k = |I_1|/2|I_2| \quad \text{and} \quad \tilde{m}_k = \tilde{h}_k/\tilde{w}_k = |I_4| \sin \frac{\pi}{n} / |I_3|.$$

As in the proof of Theorem 8.1, one checks that

$$m_k/\tilde{m}_k = |I_1||I_3|/2|I_2||I_4| \sin(\pi/n) = 1,$$

by using the geometry of the billiard table and the minimal polynomial of $2(\cos(\pi/n) + \cos(\pi/4))$ over $\mathbb{Q}(\cos(\pi/n))$. The rest of the proof is analogous to the proof of Theorem 8.1. \square

8.3. Comparison with Theorems 6.2 and 6.14. In this section we relate the billiard tables constructed in Sections 8.1 and 8.2 to the families of curves constructed in Section 6. For simplicity we suppose that $1 < m < n$ are relatively prime integers such that n is odd. This assumption avoids a case distinction. It is easy to work out the general statement.

In Theorem 6.2 we constructed a Teichmüller curve with projective affine group $\Delta(m, n, \infty)$. We constructed a concrete finite cover, C , of this Teichmüller curve. We denote by \bar{C} the corresponding projective curve. Over \bar{C} there exists a universal family $f : \mathcal{X} \rightarrow \bar{C}$ of semistable curves. In Theorem 6.14 we showed that there exist points c of \bar{C} such that the fiber $X_0 := \mathcal{X}_c$ is a smooth curve which is a $2n$ -cyclic cover of the projective line branched at $(m+3)/2$ (resp. $(m+4)/2$) points if m is odd (resp. even). There also exist fibers of \mathcal{X} which are $2m$ -cyclic covers of the projective line branched at $(n+3)/2$, but we do not regard these here since it is convenient to have as few branch points as possible, for our purposes. One may check that this is the most efficient way to represent a fiber of \mathcal{X} as an abelian cover of the projective line. This representation allows us to use Schwarz-Christoffel maps ([38, Th. C']) to represent the fiber X_0 of \mathcal{X} as the unfolding of a billiard table, under certain conditions (see below).

We first suppose that m is odd. The $2n$ -cyclic cover $X_0 \rightarrow \mathbb{P}_u^1$ of Theorem 6.14(a) is branched at the real points $2 \cos(2k\pi/m)$. The Schwarz-Christoffel map is defined as

$$\text{SC}(w) = \int_0^w (u-2)^{\frac{1}{2n}-1} \prod_{k=1}^{(m-1)/2} (u-2 \cos(2k\pi/m))^{\frac{1}{n}-1} du.$$

The integrand is the generating differential form ω_0 .

The Schwarz-Christoffel map maps the real axis to a $(m+3)/2$ -gon which we denote by $T[m, n, \infty]$. If $T[m, n, \infty]$ has no self-crossings then SC maps the upper half-plane bijectively to the interior of this $(m+3)/2$ -gon. The interior angles of $T[m, n, \infty]$ are $(m-1)/2$ times π/n and once $\pi/2n$, in this order. The remaining angle is $2\pi - m\pi/2n \bmod 2\pi$ (resp. $\pi - m\pi/2n$) if $m \equiv 1 \bmod 4$ (resp. $m \equiv 3 \bmod 4$). The number of self-crossings is therefore $(m-5)/4$ if $m \equiv 1 \bmod 4$ and $(m-3)/4$ if $m \equiv 3 \bmod 4$. In particular, this number is zero if and only if $m = 3, 5$. For $m \geq 7$ it is therefore unclear whether one can obtain (X_0, ω_0) by unfolding a billiard table. However, it follows from our results that one cannot do this via the usual theory of Schwarz-Christoffel maps. Namely, for $m \geq 7$ one cannot represent a smooth fiber of $\bar{\mathcal{X}}$ as a cyclic cover of the projective line, such that the corresponding polygon does not have self-crossings.

If $m = 3$ or 5 , Theorem C' of [38] implies that the Veech surface (X_0, ω_0) is obtained by unfolding the billiard table $T[m, n, \infty]$. For $m = 3$, we obtain Ward's family (cf. Theorem 6.14(c)). For $m = 5$, the angles of the 4-gon $T[5, n, \infty]$ coincide with those of the billiard table $T(5, n, \infty)$ which we constructed in Section 8.1. We show below that both 4-gons are similar.

The case that m even is analogous. The Schwarz-Christoffel map

$$\text{SC}(w) = \int_0^w (u-2)^{\frac{1}{2}-1} \prod_{k=1}^{m/2} (u-2 \cos((2k-1)\pi/2m))^{\frac{1}{n}-1} du$$

maps the real axis to a $(m+4)/2$ -gon $T[m, n, \infty]$.

The interior angles of $T[m, n, \infty]$ are once $\pi/2$ and $m/2$ times π/n , in this order. The remaining angle is $(3n-m)\pi/2n$ if $m \equiv 0 \bmod 4$ and $((n-m)\pi/(2n))$ if $m \equiv 2 \bmod 4$. We conclude that the number of self-crossings is $(m-4)/4$ (resp. $(m-2)/4$) if $m \equiv 0 \bmod 4$ (resp. $m \equiv 2 \bmod 4$). Therefore the number of self-crossings is zero if and only if $m = 2, 4$. The case $m = 2$ corresponds to Veech's family [35] (see §5). We show below that the case $m = 4$ corresponds to the billiards constructed in Section 8.2.

We leave it to the reader to use Theorem 6.2 and the techniques of Theorem 6.14 to construct billiard tables with projective affine group $\Delta(4, n, \infty)$ and $\Delta(5, n, \infty)$ also in the case that n is even or divisible by 5, or both.

PROPOSITION 8.4. *Let m be either 4 or 5. The billiard table $T[m, n, \infty]$ is similar to the billiard table $T(m, n, \infty)$.*

Proof. Suppose that $m = 5$. The case that $m = 4$ is similar, and left to the reader.

Recall that the interior angles of the 4-gons $T(5, n, \infty)$ and $T[5, n, \infty]$ are the same, and also occur in the same order. Therefore we only have to compare the lengths of the sides of $T[5, n, \infty]$ to those of $T(5, m, \infty)$. Since the sides of $T[5, n, \infty]$ are expressed in terms of the Schwarz-Christoffel map, it suffices to show that

$$(21) \quad \frac{|\text{SC}(2 \cos(2\pi/5)) - \text{SC}(2)|}{|\text{SC}(\infty) - \text{SC}(2)|} = \frac{|I_3|}{|I_4|} = \frac{\cos(\pi/n) + \cos(\pi/5)}{\cos(\pi/2n)}.$$

Here I_3, I_4 are the vectors corresponding to the sides of the 4-gon $T[5, n, \infty]$ as indicated in Figure 1.

We first express the length of the vector I_4 in terms of Beta integrals:

$$(22) \quad |I_4| = \int_2^\infty (u-2)^{\frac{1}{2n}-1} (u-2 \cos(2\pi/5))^{\frac{1}{n}-1} (u-2 \cos(4\pi/5))^{\frac{1}{n}-1} du \\ = \int_1^\infty z^{1-\frac{5}{2n}} (z^5-1)^{\frac{1}{n}-1} (z+1) dz.$$

Here we used the substitution $u = z + 1/z$; cf. the proof of Theorem 6.14. Substituting $z = 1/t$, we recognize this integral as the sum of two Beta integrals:

$$(23) \quad |I_4| = \frac{1}{5} \left(B\left(\frac{2}{5} - \frac{1}{2n}, \frac{1}{n}\right) + B\left(\frac{3}{5} - \frac{1}{2n}, \frac{1}{n}\right) \right).$$

Similarly, one finds that

$$(24) \quad |I_3| = \frac{1}{5} \left((-1 + \zeta_5^2 \zeta_{2n}^{-1}) B\left(\frac{2}{5} - \frac{1}{2n}, \frac{1}{n}\right) + (-1 + \zeta_5^3 \zeta_{2n}^{-1}) B\left(\frac{3}{5} - \frac{1}{2n}, \frac{1}{n}\right) \right).$$

Equation (21) follows from (23) and (24) by expressing the Beta integrals in terms of Gamma functions, using that $\Gamma(z)\Gamma(1-z) = \pi/\sin(\pi z)$, and applying the addition formulas for sines and cosines. \square

One may give an alternative proof for the statement that the 4-gons $T[5, n, \infty]$ and $T(5, n, \infty)$ are similar by showing the following. Let P be any 4-gon with the prescribed angles, and let X be the corresponding translation surface. Suppose that the affine group of X contains R and T . Then P is similar to $T(5, n, \infty)$. This can be shown by first deducing from the geometry of P that an affine diffeomorphism with derivative R has to fix each saddle connection.

Remark 8.5. Several authors ([38], [37], [16], [32]) have classified the Teichmüller curves that are obtained by unfolding a rational triangle, under certain conditions on the angles of the triangle. We have obtained the translation surfaces $X(m, n, \infty)$ for $m = 4, 5$ by unfolding 4-gons. The corresponding families of Teichmüller curves have not been found by Ward et. al. This suggests that the translation surfaces $X(m, n, \infty)$ for $m = 4, 5$ may not be obtained by unfolding triangles, but of course we have not shown this.

Remark 8.6. For $n > m \geq 6$ we have not been able to obtain the translation surface $X(m, n, \infty)$ by unfolding a billiard table, since the corresponding polygon $T[m, n, \infty]$ may not be embedded in the complex plane. However, it should in principle be possible to give a concrete description of $X(m, n, \infty)$ as obtained by gluing certain cylinders, analogous to the description in the case of $m = 4, 5$ (§§8.1 and 8.2). As for $m = 4$ and 5 , it follows from Corollary 6.9 that we would need $g(X_0)$ cylinders, which is approximately $(m - 1)(n - 1)/2$: it will be difficult to visualize the result. Therefore it seems more natural to us to represent these Teichmüller curves via the algebraic description from Section 6.

9. Lyapunov exponents

Roughly speaking, a flat normed vector bundle on a manifold with a flow, i.e. an action of \mathbb{R}^+ , can sometimes be stratified according to the growth rate of the length of vectors under parallel transport along the flow. The growth rates are then called Lyapunov exponents. In this section we will relate Lyapunov exponents to degrees of some line bundles in case that the underlying manifold is a Teichmüller curve.

For the convenience of the reader we reproduce Oseledec's theorem ([30]) that proves the existence of such exponents. We give a restatement due to Kontsevich ([18]) in a language closer to our setting.

9.1. Multiplicative ergodic theorem. We start with some definitions. A measurable vector bundle is a bundle that can be trivialized by functions which only need to be measurable. If $(V, \|\cdot\|)$ and $(V', \|\cdot\|)$ are normed vector bundles and $T : V \rightarrow V'$ is a linear map, then we let $\|T\| := \sup_{\|v\|=1} \|T(v)\|$. A reference for notions in ergodic theory is [7].

THEOREM 9.1 (Oseledec). *Let $T_t : (M, \nu) \rightarrow (M, \nu)$ be an ergodic flow on a space M with finite measure ν . Suppose that the action of $t \in \mathbb{R}^+$ lifts equivariantly to a flow S_t on some measurable real bundle V on M . Suppose there exists a (not equivariant) norm $\|\cdot\|$ on V such that for all $t \in \mathbb{R}^+$*

$$\int_M \log(1 + \|S_t\|) \nu < \infty.$$

Then there exist real constants $\lambda_1 \geq \dots \geq \lambda_k$ and a filtration

$$V = V_{\lambda_1} \supset \dots \supset V_{\lambda_k} \supset 0$$

by measurable vector subbundles such that, for almost all $m \in M$ and all $v \in V_m \setminus \{0\}$, one has

$$\|S_t(v)\| = \exp(\lambda_i t + o(t)),$$

where i is the maximal value such that $v \in (V_i)_m$.

The V_{λ_i} do not change if $\|\cdot\|$ is replaced by another norm of 'comparable' size (e.g. if one is a scalar multiple of the other).

The numbers λ_i for $i = 1, \dots, k \leq \text{rank}(V)$ are called the *Lyapunov exponents of S_t* . Note that these exponents remain unchanged if we replace M by a finite unramified covering with a lift of the flow and the pullback of V . We adopt the convention of repeating the exponents according to the rank of V_i / V_{i+1} , such that we will always have $2g$ of them, possibly some of them equal. A reference for elementary properties of Lyapunov exponents is, e.g., [1].

If the bundle V comes with a symplectic structure the Lyapunov exponents are symmetric with respect to 0 ([1, Prop. 5.1]); i.e., they are

$$1 = \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_g \geq 0 \geq -\lambda_g \geq \dots \geq -\lambda_1 = -1.$$

We specialize these concepts to the situation we are interested in. Let ΩM_g^* be the bundle of nonzero holomorphic 1-forms over the moduli space of curves. Its points are translation surfaces. The 1-forms define a flat metric on the underlying Riemann surface and we let $\Omega_1 M_g \subset \Omega M_g^*$ be the hypersurface consisting of translation surfaces of area one. As usual we replace M_g by an appropriate fine moduli space adding a level structure, but we do not indicate this in the notation. This allows us to use a universal family $f : \mathcal{X} \rightarrow M_g$.

Over $\Omega_1 M_g$, we have the local system $\mathbb{V}_{\mathbb{R}} = R^1 f_* \mathbb{R}$, whose fiber over (X, ω) is $H^1(X, \mathbb{R})$. We denote the corresponding real C^∞ -bundle by V . This bundle naturally carries the Hodge metric

$$H(\alpha, \beta) = \int_X \alpha \wedge * \beta,$$

where classes in $H^1(X, \mathbb{R})$ are represented by \mathbb{R} -valued 1-forms, and where $*$ is the Hodge star operator. We denote by $\|\cdot\| := \|\cdot\|_T$ the associated metric on V .

There is a natural $\text{SL}_2(\mathbb{R})$ -action on $\Omega_1 M_g$ obtained by post-composing the charts given by integrating the 1-form with the \mathbb{R} -linear map given by $A \in \text{SL}_2(\mathbb{R})$ to obtain a new complex structure and new holomorphic 1-form (see e.g., [24] and the references there). The geodesic flow T_t on $\Omega_1 M_g$ is the restriction of the $\text{SL}_2(\mathbb{R})$ -action to the subgroup $\text{diag}(e^t, e^{-t})$. Since V carries a flat structure, we can lift T_t by parallel transport to a flow S_t on V . This is the *Kontsevich-Zorich cocycle*. The notion ‘cocycle’ is motivated by writing the flow on a vector bundle in terms of transition matrices.

Lyapunov exponents can be studied for any finite measure ν on a subspace M of $\Omega_1 M_g$ such that T_t is ergodic with respect to ν . Starting with the work of Zorich ([43]), Lyapunov exponents have been studied for connected components of the stratification of $\Omega_1 M_g$ by the order of zeros of the 1-form. The integral structure of ΩM_g^* as an affine manifold can be used to construct a finite ergodic measure μ . Lyapunov exponents for $(\Omega_1 M_g, \mu)$ may be interpreted as deviations from ergodic averages of typical leaves of measured foliations on surfaces of genus g . The reader is referred to [18], [10], and the surveys [19] and [11] for further motivation and results.

9.2. *Lyapunov exponents for Teichmüller curves.* We want to study Lyapunov exponents in case of an arbitrary Teichmüller curve C or rather its canonical lift M to $\Omega_1 M_g$ given by providing the Riemann surfaces parametrized by C with the normalized generating differential. The lift $\pi : M \rightarrow C$ is an S^1 -bundle. We equip M with the measure ν which is induced by the Haar measure on $\mathrm{SL}_2(\mathbb{R})$, normalized such that $\nu(M) = 1$. Locally, ν is the product of the measure ν_C coming from the Poincaré volume form and the uniform measure on S^1 , both normalized to have total volume one.

We can apply Oseledec's theorem since ν_M is ergodic for the geodesic flow ([7, Th. 4.2.1]).

We start from the observation that the decomposition (2) of the VHS in Theorem 2.1 is $\mathrm{SL}_2(\mathbb{R})$ -equivariant and orthogonal with respect to Hodge metric. This implies that the Lyapunov exponents of \mathbb{V} are the union of the Lyapunov exponents of the \mathbb{L}_i with those of \mathbb{M} .

Let $\mathcal{L}_i := (\mathbb{L}_i)^{1,0}$ be the $(1,0)$ -part of the Hodge filtration of the Deligne extension of \mathbb{L}_i to \bar{C} . Denote by $d_i := \deg(\mathcal{L}_i)$ the corresponding degrees. Recall from Theorem 2.1 that precisely one of the \mathbb{L}_i , say the first one \mathbb{L}_1 is maximal Higgs. Recall that $S = \bar{C} \setminus C$ is the set of singular fibers.

THEOREM 9.2. *Let ν_M be the finite $\mathrm{SL}_2(\mathbb{R})$ -invariant measure with support in the canonical lift M of a Teichmüller curve to $\Omega_1 M_g$. Then r of the Lyapunov exponents λ_i satisfy*

$$\lambda_i = d_i / d_1 = \lambda(\mathbb{L}_i, S).$$

In particular, these exponents are rational and nonzero.

Proof. We write $(\mathbb{L}_i)_{\mathbb{R}}$ for the local subsystem of $R^1 f_* \mathbb{R}$ such that

$$(\mathbb{L}_i)_{\mathbb{R}} \otimes_{\mathbb{R}} \mathbb{C} = \mathbb{L}_i$$

and let L_i be the C^∞ -bundle attached to $(\mathbb{L}_i)_{\mathbb{R}}$. We apply Oseledec's theorem to L_i . Then

$$\lambda_i = \lim_{t \rightarrow \infty} \frac{1}{t} \log ||S_t(v_i)||,$$

for $v_i \in L_i \setminus (L_i)_{-\lambda_i}$. By averaging, we have

$$\lambda_i = \lim_{t \rightarrow \infty} \frac{1}{t} \int_{G(L_i)} \log ||S_t(v_i)|| d\nu_{G(L_i)}(v_i),$$

where $\tau : G(L_i) \rightarrow M$ is the (Grassmann) bundle of norm one vectors in L_i . This bundle is locally isomorphic to $S^1 \times M$. The measure $\nu_{G(L_i)}$ is locally the product measure of ν with the uniform measure on S^1 .

Following the idea of Kontsevich ([18]) also exploited in Forni ([10]), we estimate the growth of the length of v_i not only as a function on the T_t -ray through $\tau(v_i)$ (given as the parallel transport of the corresponding vector) but as a function on the whole (quotient by a discrete group of a) Poincaré disc $D_{\tau(v_i)}$ in M . For this purpose we write $z = e^{i\theta}r$ ($\theta \in [0, 2\pi)$) for z in the unit disc D and lift

it to $\rho_\theta \operatorname{diag}(e^t, e^{-t}) \in \operatorname{SL}_2(\mathbb{R})$, where ρ_θ is the rotation matrix by Θ and $t = (1/2) \log((1+r)/(1-r))$. Using this lift $D \rightarrow \operatorname{SL}_2(\mathbb{R})$ we obtain our disc $D_{\tau(v_i)}$ in M using the (left) $\operatorname{SL}_2(\mathbb{R})$ -action on M .

Consider the following functions

$$f_D := f_{D,i} : \begin{cases} (\pi^* L_i \setminus \{0\}) \times D & \rightarrow \mathbb{R} \\ (v_i, z) & \mapsto \log \|z \cdot v_i\|, \end{cases}$$

where $z \cdot v_i$ is the parallel transport of v_i over the disc $D_{\tau(v_i)}$. This is well-defined since the monodromy of L_i acts by matrices in $\operatorname{SL}_2(\mathbb{Z}) = \operatorname{Sp}_2(\mathbb{Z})$ and symplectic transformations do not affect the Hodge length. Note that by definition

$$(25) \quad f_D(v_i, z) = f_D(z \cdot v_i, 0).$$

On the discs $D_{\tau(v_i)}$ we may apply the (hyperbolic) Laplacian Δ_h to the functions $f_{D_{\tau(v_i)}}$ with respect to the second variable, i.e. consider

$$h_D := h_{D,i} : \begin{cases} (\pi^* L_i \setminus \{0\}) \times D & \rightarrow \mathbb{R} \\ (v_i, z) & \mapsto (\Delta_h f_D(v_i, \cdot))(z). \end{cases}$$

Using (25) and the invariance of Δ_h under isometries one deduces that there is a function $h : \pi^* L_i \setminus \{0\} \rightarrow \mathbb{R}$, such that

$$(26) \quad h_D(v_i, z) = h(z \cdot v_i).$$

Since obviously $\int_{G(L_i)} h(S_t v_i) dv_{G(L_i)}(v_i) = \int_{G(L_i)} h(v_i) dv_{G(L_i)}(v_i)$ for any t , we can apply [19, Eq. (3)] (see also [10, Lemma 3.1] to obtain

$$(27) \quad \lambda_i = \int_{G(L_i)} h(v_i) dv_{G(L_i)}(v_i).$$

We want to relate this expression to the degree d_i of the line bundles \mathcal{L}_i . Suppose $s_i(u)$ is a holomorphic section of \mathcal{L}_i over some open $U \subset C$. Recall that L_i has unipotent monodromies, by assumption. Therefore [31, Prop. 3.4] implies that the Hodge metric grows not too fast near the punctures and we have

$$(28) \quad d_i = \frac{1}{2\pi i} \int_{\bar{C}} \partial \bar{\partial} \log(\|s_i\|).$$

Here as usual, if there is no global section of \mathcal{L}_i the contributions of local holomorphic sections are added up using a partition of unity.

Instead of considering a holomorphic section s_i , we now consider a flat section $v_i(u)$ of L_i over U . Then, in $(\wedge^2(\mathbb{L}_i)_{\mathbb{C}})^{\otimes 2}(U)$ one checks the identity

$$(29) \quad (v_i \wedge * v_i) \otimes (s_i \wedge \bar{s}_i) = \frac{1}{2} (v_i \wedge s_i) \otimes (v_i \wedge \bar{s}_i).$$

We integrate this identity over the fibers \mathcal{X}_c of $f : \mathcal{X} \rightarrow C$, take logarithms and the Laplacian $\frac{1}{2\pi i} \partial \bar{\partial}$. Note that

$$(30) \quad \frac{1}{2\pi i} \partial \bar{\partial} \log \frac{1}{2} (v_i \wedge s_i) \otimes (v_i \wedge \bar{s}_i) = 0.$$

Let F be a fundamental domain for the action of the affine group Γ in a Poincaré discs $D \hookrightarrow M$. Then (28) and (30) implies that for any flat section v_i of L_i , we have

$$d_i = \frac{-1}{2\pi i} \int_F \partial \bar{\partial} \log(\|v_i\|).$$

The differential operator $\partial \bar{\partial}$ coincides, up to a scalar, with $\Delta_h(\cdot)\omega_P$, where ω_P is the Poincaré area form. Therefore we obtain for each $v_i \in (\pi \circ \tau)^*(L_i \setminus \{0\})$ that

$$d_i = \frac{1}{4\pi} \int_F \Delta_h \log \|v_i(z)\| \omega_P(z),$$

where $v_i(z)$ is obtained from v_i via parallel transport. Hence by integrating over all $G(L_i)$ and taking care of the normalization of $\nu_{G(L_i)}$, we find that

$$(31) \quad d_i = \frac{1}{4\pi} \text{vol}(C) \int_{G(L_i)} \Delta_h \log \|v_i\| \nu_{G(L_i)}(v_i).$$

The statement of the theorem now follows by comparing (31) with (27). \square

COROLLARY 9.3. *At least r of the Lyapunov exponents are nonzero.*

Proof. By Theorem 9.2, it is sufficient to show that for $\mathcal{L}_i := (\mathbb{L}_i)^{(1,0)}$ the degree $\deg(\mathcal{L}_i) \neq 0$. If $\mathcal{L}_i = 0$ then, by Simpson's correspondence ([36, Th. 1.1]), \mathbb{L}_i would be a reducible local system. But since \mathbb{L}_i is Galois conjugate to \mathbb{L}_1 , this is a contradiction. \square

Remark 9.4. If $r \geq g - 1$ all the Lyapunov exponents are known. In fact in this case we can identify the remaining Lyapunov exponent by the formula ([18], [10, Lemma 5.3])

$$\sum_{i=1}^g \lambda_i = \frac{\deg(f_* \omega_{X/C})}{2g - 2 + s}.$$

In the case of Teichmüller curves associated with triangle groups constructed in Sections 5 and 6, the proof of Theorem 9.2 yields more. Since for these curves the VHS decomposes completely into subsystems of rank 2 (Remark 6.11) we can determine all the Lyapunov exponents.

PROPOSITION 9.5. *Suppose that the local system \mathbb{M} , as in Theorem 2.1 contains a rank 2 local subsystem \mathbb{F}_i , whose $(1, 0)$ -part is a line bundle, denoted by \mathcal{F}_i . Then the Lyapunov spectrum contains (in addition to the d_i/d_1) the exponents*

$$\deg(\mathcal{F}_i)/d_1.$$

By Theorem 9.2 and Proposition 9.5 it is justified to call $\lambda(\mathbb{L}_i)$ Lyapunov exponents.

References

- [1] G. BENETTIN, I. GALGINI, A. GIORGILLI, and J.-M. STRELCYN, Lyapunov characteristic exponents for smooth dynamical systems and for Hamiltonian systems; a method for computing all of them, Part I (Meccanica, 1980), pp. 9–20.

- [2] I. I. BOUW, The p -rank of ramified covers of curves, *Compositio Math.* **126** (2001), 295–322. MR 2002e:14045 Zbl 1056.14036
- [3] ———, Reduction of the Hurwitz space of metacyclic covers, *Duke Math. J.* **121** (2004), 75–111. MR 2005d:14042 Zbl 1056.14037
- [4] ———, Pseudo-elliptic bundles, deformation data and the reduction of Galois covers, Habilitation thesis, Ph.D. thesis, Essen, 2005.
- [5] I. I. BOUW and S. WEWERS, Indigenous bundles with nilpotent p -curvature, *Internat. Math. Res. Not.* **2006** (2006), Art. ID 89254, 37. MR 2007i:14032 Zbl 1121.14026
- [6] P. COHEN and J. WOLFART, Modular embeddings for some nonarithmetic Fuchsian groups, *Acta Arith.* **56** (1990), 93–110. MR 92d:11039 Zbl 0717.14014
- [7] I. P. CORNFELD, S. V. FOMIN, and Y. G. SINAI, *Ergodic Theory*, *Grundle Math. Wissen.* **245**, Springer-Verlag, New York, 1982. MR 87f:28019 Zbl 0493.28007
- [8] P. DELIGNE, Un théorème de finitude pour la monodromie, in *Discrete Groups in Geometry and Analysis* (New Haven, Conn., 1984), *Progr. Math.* **67**, Birkhäuser, Boston, MA, 1987, pp. 1–19. MR 88h:14013 Zbl 0656.14010
- [9] P. DELIGNE, *Équations Différentielles à Points Singuliers Réguliers*, *Lecture Notes in Math.* **163**, Springer-Verlag, New York, 1970. MR 54 #5232 Zbl 0244.14004
- [10] G. FORNI, Deviation of ergodic averages for area-preserving flows on surfaces of higher genus, *Ann. of Math.* **155** (2002), 1–103. MR 2003g:37009 Zbl 1034.37003
- [11] ———, On the Lyapunov exponents of the Kontsevich-Zorich cocycle, in *Handbook of Dynamical Systems*, vol. 1B, Elsevier B. V., Amsterdam, 2006, pp. 549–580. MR 2010c:37004 Zbl 1130.37302
- [12] D. HARBATER and K. F. STEVENSON, Patching and thickening problems, *J. Algebra* **212** (1999), 272–304. MR 2000b:14004 Zbl 0944.14013
- [13] P. HOOPER, Grid graphs and lattice surfaces, preprint, 2009.
- [14] P. HUBERT and T. A. SCHMIDT, Invariants of translation surfaces, *Ann. Inst. Fourier (Grenoble)* **51** (2001), 461–495. MR 2003e:32023 Zbl 0985.32008
- [15] N. M. KATZ, Nilpotent connections and the monodromy theorem: Applications of a result of Turrittin, *Inst. Hautes Études Sci. Publ. Math.* (1970), 175–232. MR 45 #271 Zbl 0221.14007
- [16] R. KENYON and J. SMILLIE, Billiards on rational-angled triangles, *Comment. Math. Helv.* **75** (2000), 65–108. MR 2001e:37046 Zbl 0967.37019
- [17] S. KERCKHOFF, H. MASUR, and J. SMILLIE, Ergodicity of billiard flows and quadratic differentials, *Ann. of Math.* **124** (1986), 293–311. MR 88f:58122 Zbl 0637.58010
- [18] M. KONTSEVICH, Lyapunov exponents and Hodge theory, in *The Mathematical Beauty of Physics* (Saclay, 1996), *Adv. Ser. Math. Phys.* **24**, World Sci. Publ., River Edge, NJ, 1997, pp. 318–332. MR 99b:58147 Zbl 1058.37508
- [19] R. KRIKORIAN, Déviations de moyennes ergodiques, flots de Teichmüller et cocycle de Kontsevich-Zorich, in *Séminaire Bourbaki 927* (2003), *Astérisque* **299**, Soc. Math. France, Paris, 2005, pp. 59–93. MR 2006k:37109 Zbl 1094.37021
- [20] P. LOCHAK, On arithmetic curves in the moduli spaces of curves, *J. Inst. Math. Jussieu* **4** (2005), 443–508. MR 2006m:32010 Zbl 1094.14018
- [21] G. A. MARGULIS, *Discrete Subgroups of Semisimple Lie Groups*, *Ergeb. Math. Grenzgeb.* **17**, Springer-Verlag, New York, 1991. MR 92h:22021 Zbl 0732.22008
- [22] H. MASUR, On a class of geodesics in Teichmüller space, *Ann. of Math.* **102** (1975), 205–221. MR 52 #6038 Zbl 0322.32010
- [23] H. MASUR and S. TABACHNIKOV, Rational billiards and flat structures, in *Handbook of Dynamical Systems*, vol. 1A, North-Holland, Amsterdam, 2002, pp. 1015–1089. MR 2003j:37002 Zbl 1057.37034
- [24] C. T. MCMULLEN, Billiards and Teichmüller curves on Hilbert modular surfaces, *J. Amer. Math. Soc.* **16** (2003), 857–885. MR 2004f:32015 Zbl 1030.32012

- [25] ———, Prym varieties and Teichmüller curves, *Duke Math. J.* **133** (2006), 569–590. MR 2007a:32018 Zbl 1099.14018
- [26] ———, Teichmüller curves in genus two: torsion divisors and ratios of sines, *Invent. Math.* **165** (2006), 651–672. MR 2007f:14023 Zbl 1103.14014
- [27] S. MOCHIZUKI, *Foundations of p -adic Teichmüller Theory*, AMS/IP Studies in Advanced Mathematics **11**, Amer. Math. Soc., Providence, RI, 1999. MR 2000i:14025 Zbl 0969.14013
- [28] M. MÖLLER, Periodic points on Veech surfaces and the Mordell-Weil group over a Teichmüller curve, *Invent. Math.* **165** (2006), 633–649. MR 2007e:14012 Zbl 1111.14019
- [29] ———, Variations of Hodge structures of a Teichmüller curve, *J. Amer. Math. Soc.* **19** (2006), 327–344. MR 2007b:32026 Zbl 1090.32004
- [30] V. I. OSELEDEC, A multiplicative ergodic theorem. Characteristic Lyapunov exponents of dynamical systems, *Trans. Moscow Math. Soc.* **19** (1968), 197–231. MR 39 #1629 Zbl 0236.93034
- [31] C. A. M. PETERS, A criterion for flatness of Hodge bundles over curves and geometric applications, *Math. Ann.* **268** (1984), 1–19. MR 85m:14014 Zbl 0548.14004
- [32] J.-C. PUCHTA, On triangular billiards, *Comment. Math. Helv.* **76** (2001), 501–505. MR 2002f:37060 Zbl 01701563
- [33] D. SINGERMAN, Finitely maximal Fuchsian groups, *J. London Math. Soc.* **6** (1972), 29–38. MR 48 #529 Zbl 0251.20052
- [34] K. TAKEUCHI, Arithmetic triangle groups, *J. Math. Soc. Japan* **29** (1977), 91–106. MR 55 #2754 Zbl 0365.20055
- [35] W. A. VEECH, Teichmüller curves in moduli space, Eisenstein series and an application to triangular billiards, *Invent. Math.* **97** (1989), 553–583. MR 91h:58083a Zbl 0676.32006
- [36] E. VIEHWEG and K. ZUO, A characterization of certain Shimura curves in the moduli stack of abelian varieties, *J. Differential Geom.* **66** (2004), 233–287. MR 2006a:14015 Zbl 1078.11043
- [37] Y. B. VOROBETS, Plane structures and billiards in rational polygons: the Veech alternative, *Russian Math. Surveys* **51** (1996), 779–817. MR 97j:58092
- [38] C. C. WARD, Calculation of Fuchsian groups associated to billiards in a rational triangle, *Ergodic Theory Dynam. Systems* **18** (1998), 1019–1042. MR 2000b:30065 Zbl 0915.58059
- [39] S. WEWERS, Construction of Hurwitz spaces, thesis, Essen, 1998, preprint no. 21 of the IEM.
- [40] ———, Deformation of tame admissible covers of curves, in *Aspects of Galois Theory* (Gainesville, FL, 1996), *London Math. Soc. Lecture Note Ser.* **256**, Cambridge Univ. Press, Cambridge, 1999, pp. 239–282. MR 2001b:14048 Zbl 0995.14008
- [41] S. WEWERS and I. I. BOUW, Alternating groups as monodromy groups in positive characteristic, *Pacific J. Math.* **222** (2005), 185–200. MR 2006k:14050 Zbl 1111.14021
- [42] M. YOSHIDA, *Fuchsian Differential Equations*, *Aspects of Mathematics* **11**, Friedr. Vieweg & Sohn, Braunschweig, 1987. MR 90f:32025 Zbl 0618.35001
- [43] A. ZORICH, Finite Gauss measure on the space of interval exchange transformations. Lyapunov exponents, *Ann. Inst. Fourier (Grenoble)* **46** (1996), 325–370. MR 97f:58081 Zbl 0853.28007

(Received December 1, 2005)

(Revised April 21, 2007)

E-mail address: irene.bouw@uni-ulm.de

INSTITUTE OF PURE MATHEMATICS, ULM UNIVERSITY, D-89069 ULM, GERMANY

E-mail address: moeller@math.uni-frankfurt.de

INSTITUT FÜR MATHEMATIK, GOETHE-UNIVERSITÄT, ROBERT-MAYER-STR. 6-8,
D-60054 FRANKFURT AM MAIN, GERMANY

This periodical is published bimonthly by the Department of Mathematics at Princeton University with the cooperation of the Institute for Advanced Study. *Annals* is typeset in \TeX by Sarah R. Warren and produced by Mathematical Sciences Publishers. The six numbers each year are divided into two volumes of three numbers each.

Editorial correspondence

Papers submitted for publication and editorial correspondence should be addressed to Maureen Schupsky, *Annals of Mathematics*, Fine Hall-Washington Road, Princeton University, Princeton, NJ, 08544-1000 U.S.A. The e-mail address is annals@math.princeton.edu.

Preparing and submitting papers

The *Annals* requests that all papers include an abstract of about 150 words which explains to the nonspecialist mathematician what the paper is about. It should not make any reference to the bibliography. Authors are encouraged to initially submit their papers electronically and in PDF format. Please send the file to: annals@math.princeton.edu or to the Mathematics e-print arXiv: front.math.ucdavis.edu/submissions. If a paper is submitted through the arXiv, then please e-mail us with the arXiv number of the paper.

Proofs

A PDF file of the galley proof will be sent to the corresponding author for correction. If requested, a paper copy will also be sent to the author.

Offprints

Authors of single-authored papers will receive 30 offprints. (Authors of papers with one co-author will receive 15 offprints, and authors of papers with two or more co-authors will receive 10 offprints.) Extra offprints may be purchased through the editorial office.

Subscriptions

The price for a print and online subscription, or an online-only subscription, is \$390 per year for institutions. In addition, there is a postage surcharge of \$40 for print subscriptions that are mailed to countries outside of the United States. Individuals interested in subscriptions for their own personal use should contact the publisher at the address below. Subscriptions and changes of address should be sent to Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840 (e-mail: contact@mathscipub.org; phone: 1-510-643-8638; fax: 1-510-295-2608). (Checks should be made payable to "Mathematical Sciences Publishers".)

Back issues and reprints

Orders for missing issues and back issues should be sent to Mathematical Sciences Publishers at the above address. Claims for missing issues must be made within 12 months of the publication date. Online versions of papers published five or more years ago are available through JSTOR (www.jstor.org).

Microfilm

Beginning with Volume 1, microfilm may be purchased from NA Publishing, Inc., 4750 Venture Drive, Suite 400, PO Box 998, Ann Arbor, MI 48106-0998; phone: 1-800-420-6272 or 734-302-6500; email: info@napubco.com, website: www.napubco.com/contact.html.

ALL RIGHTS RESERVED UNDER THE BERNE CONVENTION AND
THE UNIVERSAL COPYRIGHT CONVENTION

TABLE OF CONTENTS

DENIS OSIN. Small cancellations over relatively hyperbolic groups and embedding theorems	1–39
HARUZO HIDA. The Iwasawa μ -invariant of p -adic Hecke L -functions.....	41–137
IRENE I. BOUW and MARTIN MÖLLER. Teichmüller curves, triangle groups, and Lyapunov exponents	139–185
IGOR RODNIANSKI and JACOB STERBENZ. On the formation of singularities in the critical $O(3)$ σ -model	187–242
YUAN-PIN LEE, HUI-WEN LIN and CHIN-LUNG WANG. Flops, motives, and invariance of quantum rings	243–290
LÁSZLÓ ERDŐS, BENJAMIN SCHLEIN and HORNG-TZER YAU. Derivation of the Gross-Pitaevskii equation for the dynamics of Bose-Einstein condensate	291–370
L. HAKAN ELIASSEN and SERGEI B. KUKSIN. KAM for the nonlinear Schrödinger equation	371–435
MIRIAM BRISKIN, NINA ROYTVARF and YOSEF YOMDIN. Center conditions at infinity for Abel differential equations.....	437–483
IGOR KRICHEVER. Characterizing Jacobians via trisecants of the Kummer variety.....	485–516
MARC BURGER, ALESSANDRA IOZZI and ANNA WIENHARD. Surface group representations with maximal Toledo invariant	517–566
TIM DOKCHITSER and VLADIMIR DOKCHITSER. On the Birch-Swinnerton-Dyer quotients modulo squares	567–596
DMITRI BELIAEV and STANISLAV SMIRNOV. Random conformal snowflakes.....	597–615
SOURAV CHATTERJEE, RON PELED, YUVAL PERES and DAN ROMIK. Gravitational allocation to Poisson points.....	617–671
RADU LAZA. The moduli space of cubic fourfolds via the period map.....	673–711
NARUTAKA OZAWA and SORIN POPA. On a class of II_1 factors with at most one Cartan subalgebra.....	713–749
MINHYONG KIM. p -adic L -functions and Selmer varieties associated to elliptic curves with complex multiplication	751–759