

ANNALS OF MATHEMATICS

A finiteness theorem for zero-cycles over p -adic fields

By SHUJI SAITO and KANETOMO SATO
APPENDIX BY UWE JANNSSEN



SECOND SERIES, VOL. 172, NO. 3

November, 2010

ANMAAH

A finiteness theorem for zero-cycles over p -adic fields

By SHUJI SAITO and KANETOMO SATO
APPENDIX BY UWE JANNSEN

Abstract

Let R be a henselian discrete valuation ring. Let X be a regular projective flat scheme over $\mathrm{Spec}(R)$ with generalized semistable reduction. We prove a bijectivity theorem for étale cycle class maps of the Chow group of 1-cycles on X . As an application, we prove a finiteness theorem for the Chow group of 0-cycles on a projective smooth variety over a p -adic field.

Introduction

1. Homology theory and the cycle map
2. Kato homology
3. The vanishing theorem
4. The Bertini theorem over a discrete valuation ring
5. Surjectivity of the cycle map
6. The blow-up formula
7. A moving lemma
8. Proof of the main theorem
9. Applications of the main theorem

Appendix A. Resolution of singularities for embedded curves, by Uwe Jannsen

References

The first named author was supported by Grant-in-Aid for Scientific Research B-18340003 and S-19104001 and JSPS Core-to-Core Program 18005.

The second named author was supported by Grant-in-Aid for Scientific Research B-20740008 and JSPS Core-to-Core Program 18005.

Introduction

Let V be a smooth projective geometrically integral variety over a field k . Let $\mathrm{CH}_0(V)$ be the Chow group of zero-cycles on V modulo rational equivalence, and let $A_0(V) \subset \mathrm{CH}_0(V)$ be the subgroup consisting of cycles of degree 0. Then we have a homomorphism

$$\phi_V : A_0(V) \longrightarrow \mathrm{Alb}_V(k)$$

induced by the Albanese map $V \rightarrow \mathrm{Alb}_V$, where Alb_V denotes the Albanese variety of V and $\mathrm{Alb}_V(k)$ denotes the group of k -rational points on Alb_V . By Abel's theorem ϕ_V is injective if $\dim(V) = 1$. It is Mumford [Mum68] who first discovered that the situation is rather chaotic in the case of higher dimension. If $\dim(V) \geq 2$ and $k = \mathbb{C}$, the kernel $\mathrm{Ker}(\phi_V)$ of ϕ_V may be too large to be understood by a standard geometric structure. On the other hand, the situation in the case that k is a field of arithmetic nature presents a striking contrast to the case $k = \mathbb{C}$. If k is finite, $\mathrm{Ker}(\phi_V)$ has been explicitly determined by geometric class field theory [KS83, Prop. 9]. One of the most fascinating and challenging conjectures in arithmetic geometry is a conjecture of Bloch and Beilinson that $\mathrm{Ker}(\phi_V)$ is torsion in case k is a number field. Combined with the conjecture that $\mathrm{CH}_0(V)$ is finitely generated for V over a number field, it would imply that $\mathrm{Ker}(\phi_V)$ is finite. Very little is known about the conjecture and we have nothing to say about it in this paper. Instead we consider its analogue over p -adic fields, more precisely, the following conjecture:

CONJECTURE 0.1. *Let V be a smooth projective geometrically integral variety over a p -adic field k . Then $\mathrm{Ker}(\phi_V)$ is the direct sum of a finite group and a divisible group.*

A result of Mattuck [Mat55] implies, by aid of [JS03, Lemma 7.8], that $\mathrm{Alb}_V(k)$ is isomorphic to the direct sum of a finite group and a p' -divisible group. Here a p' -divisible group means an abelian group which is divisible by any integer prime to p . So the above conjecture implies

CONJECTURE 0.2. *Let V be as in Conjecture 0.1. Then $A_0(V)$ is the direct sum of a finite group and a p' -divisible group.*

In essentially equivalent forms, Conjectures 0.1 and 0.2 were made by Colliot-Thélène in 1993 ([CT95]) and also in the more recent notes [CTb], [CTc]. He chose to state most of them as “questions” or “guesses”. Some finiteness results in this direction have been known for surfaces ([SS95, Th. 2.5]; see also [CT95, Th. 2.1] and some special varieties of higher dimension ([RS00], [PS95], [KS03], [CT05, Théorème and §5]). What has been known for surfaces is that for any prime $\ell \neq p$ the group $A_0(X)$ is the direct sum of a finite ℓ -primary group G_ℓ and an ℓ -divisible group. But it had not been known in general that $G_\ell = 0$ for almost all primes ℓ . That this is indeed the case is now a consequence of Corollary 0.4 below.

In this paper we will prove the following affirmative result concerning Conjecture 0.2, where \mathfrak{O}_k denotes the ring of integers in k :

THEOREM 0.3 (Theorem 9.7). *Let V be a smooth projective geometrically integral variety over a p -adic field k . Let F be the residue field of \mathfrak{O}_k . Assume that V has a regular projective flat model X over \mathfrak{O}_k on which the reduced subscheme of the divisor $X \otimes_{\mathfrak{O}_k} F$ has simple normal crossings. Then Conjecture 0.2 holds true.*

This result has not been known even in the case that $X \otimes_{\mathfrak{O}_k} F$ is smooth. See Definition 1.15 below for the definition of divisors with simple normal crossings. Jean-Louis Colliot-Thélène notices that a combination of our theorem and one of de Jong's results on alterations (see [dJ96, Th. 8.2]) implies the following result, which had been stated as a conjecture in [CT95].

COROLLARY 0.4. *Let V be a smooth projective geometrically integral variety over a p -adic field k . Then $A_0(V)$ is ℓ -divisible for almost all primes ℓ .*

Indeed the cited theorem implies that there exists a finite field extension L of k , a smooth projective variety W over L equipped with a proper generically finite morphism $\pi : W \rightarrow V$ such that W has strict semistable reduction over L . In particular, it satisfies the assumption of Theorem 0.3 over \mathfrak{O}_L . Hence Corollary 0.4 follows from Theorem 0.3 for W and the fact that the composite $A_0(V) \xrightarrow{\pi^*} A_0(W) \xrightarrow{\pi_*} A_0(V)$ is multiplication by a positive integer (cf. [Ful98, Ex. 1.7.4]).

COROLLARY 0.5. *Let the assumptions be as in Theorem 0.3. Assume further that V is rationally connected in the sense of Kollár, Miyaoka and Mori. Then $A_0(V)$ is the direct sum of a finite group and a p -primary torsion group of finite exponent.*

It is indeed known that for a smooth projective rationally connected variety V over a field, the group $A_0(V)$ is killed by some positive integer ([CT05, Prop. 11]). The corollary thus follows from Theorem 0.3. In the particular case that V is a smooth compactification of a connected linear algebraic group, Colliot-Thélène [CT05] established the conclusion of Corollary 0.5 without even assuming the existence of a regular proper model.

For a smooth integral variety V over a field k , a standard method to study $\mathrm{CH}_0(V)$ is to use the étale cycle map ([SGA $_{\frac{1}{2}}$, Cycle])

$$\rho_V : \mathrm{CH}_0(V)/n \longrightarrow H_{\mathrm{\acute{e}t}}^{2d}(V, \mathbb{Z}/n\mathbb{Z}(d)),$$

where d denotes $\dim(V)$, n is an integer prime to $\mathrm{ch}(k)$ and $\mathbb{Z}/n\mathbb{Z}(r) = \mu_n^{\otimes r}$ is the Tate twist of the étale sheaf μ_n of n -th roots of unity. The étale cohomology groups $H_{\mathrm{\acute{e}t}}^*(V, \mathbb{Z}/n\mathbb{Z}(r))$ are finite if k is a p -adic field. So by the compatibility between ρ_V and ϕ_V stated in [Ras95, Appendix], one can easily show that Conjecture 0.2 follows if ρ_V is injective for any $n \geq 1$ prime to p . We do not use this statement in what follows, because the last naive expectation concerning the injectivity of

ρ_V was shattered by an example found by Parimala-Suresh [PS95]. In this paper we shed new light on the problem by considering the cycle map for schemes over integer rings.

Let R be an excellent henselian discrete valuation ring with residue field F and fraction field k . In what follows, we do not assume that k has characteristic zero. Put $B := \operatorname{Spec}(R)$ with the closed point $s := \operatorname{Spec}(F)$ and the generic point $\eta := \operatorname{Spec}(k)$. For a scheme X over B , put $X_s := X \times_B s$ and $X_\eta := X \times_B \eta$, and let $X_{s,\text{red}}$ be the reduced subscheme of X_s . Let \mathcal{QSP} be the category of regular projective flat schemes X over B on which the reduced divisor $X_{s,\text{red}}$ has simple normal crossings. Let X be an object of \mathcal{QSP} , and let $\operatorname{CH}_1(X)$ be the Chow group of one-dimensional cycles on X (cf. Definition 1.2 below). For an integer $n \geq 1$ prime to $\operatorname{ch}(F)$, there is a cycle class map (cf. (1.9) and (1.12) below)

$$\rho_X : \operatorname{CH}_1(X)/n \longrightarrow H_{\text{ét}}^{2d}(X, \mathbb{Z}/n\mathbb{Z}(d)) \quad \text{with } d := \dim(X) - 1.$$

The main result, Theorem 0.3 will be deduced from the following result:

THEOREM 0.6 (Theorem 1.16). *If F is finite or separably closed, then ρ_X is bijective.*

There are three main ingredients in the proof of Theorem 0.6. The first one is the Bertini theorem over a discrete valuation ring proved in [JSa, Th. 1]. It affirms the existence of a nice hypersurface section $Y \subset X$ over B which belongs to $\operatorname{Ob}(\mathcal{QSP})$, whose complement $X - Y$ is affine and for which $Y \cup X_{s,\text{red}}$ is a reduced divisor with simple normal crossings on X . Such a pair (X, Y) is called an ample \mathcal{QSP} -pair. In this paper we need its refinement, obtained by a joint work of Jannsen and the first author, where it is allowed to impose an additional condition that Y contains a given regular closed subscheme $W \subset X$ with suitable conditions. We include a proof in Section 4 below. The second key ingredient is a vanishing theorem of $H_{\text{ét}}^q(X - Y, \mathbb{Z}/n\mathbb{Z}(d))$ for an ample \mathcal{QSP} -pair (X, Y) (see Theorem 3.2 below). A key to its proof is an extension of the base-change theorem of Rapoport and Zink [RZ82, Satz 2.19]. We remove the assumption on the multiplicity of the components of X_s using the absolute purity theorem of Gabber [Fuj02]. Recall that we have the affine vanishing theorem of Artin-Gabber (loc. cit. §5), which holds for more general affine schemes. Our Theorem 3.2 implies that a stronger vanishing holds in our special situation (cf. Remark 3.3). The last key ingredient of the proof of Theorem 0.6 is an arithmetic invariant associated to a quasi-projective scheme X over B . It arises from a homological complex $KC(X, \mathbb{Z}/n\mathbb{Z})$ with $(n, \operatorname{ch}(F)) = 1$ of the form

$$\begin{aligned} \bigoplus_{x \in X_0} H_{\text{ét}}^0(x, \mathbb{Z}/n\mathbb{Z}(-1)) &\leftarrow \bigoplus_{x \in X_1} H_{\text{ét}}^1(x, \mathbb{Z}/n\mathbb{Z}) \\ &\leftarrow \cdots \leftarrow \bigoplus_{x \in X_a} H_{\text{ét}}^a(x, \mathbb{Z}/n\mathbb{Z}(a-1)) \leftarrow \cdots, \end{aligned}$$

where X_a denotes the set of the points on X of dimension a , and the leftmost term is placed in degree 0. We define the group $KH_a(X, \mathbb{Z}/n\mathbb{Z})$ as the homology group of $KC(X, \mathbb{Z}/n\mathbb{Z})$ in degree a and put

$$KH_a(X, \mathbb{Q}_\ell/\mathbb{Z}_\ell) := \varinjlim_n KH_a(X, \mathbb{Z}/\ell^n\mathbb{Z}) \quad \text{for a prime } \ell \neq \text{ch}(F).$$

We construct the complex $KC(X, \mathbb{Z}/n\mathbb{Z})$ using localization sequences of a certain étale homology theory (see §§1–2 below for details). The complex $KC(X, \mathbb{Z}/n\mathbb{Z})$ is identified with the case $j = -1$ of the homological complex $C_n^j(X)$ defined by Kato [Kat86], up to signs of differential operators (cf. Definition 2.1 below):

$$\begin{aligned} \bigoplus_{x \in X_0} H_{\text{ét}}^{j+1}(x, \mathbb{Z}/n\mathbb{Z}(j)) &\leftarrow \bigoplus_{x \in X_1} H_{\text{ét}}^{j+2}(x, \mathbb{Z}/n\mathbb{Z}(j+1)) \\ &\leftarrow \cdots \leftarrow \bigoplus_{x \in X_a} H_{\text{ét}}^{a+j+1}(x, \mathbb{Z}/n\mathbb{Z}(a+j)) \leftarrow \cdots \end{aligned}$$

In loc. cit., he studied the case $j = 0$ and conjectures that $C_n^0(X)$ is exact for any $X \in \text{Ob}(\mathcal{QSP})$ if F is finite (loc. cit. Conjecture 5.1). In this paper we study the case $j = -1$ and propose the following new conjecture.

CONJECTURE 0.7 (Conjecture 2.12). *Assume that F is finite. Then for $X \in \text{Ob}(\mathcal{QSP})$, there is an isomorphism*

$$KH_a(X, \mathbb{Q}_\ell/\mathbb{Z}_\ell) \simeq \begin{cases} (\mathbb{Q}_\ell/\mathbb{Z}_\ell)^{I(X_s)} & a = 1 \\ 0 & a \neq 1, \end{cases}$$

where $I(X_s)$ denotes the set of irreducible components of $X_{s,\text{red}}$.

The vanishing of $KH_2(X, \mathbb{Q}_\ell/\mathbb{Z}_\ell)$ is equivalent to the surjectivity of the cycle map ρ_X (cf. Proof of Theorem 5.3). If $\dim(X) = 2$, $KH_2(X, \mathbb{Q}_\ell/\mathbb{Z}_\ell)$ is isomorphic to the ℓ -primary torsion part of the Brauer group of X (cf. Lemma 5.2), and its vanishing is a consequence of the base-change theorem for Brauer groups of relative curves ([Gro68, III, Th.3.1], [CTOP02, Th. 1.7 (c)]). See Theorem 5.1 below for details. If $\dim(X) = 3$, the vanishing of $KH_3(X, \mathbb{Q}_\ell/\mathbb{Z}_\ell)$ implies the injectivity of ρ_X (cf. Lemmas 8.7 and 8.8). We will prove the following result as a consequence of Theorem 0.6, where the dimension of X is arbitrary.

THEOREM 0.8 (Theorem 2.13). *Conjecture 0.7 holds in degree $a \leq 3$.*

We state some other consequences of Theorem 0.6, which will be proved in Section 9 below.

COROLLARY 0.9 (Corollary 9.2). *Assume that F is separably closed. Let X be an object of \mathcal{QSP} , and let Y_1, \dots, Y_N be the irreducible components of $X_{s,\text{red}}$. Then for any integer $n > 0$ prime to $\text{ch}(F)$, we have*

$$\text{CH}_1(X)/n \xrightarrow{\simeq} \bigoplus_{1 \leq i \leq N} \mathbb{Z}/n\mathbb{Z}, \quad \alpha \mapsto \langle \alpha, Y_i \rangle_{1 \leq i \leq N},$$

where $\langle \alpha, Y_i \rangle$ denotes the intersection number (see §9 below for its definition).

It would be interesting to compare this result with those in [Dal05] and [Mad08]. In both works some nontrivial classes in $A_0(V)$ are detected by studying intersection theory with the special fiber of a regular model over the ring of integers. In [Dal05] the residue field F is finite and in [Mad08] it is of the shape $\mathbb{C}((t))$.

COROLLARY 0.10 (Corollary 9.5). *Assume that X is smooth and projective over B and that F is finite or separably closed. Then for any integer $n > 0$ prime to $\text{ch}(F)$,*

$$\text{CH}_0(X_s)/n \xleftarrow[i^*]{\simeq} \text{CH}_1(X)/n \xrightarrow[j^*]{\simeq} \text{CH}_0(X_\eta)/n \quad (X_s \xhookrightarrow{i} X \xhookrightarrow{j} X_\eta).$$

If F is separably closed, the degree map $\text{CH}_0(X_s) \rightarrow \mathbb{Z}$ induces an isomorphism $\text{CH}_0(X_s)/n \simeq \mathbb{Z}/n$. When F is finite, $A_0(X_s)$ is a finite group computed by higher class field theory [KS83]. When moreover X_s is separably rationally connected, that computation shows $A_0(X_s) = 0$. We thus recover the prime-to- $\text{ch}(F)$ part of the main result of [KS03], which was obtained by deformation theory. One should also compare the above result with the result of [Kol04], where the base is a henselian discrete valuation ring with arbitrary perfect residue field.

COROLLARY 0.11 (Corollary 9.6). *Let (X, Y) be an ample \mathcal{LSP} -pair with $\dim(X) = d + 1 \geq 2$, and assume that F is finite (resp. separably closed). Then for any integer $n > 0$ prime to $\text{ch}(F)$, the push-forward map*

$$i_* : \text{CH}_1(Y)/n \longrightarrow \text{CH}_1(X)/n \quad (i : Y \hookrightarrow X)$$

is bijective for $d \geq 3$ and surjective for $d = 2$ (resp. bijective for $d \geq 2$ and surjective for $d = 1$).

This result can be viewed as a hard Lefschetz theorem for $\text{CH}_1(X)/n$. Theorem 0.3 is an easy consequence of it (see §9 below).

1. Homology theory and the cycle map

Unless indicated otherwise, all cohomology groups of schemes are taken over the étale topology. Throughout this paper, we fix a discrete valuation ring R with residue field F and quotient field k . We put $B := \text{Spec}(R)$, $s := \text{Spec}(F)$ and $\eta := \text{Spec}(k)$. Let Sch_B^{qp} be the category of quasi-projective B -schemes and B -morphisms. For $X \in \text{Ob}(\text{Sch}_B^{qp})$, we put $X_s := X \times_B s$ and $X_\eta := X \times_B \eta$, and write $X_{s,\text{red}}$ for the reduced subscheme of X_s . We define $\dim'(X)$ to be the Krull dimension of a compactification of X over B , which does not depend on the choice of compactifications. For an integer $q \geq 0$, we define X_q as the set of all points x on X with $\dim'(\overline{\{x\}}) = q$, where $\overline{\{x\}}$ is the closure of x in X . Note that a point x on X_η (resp. on X_s) belongs to X_q if and only if the residue field $\kappa(x)$ satisfies

$$\text{trdeg}_k(\kappa(x)) = q - 1 \quad (\text{resp. } \text{trdeg}_F(\kappa(x)) = q).$$

One can easily check that

$$(1.1) \quad X_q \cap Y = Y_q \quad \text{for } Y \text{ locally closed in } X.$$

Let \mathcal{C} be the full subcategory of Sch_B^{qp} consisting of objects whose structural morphisms do not factor through η . For $X \in \text{Ob}(\mathcal{C})$, $\dim'(X)$ agrees with the Krull dimension of X , and we simply denote $\dim'(X)$ by $\dim(X)$.

Definition 1.2. For $X \in \text{Ob}(\mathcal{C})$ and $q \in \mathbb{Z}_{\geq 0}$, let $\mathcal{Z}_q(X)$ be the free abelian group over the set X_q . We define the rational equivalence relation on $\mathcal{Z}_q(X)$ in the same way as in [Ful98, §1.3], and define the group $\text{CH}_q(X)$ as the quotient of $\mathcal{Z}_q(X)$ by this equivalence relation. We call $\text{CH}_q(X)$, the Chow group of q -cycles modulo rational equivalence. This group is identified with the cokernel of the divisor map defined in [Ful98, §1.2]

$$\text{div} : \bigoplus_{x \in X_{q+1}} \kappa(x)^\times \longrightarrow \bigoplus_{x \in X_q} \mathbb{Z}.$$

Remark 1.3. For a scheme $X \in \text{Ob}(\mathcal{C})$ whose structural morphism factors through $s \hookrightarrow B$, the above definition of the Chow groups agrees with that in [Ful98, §1.3]. We did not include schemes over η in the category \mathcal{C} to avoid confusion caused by the fact that for $X \in \text{Ob}(\text{Sch}_B^{qp})$ lying over η , our notation $\text{CH}_q(X)$ corresponds to the usual Chow group of $(q-1)$ -cycles. We will not deal with Chow groups of any schemes which do not belong to \mathcal{C} until the end of Section 8.

Throughout this paper we fix a prime number ℓ different from $\text{ch}(F)$ and write

$$\Lambda_n = \mathbb{Z}/\ell^n \mathbb{Z} \quad \text{for an integer } n \geq 1 \quad \text{and} \quad \Lambda_\infty = \mathbb{Q}_\ell/\mathbb{Z}_\ell = \varinjlim_m \Lambda_n.$$

Let Λ denote either Λ_n or Λ_∞ . For $X \in \text{Ob}(\text{Sch}_B^{qp})$ and $q \in \mathbb{Z}$, we define

$$(1.4) \quad H_q(X, \Lambda) := H^{2-q}(X, Rf^! \Lambda)$$

where f denotes the structural morphism $X \rightarrow B$ and $Rf^!$ denotes Deligne's twisted inverse image functor [SGA4, XVIII, Th. 3.1.4]. This assignment gives a homology theory on \mathcal{C} in the sense of [JS03, Def. 2.1]. More precisely, we have the following proposition:

PROPOSITION 1.5. *Let Sch_{B*}^{qp} be the category whose objects are the same as Sch_B^{qp} and whose morphisms are proper B -morphisms. Then $H_q(-, \Lambda)$ is a covariant functor from Sch_{B*}^{qp} to the category of abelian groups for each $q \in \mathbb{Z}$, and satisfies the following (1)–(3):*

- (1) $H_q(-, \Lambda)$ is contravariantly functorial with respect to étale morphisms in Sch_B^{qp} .
- (2) If $i : Y \hookrightarrow X$ is a closed immersion in Sch_B^{qp} with open complement $j : U \hookrightarrow X$, then there is a long exact sequence

$$\cdots \xrightarrow{\delta} H_q(Y, \Lambda) \xrightarrow{i_*} H_q(X, \Lambda) \xrightarrow{j^*} H_q(U, \Lambda) \xrightarrow{\delta} H_{q-1}(Y, \Lambda) \longrightarrow \cdots,$$

which we call a localization sequence in what follows.

- (3) *The localization sequence in (2) is functorial with respect to proper morphisms and étale morphisms in the following sense. Suppose we are given cartesian squares in Sch_B^{qp}*

$$\begin{array}{ccccc} U' & \xrightarrow{j'} & X' & \xleftarrow{i'} & Y' \\ f_U \downarrow & \square & f \downarrow & \square & f_Y \downarrow \\ U & \xrightarrow{j} & X & \xleftarrow{i} & Y, \end{array}$$

where i (resp. j) is a closed (resp. open) immersion and $U = X - Y$. If f is proper, then there is a commutative diagram with exact rows

$$\begin{array}{ccccccc} \cdots & \xrightarrow{\delta} & H_q(Y', \Lambda) & \xrightarrow{i'_*} & H_q(X', \Lambda) & \xrightarrow{(j')^*} & H_q(U', \Lambda) \xrightarrow{\delta} H_{q-1}(Y', \Lambda) \longrightarrow \cdots \\ & & f_{Y*} \downarrow & & f_* \downarrow & & f_{U*} \downarrow & f_{Y*} \downarrow \\ \cdots & \xrightarrow{\delta} & H_q(Y, \Lambda) & \xrightarrow{i_*} & H_q(X, \Lambda) & \xrightarrow{j^*} & H_q(U, \Lambda) \xrightarrow{\delta} H_{q-1}(Y, \Lambda) \longrightarrow \cdots \end{array}$$

If f is étale, then there is a similar commutative diagram with vertical arrows f_Y^*, f^*, f_U^* in the opposite direction.

Proof. An outline of the proof of these statements is given in [JS03, Ex. 2.2]. The details are straightforward and left to the reader. We recall here only the definition of the map $g_* : H_q(Y', \Lambda) \rightarrow H_q(Y, \Lambda)$ for a proper morphism $g : Y' \rightarrow Y$ in Sch_B^{qp} , which will be useful later in Lemma 1.8. Let β and β' be the structural morphisms $Y \rightarrow B$ and $Y' \rightarrow B$, respectively. Then g_* is defined as the composite

$$\begin{aligned} g : H_q(Y', \Lambda) &= H^{2-q}(Y', R(\beta')^! \Lambda_B) = H^{2-q}(Y, Rg_* R(\beta')^! \Lambda_B) \\ (1.6) \quad &= H^{2-q}(Y, Rg_* Rg^! R\beta^! \Lambda_B) \xrightarrow{\text{ad}_g} H^{2-q}(Y, R\beta^! \Lambda_B) = H_q(Y, \Lambda), \end{aligned}$$

where ad_g denotes the adjunction map $Rg_* Rg^! \rightarrow \text{id}$ obtained by the properness of g (i.e., $Rg_* = Rg_!$) and the adjointness between $Rg_!$ and $Rg^!$. \square

Remark 1.7. For $X \in \text{Ob}(\mathcal{C})$ whose structural morphism f factors as $X \xrightarrow{g} s \xrightarrow{i} B$, we have

$$H_q(X, \Lambda) = H^{2-q}(X, Rf^! \Lambda_B) = H^{2-q}(X, Rg^! Ri^! \Lambda_B) \simeq H^{-q}(X, Rg^! \Lambda(-1)_s),$$

where the last isomorphism follows from the purity for discrete valuation rings [SGA5, I, Th. 5.1]. The last group is a special case of the Bloch-Ogus homology [BO74, (2.1)].

We next provide the following lemma:

LEMMA 1.8. *Assume that $X \in \text{Ob}(\text{Sch}_B^{qp})$ is integral and regular, and put $d := \dim'(X) - 1$. For $m \in \mathbb{Z}$, let $\Lambda(m)$ be the Tate twist. Then for a closed*

subscheme $Y \subset X$, there is a canonical isomorphism

$$(1.9) \quad \tau_{Y,X} : H_Y^{2d-q+2}(X, \Lambda(d)) \xrightarrow{\simeq} H_q(Y, \Lambda)$$

which satisfies the following functorial properties:

- (1) Let $f : X' \rightarrow X$ be a proper morphism (resp. étale morphism) in Sch_B^{qp} with X' integral and regular, and let Y' be a closed subscheme of $f^{-1}Y = X' \times_X Y$ (resp. put $Y' := f^{-1}Y$). Let g be the composite map $Y' \rightarrow f^{-1}Y \rightarrow Y$. Then there is a commutative diagram

$$\begin{array}{ccc} H_{Y'}^{2d'-q+2}(X', \Lambda(d')) & \xrightarrow[\simeq]{\tau_{Y',X'}} & H_q(Y', \Lambda) \\ f_* \downarrow & & \downarrow g_* \\ H_Y^{2d-q+2}(X, \Lambda(d)) & \xrightarrow[\simeq]{\tau_{Y,X}} & H_q(Y, \Lambda) \end{array}$$

$$\left(\begin{array}{ccc} H_{Y'}^{2d-q+2}(X', \Lambda(d)) & \xrightarrow[\simeq]{\tau_{Y',X'}} & H_q(Y', \Lambda) \\ \text{resp.} & f^* \uparrow & \uparrow g^* \\ H_Y^{2d-q+2}(X, \Lambda(d)) & \xrightarrow[\simeq]{\tau_{Y,X}} & H_q(Y, \Lambda) \end{array} \right),$$

where d' in the left square denotes $\dim'(X') - 1$ and the arrow f_* in the left square is the push-forward map of cohomology groups with supports (see the proof below).

- (2) Let i be the closed immersion $Y \hookrightarrow X$ and let j be the open complement $U \hookrightarrow X$. Then there is a commutative diagram with exact rows

$$\begin{array}{ccccccc} \cdots & \xrightarrow{\delta} & H_Y^r(X, \Lambda(d)) & \xrightarrow{i_*} & H^r(X, \Lambda(d)) & \xrightarrow{j^*} & H^r(U, \Lambda(d)) \xrightarrow{\delta} H_Y^{r+1}(X, \Lambda(d)) \longrightarrow \cdots \\ & & \tau_{Y,X} \downarrow \simeq & & \tau_{X,X} \downarrow \simeq & & \tau_{U,U} \downarrow \simeq & & \tau_{Y,X} \downarrow \\ \cdots & \xrightarrow{\delta} & H_q(Y, \Lambda) & \xrightarrow{i_*} & H_q(X, \Lambda) & \xrightarrow{j^*} & H_q(U, \Lambda) & \xrightarrow{\delta} & H_{q-1}(Y, \Lambda) \longrightarrow \cdots, \end{array}$$

where we put $r := 2d - q + 2$, for simplicity.

Proof. Let $D(X_{\text{ét}})$ be the derived category of complexes of étale sheaves on X , and let α be the structural morphism $X \rightarrow B$. We will prove

- (3) There is a canonical isomorphism $T_\alpha : \Lambda(d)_X[2d] \xrightarrow{\simeq} R\alpha^! \Lambda_B$ in $D(X_{\text{ét}})$.

We first finish the proof of the lemma admitting this statement. We define the isomorphism $\tau_{Y,X}$ in (1.9) as that induced by T_α . The right square in (1) is obviously commutative. We obtain the commutative diagram in (2) by applying a standard localization argument to T_α (cf. [JSS, (0.4.2)]). For a proper morphism $f : X' \rightarrow X$ as in (1), we define the morphism

$$\text{Tr}_f : Rf_* \Lambda(d')_X[2d'] \rightarrow \Lambda(d)_X[2d] \quad \text{in } D(X_{\text{ét}})$$

by the commutative square

$$\begin{array}{ccc} Rf_* \Lambda(d')_X[2d'] & \xrightarrow[\simeq]{T_{\alpha'}} & Rf_* R(\alpha')^! \Lambda_B \\ \text{Tr}_f \downarrow & & \downarrow \text{ad}_f \\ \Lambda(d)_X[2d] & \xrightarrow[\simeq]{T_\alpha} & R\alpha^! \Lambda_B, \end{array}$$

where α' denotes the structural morphism $X' \rightarrow B$, and ad_f denotes the adjunction map $Rf_* Rf^! \rightarrow \text{id}$ (see ad_g in (1.6)). We define the arrow f_* in the left square of (1) as the map induced by Tr_f and the cobase-change map $H_{Y'}^r(X', -) \rightarrow H_Y^r(X, Rf_*(-))$ [SGA4, XVIII, (3.1.13.2)]. In view of the definition of g_* in (1.6), f_* makes the left square of (1) commutative. Thus it remains to show (3), which we prove in what follows.

Since B is regular of dimension one, α is flat or otherwise factors as $X \xrightarrow{\beta} s \xrightarrow{i} B$. In the flat case, we define T_α as the adjoint of the trace morphism $\text{Tr}_\alpha : R\alpha^! \Lambda(d)_X[2d] \rightarrow \Lambda_B$ due to Deligne ([SGA4, XVIII, Th. 2.9]). In the latter case, we define T_α as the adjoint of the composite morphism

$$R\alpha^! \Lambda(d)_X[2d] = Ri_* R\beta^! \Lambda(d)_X[2d] \xrightarrow{\text{Tr}_\beta} Ri_* \Lambda(-1)_s[-2] \xrightarrow{\text{Gys}_i} \Lambda_B,$$

where the last arrow is the Gysin map for i ([Fuj02, §1.2]). We show that T_α is an isomorphism. Fix a locally closed embedding $\gamma : X \hookrightarrow \mathbb{P} = \mathbb{P}_B^N$ into a projective space over B . Let π be the natural projection $\mathbb{P} \rightarrow B$. Then by [SGA $\frac{1}{2}$, Cycle, Th. 2.3.8 (i)], T_α factors as

$$\Lambda(d)_X[2d] \xrightarrow{\text{Gys}_\gamma} R\gamma^! \Lambda(N)_\mathbb{P}[2N] \xrightarrow{T_\pi} R\gamma^! R\pi^! \Lambda_B = R\alpha^! \Lambda_B.$$

Here the arrow Gys_γ denotes the Gysin morphism for γ ([Fuj02, §1.2]), which is an isomorphism by the absolute purity (loc. cit. Theorem 2.2.1). The second arrow is an isomorphism by [SGA4, XVIII, Th. 3.2.5]. Hence T_α is an isomorphism and we obtain the assertion (3). This completes the proof of Lemma 1.8. \square

COROLLARY 1.10. *For $X \in \text{Ob}(\text{Sch}_B^{qp})$, there is a homologically graded spectral sequence*

$$E_{a,b}^1(X, \Lambda) = \bigoplus_{x \in X_a} H^{a-b}(x, \Lambda(a-1)) \implies H_{a+b}(X, \Lambda),$$

which we call the niveau spectral sequence. This spectral sequence is covariant with respect to proper morphisms in Sch_B^{qp} and contravariant with respect to étale morphisms in Sch_B^{qp} . See Figure 1 below for a table of E^1 -terms of this spectral sequence.

Proof. By Proposition 1.5 and [JS03, Prop. 2.7], there is a spectral sequence

$$E_{a,b}^1(X, \Lambda) = \bigoplus_{x \in X_a} H_{a+b}(x, \Lambda) \implies H_{a+b}(X, \Lambda),$$

Definition 1.2, by Lemma 1.8 (2) and [JSS, Th. 1.1.1]. We define the cycle class mapping

$$(1.12) \quad \rho_X : \mathrm{CH}_1(X) \otimes \Lambda \rightarrow H_2(X, \Lambda)$$

to be the edge homomorphism $E_{1,1}^2 \rightarrow H_2(X, \Lambda)$ of the niveau spectral sequence. By the functorial properties of the niveau spectral sequences, we obtain the following proposition, which will be frequently used in later sections:

PROPOSITION 1.13. (1) *The cycle map (1.12) is covariantly functorial with respect to proper morphisms in \mathcal{C} and contravariantly functorial with respect to étale morphisms in \mathcal{C} .*

(2) *For a closed immersion $i : Y \hookrightarrow X$ in \mathcal{C} whose open complement $j : U \hookrightarrow X$ belongs to \mathcal{C} , there is a commutative diagram with exact rows*

$$\begin{array}{ccccccc} \mathrm{CH}_1(Y) \otimes \Lambda & \xrightarrow{i_*} & \mathrm{CH}_1(X) \otimes \Lambda & \xrightarrow{j^*} & \mathrm{CH}_1(U) \otimes \Lambda & \longrightarrow & 0 \\ \downarrow \rho_Y & & \downarrow \rho_X & & \downarrow \rho_U & & \\ H_2(Y, \Lambda) & \xrightarrow{i_*} & H_2(X, \Lambda) & \xrightarrow{j^*} & H_2(U, \Lambda), & & \end{array}$$

where the lower row is the localization exact sequence in Proposition 1.5 (2).

The exactness of the upper row in the above diagram follows from the same argument as for [Ful98, Prop. 1.8]. The details are straightforward and left to the reader.

Remark 1.14. The results in Proposition 1.5 – Corollary 1.10 are covariantly functorial in Λ . In particular, for $X \in \mathrm{Ob}(\mathcal{C})$ and $m, n \geq 1$ there is a commutative diagram with exact rows

$$\begin{array}{ccccccc} \mathrm{CH}_1(X) \otimes \Lambda_m & \xrightarrow{\times \ell^n} & \mathrm{CH}_1(X) \otimes \Lambda_{m+n} & \xrightarrow{\mathrm{mod} \ell^n} & \mathrm{CH}_1(X) \otimes \Lambda_n & \longrightarrow & 0 \\ \downarrow \rho_X & & \downarrow \rho_X & & \downarrow \rho_X & & \\ H_2(X, \Lambda_m) & \xrightarrow{\times \ell^n} & H_2(X, \Lambda_{m+n}) & \xrightarrow{\mathrm{mod} \ell^n} & H_2(X, \Lambda_n). & & \end{array}$$

where the lower row is a part of the long exact sequence of homology groups associated with the short exact sequence of étale sheaves on B :

$$0 \longrightarrow \Lambda_m \xrightarrow{\times \ell^n} \Lambda_{m+n} \xrightarrow{\mathrm{mod} \ell^n} \Lambda_n \longrightarrow 0.$$

We introduce here the following terminology, in order to state a main result of this paper.

Definition 1.15. (1) For a regular scheme X and a reduced divisor D on X , we say that D has *simple normal crossings* if it satisfies the following conditions:

(1a) *The irreducible components D_i ($i \in I$) of D are regular.*

- (1b) For an arbitrary nonempty subset $J \subset I$, the scheme-theoretical intersection $\bigcap_{i \in J} D_i$ is regular of codimension $\sharp(J)$ in X , or otherwise empty.
- (2) We call an object $X \in \mathcal{C}$ *quasi-semistable*, if it satisfies the following conditions:
- (2a) X is pure-dimensional, regular and flat of finite type over B .
- (2b) The reduced divisor $X_{s,\text{red}}$ on X has simple normal crossings.
- Note that $X_{s,\text{red}}$ is nonempty by the definition of the category \mathcal{C} .
- (3) We define the category \mathcal{QS} (resp. \mathcal{QSP}) as the full subcategory of \mathcal{C} consisting of objects which are quasi-semistable (resp. quasi-semistable and projective over B).

The following theorem is one of the main results of this paper:

THEOREM 1.16 (Theorem 0.6). *Assume that R is excellent and henselian and that F is finite or separably closed. Then for $X \in \text{Ob}(\mathcal{QSP})$ and $n \geq 1$, the cycle map $\rho_X : \text{CH}_1(X)/\ell^n \rightarrow H_2(X, \Lambda_n)$ is bijective.*

The proof of Theorem 1.16 will be completed in Section 8. We note here the following lemma, where R may not be excellent or henselian and F may be arbitrary:

LEMMA 1.17. *If $X \in \text{Ob}(\mathcal{C})$ has dimension ≤ 2 , then the map $\rho_X : \text{CH}_1(X)/\ell^n \rightarrow H_2(X, \Lambda_n)$ is injective for any $n \geq 1$.*

Proof. This follows from the definition of ρ_X and the fact that $E_{a,b}^1(X) = 0$ for $a \geq 3$. \square

2. Kato homology

The notation remains as in Section 1. In this section, we assume that R is henselian, and introduce an arithmetic invariant associated to each object of \mathcal{C} , which plays a crucial role in the proof of Theorem 1.16.

Definition 2.1. Let X be an object of \mathcal{C} . Let Λ denote Λ_n or Λ_∞ . Let us recall the niveau spectral sequence obtained in Corollary 1.10:

$$(2.2) \quad E_{a,b}^1(X, \Lambda) = \bigoplus_{x \in X_a} H^{a-b}(x, \Lambda(a-1)) \implies H_{a+b}(X, \Lambda).$$

- (1) We define the complex $KC(X, \Lambda)$ as the following sequence of E^1 -terms:

$$\begin{array}{ccccccc} E_{0,0}^1(X, \Lambda) & \xleftarrow{d_{1,0}^1} & E_{1,0}^1(X, \Lambda) & \xleftarrow{d_{2,0}^1} \cdots \xleftarrow{d_{a,0}^1} & E_{a,0}^1(X, \Lambda) & \xleftarrow{d_{a+1,0}^1} \cdots, \\ \parallel & & \parallel & & \parallel & & \\ \bigoplus_{x \in X_0} H^0(x, \Lambda(-1)) & & \bigoplus_{x \in X_1} H^1(x, \Lambda) & & \bigoplus_{x \in X_a} H^a(x, \Lambda(a-1)) & & \end{array}$$

where $E_{a,0}^1(X, \Lambda)$ is placed in degree a . By Lemma 1.8 (2) and [JSS, Th. 1.1.1], the differential maps agree with the negative of the boundary maps of Galois

cohomology groups defined in [Kat86, §1]. In particular, $KC(X, \Lambda)$ is a special case of the Bloch-Ogus-Kato complexes defined in loc. cit., up to signs of differential maps.

- (2) We define the group $KH_a(X, \Lambda)$, the Kato homology of X , to be $E_{a,0}^2(X, \Lambda)$, which is by definition the homology group of the complex $KC(X, \Lambda)$ in degree a .

By the functorial properties of the niveau spectral sequences (cf. Corollary 1.10), the complex $KC(X, \Lambda)$ is covariantly functorial for proper morphisms in \mathcal{C} and contravariantly functorial for étale morphisms in \mathcal{C} . In particular, for a closed immersion $i : Y \hookrightarrow X$ in \mathcal{C} whose open complement $j : U \hookrightarrow X$ belongs to \mathcal{C} , there is an exact sequence of complexes (cf. (1.1))

$$(2.3) \quad 0 \longrightarrow KC(Y, \Lambda) \xrightarrow{i_*} KC(X, \Lambda) \xrightarrow{j^*} KC(U, \Lambda) \longrightarrow 0.$$

By Lemma 2.6 below, there is an edge homomorphism

$$(2.4) \quad \epsilon_a : H_a(X, \Lambda_\infty) \longrightarrow KH_a(X, \Lambda_\infty) \quad \text{if } F \text{ is finite,}$$

$$(2.5) \quad \epsilon_a : H_a(X, \Lambda_n) \longrightarrow KH_a(X, \Lambda_n) \quad \text{if } F \text{ is separably closed.}$$

LEMMA 2.6. *Assume that R is henselian. Let X be an object of \mathcal{C} .*

- (1) *If F is separably closed, then $E_{a,b}^1(X, \Lambda_n) = 0$ for $b \leq -1$.*
 (2) *If F is finite, then $E_{a,b}^1(X, \Lambda_n) = 0$ for $b \leq -2$ and $E_{a,-1}^1(X, \Lambda_\infty) = 0$.*

Proof. (1) and the first vanishing of (2) follow from the fact $\text{cd}(x) = a + \text{cd}(F)$ for $x \in X_a$. The second vanishing of (2) follows from the next lemma, which is an analogue of a result of Kahn [Kah93, p. 61, Prop. 4]. \square

LEMMA 2.7. *Assume that R is henselian and that F is finite. Let $r \geq 0$ be an integer. Let L be a field which is either finitely generated over k with $\text{trdeg}_k(L) = r - 1$ or finitely generated over F with $\text{trdeg}_F(L) = r$. Then we have $H^{r+1}(L, \Lambda_\infty(n)) = 0$ for any $n \neq r$. Here for a field K , $H^*(K, -)$ denotes the étale cohomology of $\text{Spec}(K)$, which agrees with the Galois cohomology of the absolute Galois group G_K (cf. [Mil80, III, Ex. 1.7]).*

Proof. We prove the case that L is over k . The case that L is over F is similar and left to the reader (see also Lemma 3.10 below). If L is a finite extension of k , then the assertion follows from the Tate duality for Galois cohomology of local fields (cf. [Ser92, II, §5.2, Th. 2]). We continue the proof by induction on $r = \text{trdeg}_k(L) + 1$. Assume $r \geq 2$. There is a subfield $L_{r-1} \subset L$ which is purely transcendental over k and for which L/L_{r-1} is finite. Take a subfield $L_{r-2} \subset L_{r-1}$ which is purely transcendental over k of degree $r - 2$, and let K be the algebraic separable closure of L_{r-2} in L . Note that $\text{trdeg}_K(L) = 1$ and that $\text{trdeg}_k(K) = r - 2$. Fix a separable closure \bar{K} of K . There is a Hochschild-Serre spectral sequence

$$E_2^{a,b} = H^a(K, H^b(L \otimes_K \bar{K}, \Lambda_\infty(n))) \Rightarrow H^{a+b}(L, \Lambda_\infty(n)).$$

Since $\text{cd}(K) \leq \text{trdeg}_k(K) + 2 = r$ ([Ser92, II, §§4.2–4.3]) and $\text{cd}(L \otimes_K \bar{K}) = 1$ (loc. cit. II, §§3.2–3.3), we have $E_2^{a,b} = 0$ if $b \geq 2$ or $a \geq r + 1$. Hence it suffices to show

$$E_2^{r,1} = H^r(K, H^1(L \otimes_K \bar{K}, \Lambda_\infty(n))) = 0.$$

By Hilbert's Theorem 90 (loc. cit. II, §1.2, Prop. 1), we have

$$\begin{aligned} H^r(K, H^1(L \otimes_K \bar{K}, \Lambda_\infty(n))) &= H^r(K, (L \otimes_K \bar{K})^\times \otimes \Lambda_\infty(n-1)) \\ &= \varinjlim_{K' \subset \bar{K}} H^r(K, (L \otimes_K K')^\times \otimes \Lambda_\infty(n-1)), \end{aligned}$$

where $K' \subset \bar{K}$ ranges over all finite Galois extensions of K . See [Ser92, II, §2.2, Prop. 8] for the last equality. We show that the last group is zero. Fix a finite Galois extension K' of K . Since $\text{cd}(K) \leq r$, the corestriction map

$$H^r(K', (L \otimes_K K')^\times \otimes \Lambda_\infty(n-1)) \longrightarrow H^r(K, (L \otimes_K K')^\times \otimes \Lambda_\infty(n-1))$$

is surjective. Because $G_{K'}$ acts on $(L \otimes_K K')^\times$ trivially, the natural map induced by cup-product

$$(L \otimes_K K')^\times \otimes H^r(K', \Lambda_\infty(n-1)) \longrightarrow H^r(K', (L \otimes_K K')^\times \otimes \Lambda_\infty(n-1))$$

is bijective. Here the left-hand side is zero by the induction hypothesis. Thus we obtain Lemmas 2.7 and 2.6. \square

Assume that F is finite and put $\Lambda := \Lambda_\infty$ until we state Conjecture 2.11. Let $D(\mathbb{Z}_\ell)$ be the derived category of complexes of \mathbb{Z}_ℓ -modules. For $X \in \text{Ob}(\mathcal{QSP})$, we construct a morphism

$$(2.8) \quad KC(X, \Lambda) \longrightarrow \Lambda^{I(X_s)}[-1] \quad \text{in } D(\mathbb{Z}_\ell).$$

Here $I(X_s)$ denotes the set of the irreducible components of $X_{s,\text{red}}$ and $\Lambda^{I(X_s)}[-1]$ is the homological complex with $\Lambda^{I(X_s)}$ placed in degree 1 and with 0 in the other degrees. We show that the first boundary map of $KC(X, \Lambda)$ is surjective:

$$d_{1,0}^1: \bigoplus_{y \in X_1} H^1(y, \Lambda) \twoheadrightarrow \bigoplus_{x \in X_0} H^0(x, \Lambda(-1)).$$

Indeed, for a given point $x \in X_0$ there exists a closed point y on X_η whose closure $\overline{\{y\}} \subset X$ contains x and is regular at x by Lemma 7.2 below. Because R is henselian, $\{y\}$ is the spectrum of a henselian discrete valuation ring. By Hensel's lemma, all roots of unity of order prime to $\text{ch}(\kappa(x))$ in $\kappa(x)$ lift to $\kappa(y)$, and one can easily check the (y, x) -factor of $d_{1,0}^1$ is surjective by [JSS, Th. 1.1.1]. Thus $d_{1,0}^1$ is surjective, and we get a natural morphism

$$(2.9) \quad KC(X, \Lambda) \longrightarrow H_1(KC(X, \Lambda))[-1] \simeq H_1(X, \Lambda)[-1] \quad \text{in } D(\mathbb{Z}_\ell),$$

where the second isomorphism is given by the map (2.4) with $a = 1$ (cf. Figure 1 after Corollary 1.10 and Lemma 2.6). On the other hand, there is a composite map (2.10)

$$H_1(X, \Lambda) \xrightarrow{(1.9)} H^{2d+1}(X, \Lambda(d)) \xrightarrow{(*)} \bigoplus_{1 \leq j \leq m} H^{2d+1}(Y_j, \Lambda(d)) \xrightarrow{\simeq} \Lambda^{I(X_s)},$$

where d denotes $\dim(X) - 1 = \dim(X_s)$ and Y_1, \dots, Y_m are the irreducible components of $X_{s,\text{red}}$. The arrow $(*)$ is the natural pull-back map, and the last arrow is the trace isomorphisms for projective smooth varieties over F (cf. [JS03, Lemma 5.3 (2)]). We define the morphism (2.8) as the composite of (2.9) and (2.10).

We now propose the following conjectures:

CONJECTURE 2.11. *If R is excellent and henselian and F is separably closed, then $KC(X, \Lambda_n)$ is exact for any $X \in \text{Ob}(\mathcal{QSP})$.*

CONJECTURE 2.12 (Conjecture 0.7). *If R is excellent and henselian and F is finite, then for any $X \in \text{Ob}(\mathcal{QSP})$, the morphism (2.8) is an isomorphism, i.e., we have*

$$KH_a(X, \Lambda_\infty) \simeq \begin{cases} \Lambda_\infty^{I(X_s)} & a = 1 \\ 0 & a \neq 1. \end{cases}$$

Recall that $KC(X, \Lambda)$ is identified with the case $j = -1$ of the Bloch-Ogus-Kato complex

$$\begin{aligned} \bigoplus_{x \in X_0} H^{j+1}(x, \Lambda(j)) &\leftarrow \bigoplus_{x \in X_1} H^{j+2}(x, \Lambda(j+1)) \\ &\leftarrow \dots \leftarrow \bigoplus_{x \in X_a} H^{a+j+1}(x, \Lambda(a+j)) \leftarrow \dots \end{aligned}$$

up to signs of differential maps (cf. Definition 2.1). In [Kat86, Conj. 5.1], he conjectures that this complex with $j = 0$ is exact for both $\Lambda = \Lambda_n$ and Λ_∞ , if F is finite and X is a regular scheme which is proper flat over B . It is interesting to ask if Conjecture 2.12 is valid for Λ_n . We will prove the following affirmative result on these conjectures using Theorem 1.16.

THEOREM 2.13 (Theorem 0.8). *Conjectures 2.11 and 2.12 hold in degree $a \leq 3$.*

The proof of this theorem will be given in Section 9 below. In the rest of this section, we prove the following proposition, where R may not be excellent:

PROPOSITION 2.14. *Conjectures 2.11 and 2.12 hold in degree $a \leq 1$.*

Proof. Let Λ denote Λ_n if F is separably closed and Λ_∞ if F is finite. We have

$$KH_a(X, \Lambda) = E_{a,0}^2(X, \Lambda) = E_{a,0}^\infty(X, \Lambda) = H_a(X, \Lambda) \quad \text{for } a \leq 1$$

by Lemma 2.6. Hence the proposition follows from the following lemma.

LEMMA 2.15. Assume R is henselian, and let X be an object of \mathcal{QSP} .

- (1) If F is separably closed, then $H_a(X, \Lambda)$ is zero for $a \leq 1$.
- (2) If F is finite, then the composite map (2.10) is bijective and $H_a(X, \Lambda)$ is zero for $a \leq 0$.

Proof of Lemma 2.15. We prove only the case $a = 1$ of (2), i.e., the bijectivity of the arrow $(*)$ in (2.10). The other assertions follow from arguments similar to those below and are left to the reader. The map $(*)$ in question factors as follows

$$\begin{aligned} H^{2d+1}(X, \Lambda(d)) &\longrightarrow H^{2d+1}(X_s, \Lambda(d)) \\ &\longrightarrow H^{2d+1}(X_{s,\text{red}}, \Lambda(d)) \xrightarrow{(**)} \bigoplus_{1 \leq j \leq m} H^{2d+1}(Y_j, \Lambda(d)), \end{aligned}$$

where all arrows are pull-back maps. Since R is henselian, the first arrow is bijective by the proper base-change theorem [SGA4, XII Cor. 5.5 (iii)]. The second arrow is bijective by the invariance of étale topology for the nilpotent closed immersion $X_{s,\text{red}} \hookrightarrow X_s$ (loc. cit. VIII, Cor. 1.2). We show that the last arrow $(**)$ is bijective. Let ν be the canonical map $\coprod_{i=1}^m Y_j \rightarrow X_{s,\text{red}}$, which is finite because each component $Y_j \rightarrow X_{s,\text{red}}$ is a closed immersion. Let Σ be the singular locus of $X_{s,\text{red}}$. As ν is an isomorphism over $X_{s,\text{red}} - \Sigma$, the adjunction map $\alpha : \Lambda(d) \rightarrow \nu_* \nu^* \Lambda(d)$ of étale sheaves on $X_{s,\text{red}}$ is injective, and the support of $\text{Coker}(\alpha)$ is contained in Σ . Hence we have a long exact sequence

$$\begin{aligned} \cdots \longrightarrow H^{2d}(\Sigma, \text{Coker}(\alpha)) &\longrightarrow H^{2d+1}(X_{s,\text{red}}, \Lambda(d)) \xrightarrow{(**)} \bigoplus_{1 \leq j \leq m} H^{2d+1}(Y_j, \Lambda(d)) \\ &\longrightarrow H^{2d+1}(\Sigma, \text{Coker}(\alpha)) \longrightarrow \cdots, \end{aligned}$$

and the map $(**)$ is bijective by the fact $\text{cd}_\ell(\Sigma) \leq 2 \dim(\Sigma) + \text{cd}_\ell(F) = 2d - 1$ (loc. cit. X, Cor. 4.3). Thus we obtain the lemma and Proposition 2.14. \square

3. The vanishing theorem

Let the notation be as in Section 1. For a scheme X , $D(X_{\text{ét}})$ denotes the derived category of complexes of étale sheaves on X .

Definition 3.1. See Definition 1.15 (3) for the categories \mathcal{QS} and \mathcal{QSP} .

- (1) A \mathcal{QS} -pair is a pair (X, Y) consisting of schemes $X, Y \in \text{Ob}(\mathcal{QS})$ such that Y is a divisor on X and such that the reduced divisor $X_{s,\text{red}} \cup Y$ on X has simple normal crossings. Note that no component of Y is a component of $X_{s,\text{red}}$, because Y is flat over B by assumption.
- (2) A \mathcal{QSP} -pair is a \mathcal{QS} -pair (X, Y) such that $X, Y \in \text{Ob}(\mathcal{QSP})$. For a \mathcal{QSP} -pair (X, Y) , we call $U := X - Y$ the complement of (X, Y) , and often denote it by $(X, Y; U)$.
- (3) An *ample* \mathcal{QSP} -pair is a \mathcal{QSP} -pair $(X, Y; U)$ such that U is affine.

The aim of this section is to prove the following vanishing result:

THEOREM 3.2. *Let $(X, Y; U)$ be an ample \mathcal{LSP} -pair with $\dim(X) = d + 1 \geq 2$, and assume that R is henselian.*

- (1) *If F is separably closed, then $H_q(U, \Lambda_n) = 0$ for $q \leq d + 1$.*
- (2) *If F is finite, then $H_q(U, \Lambda_n) = 0$ for $q \leq d$. Furthermore if $d \geq 2$, then $H_{d+1}(U, \Lambda_\infty) = 0$.*

Remark 3.3. For a regular scheme U which is flat of relative dimension d over B , we have

$$H_q(U, \Lambda) \simeq H^{2d+2-q}(U, \Lambda(d))$$

by (1.9). When U is affine and R is henselian, the affine Lefschetz theorems of Artin-Gabber (cf. [Fuj02, §5]) imply that the last group vanishes if F is separably closed and $q \leq d$, or if F is finite and $q \leq d - 1$. Theorem 3.2 asserts a stronger vanishing result, assuming that U is the complement of an ample \mathcal{LSP} -pair (X, Y) . For example, assume F is separably closed, and consider the case that $X = \text{Proj}(R[T_0, T_1])$ and that $Y \subset X$ is the ample divisor $T_1^m - \pi \cdot T_0^m = 0$ ($m \geq 1$), where π is a prime element of R . We compute the natural map α in the exact sequence

$$H_Y^2(X, \Lambda_n(1)) \xrightarrow{\alpha} H^2(X, \Lambda_n(1)) \longrightarrow H^2(U, \Lambda_n(1)) \quad (U := X - Y).$$

The group $H_Y^2(X, \Lambda_n(1))$ is generated by the divisor class $\text{cl}_X(Y)$ [SGA $_{\frac{1}{2}}$, Cycle Prop. 2.1.4]. Consider the isomorphism

$$\beta : H^2(X, \Lambda_n(1)) \xrightarrow{\sim} H^2(X_s, \Lambda_n(1)) \xrightarrow{\sim} \Lambda_n,$$

where the first isomorphism is obtained by the proper base change theorem for étale cohomology and the second is the trace map for $X_s \rightarrow s$ (cf. (9.3) below). Then we have $\beta\alpha(\text{cl}_X(Y)) = m \in \Lambda_n$ by Lemma 9.4 below. Hence $H^2(U, \Lambda_n(1))$ is nonzero if ℓ divides m . On the other hand, Theorem 3.2 asserts that $H^2(U, \Lambda_n(1))$ is zero if (X, Y) is a \mathcal{LSP} -pair, i.e., $m = 1$.

The following lemma is crucial to the proof of Theorem 3.2, which extends the base-change theorem of Rapoport-Zink ([RZ82, Satz 2.19]) to \mathcal{LSP} -pairs by the absolute purity theorem of Gabber [Fuj02, Th. 2.1.1]. For a scheme $X \in \text{Ob}(\mathcal{C})$ and an effective Cartier divisor $Y \subset X$, let $\text{cl}_X(Y) \in H_Y^2(X, \Lambda_n(1))$ be the divisor class of Y ([SGA $_{\frac{1}{2}}$, Cycle §2.1], [Fuj02, §1.1]), which induces a morphism $\Lambda_n(-1)_Y[-2] \rightarrow R\kappa^! \Lambda_{n,X}$ in $D(Y_{\text{ét}})$ ($\kappa : Y \hookrightarrow X$) by the natural identification (cf. [SGA $_{\frac{1}{2}}$, Catégories Dérivées II.2.3 (4)])

$$H_Y^2(X, \Lambda_n(1)) = \text{Hom}_{D(Y_{\text{ét}})}(\Lambda_n(-1)_Y[-2], Ri^! \Lambda_{n,X}).$$

LEMMA 3.4. *Let (X, Y) be a \mathcal{LSP} -pair over B . Put $Z := X_{s,\text{red}}$ and $W := Y_{s,\text{red}}$. Then:*

- (1) W is an effective Cartier divisor on Z and the divisor class

$$\mathrm{cl}_Z(W) \in H_W^2(Z, \Lambda_n(1))$$

induces an isomorphism $H^{q-2}(W, \Lambda_n(j-1)) \simeq H_W^q(Z, \Lambda_n(j))$ for any $q, j \in \mathbb{Z}$.

- (2) Assume further that R is henselian and that $(X, Y; U)$ is a \mathcal{QSP} -pair over B . Put $V := U_{s, \text{red}}$. Then the pull-back map $H^q(U, \Lambda_n(j)) \rightarrow H^q(V, \Lambda_n(j))$ is bijective for any $q, j \in \mathbb{Z}$.

Proof. Put $\Lambda := \Lambda_n$. We first show (1). Since (X, Y) is a \mathcal{QS} -pair by assumption, the following square consisting of canonical closed immersions is cartesian:

$$(3.5) \quad \begin{array}{ccc} W & \xrightarrow{f} & Z \\ i_Y \downarrow & \square & \downarrow i_X \\ Y & \xrightarrow{\kappa} & X. \end{array}$$

In particular, W is an effective Cartier divisor on Z and the divisor class $\mathrm{cl}_Z(W)$ is defined in $H_W^2(Z, \Lambda(1))$. Our task is to show that the morphism

$$\mathrm{Gys}_f : \Lambda(-1)_W[-2] \longrightarrow Rf^! \Lambda_Z \quad \text{in } D(W_{\text{ét}})$$

induced by $\mathrm{cl}_Z(W)$ is an isomorphism. For $q \geq 1$, let $Z^{(q)}$ be the disjoint union of q -fold intersections of distinct irreducible components of Z . We define $W^{(q)}$ ($q \geq 1$) in a similar way. Since X and Y belong to $\mathrm{Ob}(\mathcal{QS})$, each of $Z^{(q)}$ and $W^{(q)}$ is empty or otherwise regular of pure dimension $\dim(Z) + 1 - q$ and $\dim(W) + 1 - q$, respectively. Since (X, Y) is a \mathcal{QS} -pair, one can easily check

$$(3.6) \quad W^{(q)} = Z^{(q)} \times_Z W \quad \text{for any } q \geq 1.$$

Let $u_q : Z^{(q)} \rightarrow Z$ and $v_q : W^{(q)} \rightarrow W$ be the canonical finite morphisms. There is an exact sequence of étale sheaves on Z

$$(3.7) \quad 0 \longrightarrow \Lambda_Z \xrightarrow{d^0} u_{1*} \Lambda_{Z^{(1)}} \xrightarrow{d^1} u_{2*} \Lambda_{Z^{(2)}} \xrightarrow{d^2} \cdots \xrightarrow{d^{q-1}} u_{q*} \Lambda_{Z^{(q)}} \xrightarrow{d^q} \cdots,$$

where d^0 is the adjunction map and d^q for $q \geq 1$ is an alternating sum of adjunction maps. The signs of these alternating sums are defined as follows. Let Z_1, \dots, Z_m be the irreducible components of Z , and let S and T be the components of $Z^{(q)}$ and $Z^{(q+1)}$ given by

$$S = Z_{i_1} \cap Z_{i_2} \cap \cdots \cap Z_{i_q} \quad \text{and} \quad T = S \cap Z_{i_{q+1}},$$

where i_j 's ($1 \leq j \leq q+1$) are pairwise distinct and we assumed $i_1 < i_2 < \cdots < i_q$. Then the sign of the (S, T) -factor of d^q is defined as $(-1)^{q-r}$, where r is the integer with $i_r < i_{q+1} < i_{r+1}$. The exact sequence (3.7) yields a spectral sequence of sheaves on W (cf. [Mil80, App. C (g)])

$$E_1^{a,b} = R^b f^! u_{a*} \Lambda_{Z^{(a)}} \implies R^{a+b-1} f^! \Lambda_Z.$$

We compute the sheaf $E_1^{a,b}$. Since u_a and v_a are finite, $R^b f^! u_{a*} = v_{a*} R^b (f^{(a)})^!$ by (3.6), where $f^{(a)}$ denotes the closed immersion $W^{(a)} \hookrightarrow Z^{(a)}$. Hence,

$$E_1^{a,b} \simeq \begin{cases} v_{a*} \Lambda(-1)_{W^{(a)}} & (\text{if } b = 2 \text{ and } a \geq 1) \\ 0 & (\text{otherwise}) \end{cases}$$

by the absolute purity [Fuj02, Th. 2.1.1], where the isomorphism for $b = 2$ arises from the isomorphism $\Lambda(-1)_{W^{(a)}} \xrightarrow{\cong} R^2 (f^{(a)})^! \Lambda_{Z^{(a)}}$ induced by the divisor class $\text{cl}_{Z^{(a)}}(W^{(a)}) \in H_{W^{(a)}}^2(Z^{(a)}, \Lambda(1))$. In particular, $R^q f^! \Lambda_Z = E_2^{q-1,2}$ for any $q \in \mathbb{Z}$. Finally we compute the sheaves $E_2^{*,2}$ by the following diagram of sheaves:

$$\begin{array}{ccccccc} 0 \longrightarrow & R^2 f^! \Lambda_Z \longrightarrow & E_1^{1,2} & \xrightarrow{d_1^{1,2}} & E_1^{2,2} & \xrightarrow{d_1^{2,2}} & \cdots \\ & \text{Gys}_f \uparrow & \parallel & & \parallel & & \\ 0 \longrightarrow & \Lambda(-1)_W \xrightarrow{d^0} v_{1*} \Lambda(-1)_{W^{(1)}} \xrightarrow{d^1} v_{2*} \Lambda(-1)_{W^{(2)}} \xrightarrow{d^2} \cdots & & & & & \\ & & & \cdots \xrightarrow{d_1^{q-1,2}} & E_1^{q,2} & \xrightarrow{d_1^{q,2}} & \cdots \\ & & & & \parallel & & \\ & & & \cdots \xrightarrow{d^{q-1}} & v_{q*} \Lambda(-1)_{W^{(q)}} \xrightarrow{d^q} \cdots & & \end{array}$$

The lower row is an exact sequence analogous to (3.7) and the signs of the alternating sum in d^q ($q \geq 1$) are defined by those in (3.7) and the isomorphism (3.6). We prove these squares are commutative, which implies Lemma 3.4 (1). Indeed, for any $q \geq 1$ the pull-back map $H_W^2(Z, \Lambda(1)) \rightarrow H_{W^{(q)}}^2(Z^{(q)}, \Lambda(1))$ sends $\text{cl}_Z(W)$ to $\text{cl}_{Z^{(q)}}(W^{(q)})$ by (3.6) and [Fuj02, Prop. 1.1.3], which shows the commutativity in question. Thus we obtain Lemma 3.4 (1).

We prove Lemma 3.4 (2). Since $W = Y \times_X Z$ by (3.5), the pull-back map $H_Y^2(X, \Lambda(1)) \rightarrow H_W^2(Z, \Lambda(1))$ sends $\text{cl}_X(Y)$ to $\text{cl}_Z(W)$ by [Fuj02, Prop. 1.1.3], which yields a commutative diagram in $D(W_{\text{ét}})$

$$(3.8) \quad \begin{array}{ccc} i_Y^* R\kappa^! \Lambda_X & \longleftarrow & i_Y^* \Lambda(-1)_Y[-2] \\ \downarrow & & \parallel \\ Rf^! \Lambda_Z & \xleftarrow{\text{Gys}_f} & \Lambda(-1)_W[-2]. \end{array}$$

Here the left vertical arrow is the base-change morphism associated to the square (3.5) (cf. [SGA4, XVIII, 3.1.14.2]). The top arrow is the morphism induced by $\text{cl}_X(Y)$, which is an isomorphism by the absolute purity [Fuj02, Th. 2.1.1]. The bottom arrow is an isomorphism by (1). Hence the left vertical arrow is an isomorphism as well. Since R is henselian, this fact implies Lemma 3.4 (2) by similar arguments as for [RZ82, Lemma 2.18] and the invariance of étale topology for

nilpotent closed immersions [SGA4, VIII, Cor. 1.2]. The details are straightforward and left to the reader. \square

We start the proof of Theorem 3.2. Put $Z := X_{s,\text{red}}$, $W := Y_{s,\text{red}}$ and $V := U_{s,\text{red}}$. By (1.9) and Lemma 3.4 (2), we have

$$H_q(U, \Lambda_n) \simeq H^{2d+2-q}(U, \Lambda_n(d)) \simeq H^{2d+2-q}(V, \Lambda_n(d)).$$

By the ampleness assumption, the F -variety V is affine of dimension d . By the affine Lefschetz theorem ([SGA4, XIV, Cor. 3.2]), we have

$$(3.9) \quad H^{2d+2-q}(V, \Lambda_n(d)) = 0 \quad \begin{cases} \text{if } F \text{ is separably closed and } q \leq d+1 \\ \text{or if } F \text{ is finite and } q \leq d. \end{cases}$$

It remains to show $H^{d+1}(V, \Lambda_\infty(d))$ is zero when we assume that F is finite and $d \geq 2$. There is a long exact sequence of étale cohomology groups

$$\begin{aligned} \cdots \rightarrow H^r(V, \Lambda_m(d)) &\rightarrow H^r(V, \Lambda_{m+n}(d)) \\ &\rightarrow H^r(V, \Lambda_n(d)) \rightarrow H^{r+1}(V, \Lambda_m(d)) \rightarrow \cdots \end{aligned}$$

Because these groups are finite by [SGA $\frac{1}{2}$, Th. finitude, Cor. 1.10], we obtain the following long exact sequence by standard projective and inductive limit arguments:

$$\begin{aligned} \cdots \rightarrow H^r(V, \mathbb{Z}_\ell(d)) &\rightarrow H^r(V, \mathbb{Q}_\ell(d)) \\ &\rightarrow H^r(V, \Lambda_\infty(d)) \rightarrow H^{r+1}(V, \mathbb{Z}_\ell(d)) \rightarrow \cdots \end{aligned}$$

Since $H^{d+2}(V, \mathbb{Z}_\ell(d)) = 0$ by (3.9), the map

$$H^{d+1}(V, \mathbb{Q}_\ell(d)) \rightarrow H^{d+1}(V, \Lambda_\infty(d))$$

is surjective. Therefore it is enough to show $H^{d+1}(V, \mathbb{Q}_\ell(d))$ is zero, which we prove in what follows.

LEMMA 3.10. *For any integer $n \in \mathbb{Z}$, there is an isomorphism*

$$H^{d+1}(V, \mathbb{Q}_\ell(n)) \simeq \text{Coker}(1 - \varphi_s : H^d(\bar{V}, \mathbb{Q}_\ell(n)) \rightarrow H^d(\bar{V}, \mathbb{Q}_\ell(n))).$$

Here φ_s denotes the arithmetic Frobenius operator, i.e., the element of the absolute Galois group G_F that corresponds to the Frobenius substitution on a fixed algebraic closure \bar{F} of F . For a scheme $X \in \text{Ob}(\mathcal{C})$ which lies over $s = \text{Spec}(F)$, \bar{X} denotes the scalar extension $X \otimes_F \bar{F}$, which we regard as a G_F -scheme by its natural action on \bar{F} .

Proof of Lemma 3.10. There is a short exact sequence (cf. [CTSS83, p. 780, (28)])

$$0 \longrightarrow H^d(\bar{V}, \mathbb{Q}_\ell(n))_{G_F} \longrightarrow H^{d+1}(V, \mathbb{Q}_\ell(n)) \longrightarrow H^{d+1}(\bar{V}, \mathbb{Q}_\ell(n))^{G_F} \longrightarrow 0,$$

where for a G_F -module M , M^{G_F} (resp. M_{G_F}) denotes the maximal subgroup (resp. maximal quotient group) of M on which G_F acts trivially. Since $H^{d+1}(\bar{V}, \mathbb{Q}_\ell(n))$

$= 0$ by (3.9), we have $H^{d+1}(V, \mathbb{Q}_\ell(n)) \simeq H^d(\bar{V}, \mathbb{Q}_\ell(n))_{G_F}$. The last group is identified with the cokernel of the endomorphism $1 - \varphi_s$ on $H^d(\bar{V}, \mathbb{Q}_\ell(n))$, because G_F is topologically generated by φ_s . \square

By Lemma 3.10, it suffices to show that the eigenvalues of φ_s on $H^d(\bar{V}, \mathbb{Q}_\ell(d))$ are different from 1. By Deligne's theorem on weights ([Del80, Th. I]) and the assumption that $d \geq 2$, the eigenvalues of φ_s on

$$H^d(\bar{Z}, \mathbb{Q}_\ell(d)) \quad \text{and} \quad H^{d-1}(\bar{W}, \mathbb{Q}_\ell(d-1))$$

are different from 1. It remains to show that there is a long exact sequence of G_F - \mathbb{Q}_ℓ -vector spaces

$$\begin{aligned} \cdots \rightarrow H^{r-2}(\bar{W}, \mathbb{Q}_\ell(d-1)) \rightarrow H^r(\bar{Z}, \mathbb{Q}_\ell(d)) \rightarrow H^r(\bar{V}, \mathbb{Q}_\ell(d)) \\ \rightarrow H^{r-1}(\bar{W}, \mathbb{Q}_\ell(d-1)) \rightarrow \cdots \end{aligned}$$

Indeed we have

$$H^{r-2}(\bar{W}, \mathbb{Q}_\ell(d-1)) \simeq H_{\bar{W}}^r(\bar{Z}, \mathbb{Q}_\ell(d))$$

by Lemma 3.4(1), which is G_F -equivariant because $\text{cl}_{\bar{Z}}(\bar{W}) \in H_{\bar{W}}^2(\bar{Z}, \mathbb{Z}_\ell(1))$ arises from $\text{cl}_Z(W) \in H_W^2(Z, \mathbb{Z}_\ell(1))$. On the other hand, there is a localization long exact sequence consisting of finite Λ_n -modules for each $n \geq 1$,

$$\begin{aligned} \cdots \rightarrow H_{\bar{W}}^r(\bar{Z}, \Lambda_n(d)) \rightarrow H^r(\bar{Z}, \Lambda_n(d)) \rightarrow H^r(\bar{V}, \Lambda_n(d)) \\ \rightarrow H_{\bar{W}}^{r+1}(\bar{Z}, \Lambda_n(d)) \rightarrow \cdots \end{aligned}$$

Hence we obtain the desired long exact sequence by taking the projective limit of these localization long exact sequences with respect to $n \geq 1$ and then taking $\otimes_{\mathbb{Z}_\ell} \mathbb{Q}_\ell$. Thus $H^{d+1}(V, \mathbb{Q}_\ell(d))$ and $H^{d+1}(V, \Lambda_\infty(d))$ are zero, and we obtain Theorem 3.2. \square

4. The Bertini theorem over a discrete valuation ring

Let the notation be as in Section 1. The discrete valuation ring R is not necessarily henselian, and the residue field F of R is arbitrary in this section. For a free R -module E of finite rank, let

$$\mathbb{P}_B(E) \longrightarrow B = \text{Spec}(R)$$

be the associated projective bundle (cf. [EGAII, 4.1.1]). Put $\mathbb{P}_s(E) := \mathbb{P}_B(E) \otimes_R F$, which is the projective bundle over F associated to $E_s := E \otimes_R F$. Let $G_B(E)$ be the set of the invertible (i.e., rank one) R -submodules $N \subset E$ such that E/N is free. Such N induces a closed immersion

$$H(N) := \mathbb{P}_B(E/N) \hookrightarrow \mathbb{P}_B(E),$$

which we call a hyperplane in $\mathbb{P}_B(E)$ corresponding to N . There is a specialization map

$$\text{sp}_E : G_B(E) \longrightarrow G_s(E), \quad N \mapsto N \otimes_R F,$$

where $G_s(E)$ is the set of the one-dimensional F -subspaces of $E \otimes_R F$. This map is surjective. In terms of hyperplanes in projective bundles, sp_E assigns to a hyperplane $H(N) \subset \mathbb{P}_B(E)$ over B , a hyperplane $H_s(N) := H(N) \otimes_R F \subset \mathbb{P}_s(E)$ over s . We quote here the following fact proved in [JSa, Th. 1].

THEOREM 4.1. *Let $X \in \mathrm{Ob}(\mathcal{QSP})$ with $\dim(X) \geq 2$. Then there exist a free R -module E of finite rank, an embedding $X \hookrightarrow \mathbb{P}_B(E)$ and an invertible R -module $N \in G_B(E)$ such that $X \cdot H(N) := X \times_{\mathbb{P}_B(E)} H(N)$ lies in \mathcal{QSP} and such that $(X, X \cdot H(N))$ is a \mathcal{QSP} -pair.*

In this paper, we need the following stronger result, which was obtained also in a joint work of Uwe Jannsen and the first author:

THEOREM 4.2 (Jannsen-Saito). *Let $X \in \mathrm{Ob}(\mathcal{QSP})$ with $\dim(X) = d + 1 \geq 2$. Let Y_1, \dots, Y_r be the irreducible components of $Y := X_{s, \mathrm{red}}$, which are smooth of dimension d over F . For integers $1 \leq a \leq d$ and $1 \leq i_1 < \dots < i_a \leq r$, we put $Y_{i_1, \dots, i_a} := Y_{i_1} \cap \dots \cap Y_{i_a}$. Let $W \subset X$ be a closed subscheme satisfying the following three conditions:*

- (i) *W is the disjoint union of integral regular schemes W_1, \dots, W_m .*
- (ii) *For integers $1 \leq a \leq d$, $1 \leq i_1 < \dots < i_a \leq r$ and $1 \leq v \leq m$, if $W_v \not\subset Y_{i_1, \dots, i_a}$, then $W_v \times_X Y_{i_1, \dots, i_a}$ is empty or regular of dimension $< \frac{1}{2} \dim(Y_{i_1, \dots, i_a})$.*
- (iii) *For integers $1 \leq a \leq d$, $1 \leq i_1 < \dots < i_a \leq r$ and $1 \leq v \leq m$, if $W_v \subset Y_{i_1, \dots, i_a}$, then $\dim(W_v) < \frac{1}{2} \dim(Y_{i_1, \dots, i_a})$.*

Then there exist a free R -module E of finite rank, an embedding $X \hookrightarrow \mathbb{P}_B(E)$ and an invertible R -module $N \in G_B(E)$ satisfying the conditions:

- (1) *$X \cdot H(N) := X \times_{\mathbb{P}_B(E)} H(N)$ lies in \mathcal{QSP} and $(X, X \cdot H(N))$ is a \mathcal{QSP} -pair.*
- (2) *$H(N)$ contains W .*

Remark 4.3. (1) In the original form of Theorem 4.1, it is assumed that F is infinite. This assumption is removed by virtue of a theorem of Poonen [Poo04, Th. 1.1]. By the same token, another theorem due to him [Poo08, Th. 1.1], enables us to show Theorem 4.2 in the case that F is finite (cf. Theorem 4.7 below).

- (2) Theorem 4.2 will play a crucial role in the proof of Theorem 1.16 in Section 8 below. In fact, in the proof of Theorem 1.16 for the case that F is finite, a standard norm argument enables one to use Theorem 4.2 just in the case that the residue field F is infinite.
- (3) Put $H_s(N) := H(N) \otimes_R F \subset \mathbb{P}_s(E)$. Theorem 4.2 (i) implies that $H_s(N) \times_{\mathbb{P}_s(E)} Y_{i_1, \dots, i_a}$ is regular for any $1 \leq i_1 < \dots < i_a \leq m$ and that it is empty if $\dim(Y_{i_1, \dots, i_a}) = 0$. We also note that $H_s(N) \times_{\mathbb{P}_s(E)} Y_{i_1, \dots, i_a}$ is not empty if $\dim(Y_{i_1, \dots, i_a}) \geq 1$, because it is a hyperplane section of $Y_{i_1, \dots, i_a} \subset \mathbb{P}_s(E)$.

For the proof of Theorem 4.2, we need the following fact proved in [JSa, Lemma 1].

LEMMA 4.4. *Let $X \in \text{Ob}(\mathcal{QSP})$ be endowed with an embedding $X \hookrightarrow \mathbb{P}_B(E)$. Let $H \hookrightarrow \mathbb{P}_B(E)$ be a hyperplane. Assume that $H_s := H \otimes_R F$ and Y_{i_1, \dots, i_a} for any i_1, \dots, i_a intersect transversally on $\mathbb{P}_s(E)$. Then $X \cdot H (= X \times_{\mathbb{P}_B(E)} H)$ lies in \mathcal{QSP} and $(X, X \cdot H)$ is a \mathcal{QSP} -pair.*

Theorem 4.1 is a rather immediate consequence of this lemma (cf. loc. cit. §1). We deduce Theorem 4.2 from Lemma 4.4 by a refined argument.

Proof of Theorem 4.2. Let $W^f \subset W$ (resp. W^{nf}) be the (disjoint) union of those W_v 's which are not contained (resp. contained) in Y . By the projectivity assumption on X , take a free R -module E_0 of finite rank and an embedding $i : X \hookrightarrow \mathbb{P}_B(E_0)$. Put $\mathcal{O}_X(n) = i^* \mathcal{O}_{\mathbb{P}_B(E_0)}(n)$ for an integer $n \geq 1$. By the Serre vanishing theorem (e.g. [Har77, III, Th. 5.2]),

$$H^1(X, \mathcal{O}_X(n)) = H^1(X, \mathcal{O}_X(n) \otimes I_X(W^f)) = 0$$

for a sufficiently large $n > 0$, where $I_X(W^f) \subset \mathcal{O}_X$ is the ideal sheaf for $W^f \subset X$. We fix such n in what follows. The R -modules

$$E_n := H^0(X, \mathcal{O}_X(n)) \supset \tilde{E}_n := H^0(X, \mathcal{O}_X(n) \otimes_{\mathcal{O}_X} I_X(W^f))$$

are free of finite rank and

$$E_n \otimes_R F = H^0(X_s, \mathcal{O}_{X_s}(n)) \quad \text{and} \quad \tilde{E}_n \otimes_R F = H^0(X_s, \mathcal{O}_{X_s}(n) \otimes_{\mathcal{O}_X} I_{X_s}(W_s^f)).$$

Here we have used the flatness of W^f over B . The last flatness also implies that $I_{X_s}(W_s^f) = I_X(W^f) \otimes_{\mathcal{O}_X} \mathcal{O}_{X_s}$, so that there is a short exact sequence

$$\begin{aligned} 0 \rightarrow \mathcal{O}_X(n) \otimes_{\mathcal{O}_X} I_X(W^f) &\xrightarrow{\times \pi} \mathcal{O}_X(n) \otimes_{\mathcal{O}_X} I_X(W^f) \\ &\rightarrow \mathcal{O}_{X_s}(n) \otimes_{\mathcal{O}_{X_s}} I_{X_s}(W_s^f) \rightarrow 0, \end{aligned}$$

where π denotes a prime element of R . By this short exact sequence, the quotient E_n/\tilde{E}_n is free, and moreover, for an invertible R -submodule $N \subset \tilde{E}_n$ with \tilde{E}_n/N free, the quotient E_n/N is again free. Thus we obtain a commutative diagram

$$\begin{array}{ccc} G_B(\tilde{E}_n) & \xhookrightarrow{\iota} & G_B(E_n) \\ \text{sp}_{\tilde{E}_n} \downarrow & & \downarrow \text{sp}_{E_n} \\ G_s(\tilde{E}_n) & \xhookrightarrow{\iota} & G_s(E_n). \end{array}$$

The images $\iota(G_B(\tilde{E}_n))$ and $\iota(G_s(\tilde{E}_n))$ are identified with the sets

$$\Phi := \{N \in G_B(E_n) \mid W^f \subset H(N)\} \quad \text{and} \quad \Phi_s := \{M \in G_s(E_n) \mid W_s^f \subset H_s(M)\},$$

respectively. Here $H(N)$ denotes the hyperplane $\mathbb{P}_B(E/N) \hookrightarrow \mathbb{P}_B(E)$ and $H_s(M) \subset \mathbb{P}_s(E_n)$ is the hyperplane corresponding to $(E_n \otimes_R F)/M$. We have used the

embedding $X \hookrightarrow \mathbb{P}_B(E_n)$ associated to $\mathbb{O}_X(n)$. Consider the set

$$\begin{aligned}\Phi_{s,\text{red}} &:= \{M \in G_s(E_n) \mid W_{s,\text{red}}^f = W^f \cap X_{s,\text{red}} \subset H_s(M)\} \\ &= \{M \in G_s(E_n) \mid W^f \cap Y_{i_1, \dots, i_a} \subset H_s(M) \text{ for any } i_1, \dots, i_a\}.\end{aligned}$$

By definition we have $\Phi_s \subset \Phi_{s,\text{red}}$. We show the following:

LEMMA 4.5. *Put*

$$\mathcal{F} := \text{Ker}(I_{X_s}(X_{s,\text{red}}) \rightarrow I_{X_s}(X_{s,\text{red}}) \cdot \mathbb{O}_{W_s^f}) \simeq I_{X_s}(X_{s,\text{red}}) \otimes I_{X_s}(W_s^f),$$

where the tensor products of sheaves are taken over \mathbb{O}_{X_s} . Replacing n with a sufficiently larger one if necessary, assume that $H^1(X_s, \mathbb{O}_{X_s}(n) \otimes \mathcal{F}) = 0$. Then for a given $M \in \Phi_{s,\text{red}}$, there exists an $M' \in \Phi_s$ such that

$$H_s(M) \cap X_{s,\text{red}} = H_s(M') \cap X_{s,\text{red}} \subset \mathbb{P}_s(E_n).$$

Proof of Lemma 4.5. By definition we have

$$(4.6) \quad \Phi_s = (\text{Ker}(\tau) - \{0\})/F^\times, \quad \Phi_{s,\text{red}} = (\text{Ker}(\sigma\tau) - \{0\})/F^\times,$$

where σ and τ are as follows:

$$\begin{aligned}H^0(X_s, \mathbb{O}_{X_s}(n)) &\xrightarrow{\tau} H^0(X_s, \mathbb{O}_{X_s}(n) \otimes \mathbb{O}_{W_s^f}) \\ &\xrightarrow{\sigma} H^0(X_s, \mathbb{O}_{X_s}(n) \otimes \mathbb{O}_{W_{s,\text{red}}^f}).\end{aligned}$$

The assumption of the claim implies that the map

$$H^0(X_s, \mathbb{O}_{X_s}(n) \otimes I_{X_s}(X_{s,\text{red}})) \rightarrow H^0(X_s, \mathbb{O}_{X_s}(n) \otimes (I_{X_s}(X_{s,\text{red}}) \cdot \mathbb{O}_{W_s^f})) = \text{Ker}(\sigma)$$

is surjective. Hence we have

$$\text{Ker}(\sigma\tau) = \text{Ker}(\tau) + H^0(X_s, \mathbb{O}_{X_s}(n) \otimes I_{X_s}(X_{s,\text{red}})).$$

In view of (4.6), this equality implies the desired assertion. \square

We turn to the proof of Theorem 4.2. Put

$$\begin{aligned}\Phi_{s,\text{red}} \supset \Psi &:= \{M \in G_s(E_n) \mid W^{nf} \cup W_{s,\text{red}}^f \subset H_s(M)\} \\ &= \{M \in G_s(E_n) \mid W \cap Y_{i_1, \dots, i_a} \subset H_s(M) \text{ for any } i_1, \dots, i_a\}\end{aligned}$$

and let $\Psi_{\text{reg}} \subset \Psi$ be the subset of such M that $H_s(M)$ and Y_{i_1, \dots, i_a} intersect transversally on $\mathbb{P}_s(E_n)$ for any i_1, \dots, i_a . We use here the following fact, which is due to Altman-Kleiman ([KA79, Th. 7]) if F is infinite, and due to Poonen ([Poo04, Th. 1.1], [Poo08, Th. 1.1]) if F is finite.

THEOREM 4.7 (Altman-Kleiman/Poonen). *Let Y be a smooth projective scheme over F and let $Z \subset Y$ be a smooth closed subscheme. Assume $\dim(Z) < \frac{1}{2} \dim(Y)$. Then for any integer $n \geq 0$ large enough, there exists a smooth hypersurface section $H \subset X$ of degree n containing Z .*

By this theorem and the assumptions of Theorem 4.2, we may suppose that Ψ_{reg} is nonempty, replacing n again with a sufficiently larger one if necessarily. By the surjectivity of $\text{sp}_{\tilde{E}_n}$ and by Lemma 4.5, there exists an $N \in \Phi \subset G_B(E_n)$ satisfying $\text{sp}_{E_n}(N) \in \Psi_{\text{reg}}$. Now Theorem 4.2 follows from Lemma 4.4 for the embedding $X \hookrightarrow G_B(E_n)$. \square

5. Surjectivity of the cycle map

Let the notation be as in Section 1. In this section, we prove the following:

THEOREM 5.1. *Assume that R is henselian and that F is finite or separably closed. Let X be a scheme over $\text{Spec}(R)$ which belongs to \mathfrak{QSP} . Then the cycle map defined in (1.12)*

$$\rho_X : \text{CH}_1(X) \otimes \Lambda \rightarrow H_2(X, \Lambda)$$

is surjective for both $\Lambda = \Lambda_n$ and Λ_∞ .

Proof. It suffices to prove the case that $\Lambda = \Lambda_n$. Without loss of generality, we may assume that X is integral. The case $\dim(X) = 1$ follows from the definition of ρ_X . We prove the case $\dim(X) = 2$. Since $\dim(X) = 2$, the niveau spectral sequence (2.2) gives rise to an exact sequence

$$0 \longrightarrow \text{CH}_1(X)/\ell^n \xrightarrow{\rho_X} H_2(X, \Lambda_n) \longrightarrow KH_2(X, \Lambda_n) \longrightarrow 0.$$

We will prove the following:

LEMMA 5.2. *If $\dim(X) = 2$, then $KH_2(X, \Lambda_n)$ is canonically isomorphic to $\text{Br}(X)[\ell^n]$, where for a scheme Z , $\text{Br}(Z)$ denotes the Grothendieck-Brauer group $H^2(Z, \mathbb{G}_m)$.*

Since R is henselian and F is finite or separably closed, $\text{Br}(X)\{\ell\} = 0$ by [CTOP02, Cors. 1.10 (b) and 1.11 (b)], which is a consequence the base-change theorem for Brauer groups of relative curves; see [Gro68, III, Th. 3.1] and [CTOP02, Th. 1.7 (c)]. Hence Theorem 5.1 for the case $\dim(X) = 2$ follows from this lemma.

Proof of Lemma 5.2. Let y be the generic point of X . Since $\dim(X) = 2$, we have

$$\begin{aligned} KH_2(X, \Lambda_n) &= \text{Ker}(d_{2,0}^1 : H^2(y, \Lambda_n(1)) \rightarrow \bigoplus_{x \in X_1} H^1(x, \Lambda_n)) \\ &= \bigcap_{x \in X_1} \text{Ker}(d_x : H^2(y, \Lambda_n(1)) \rightarrow H^1(x, \Lambda_n)) \\ &= \bigcap_{x \in X_1} \text{Br}(\mathbb{O}_{X,x})[\ell^n] = \text{Br}(X)[\ell^n], \end{aligned}$$

where $d_{2,0}^1$ denotes the differential map of the niveau spectral sequence (2.2), and d_x ($x \in X_1$) denotes the x -factor of $d_{2,0}^1$. The last equality follows from the assumption that X is regular of dimension 2 and theorems of Auslander-Goldman and Grothendieck [Gro68, II, Prop. 2.3, Cor. 2.2]. We show the third equality. Fix

a point $x \in X^1$, and put $Z := \text{Spec}(\mathbb{O}_{X,x})$. By Lemma 1.8 (2), d_x is identified with the arrow δ^2 in the localization exact sequence of étale cohomology groups

$$\begin{aligned} H^1(y, \Lambda_n(1)) &\xrightarrow{\delta^1} H_x^2(Z, \Lambda_n(1)) \rightarrow H^2(Z, \Lambda_n(1)) \\ &\rightarrow H^2(y, \Lambda_n(1)) \xrightarrow{\delta^2} H_x^3(Z, \Lambda_n(1)), \end{aligned}$$

where $H^1(y, \Lambda_n(1)) \simeq \kappa(y)^\times / \ell^n$ by Hilbert's theorem 90, and $H_x^2(Z, \Lambda_n(1)) \simeq \Lambda_n$ by [SGA $\frac{1}{2}$, Cycle Prop. 2.1.4]. By [JSS, Th. 1.1.1], δ^1 agrees with the negative of the divisor map $\kappa(y)^\times / \ell^n \rightarrow \Lambda_n$. Because $\mathbb{O}_{X,x}$ is a principal ideal domain, δ^1 is surjective and we have $H^2(Z, \Lambda_n(1)) = \text{Ker}(\delta^2)$. On the other hand, the short exact sequence $0 \rightarrow \Lambda_n(1) \rightarrow \mathbb{G}_m \xrightarrow{\times \ell^n} \mathbb{G}_m \rightarrow 0$ on $Z_{\text{ét}}$ gives rise to a short exact sequence

$$0 \longrightarrow \text{Pic}(\mathbb{O}_{X,x}) / \ell^n \longrightarrow H^2(Z, \Lambda_n(1)) \longrightarrow \text{Br}(\mathbb{O}_{X,x})[\ell^n] \longrightarrow 0,$$

and $\text{Pic}(\mathbb{O}_{X,x})$ is zero because $\mathbb{O}_{X,x}$ is local. Thus we have $\text{Ker}(d_x) = \text{Ker}(\delta^2) = H^2(Z, \Lambda_n(1)) = \text{Br}(\mathbb{O}_{X,x})[\ell^n]$, which completes the proof of the lemma. \square

Finally we prove the case that $\dim(X) \geq 3$ by induction on $\dim(X)$. By Theorem 4.2 we take an ample divisor $Y \subset X$ such that (X, Y) is an ample \mathcal{LSP} -pair, and put $U := X - Y$. By Proposition 1.13 (2), there is a commutative diagram whose lower row is exact:

$$\begin{array}{ccccc} \text{CH}_1(Y) / \ell^n & \longrightarrow & \text{CH}_1(X) / \ell^n & & \\ \rho_Y \downarrow & & \rho_X \downarrow & & \\ H_2(Y, \Lambda_n) & \longrightarrow & H_2(X, \Lambda_n) & \longrightarrow & H_2(U, \Lambda_n). \end{array}$$

By the induction hypothesis, ρ_Y is surjective. Since R is henselian, we have $H_2(U, \Lambda_n) = 0$ by Theorem 3.2, which implies the surjectivity of ρ_X . Thus we obtain Theorem 5.1. \square

THEOREM 5.3. *Let the assumptions be as in Theorem 5.1. Let Λ denote Λ_n if F is separably closed and Λ_∞ if F is finite. Then $KH_2(X, \Lambda) = 0$.*

Proof. The niveau spectral sequence (2.2) and Lemma 2.6 yield isomorphisms

$$KH_2(X, \Lambda) = E_{2,0}^2(X) = E_{2,0}^\infty(X) = \text{Coker}(\rho_X : \text{CH}_1(X) \otimes \Lambda \rightarrow H_2(X, \Lambda)).$$

Hence Theorem 5.3 follows from Theorem 5.1. \square

6. The blow-up formula

Let the notation be as in Section 1. We prove here some lemmas which explain how homology groups behave under blow-ups at closed points. The goal of this section is to prove Lemma 6.4 below, which will be used in the proof of Theorem 1.16. We do not assume R is henselian in this section. The residue field F of R is arbitrary in the following lemma.

LEMMA 6.1. *Let $X \in \text{Ob}(\mathcal{C})$ be regular of dimension $d + 1$. Fix a closed point x on X_S and let $\pi : \tilde{X} \rightarrow X$ be the blow-up of X at x and let $\iota : E \hookrightarrow \tilde{X}$ be the exceptional divisor. For any integer q there is a split short exact sequence*

$$0 \longrightarrow \tilde{H}_q(E, \Lambda) \xrightarrow{\iota_*} H_q(\tilde{X}, \Lambda) \xrightarrow{\pi_*} H_q(X, \Lambda) \longrightarrow 0.$$

Here $\tilde{H}_q(E, \Lambda)$ is defined as

$$\tilde{H}_q(E, \Lambda) := \text{Ker}(\pi_* : H_q(E, \Lambda) \longrightarrow H_q(x, \Lambda)),$$

which agrees with $H_q(E, \Lambda)$ for $q \geq 1$ because $H_q(x, \Lambda)$ is zero for $q \geq 1$ (see Remark 1.7 or (1.9)).

Proof. By Proposition 1.5 (3), there is a commutative diagram with exact rows

$$\begin{array}{ccccccc} \cdots & \longrightarrow & H_q(E, \Lambda) & \longrightarrow & H_q(\tilde{X}, \Lambda) & \longrightarrow & H_q(\tilde{X} - E, \Lambda) \longrightarrow H_{q-1}(E, \Lambda) \longrightarrow \cdots \\ & & \downarrow & & \pi_* \downarrow & & \parallel & & \downarrow \\ \cdots & \longrightarrow & H_q(x, \Lambda) & \longrightarrow & H_q(X, \Lambda) & \longrightarrow & H_q(X - x, \Lambda) \longrightarrow H_{q-1}(x, \Lambda) \longrightarrow \cdots \end{array}$$

It is enough to show that $\pi_* : H_q(\tilde{X}, \Lambda_n) \rightarrow H_q(X, \Lambda_n)$ is split surjective. We define the map $\pi^* : H_q(X, \Lambda_n) \rightarrow H_q(\tilde{X}, \Lambda_n)$ as follows:

$$\begin{aligned} \pi^* : H_q(X, \Lambda_n) &\stackrel{(1.9)}{\simeq} H^{2d+2-q}(X, \Lambda_n(d)) \\ &\xrightarrow{\pi^*} H^{2d+2-q}(\tilde{X}, \Lambda_n(d)) \stackrel{(1.9)}{\simeq} H_q(\tilde{X}, \Lambda_n). \end{aligned}$$

We show $\pi_* \pi^* = 1$ as endomorphisms on $H_q(X, \Lambda)$. By Lemma 1.8 (1), π_* is identified with the push-forward map of étale cohomology groups. By the projection formula, i.e., the commutative diagram of cup-product pairings (cf. Proof of [Mil80, VI, Cor. 6.4 (a)])

$$\begin{array}{ccccc} H^0(\tilde{X}, \Lambda_n) \times H^{2d+2-q}(\tilde{X}, \Lambda_n(d)) & \xrightarrow{\cup} & H^{2d+2-q}(\tilde{X}, \Lambda_n(d)) \\ \pi_* \downarrow & & \pi^* \uparrow & & \pi_* \downarrow \\ H^0(X, \Lambda_n) \times H^{2d+2-q}(X, \Lambda_n(d)) & \xrightarrow{\cup} & H^{2d+2-q}(X, \Lambda_n(d)), \end{array}$$

we have only to check the left vertical π_* maps 1-to-1, which follows from the fact that π is an isomorphism outside of E . Thus we obtain the lemma. \square

Let $\pi : \tilde{X} \rightarrow X$ and $\iota : E \hookrightarrow \tilde{X}$ be as in Lemma 6.1. Let $W \subset \tilde{X}$ be an integral closed subscheme of dimension one. For $\Lambda = \Lambda_n$, let

$$\Lambda \langle W \rangle \subset H_2(\tilde{X}, \Lambda)$$

be the subgroup generated by $\rho_X(W) \in H_2(\tilde{X}, \Lambda)$. For $\Lambda = \Lambda_\infty$, we put

$$\Lambda \langle W \rangle := \varinjlim_{n \in \mathbb{N}} \Lambda_n \langle W \rangle \subset H_2(\tilde{X}, \Lambda_\infty).$$

See Remark 1.14 for the compatibility of cycle class maps with respect to the multiplication by ℓ from Λ_n to Λ_{n+1} . Recall that $E \simeq \mathbb{P}_x^d$. For an integral closed subscheme $W \subset E$ of dimension one, we define $\deg_{E/x}(W) \in \mathbb{Z}$ to be the degree over x (cf. [Ful98, Def. 1.4]) of the intersection of W and a hyperplane $H \subset E$ with $W \not\subset H$. Under these settings, we prove:

COROLLARY 6.2. *Assume $\text{cd}_\ell(F) \leq 1$ (e.g., F is finite or separably closed). Let $W \subset E$ be a closed integral subscheme of dimension one, and assume either that $\Lambda = \Lambda_\infty$ or that $\deg_{E/x}(W)$ is prime to ℓ . Then*

$$\text{Ker}(\pi_* : H_2(\tilde{X}, \Lambda) \rightarrow H_2(X, \Lambda)) = \Lambda\langle W \rangle.$$

Proof. Lemma 6.1 yields an exact sequence

$$0 \longrightarrow H_2(E, \Lambda) \xrightarrow{\iota_*} H_2(\tilde{X}, \Lambda) \xrightarrow{\pi_*} H_2(X, \Lambda) \longrightarrow 0.$$

Since $E \simeq \mathbb{P}_x^d$, $\text{CH}_1(E)$ is generated by a line $L \subset E$ and we have

$$\deg_{E/x} : \text{CH}_1(E) \xrightarrow{\sim} \mathbb{Z}.$$

We show that the map $\rho_E : \text{CH}_1(E) \otimes \Lambda \rightarrow H_2(E, \Lambda)$ is bijective. Indeed, we have

$$H_2(E, \Lambda_n) \underset{(1.9)}{\simeq} H^{2d-2}(E, \Lambda_n(d-1)) \underset{\text{cd}_\ell(F) \leq 1}{\simeq} \Lambda_n$$

and $H_2(E, \Lambda_n)$ is generated by $\rho_E(L)$, which implies the bijectivity of ρ_E for both $\Lambda = \Lambda_n$ and Λ_∞ . Corollary 6.2 follows easily from these facts and Proposition 1.13 (1) for $E \hookrightarrow \tilde{X}$. \square

Setting 6.3. We consider the following situation:

$$X = X^{(0)} \xleftarrow{\pi_1} X^{(1)} \xleftarrow{\pi_2} X^{(2)} \xleftarrow{\pi_3} \dots \xleftarrow{\pi_N} X^{(N)} = \tilde{X}.$$

Here $X \in \text{Ob}(\mathcal{C})$ is regular of dimension $d+1$, and $\pi_i : X^{(i)} \rightarrow X^{(i-1)}$ ($1 \leq i \leq N$) is the blow-up at a single closed point x_{i-1} on $(X^{(i-1)})_s$. Let $\pi : \tilde{X} \rightarrow X$ be the composite of the above maps. Let $E_i \subset X^{(i)}$ be the exceptional divisor of π_i and let $\tilde{E}_i \subset \tilde{X}$ be its strict transform (cf. Appendix A below for the definition). Note that $E_i \simeq \mathbb{P}_{x_{i-1}}^d$. Let $\tau_i : \tilde{E}_i \rightarrow E_i$ be the natural map. We have the following facts (see Proposition A.4 below for (1)–(3)):

- (1) $\tilde{E}_1, \dots, \tilde{E}_N$ are regular irreducible components of $\tilde{X}_{s, \text{red}}$.
- (2) The reduced divisor $\bigcup_{i=1}^N \tilde{E}_i$ on \tilde{X} has simple normal crossings.
- (3) If $X \in \text{Ob}(\mathcal{Q}S)$, then $\tilde{X} \in \text{Ob}(\mathcal{Q}S)$.
- (4) For $1 \leq i \leq N$, there exists a unique finite set S_i of closed points on E_i such that

$$\tilde{S}_i := \tau_i^{-1}(S_i) = \tilde{E}_i \cap \left(\bigcup_{1 \leq j \leq N, j \neq i} \tilde{E}_j \right)$$

and such that $\tilde{E}_i - \tilde{S}_i \simeq E_i - S_i$ via τ_i . Note that S_i may be empty.

These facts and the following lemma will be used later in Section 8 below.

LEMMA 6.4. Assume $\mathrm{cd}_\ell(F) \leq 1$. For $1 \leq i \leq N$, let $W_i \subset \tilde{E}_i$ be a closed integral subscheme of dimension one such that $W_i \not\subset \tilde{S}_i$. Assume either that $\Lambda = \Lambda_\infty$ or that $\deg_{E_i/x_{i-1}}(\tau_i(W_i))$ is prime to ℓ for all $1 \leq i \leq N$. Then we have

$$\mathrm{Ker}(\pi_* : H_2(\tilde{X}, \Lambda) \rightarrow H_2(X, \Lambda)) = \sum_{1 \leq i \leq N} \Lambda \langle W_i \rangle.$$

Proof. Let p be the projection $\tilde{X} \rightarrow X^{(1)}$. Since $\pi(W_i)$ is a closed point of X_s , the right-hand side is contained in the left-hand side by Proposition 1.13 (1). There is an exact sequence

$$0 \longrightarrow \mathrm{Ker}(p_* : H_2(\tilde{X}, \Lambda) \rightarrow H_2(X^{(1)}, \Lambda)) \longrightarrow \mathrm{Ker}(\pi_* : H_2(\tilde{X}, \Lambda) \rightarrow H_2(X, \Lambda)) \xrightarrow{p_*} \mathrm{Ker}(\pi_{1*} : H_2(X^{(1)}, \Lambda) \rightarrow H_2(X, \Lambda)).$$

By induction on N , we may suppose that the first term is equal to $\sum_{2 \leq i \leq N} \Lambda \langle W_i \rangle$. The last group is equal to $\Lambda \langle p_* W_1 \rangle$ by Corollary 6.2, which agrees with $p_*(\Lambda \langle W_1 \rangle)$ by Proposition 1.13 (1) for the proper map $p : \tilde{X} \rightarrow X^{(1)}$. Lemma 6.4 follows easily from this fact. \square

7. A moving lemma

Let the notation be as in Section 1. The discrete valuation ring R is not necessarily henselian, and the residue field F of R is arbitrary. In this section we prove the following moving lemma for cycles on $X \in \mathrm{Ob}(\mathcal{C})$, which will be used in the proof of Theorem 1.16.

PROPOSITION 7.1. *Let X be an integral regular scheme which belongs to $\mathrm{Ob}(\mathcal{C})$. Let $Y \subset X$ be a proper closed subscheme, and put $U := X - Y$. Then for $q \geq 0$, the natural map $\bigoplus_{x \in X_q \cap U} \mathbb{Z} \rightarrow \mathrm{CH}_q(X)$ is surjective.*

A slight modification of the following argument will show that the assertion of Proposition 7.1 is valid even if the base scheme B is the spectrum of a Dedekind ring. Since this generalization is not used in this paper, we do not pursue it here.

For the proof of the proposition, we need the following lemmas.

LEMMA 7.2. *Let X be an integral regular scheme and let $Y \subset X$ be a proper closed subscheme with $U = X - Y$. Let y be a point on Y , and put $c := \mathrm{codim}_X(y)$. Then there exists an integral closed subscheme $Z \subset X$ of codimension $c - 1$ satisfying the conditions (1) $Z \cap U \neq \emptyset$, (2) $y \in Z$, (3) Z is regular at y .*

LEMMA 7.3. *Let Z be a quasi-projective scheme over B and assume given a finite number of points z_1, \dots, z_r on Z . Then there exist a dense open subset $V \subset Z$ and a dense open immersion $V \hookrightarrow U$ such that V contains all z_1, \dots, z_r and such that U is affine.*

We first prove Proposition 7.1 admitting these lemmas. Take an arbitrary $y \in Y_q$. Let $Z \subset X$ be as in Lemma 7.2. Then $\dim(Z \cap Y) \leq q$ and $Z_q \cap Y$ is finite. Let $\pi : \tilde{Z} \rightarrow Z$ be the normalization of Z . The assumption that X

is quasi-projective over B implies that \tilde{Z} is quasi-projective over B , and hence by Lemma 7.3, there exists an open subset $V \subset \tilde{Z}$ and a dense open immersion $V \hookrightarrow U = \operatorname{Spec}(A)$ such that V contains $\pi^{-1}(Z_q \cap Y)$. Let A' be the semi-localization of A at the ideals corresponding to $\pi^{-1}(Z_q \cap Y) \subset U$. It is a regular semi-local domain of dimension one, and hence a principal ideal domain. Since Z is regular at y , there is a unique point $\tilde{y} \in \tilde{Z}$ lying over y . Suppose $f \in A'$ defines the principal divisor \tilde{y} on $\operatorname{Spec}(A')$, which we regard as a rational function on \tilde{Z} . Then the divisor $\operatorname{div}_{\tilde{Z}}(f)$ on \tilde{Z} has the form

$$\operatorname{div}_{\tilde{Z}}(f) = [\tilde{y}] + \sum_{i=1}^N m_i [x_i]$$

for some points $x_i \in \tilde{Z}_q \setminus (\{\tilde{y}\} \cup \pi^{-1}(Z_q \cap Y))$ and some integers m_i for $1 \leq i \leq N$. Applying π_* to both sides, we get the following formula for the divisor $\operatorname{div}_Z(f) = \pi_* \operatorname{div}_{\tilde{Z}}(f)$ on Z :

$$\operatorname{div}_Z(f) = [y] + \sum_{i=1}^N m'_i [x'_i] \quad \text{for some integers } m'_1, \dots, m'_N,$$

where $x'_i = \pi(x_i) \in Z_q \setminus (\{y\} \cup (Z_q \cap Y))$. This proves the desired assertion. \square

Proof of Lemma 7.2. We have only to show the case that Y is a divisor on X , and we may replace X with $\operatorname{Spec}(\mathbb{O}_{X,y})$. Suppose that Y is defined by a nonzero element $\pi \in \mathbb{O}_{X,y}$ on $X = \operatorname{Spec}(\mathbb{O}_{X,y})$. It suffices to show the following:

SUBLEMMA 7.4. *Let (A, \mathfrak{m}) be a regular local ring of dimension $c \geq 1$ and let $\pi \in \mathfrak{m}$ be a nonzero element. Then there exists a regular system of parameters $\{a_1, \dots, a_{c-1}, a_c\}$ of A for which π does not belong to the ideal $(a_1, \dots, a_{c-1}) \subset A$.*

We prove this sublemma by induction on $c \geq 1$. The case $c = 1$ is clear. Suppose $c \geq 2$. Because A is a unique factorization domain and $\mathfrak{m}/(\mathfrak{m}^2 + (\pi))$ is nonzero by the assumption $c \geq 2$, one can find a prime element a_1 which does not divide π and does not belong to \mathfrak{m}^2 . Then $A' := A/(a_1)$ is a regular local ring of dimension $c - 1$ such that the residue class $\bar{\pi} \in A'$ of π is nonzero. By the induction hypothesis, there is a regular system of parameters $\{b_2, \dots, b_{c-1}, b_c\}$ of A' such that $\bar{\pi} \notin (b_2, \dots, b_{c-1})$. Taking lifts $a_2, \dots, a_{c-1}, a_c \in A$ of b_2, \dots, b_{c-1}, b_c , we get a regular system of parameters $\{a_1, \dots, a_{c-1}, a_c\}$ of A such that $\pi \notin (a_1, \dots, a_{c-1})$. Thus we obtain the sublemma and Lemma 7.2. \square

Proof of Lemma 7.3. By assumption there is a locally closed immersion $i : Z \hookrightarrow \mathbb{P}_B^N$, where \mathbb{P}_B^N denotes the projective space $\operatorname{Proj}(R[T_0, \dots, T_N])$ over $B = \operatorname{Spec}(R)$. Replacing Z by the closure of $i(Z)$ in \mathbb{P}_B^N , we may assume Z is a closed subscheme of \mathbb{P}_B^N . Then, for any affine open subscheme $U \subset \mathbb{P}_B^N$, $U \cap Z$ is affine, and hence we may assume $X = \mathbb{P}_B^N$. If an open subset $U \subset \mathbb{P}_B^N$ contains a point in the closure of $z_i \in \mathbb{P}_B^N$, then it should contain z_i as well. Therefore we may

assume that z_1, \dots, z_r are closed points of $\mathbb{P}_s^N = \mathbb{P}_B^N \times_B s = \text{Proj}(F[T_0, \dots, T_N])$. Let $\mathfrak{m}_i \subset A := F[T_0, \dots, T_N]$ be the homogeneous ideal corresponding to z_i for $1 \leq i \leq r$. We will prove the following sublemma:

SUBLEMMA 7.5. *There exists a nonzero homogeneous polynomial $f \in A$ which is not contained in any of $\mathfrak{m}_1, \dots, \mathfrak{m}_r$, or equivalently, for which the hypersurface $H_f \subset \mathbb{P}_s^N$ defined by f does not contain any of z_1, \dots, z_r .*

We first finish the proof of Lemma 7.3 admitting this sublemma. Let $f' \in R[T_0, \dots, T_N]$ be a homogeneous element with $f' \bmod (\pi) = f$, where f is taken as in the claim. Then the open subset $U = \{f' \neq 0\} \subset \mathbb{P}_B^N$ is affine and contains all z_1, \dots, z_r .

Proof of Sublemma 7.5. We first assume that F is infinite. Then the assertion follows from the following fact: In the dual projective space $(\mathbb{P}_s^N)^\vee$ (i.e. the Grassmannian of hyperplanes in \mathbb{P}_s^N), the locus S_i of those hypersurfaces that contain z_i is a proper closed subset, and hence $(\mathbb{P}_s^N)^\vee - (\bigcup_{1 \leq i \leq r} S_i)$ is dense open in $(\mathbb{P}_s^N)^\vee$. We next prove the case that F is finite. By the previous case, there exists a finite Galois extension F'/F and a linear form $L = \sum_{0 \leq j \leq N} a_j T_j \in A \otimes_F F'$ such that $L \notin \bigcup_{1 \leq i \leq r} \mathfrak{m}_i \otimes_F F'$. It is easy to see that for any $\sigma \in \text{Gal}(F'/F)$ and any i with $1 \leq i \leq r$, the linear form $L^\sigma = \sum_{0 \leq j \leq N} a_j^\sigma T_j$ does not belong to $\mathfrak{m}_i \otimes_F F'$. Hence the homogeneous polynomial $f := \prod_{\sigma \in \text{Gal}(F'/F)} L^\sigma \in A$ satisfies the condition in Sublemma 7.5. Thus we obtain the sublemma, Lemma 7.3 and Proposition 7.1. \square

8. Proof of the main theorem

In this section we complete the proof of Theorem 1.16, which consists of three steps. Assume that R is excellent and henselian and that F is finite or separably closed. We first introduce the following construction, which will be useful later:

Construction 8.1. Let X be an object of \mathcal{QSP} , and suppose we are given integral closed subschemes $C_1, \dots, C_m \subset X$ of dimension one such that $C_j \not\subset X_s$ for any $1 \leq j \leq m$. Then by the resolution of singularities for embedded curves (see Theorem A.1 in Appendix A below), there exists a sequence of blow-ups at single closed points as in Setting 6.3

$$(8.2) \quad X = X^{(0)} \xleftarrow{\pi_1} X^{(1)} \xleftarrow{\pi_2} X^{(2)} \xleftarrow{\pi_3} \dots \xleftarrow{\pi_N} X^{(N)} = \tilde{X}$$

such that the strict transforms $\tilde{C}_1, \dots, \tilde{C}_m \subset \tilde{X}$ of C_1, \dots, C_m are disjoint and satisfy the condition (ii) of Theorem 4.2. For each $1 \leq i \leq N$, let $E_i \subset X^{(i)}$ be the exceptional divisor of π_i and let $\tilde{E}_i \subset \tilde{X}$ be its strict transform. Let $\tau_i : \tilde{E}_i \rightarrow E_i$ be the natural map. Note that \tilde{X} belongs to $\text{Ob}(\mathcal{QSP})$ by Setting 6.3 (3).

Step 1. The aim of this step is to prove the following important lemma:

LEMMA 8.3. $\rho_X : \text{CH}_1(X) \otimes \Lambda_\infty \rightarrow H_2(X, \Lambda_\infty)$ is injective for $X \in \mathcal{QSP}$ with $\dim(X) = 3$.

Proof. Take $\alpha \in \mathrm{CH}_1(X) \otimes \Lambda_\infty$ and assume $\rho_X(\alpha) = 0$. By Proposition 7.1, we have $\alpha = \sum_{1 \leq j \leq m} [C_j] \otimes \lambda_j$ ($\lambda_j \in \Lambda_\infty$) for some integral closed subschemes $C_1, \dots, C_m \subset X$ of dimension one such that $C_j \not\subset X_s$ for any $1 \leq j \leq m$. Apply Construction 8.1 to C_1, \dots, C_m , and let $\pi : \tilde{X} \rightarrow X$ be the composite of (8.2). Applying Theorem 4.2 to $\tilde{C} := \tilde{C}_1 \cup \dots \cup \tilde{C}_m \hookrightarrow \tilde{X}$, we obtain an ample divisor $\iota : Y \hookrightarrow \tilde{X}$ which contains \tilde{C} and for which (\tilde{X}, Y) is an ample \mathcal{LSP} -pair. We have

$$\alpha = \pi_* \iota_* \beta \quad \text{with} \quad \beta := \sum_{1 \leq j \leq m} [\tilde{C}_j] \otimes \lambda_j \in \mathrm{CH}_1(Y) \otimes \Lambda_\infty.$$

For each $1 \leq i \leq N$, let $\tilde{E}_i \subset \tilde{X}$ be as in Construction 8.1, and put $W_i := \tilde{E}_i \cap Y$. Each W_i is nonempty (cf. Remark 4.3 (3)), integral and regular of dimension one by the assumption $\dim(X) = 3$, and not contained in $\tilde{E}_{i'}$ for any $i' \neq i$ by Setting 6.3 (1) and (2). Hence by Lemma 6.4 we have

$$(8.3) \quad \mathrm{Ker}(\pi_* : H_2(\tilde{X}, \Lambda_\infty) \rightarrow H_2(X, \Lambda_\infty)) = \sum_{1 \leq i \leq N} \Lambda_\infty \langle W_i \rangle.$$

Now we consider the following diagram whose middle row is exact and whose square is commutative (cf. Propositions 1.5 (2) and 1.13 (1)):

$$(8.4) \quad \begin{array}{ccccc} & & \mathrm{CH}_1(Y) \otimes \Lambda_\infty & \xrightarrow{\iota_*} & \mathrm{CH}_1(\tilde{X}) \otimes \Lambda_\infty \\ & & \downarrow \rho_Y & & \downarrow \rho_{\tilde{X}} \\ H_3(\tilde{X} - Y, \Lambda_\infty) & \longrightarrow & H_2(Y, \Lambda_\infty) & \xrightarrow{\iota_*} & H_2(\tilde{X}, \Lambda_\infty) \\ \text{(Theorem 3.2)} \parallel & & & & \downarrow \pi_* \\ & & 0 & & H_2(X, \Lambda_\infty). \end{array}$$

Let us recall the assumption that $\rho_X(\alpha) = 0$. Since $\pi_* \iota_* \beta = \alpha$, we have $\pi_* \iota_* \rho_Y(\beta) = \rho_X(\pi_* \iota_* \beta) = \rho_X(\alpha) = 0$ by Proposition 1.13 (1) for the proper map $\pi \circ \iota : Y \rightarrow X$. Hence $\iota_* \rho_Y(\beta)$ belongs to $\mathrm{Ker}(\pi_*)$, and there exist $\mu_1, \dots, \mu_N \in \Lambda_\infty$ for which the 1-cycle

$$\gamma := \sum_{1 \leq i \leq N} [W_i] \otimes \mu_i \in \mathrm{CH}_1(Y) \otimes \Lambda_\infty$$

satisfies $\iota_* \rho_Y(\gamma) = \rho_{\tilde{X}}(\iota_* \gamma) = \iota_* \rho_Y(\beta)$, by (8.3) and the commutativity of the square in (8.4). Then we have $\beta = \gamma$, because $\iota_* \rho_Y$ is injective by Lemma 1.17 and Theorem 3.2. Since $\pi(W_i)$ is a closed point of X , we get $\alpha = \pi_* \iota_* \beta = \pi_* \iota_* \gamma = 0$, and the proof of Lemma 8.3 is complete. \square

Step 2. In this step we prove Theorem 1.16, assuming $\dim(X) = 3$. We need the following general lemmas:

LEMMA 8.7. *For $X \in \mathrm{Ob}(\mathcal{C})$ of dimension ≤ 3 and $\Lambda = \Lambda_n$ and Λ_∞ , there is an exact sequence*

$$H_3(X, \Lambda) \longrightarrow KH_3(X, \Lambda) \longrightarrow \mathrm{CH}_1(X) \otimes \Lambda \xrightarrow{\rho_X} H_2(X, \Lambda).$$

Proof. The assertion follows from the niveau spectral sequence (2.2) and (1.11). \square

LEMMA 8.8. *For $X \in \text{Ob}(\mathcal{C})$ of dimension ≤ 3 , we have $KH_3(X, \Lambda_n) = KH_3(X, \Lambda_\infty)[\ell^n]$.*

Proof. Since $\dim(X) \leq 3$, we have

$$KH_3(X, \Lambda) = \text{Ker} \left(d_{3,0}^1 : \bigoplus_{x \in X_3} H^3(x, \Lambda(2)) \rightarrow \bigoplus_{x \in X_2} H^2(x, \Lambda(1)) \right)$$

for $\Lambda = \Lambda_n$ and Λ_∞ , where $d_{3,0}^1$ denotes the differential map $E_{3,0}^1 \rightarrow E_{2,0}^1$ of the niveau spectral sequence (2.2). The assertion follows from the isomorphisms

$$H^3(x, \Lambda_n(2)) \simeq H^3(x, \Lambda_\infty(2))[\ell^n] \text{ and } H^2(x, \Lambda_n(1)) \simeq H^2(x, \Lambda_\infty(1))[\ell^n],$$

which are consequences of the divisibility of the groups $H^2(x, \Lambda_\infty(2))$ (Merkur'ev-Suslin [MS82]) and $H^1(x, \Lambda_\infty(1))$ (Hilbert's theorem 90), respectively. \square

We prove Theorem 1.16 assuming $\dim(X) = 3$. By Theorem 5.1, it is enough to prove the injectivity of $\rho_X : \text{CH}_1(X)/\ell^n \rightarrow H_2(X, \Lambda_n)$. By Lemmas 8.7 and 8.8, the problem is reduced to showing $KH_3(X, \Lambda_\infty) = 0$. By Theorem 4.2, we take an ample divisor $Y \subset X$ for which (X, Y) is an ample \mathcal{QSP} -pair. Put $U := X - Y$. Since $\dim(X) = 3$, we have $KH_3(X, \Lambda_\infty) \hookrightarrow KH_3(U, \Lambda_\infty)$ by (2.3). We prove $KH_3(U, \Lambda_\infty) = 0$. By Proposition 1.13 (2), there is a commutative diagram with exact rows

$$\begin{array}{ccccccc} \text{CH}_1(Y) \otimes \Lambda_\infty & \longrightarrow & \text{CH}_1(X) \otimes \Lambda_\infty & \longrightarrow & \text{CH}_1(U) \otimes \Lambda_\infty & \longrightarrow & 0 \\ \downarrow \rho_Y & & \downarrow \rho_X & & \downarrow \rho_U & & \\ H_2(Y, \Lambda_\infty) & \longrightarrow & H_2(X, \Lambda_\infty) & \longrightarrow & H_2(U, \Lambda_\infty) & & \end{array}$$

By Lemma 8.3 and Theorem 5.1, ρ_X is injective and ρ_Y is surjective. Hence ρ_U is injective and $\text{CH}_1(U) \otimes \Lambda_\infty$ is zero, because $H_2(U, \Lambda_\infty)$ is zero by Theorem 3.2. Finally $H_3(U, \Lambda_\infty)$ is zero by Theorem 3.2, and $KH_3(U, \Lambda_\infty)$ is zero as well by Lemma 8.7. This completes Step 2. \square

Step 3. We prove Theorem 1.16 by induction on $\dim(X) = d + 1 \geq 3$. The case $d = 2$ has been shown in Step 2. Assume $d \geq 3$. By Theorem 5.1, it remains to prove the injectivity of $\rho_X : \text{CH}_1(X)/\ell^n \rightarrow H_2(X, \Lambda_n)$. Take $\alpha \in \text{CH}_1(X)/\ell^n$ and assume $\rho_X(\alpha) = 0$. We have to show $\alpha = 0$. By Proposition 7.1, we have $\alpha = \sum_{1 \leq j \leq m} [C_j] \otimes \lambda_j$ ($\lambda_j \in \Lambda_n$) for some integral closed subschemes $C_1, \dots, C_m \subset X$ of dimension one such that $C_j \not\subset X_s$ for any $1 \leq j \leq m$. Apply Construction 8.1 to C_1, \dots, C_m , and let $\tau_i : \tilde{E}_i \rightarrow E_i$ ($1 \leq i \leq N$) be as in Construction 8.1. We need the following lemma, where F is arbitrary.

LEMMA 8.9. *For each $1 \leq i \leq N$, there is an integral regular closed subscheme $W_i \subset \tilde{E}_i$ of dimension one which satisfies the following three conditions:*

- (1) $\tau_i(W_i) (\subset E_i)$ has dimension one and degree prime to ℓ in the sense of Lemma 6.4.

- (2) $W_i \cap \tilde{C}$ is empty and $W_i \cap \tilde{E}_j$ is empty for any $j \neq i$.
- (3) W_i meets the strict transform of $X_{s,\text{red}}$ (in \tilde{X}) only in its regular locus and transversally.

Proof. Fix an i with $1 \leq i \leq N$. We first prove the case that F is infinite. Let $S_i \subset E_i$ be the finite subset as in Setting 6.3 (4), which may be empty. Let $C^{(i)}$ be the union of the strict transforms in $X^{(i)}$ of C_j 's with $j = 1, \dots, m$. Note that $\Sigma_i := C^{(i)} \cap E_i$ is finite, because $E_i \subset (X^{(i)})_s$ and any irreducible component of $C^{(i)}$ is not contained in $(X^{(i)})_s$. Let Z_1, \dots, Z_r be the irreducible components of $X_{s,\text{red}}$ whose strict transforms $Z_1^{(i)}, \dots, Z_r^{(i)} \subset X^{(i)}$ meet E_i . For $1 \leq \lambda \leq r$, put $H_\lambda := Z_\lambda^{(i)} \cap E_i$, which is a hyperplane on E_i . Since F is infinite by assumption, there is a line $l_i \subset E_i$ which does not meet $S_i \cup \Sigma_i$ and intersects H_λ transversally for any $1 \leq \lambda \leq r$. Then $W_i := \tau_i^{-1}(l_i) \subset \tilde{E}_i$ is the strict transform of l_i in \tilde{X} and we have $\tau_i|_{W_i} : W_i \xrightarrow{\sim} l_i$. One can easily check that W_i satisfies the conditions (1)–(3) in the lemma by the choice of $l_i \subset E_i$.

We next prove the case that F is finite. Put $F_i := \Gamma(E_i, \mathbb{O}_{E_i})$, the constant field of E_i . Fix a prime number ℓ' different from ℓ , and let F'_i/F_i be the maximal pro- ℓ' field extension. Since F'_i is infinite, there is a line $l_i \subset E_i \otimes_{F_i} F'_i$ which satisfies the same conditions as in the previous case with $(S_i, \Sigma_i, H_\lambda)$ replaced by $(S_i \otimes_{F_i} F'_i, \Sigma_i \otimes_{F_i} F'_i, H_\lambda \otimes_{F_i} F'_i)$. Let $l'_i \subset E_i$ be the image of l_i . Then $W_i := \tau_i^{-1}(l'_i) \subset \tilde{E}_i$ is the strict transform of l'_i in \tilde{X} and we have $\tau_i|_{W_i} : W_i \xrightarrow{\sim} l'_i$. The degree of l'_i in the sense of Lemma 6.4 is a power of ℓ' , which implies the condition (1). The conditions (2) and (3) follow from the choice of $l_i \subset E_i \otimes_{F_i} F'_i$. \square

We turn to the proof of Theorem 1.16 and fix W_1, \dots, W_N as in Lemma 8.9. By the condition (1) in Lemma 8.9 and Lemma 6.4 we have

$$\text{Ker}(\pi_* : H_2(\tilde{X}, \Lambda_n) \rightarrow H_2(X, \Lambda_n)) = \sum_{1 \leq i \leq N} \Lambda_n \langle W_i \rangle.$$

By the conditions (2) and (3) in Lemma 8.9 and Theorem 4.2, there exists an ample divisor $\iota : Y \hookrightarrow \tilde{X}$ which contains \tilde{C} and $\bigcup_{i=1}^N W_i$ and for which (\tilde{X}, Y) is an ample \mathcal{QSP} -pair. Here we have used the assumption $d \geq 3$ to ensure that $\dim(W_i) < \frac{1}{2} \dim(\tilde{E}_i)$ (cf. condition (iii) in Theorem 4.2). By the induction hypothesis, $\rho_Y : \text{CH}_1(Y)/\ell^n \rightarrow H_2(Y, \Lambda_n)$ is injective. Since $d \geq 3$, we have $H_3(\tilde{X} - Y, \Lambda_n) = 0$ by Theorem 3.2. Therefore we see that $\alpha = 0$ by the same arguments as in Step 1 using the diagram (8.4) with Λ_∞ replaced by Λ_n . This completes the proof of Theorem 1.16. \square

9. Applications of the main theorem

In this section we prove Corollaries 0.9–0.11 and Theorems 0.3 and 2.13. Throughout this section, ℓ denotes a prime number prime to $p := \text{ch}(F)$. For a scheme $X \in \text{Ob}(\mathcal{QS})$ and an effective Cartier divisor Y on X which belongs to

$\text{Ob}(\mathcal{C})$, we define the pull-back homomorphism

$$i^* : \text{CH}_q(X) \longrightarrow \text{CH}_{q-1}(Y) \quad (i : Y \hookrightarrow X)$$

by the same arguments as in [Ful98, §2.3 and Prop. 2.6(a)]. This map sends an integral closed subscheme $f : Z \hookrightarrow X$ of dimension q with $Z \not\subset Y$ to the Cartier divisor f^*Y on Z , regarded as a $(q-1)$ -cycle class on Y . If, further, X belongs to $\text{Ob}(\mathcal{QSP})$ and Y is over $s = \text{Spec}(F)$, then we define the intersection number $\langle \alpha, Y \rangle$ for $\alpha \in \text{CH}_1(X)$ to be the value of α under the composite map

$$\text{CH}_1(X) \xrightarrow{i^*} \text{CH}_0(Y) \xrightarrow{\deg_s} \mathbb{Z},$$

where the last arrow is the degree map over s (loc. cit. Def. 1.4). Note that the degree map is well-defined by the properness of Y . We first show

PROPOSITION 9.1. *Let X be an object of \mathcal{QSP} , and let $C \subset X$ be an integral closed subscheme of dimension one which is flat over B . Let x be the closed point of C . Let Y be an irreducible component of $X_{s,\text{red}}$ which contains x , and let i be the closed immersion $Y \hookrightarrow X$. Put $d := \dim(X) - 1$. Let $\text{cl}_X(-) \in H^{2d}(X, \Lambda_n(d))$ and $\text{cl}_Y(-) \in H^{2d}(Y, \Lambda_n(d))$ be the cycle class of 1-cycles on X and 0-cycles on Y , respectively ([Fuj02, Def. 1.1.2; see also loc. cit. Cor. 1.1.5]). Then:*

- (1) *We have $i^* \text{cl}_X(C) = \text{cl}_Y(i^*C)$ in $H^{2d}(Y, \Lambda_n(d))$.*
- (2) *$\langle C, Y \rangle$ is divisible by $\deg_s(x)$ and we have*

$$i^* \text{cl}_X(C) = \frac{\langle C, Y \rangle}{\deg_s(x)} \cdot \text{cl}_Y(x) \quad \text{in } H^{2d}(Y, \Lambda_n(d)).$$

Proof. Put $\varepsilon := C \times_X Y$, which is the spectrum of an artinian local ring A and represents the 0-cycle i^*C on Y . We have $\langle C, Y \rangle = \text{length}_A(A) \cdot \deg_s(x)$ by the definition of the intersection number $\langle -, Y \rangle$. We have $i^* \text{cl}_X(C) = \text{cl}_Y(\varepsilon)$ by [Fuj02, Prop. 1.1.9], and $\text{cl}_Y(\varepsilon) = \text{length}_A(A) \cdot \text{cl}_Y(x)$ by [SGA $_{\frac{1}{2}}$, Cycle Th. 2.3.8(i)]. The assertions follow from these facts. \square

We prove Corollaries 0.9–0.11.

COROLLARY 9.2 (Corollary 0.9). *Assume that F is separably closed. Let X be an object of \mathcal{QSP} , and let Y_1, \dots, Y_N be the irreducible components of $X_{s,\text{red}}$. Then the assignment $\alpha \in \text{CH}_1(X) \mapsto \langle \alpha, Y_j \rangle_{1 \leq j \leq N} \in \mathbb{Z}^N$ induces an isomorphism*

$$\text{CH}_1(X)/\ell^n \xrightarrow{\cong} \bigoplus_{1 \leq j \leq N} \mathbb{Z}/\ell^n \mathbb{Z} \quad \text{for any } n \geq 1.$$

Proof. Put $d := \dim(X) - 1$. We note the following isomorphisms analogous to (2.10):

(9.3)

$$H_2(X, \Lambda_n) \stackrel{(1.9)}{\cong} H^{2d}(X, \Lambda_n(d)) \xrightarrow{\cong} \bigoplus_{1 \leq j \leq N} H^{2d}(Y_j, \Lambda_n(d)) \xrightarrow{\cong} \bigoplus_{1 \leq j \leq N} \Lambda_n,$$

where the second arrow is bijective by a similar argument as for Lemma 2.15 (2). The last arrow is the direct sum of the trace maps for $Y_j \rightarrow s$ ([SGA4, XVIII, Th. 2.9]), which is bijective by [Mil80, VI, Lemma 11.3]. Hence the corollary follows from Theorem 1.16, Proposition 7.1 and the following lemma:

LEMMA 9.4. *Let $C \subset X$ be an integral closed subscheme of dimension one which is flat over B . Then the composite map (9.3) sends the cycle class $\rho_X(C)$ to $\langle C, Y_j \rangle_{1 \leq j \leq N}$.*

Proof of Lemma 9.4 Let x be the closed point of C . By Lemma 1.8 (1), the first isomorphism in (9.3) sends $\rho_X(C)$ to the cycle class $\text{cl}_X(C)$. By Proposition 9.1 (2), we have

$$i_j^* \text{cl}_X(C) = \frac{\langle C, Y_j \rangle}{\deg_s(x)} \cdot \text{cl}_{Y_j}(x) \quad \text{in } H^{2d}(Y_j, \Lambda_n(d))$$

for $1 \leq j \leq N$. Here i_j denotes the closed immersion $Y_j \hookrightarrow X$, and $\text{cl}_{Y_j}(x)$ means zero if $x \notin Y_j$. The assertion of the lemma follows from this formula. Indeed if $x \in Y_j$, then the trace map $H^{2d}(Y_j, \Lambda_n(d)) \rightarrow \Lambda_n$ sends $\text{cl}_{Y_j}(x)$ to $\deg_s(x)$ by [SGA $\frac{1}{2}$, Cycle Th. 2.3.8 (i)]. This completes the proofs of Lemma 9.4 and Corollary 9.2. \square

COROLLARY 9.5 (Corollary 0.10). *Assume that X is smooth and projective over B and that F is finite or separably closed. Then for any $n \geq 1$,*

$$s\text{CH}_0(X_s)/\ell^n \xleftarrow{i^*} \text{CH}_1(X)/\ell^n \xrightarrow{j^*} \text{CH}_0(X_\eta)/\ell^n \quad (X_s \xrightarrow{i} X \xleftarrow{j} X_\eta).$$

Here $\text{CH}_0(X_\eta)$ denotes the Chow group of 0-cycles on X_η in the usual sense (cf. Remark 1.3).

Proof. By Lemma 7.2 and Proposition 9.1 (1), there is a commutative diagram

$$\begin{array}{ccc} \text{CH}_1(X)/\ell^n & \xrightarrow{\text{cl}_X} & H^{2d}(X, \Lambda_n(d)) \\ i^* \downarrow & & \downarrow i^* \\ \text{CH}_0(X_s)/\ell^n & \xrightarrow{\text{cl}_{X_s}} & H^{2d}(X_s, \Lambda_n(d)). \end{array}$$

where cl_X and cl_{X_s} send cycles to their cycle classes. See [JSS, Th. 1.1.1] for why these assignments factor through the Chow groups. The right vertical i^* is bijective by [SGA4, XII Cor. 5.5 (iii)], and cl_X is bijective by Theorem 1.16 and Lemma 1.8. The arrow cl_{X_s} is obviously bijective if F is separably closed. If F is finite, then cl_{X_s} is bijective by the unramified class field theory for the smooth and projective F -variety X_s ([KS83], [CTSS83, p. 792, Th. 5; see also p.793, Rem. 3]). Hence the left vertical i^* is bijective.

To show the bijectivity of j^* in the corollary, we note the following exact sequence, which follows from the same argument as for [Ful98, Prop. 1.8]:

$$\text{CH}_1(X_s)/\ell^n \xrightarrow{i^*} \text{CH}_1(X)/\ell^n \xrightarrow{j^*} \text{CH}_0(X_\eta)/\ell^n \longrightarrow 0.$$

We show the map i_* is zero. Because X_s is a principal divisor on X , the composite map

$$\mathrm{CH}_1(X_s)/\ell^n \xrightarrow{i_*} \mathrm{CH}_1(X)/\ell^n \xrightarrow[\simeq]{i^*} \mathrm{CH}_0(X_s)/\ell^n$$

is zero by [Ful98, Prop. 2.6 (c)]. Hence the bijectivity of i^* implies that i_* is zero. Thus we obtain Corollary 9.5. \square

COROLLARY 9.6 (Corollary 0.11). *Let (X, Y) be an ample \mathcal{QSP} -pair with $\dim(X) = d + 1 \geq 2$, and let $i : Y \hookrightarrow X$ be the natural closed immersion. Assume that F is finite (resp. separably closed). Then for any $n \geq 1$, the push-forward map $i_* : \mathrm{CH}_1(Y)/\ell^n \rightarrow \mathrm{CH}_1(X)/\ell^n$ is bijective for $d \geq 3$ and surjective for $d = 2$ (resp. bijective for $d \geq 2$ and surjective for $d = 1$).*

Proof. Put $\Lambda := \Lambda_n$. By Propositions 1.5 (2) and 1.13 (2), there is a commutative diagram with exact lower row

$$\begin{array}{ccccccc} \mathrm{CH}_1(Y)/\ell^n & \xrightarrow{i_*} & \mathrm{CH}_1(X)/\ell^n & & & & \\ \rho_Y \downarrow \simeq & & \rho_X \downarrow \simeq & & & & \\ H_3(U, \Lambda_n) & \longrightarrow & H_2(Y, \Lambda_n) & \xrightarrow{i_*} & H_2(X, \Lambda_n) & \longrightarrow & H_2(U, \Lambda_n), \end{array}$$

where the vertical arrows are bijective by Theorem 1.16. Corollary 0.11 follows from this diagram and the vanishing results in Theorem 3.2. \square

Next we prove the following result:

THEOREM 9.7 (Theorem 0.3). *Assume that R is a p -adic integer ring. Let X be an object of \mathcal{QSP} , and put $V := X_\eta$. Then:*

- (1) *The group $A_0(V) \hat{\otimes} \mathbb{Z}_\ell := \varprojlim_{n \in \mathbb{N}} A_0(V)/\ell^n$ is finite for any prime $\ell \neq p$ and is zero for almost all primes ℓ .*
- (2) *Conjecture 0.2 holds true for V .*

Proof. We first show (1). By Theorem 4.1 and Corollary 9.6, the problem is reduced to the case $\dim(V) = 1$. Let J be the Jacobian variety of V . Then $A_0(V)$ is a subgroup of $J(k)$ of finite index, and the assertion follows from a theorem of Mattuck [Mat55] and [JS03, Lemma 7.8].

The assertion (2) follows from (1) and loc. cit. Lemma 7.7 applied to

$$\begin{aligned} & (A, \{B_n\}_{n \in \mathbb{N}, (n,p)=1}, \{\varphi_n : A/n \rightarrow B_n\}_{n \in \mathbb{N}, (n,p)=1}) \\ &= (A_0(V), \{A_0(V)/n\}_{n \in \mathbb{N}, (n,p)=1}, \{\mathrm{id} : A_0(V)/n \rightarrow A_0(V)/n\}_{n \in \mathbb{N}, (n,p)=1}). \end{aligned}$$

This completes the proof of Theorem 9.7. \square

Finally we prove Theorem 2.13. The case $a \leq 2$ has been shown in Proposition 2.14 and Theorem 5.3. Let Λ denote Λ_n if F is separably closed, and denote Λ_∞ if F is finite. Let X be an object of \mathcal{QSP} . We have to show $KH_3(X, \Lambda) = 0$. We proceed to the proof by induction on $\dim(X)$. The assertion is clear if $\dim(X) \leq 2$.

Assume $\dim(X) \geq 3$. By Theorem 4.2 we take an ample divisor $Y \subset X$ such that (X, Y) is a \mathcal{LSP} -pair, and put $U := X - Y$. By the exact sequence of complexes in (2.3), there is a long exact sequence

$$\cdots \longrightarrow KH_3(Y, \Lambda) \longrightarrow KH_3(X, \Lambda) \longrightarrow KH_3(U, \Lambda) \longrightarrow \cdots.$$

By the induction hypothesis, it is enough to show $KH_3(U, \Lambda)$ is zero. By (1.11) and Lemma 2.6, the niveau spectral sequence (2.2) yields an exact sequence

$$H_3(U, \Lambda) \longrightarrow KH_3(U, \Lambda) \longrightarrow \mathrm{CH}_1(U) \otimes \Lambda.$$

By Theorem 3.2 we have $H_3(U, \Lambda) = 0$. By Corollary 9.6 and Proposition 1.13 (2), we have $\mathrm{CH}_1(U) \otimes \Lambda = 0$. Hence $KH_3(U, \Lambda)$ is zero. This completes the proof of Theorem 2.13. \square

Appendix A. Resolution of singularities for embedded curves

by Uwe Jannsen

The purpose of this appendix is to show the following form of resolution of singularities for curves embedded in a regular scheme, which has been used in the proof of Theorem 1.16 of the main body (cf. §8). We believe that it has been known to experts but there is no written proof of it as far as we know.

For a noetherian scheme X and a closed subscheme $Z \subset X$, let

$$\pi : \mathrm{Bl}_Z(X) = \mathrm{Proj} \left(\bigoplus_{n \geq 0} \mathcal{I}_Z^n \right) \longrightarrow X$$

denote the blow-up of X in Z , i.e., the blow-up of the ideal sheaf $\mathcal{I}_Z \subset \mathcal{O}_X$ defining Z , and let $E_Z(X) = Z \times_X \mathrm{Bl}_Z(X)$ denote the exceptional divisor. If $i : Y \hookrightarrow X$ is another closed immersion, then the strict transform of Y in $\mathrm{Bl}_Z(X)$ is by definition $\mathrm{Bl}_{Z \times_X Y}(Y)$, the blow-up of the inverse image ideal sheaf $i^{-1} \mathcal{I}_Z \cdot \mathcal{O}_Y \subset \mathcal{O}_Y$. By the universal property of blow-ups there is a canonical morphism $\mathrm{Bl}_{Z \times_X Y}(Y) \rightarrow \mathrm{Bl}_Z(X)$, which is a closed immersion.

THEOREM A.1. *Let X be a regular excellent scheme and let Y be a reduced divisor with simple normal crossings on X (cf. Definition 1.15 (1)). Let W be a one-dimensional reduced closed subscheme of X such that any irreducible component of W is not contained in Y . Then there exists a proper morphism $\pi : X' \rightarrow X$ which satisfies the following conditions:*

- (i) *π is the composite of a sequence of blow-ups $X' = X^{(n)} \rightarrow \cdots \rightarrow X^{(1)} \rightarrow X$ at closed points on W and closed points on the strict transforms of W in $X^{(i)}$ ($1 \leq i \leq n-1$).*
- (ii) *The closed subset $\pi^{-1}(Y)$ with reduced subscheme structure is a reduced divisor with simple normal crossings.*
- (iii) *The strict transform $W' \subset X'$ of W is regular and intersects Y' transversally, which means that W' intersects Y' only in the regular locus of Y' and transversally.*

By Proposition A.4 (3) below, the condition (ii) in Theorem A.1 is a consequence of the condition (i). The proof of the theorem is achieved in several steps. The following statement should be well-known, but we include a proof for lack of a suitable reference.

PROPOSITION A.2. *Let X be a reduced excellent scheme of dimension one. Then after a sequence of blow-ups $X^{(n)} \rightarrow \cdots \rightarrow X^{(1)} \rightarrow X$ at singular closed points, the resulting scheme $X^{(n)}$ is regular.*

Proof. Since X is reduced and excellent, it has a regular dense open subset by [EGAIV, 7.8.6 (iii)]. Because blow-ups are compatible with Zariski localization, we may assume that X is an affine scheme $\text{Spec}(R)$. Let x be a singular closed point of X , and let $\mathfrak{m} \subset R$ be the maximal ideal that corresponds to x . Let π be the blow-up of X at x :

$$\pi : X' := \text{Bl}_x(X) = \text{Proj}(S) \longrightarrow X \quad \text{with} \quad S := \bigoplus_{n \geq 0} \mathfrak{m}^n.$$

We first show

LEMMA A.3. *π is a finite morphism. In particular, X' is affine by [EGAII, 6.7.1].*

Proof of Lemma A.3. Since π is projective, it is enough to show that π is quasi-finite (cf. [Mil80, I, Th. 1.8]). Since π is an isomorphism over $X \setminus \{x\}$, it suffices to show that the fiber

$$\pi^{-1}(x) = \text{Proj}(\bar{S}) \quad \text{with} \quad \bar{S} := S \otimes_R R/\mathfrak{m} = \bigoplus_{n \geq 0} \mathfrak{m}^n / \mathfrak{m}^{n+1}$$

consists of a finite number of points. Let $R_{\mathfrak{m}}$ be the local ring $\mathcal{O}_{X,x}$, and let $\mathfrak{m}R_{\mathfrak{m}}$ be its maximal ideal. Put $k := R/\mathfrak{m} = R_{\mathfrak{m}}/\mathfrak{m}R_{\mathfrak{m}}$. There is an isomorphism of finitely generated k -algebras

$$\bar{S} \simeq \bigoplus_{n \geq 0} (\mathfrak{m}R_{\mathfrak{m}})^n / (\mathfrak{m}R_{\mathfrak{m}})^{n+1} =: \text{gr}^{\bullet} R_{\mathfrak{m}}.$$

Because $R_{\mathfrak{m}}$ is one-dimensional by assumption, $\text{gr}^{\bullet} R_{\mathfrak{m}}$ has dimension one as well by the theory of noetherian local rings. Hence $\pi^{-1}(x) = \text{Proj}(\text{gr}^{\bullet} R_{\mathfrak{m}})$ has dimension zero by [Kun85, II, Prop. 4.4 (b)], and the lemma follows. \square

We turn to the proof of Proposition A.2. Let X_1, \dots, X_N be the irreducible components of $X = \text{Spec}(R)$. Since X is reduced, we may write $X_i = \text{Spec}(R_i)$ ($1 \leq i \leq N$), where R_i is an integral domain of dimension one. Let \tilde{R}_i be the normalization of R_i in its fraction field K_i . Put $K := K_1 \times \cdots \times K_N$ and $\tilde{R} := \tilde{R}_1 \times \cdots \times \tilde{R}_N$. We want to show that after a sequence of blow-ups $X^{(n)} \rightarrow \cdots \rightarrow X^{(1)} = X' \rightarrow X$ at singular points, we get $X^{(n)} = \text{Spec}(\tilde{R})$. By Lemma A.3 we have $X' = \text{Spec}(R_1)$ for some finite R -algebra R_1 . Since X is reduced by assumption, R_1 is reduced. Hence we have inclusions $R \subset R_1 \subset \tilde{R} \subset K$. As R

is excellent, \tilde{R} is finite over R by [EGAIV, 7.8.6(ii)]. As $\dim(R) = 1$, \tilde{R}/R has finite length over R (it is supported in the finitely many singular points of $\operatorname{Spec}(R)$). By [EGAII, 8.1.7], the ideal $\mathfrak{m}R_1 \subset R_1$ is invertible, while $\mathfrak{m} \subset R$ is not invertible because x is a singular point of $\operatorname{Spec}(R)$ and $\mathfrak{m}R_{\mathfrak{m}} \subset R_{\mathfrak{m}}$ is not principal. Therefore $R_1 \neq R$, and $\ell_{R_1}(\tilde{R}/R_1) \leq \ell_R(\tilde{R}/R_1) < \ell_R(\tilde{R}/R)$, where $\ell_R(-)$ denotes the length over R . Hence we obtain Proposition A.2 by induction on $\ell_R(\tilde{R}/R)$. \square

PROPOSITION A.4. *Let X be an integral regular noetherian scheme of dimension d , and let $Z \subset X$ be an integral regular closed subscheme with $Z \neq X$. Then:*

- (1) $\operatorname{Bl}_Z(X)$ is again integral and regular of dimension d , and the exceptional divisor $E_Z(X)$ is integral and regular.
- (2) Let $Y_1, Y_2 \subset X$ be regular closed subschemes containing Z . Assume that Y_1 and Y_2 intersect transversally. Then the canonical morphism

$$\operatorname{Bl}_Z(Y_1 \times_X Y_2) \longrightarrow \operatorname{Bl}_Z(Y_1) \times_{\operatorname{Bl}_Z(X)} \operatorname{Bl}_Z(Y_2)$$

is an isomorphism.

- (3) Let $Y \subset X$ be a reduced divisor with simple normal crossings on X and let Y_1, \dots, Y_r be the irreducible components of Y . Assume that all Y_i contain Z . Then the union of $E_Z(X)$ and the strict transforms of Y_i in $\operatorname{Bl}_Z(X)$ for $1 \leq i \leq r$ is a reduced divisor with simple normal crossings on $\operatorname{Bl}_Z(X)$.

Proof. We first show (1). Since X is integral and $Z \neq X$, $\operatorname{Bl}_Z(X)$ is integral of dimension d by [EGAII, 8.1.4]. Since X and Z are regular, $\operatorname{Bl}_Z(X)$ is regular by [EGAIV, 19.4.3 and 19.4.4]. It follows from loc. cit. 19.4.2, that $E_Z(X)$ is isomorphic to the projective bundle over Z associated to the conormal sheaf of Z in X . Hence $E_Z(X)$ is integral and regular. To prove Proposition A.4 (2), we need the following standard fact (cf. [Ful98, Rem. A.6]).

LEMMA A.5. *Let A be a commutative ring with unity, and let $a_1, \dots, a_m \in A$ be a regular sequence. Put $I := (a_1, \dots, a_m) \subset A$. Then there is an isomorphism of graded A -algebras*

$$A[T_1, \dots, T_m]/(a_i T_j - a_j T_i \mid 1 \leq i < j \leq m) \xrightarrow{\cong} \bigoplus_{n \geq 0} I^n, \quad T_i \mapsto a_i \in I.$$

We start the proof of Proposition A.4 (2). Since the problems are local in the Zariski topology, we may assume that $X = \operatorname{Spec}(A)$ for a noetherian regular local ring A with maximal ideal \mathfrak{m} . Put $d := \dim(A)$. By the assumptions on Y_1, Y_2 and Z , there is a regular system of parameters $\{a_1, \dots, a_d\}$ of A such that $Z = \operatorname{Spec}(A/I)$ and $Y_i = \operatorname{Spec}(A/I_i)$ ($i = 1, 2$), where

$$I = (a_1, \dots, a_m), \quad I_1 = (a_1, \dots, a_\mu), \quad I_2 = (a_{\mu+1}, \dots, a_{\mu+\nu})$$

with $\mu + \nu \leq m := \text{codim}_X(Z)$. By Lemma A.5, we have

$$\text{Bl}_Z(X) \simeq \text{Proj}(B), \quad \text{Bl}_Z(Y_i) \simeq \text{Proj}(B_i) \quad (i = 1, 2), \quad \text{Bl}_Z(Y_1 \times Y_2) \simeq \text{Proj}(B_{12}),$$

where $B := A[T_1, \dots, T_m]/(a_i T_j - a_j T_i \mid 1 \leq i < j \leq m)$ and

$$\begin{aligned} B_1 &:= (A/I_1)[T_{\mu+1}, \dots, T_m]/(a_i T_j - a_j T_i \mid \mu+1 \leq i < j \leq m) \\ &= B/(T_1, \dots, T_\mu, a_1, \dots, a_\mu), \\ B_2 &:= (A/I_2)[T_1, \dots, T_\mu, T_{\mu+\nu+1}, \dots, T_m]/(a_i T_j - a_j T_i \mid i, j \in [1, \mu] \cup [\mu+\nu+1, m]) \\ &= B/(T_{\mu+1}, \dots, T_{\mu+\nu}, a_{\mu+1}, \dots, a_{\mu+\nu}), \\ B_{12} &:= (A/(I_1 + I_2))[T_{\mu+\nu+1}, \dots, T_m]/(a_i T_j - a_j T_i \mid \mu+\nu+1 \leq i < j \leq m) \\ &= B/(T_1, \dots, T_{\mu+\nu}, a_1, \dots, a_{\mu+\nu}). \end{aligned}$$

The isomorphism in question is a direct consequence of these descriptions.

Finally we prove Proposition A.4 (3). By (1), the exceptional divisor $E_Z(X) \subset \text{Bl}_Z(X)$ and the strict transforms $\text{Bl}_Z(Y_i)$ of Y_i ($1 \leq i \leq r$) are regular divisors on $\text{Bl}_Z(X)$. For $p \geq 1$ and $1 \leq i_1 < i_2 < \dots < i_p \leq s$, let

$$Y_{i_1, \dots, i_p} = Y_{i_1} \times_X \dots \times_X Y_{i_p}$$

be the p -fold intersection of irreducible components of Y . Applying (2) repeatedly, we have

$$\text{Bl}_Z(Y_{i_1}) \times_{\text{Bl}_Z(X)} \dots \times_{\text{Bl}_Z(X)} \text{Bl}_Z(Y_{i_p}) \simeq \text{Bl}_Z(Y_{i_1, \dots, i_p}),$$

where the scheme on the right-hand side is regular by (1). This shows that the reduced divisor $\bigcup_{i=1}^r \text{Bl}_Z(Y_i)$ has simple normal crossings. As for the intersection of $\text{Bl}_Z(Y_{i_1, \dots, i_p})$ and $E_Z(X) = Z \times_X \text{Bl}_Z(X)$, we have

$$\begin{aligned} E_Z(X) \times_{\text{Bl}_Z(X)} \text{Bl}_Z(Y_{i_1, \dots, i_p}) &= Z \times_X \text{Bl}_Z(X) \times_{\text{Bl}_Z(X)} \text{Bl}_Z(Y_{i_1, \dots, i_p}) \\ &= Z \times_X \text{Bl}_Z(Y_{i_1, \dots, i_p}) = Z \times_{Y_{i_1, \dots, i_p}} \text{Bl}_Z(Y_{i_1, \dots, i_p}), \end{aligned}$$

which is the exceptional divisor of the blow-up $\text{Bl}_Z(Y_{i_1, \dots, i_p}) \rightarrow Y_{i_1, \dots, i_p}$ and is a regular divisor on $\text{Bl}_Z(Y_{i_1, \dots, i_p})$ by (1). This shows that $E_Z(X)$ intersects all Y_{i_1, \dots, i_p} 's transversally. This completes the proof of Proposition A.4. \square

PROPOSITION A.6. *Let X be a regular noetherian scheme, let D be a union of finitely many regular divisors on X , and let $C \subset X$ be an integral regular closed subscheme of dimension one. Assume that C is not contained in D . Let $X^{(1)}$ be the blow-up of X at the (closed) points of $C \cap D$. Let $C^{(1)}$ be the strict transform of C in $X^{(1)}$, and let $D^{(1)}$ be the union of the strict transform of D and the exceptional divisors of the blow-up $X^{(1)} \rightarrow X$. For $n \geq 2$, define $(X^{(n)}, C^{(n)}, D^{(n)})$ inductively by applying this procedure to $(X^{(n-1)}, C^{(n-1)}, D^{(n-1)})$. Then $C^{(n)}$ intersects $D^{(n)}$ only in the regular locus of $D^{(n)}$ and transversally, for n large enough.*

Proof. Because $C \cap D$ is discrete and the question is local around the points $x \in C \cap D$, we may localize X to assume $X = \operatorname{Spec}(A)$ for a noetherian regular local ring A and that x corresponds to the maximal ideal \mathfrak{m} of A . Suppose that A has dimension d , and take a regular system of parameters $\{a_1, \dots, a_d\}$ of A for which $C = \operatorname{Spec}(\bar{A})$ with $\bar{A} := A/(a_1, \dots, a_{d-1})$. Let D_1, \dots, D_s be the irreducible components of D . For $1 \leq i \leq s$, we have $D_i = \operatorname{Spec}(A/(f_i))$ for some $f_i \in A$. The assumption that C is not contained in D implies that $f_i \notin (a_1, \dots, a_{d-1})$ for any i . We now fix $i \in [1, s]$. By assumption, $\bar{A} = A/(a_1, \dots, a_{d-1})$ is a discrete valuation ring with a prime element $a_d \bmod (a_1, \dots, a_{d-1}) \in \bar{A}$. We define an integer $r_x(C, D_i)$ as the (normalized) valuation of $f_i \bmod (a_1, \dots, a_{d-1}) \in \bar{A}$. It depends only on the triple (C, D_i, x) and not on the choice of a regular system of parameters. We will prove

LEMMA A.7. *Let $\pi : \tilde{X} = \operatorname{Bl}_x(X) \rightarrow X$ be the blow-up of $X = \operatorname{Spec}(A)$ at x , and let $E \subset \tilde{X}$ be the exceptional divisor. Let $\tilde{C} \subset \tilde{X}$ (resp. $\tilde{D}_i \subset \tilde{X}$) be the strict transform of C (resp. D_i). Then:*

- (1) \tilde{C} and E meet transversally.
- (2) If $r_x(C, D_i) = 1$, then $\tilde{C} \cap \tilde{D}_i$ is empty.
- (3) If $r_x(C, D_i) \geq 2$, then $r_y(\tilde{C}, \tilde{D}_i) < r_x(C, D_i)$ for any point $y \in \tilde{C} \cap \tilde{D}_i$.

These statements imply that for a sufficiently large n , the strict transform $C^{(n)}$ of C in $X^{(n)}$ meets only the exceptional divisor $E^{(n)}$ of the last blow-up among the irreducible components of the inverse image of D . Moreover, $C^{(n)}$ intersects $E^{(n)}$ transversally. This implies Proposition A.6. It remains to prove the lemma.

Proof of Lemma A.7. We show (1). By Lemma A.5 we have

$$\begin{aligned} \tilde{X} &= \operatorname{Proj}(B) \quad \text{with} \quad B = A[T_1, \dots, T_d]/(a_i T_j - a_j T_i \mid 1 \leq i < j \leq d) \\ \tilde{C} &= \operatorname{Proj}(B') \quad \text{with} \quad B' = \bar{A}[T_d], \end{aligned}$$

where \tilde{C} is of course isomorphic to C . Moreover, the closed immersion $\tilde{C} \hookrightarrow \tilde{X}$ is induced by the ring homomorphism $B \rightarrow B'$ sending $T_i \mapsto 0$ for $i \neq d$. This shows that \tilde{C} lies in the chart $V_+(T_d) = \{T_d \neq 0\} \subset \operatorname{Proj}(B)$, which is the spectrum of the ring

$$(A.8) \quad A[t_1, \dots, t_{d-1}]/(a_d t_i - a_i \mid i = 1, \dots, d-1) \quad (t_i = T_i/T_d)$$

In this chart, $E \cap V_+(T_d)$ (resp. $\tilde{C} \cap V_+(T_d)$) is described as the locus where $a_d = 0$ (resp. $t_1 = \dots = t_{d-1} = 0$). This immediately implies Lemma A.7 (1).

To show Lemma A.7 (2) and (3), we need to compute the strict transform \tilde{D}_i of D_i . For simplicity we put $D := D_i$, $f := f_i$ and $r := r_x(C, D)$. We normalize the regular system $\{a_1, \dots, a_d\}$ as follows. We have $f = u \cdot a_d^r + g$ for some $g \in (a_1, \dots, a_{d-1})$ and some $u \in A^\times$. Replacing f by $u^{-1}f$, we suppose $f = a_d^r + g$. We distinguish two cases. If $r = 1$, then D intersects C transversally.

Replacing a_d with f , we suppose

$$f = a_d.$$

If $r \geq 2$, then we have $f = a_d^r + a_1 g_1 + \cdots + a_{d-1} g_{d-1}$ for some $g_j \in A$. Note that not all g_j 's are contained in \mathfrak{m} since $f \notin \mathfrak{m}^2$ by regularity of D . If $g_k \notin \mathfrak{m}$, then g_k is a unit and we replace a_k with $a_1 g_1 + \cdots + a_{d-1} g_{d-1}$. By renumbering, if necessary, we suppose

$$f = a_d^r + a_1.$$

Because \tilde{C} is contained in the affine chart $V_+(T_d) \subset \tilde{X}$, we compute $\tilde{D} \cap V_+(T_d)$. Note that \tilde{D} is an integral closed subscheme of $\pi^{-1}(D) = D \times_X \tilde{X}$, and that $\pi^{-1}(D) \cap V_+(T_d)$ is defined by f , regarded as an element of the ring (A.8). When $r = 1$, we assumed $f = a_d$, which defines E on $V_+(T_d)$. Since $\tilde{D} \not\subset E$, this implies $\tilde{D} \cap V_+(T_d) = \emptyset$ and hence \tilde{D} does not meet \tilde{C} . This shows Lemma A.7 (2). Now consider the case $r \geq 2$, where we assumed $f = a_d^r + a_1$. In the ring (A.8), we have a factorization

$$f = a_d(a_d^{r-1} + t_1),$$

which implies that the divisor $\tilde{D} \cap V_+(T_d) \subset V_+(T_d)$ is defined by $a_d^{r-1} + t_1$. Since \tilde{C} is defined by the equations $t_1 = \cdots = t_{d-1} = 0$, we see that \tilde{D} meets \tilde{C} only at the point $y \in V_+(T_d)$ corresponding to the ideal $(a_d, t_1, \dots, t_{d-1})$, and that $r_y(\tilde{C}, \tilde{D}) = r - 1$. This shows Lemma A.7 (3), and the proof of Proposition A.6 is complete. \square

We deduce Theorem A.1 from Propositions A.2, A.4 and A.6. Let $Y \subset X$ and $W \subset X$ be as in the theorem. Since X is excellent by assumption, Z is excellent as well by [EGAIV, 7.8.6 (i)]. After a sequence of blow-ups $X^{(n)} \rightarrow \cdots \rightarrow X^{(1)} \rightarrow X$ at singular points of W and singular points of the strict transform $W^{(i)} \subset X^{(i)}$ of W ($1 \leq i \leq n-1$), the resulting strict transform $W^{(n)} \subset X^{(n)}$ of W is regular by Proposition A.2. The inverse image $Y^{(n)} \subset X^{(n)}$ of Y endowed with reduced subscheme structure remains a reduced divisor with simple normal crossings by Proposition A.4. Finally applying Proposition A.6 to $(X, D, C) = (X^{(n)}, Y^{(n)}, W^{(n)})$, we obtain a desired proper morphism $\pi : X' \rightarrow X$. This completes the proof of Theorem A.1. \square

Acknowledgments. The first author thanks Prof. Uwe Jannsen for inspiration on this work obtained from joint works [JS03], [JSa], and [JSb] on the Kato conjecture and its relation with motivic cohomology of arithmetic schemes. The authors thank him also for his generous permission to include his note on resolution of singularities for embedded curves as the appendix in our paper. They are grateful to Prof. Jean-Louis Colliot-Thélène for his helpful comments on the first version of this paper and his lucid expository paper [CTa] on the subject of this paper based on his Bourbaki talk. In particular, Corollary 0.4 was suggested by him. Thanks are also due to Tamás Szamuely for his valuable comments. Finally the authors are grateful to the referee for offering numerous constructive comments to improve greatly the presentation of our paper.

References

- [SGA4] M. ARTIN, A. GROTHENDIECK, and J. L. VERDIER (eds.), *Théorie des topos et cohomologie étale des schémas. Tome 2, Lecture Notes in Math.* **270**, Springer-Verlag, New York, 1972, Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4), Dirigé par M. Artin, A. Grothendieck et J. L. Verdier. Avec la collaboration de N. Bourbaki, P. Deligne et B. Saint-Donat. MR 50 #7131 Zbl 0237.00012
- [BO74] S. BLOCH and A. OGUS, Gersten’s conjecture and the homology of schemes, *Ann. Sci. École Norm. Sup.* **7** (1974), 181–201 (1975). MR 54 #318 Zbl 0307.14008
- [CTa] J.-L. COLLIOT-THÉLÈNE, Groupe de Chow des zéro-cycles sur les variétés p -adiques (d’après S. Saito, K. Sato et al.), *Séminaire N. Bourbaki* 2009–2010, n° 1012. arXiv 1004.1371 arXiv
- [CTb] ———, R -equivalence and rational equivalence on varieties over p -adic fields, with special regards to rationally connected varieties, notes for the Bangalore conference (2003) and the Sestri Levante summer school (2004).
- [CTc] ———, Zéro-cycles sur les surfaces sur un corps p -adique: quelques résultats obtenus au moyen de la K -théorie algébrique, notes for the Bangalore conference (2003) and the Sestri Levante summer school (2004).
- [CT95] ———, L’arithmétique du groupe de Chow des zéro-cycles, *J. Théor. Nombres Bordeaux* **7** (1995), 51–73. MR 97i:14006 Zbl 0870.14002
- [CT05] ———, Un théorème de finitude pour le groupe de Chow des zéro-cycles d’un groupe algébrique linéaire sur un corps p -adique, *Invent. Math.* **159** (2005), 589–606. MR 2005m:14009 Zbl 1080.14012
- [CTOP02] J.-L. COLLIOT-THÉLÈNE, M. OJANGUREN, and R. PARIMALA, Quadratic forms over fraction fields of two-dimensional Henselian rings and Brauer groups of related schemes, in *Algebra, Arithmetic and Geometry, Part I* (Mumbai, 2000), *Tata Inst. Fund. Res. Stud. Math.* **16**, Tata Inst. Fund. Res., Bombay, 2002, pp. 185–217. MR 2004c:14031 Zbl 1055.14019
- [CTSS83] J.-L. COLLIOT-THÉLÈNE, J.-J. SANSUC, and C. SOULÉ, Torsion dans le groupe de Chow de codimension deux, *Duke Math. J.* **50** (1983), 763–801. MR 85d:14010 Zbl 0574.14004
- [Dal05] C. S. DALAWAT, Le groupe de Chow d’une surface rationnelle sur un corps local, *Compos. Math.* **141** (2005), 344–358. MR 2006a:14005 Zbl 1131.14009
- [dJ96] A. J. DE JONG, Smoothness, semi-stability and alterations, *Inst. Hautes Études Sci. Publ. Math.* (1996), 51–93. MR 98e:14011 Zbl 0916.14005
- [Del80] P. DELIGNE, La conjecture de Weil. II, *Inst. Hautes Études Sci. Publ. Math.* (1980), 137–252. MR 83c:14017 Zbl 0456.14014
- [SGA $\frac{1}{2}$] P. DELIGNE, J.-F. BOUTOT, A. GROTHENDIECK, L. ILLUSIE, and J.-L. VERIDER, *Cohomologie Étale, Lecture Notes in Math.* **569**, Springer-Verlag, New York, 1977.
- [Fuj02] K. FUJIWARA, A proof of the absolute purity conjecture (after Gabber), in *Algebraic Geometry 2000, Azumino (Hotaka)*, *Adv. Stud. Pure Math.* **36**, Math. Soc. Japan, Tokyo, 2002, pp. 153–183. MR 2004d:14015 Zbl 1059.14026
- [Ful98] W. FULTON, *Intersection Theory*, second ed., *Ergeb. Math. Grenzgeb.* **2**, Springer-Verlag, New York, 1998. MR 99d:14003 Zbl 0885.14002
- [Gro68] A. GROTHENDIECK, Le groupe de Brauer. I. Algèbres d’Azumaya et interprétations diverses, in *Dix Exposés sur la Cohomologie des Schémas*, North-Holland, Amsterdam, 1968, pp. 46–66. MR 39 #5586a Zbl 0193.21503

- [SGA5] A. GROTHENDIECK, I. BUCUR, C. HOUZEL, L. ILLUSIE, J.-P. JOUANLOU, and J.-P. SERRE, *Cohomologie ℓ -adique et Fonctions L* , *Lecture Notes in Math.* **589**, Springer-Verlag, New York, 1977. Zbl 0345.00011
- [EGAII] A. GROTHENDIECK and J. DIEUDONNÉ, Étude globale élémentaire de quelques classes de morphismes, *Inst. Hautes Études Sci. Publ. Math.* **8** (1961).
- [EGAIV] ———, Étude locale des schémas et des morphismes de schémas, *Inst. Hautes Études Sci. Publ. Math.* **20** (1964), **24** (1965), **28** (1966), **32** (1967).
- [Har77] R. HARTSHORNE, *Algebraic Geometry*, *Grad. Texts Math.* **52**, Springer-Verlag, New York, 1977. MR 57 #3116 Zbl 0367.14001
- [JSa] U. JANNSEN and S. SAITO, Bertini theorems and Lefschetz pencils over discrete valuation rings, with applications to higher class field theory, to appear in *J. Algebraic Geom.*
- [JSb] ———, Kato conjecture and motivic cohomology over finite fields, preprint. Available at www.mathematik.uni-regensburg.de/Jannsen
- [JS03] ———, Kato homology of arithmetic schemes and higher class field theory over local fields, *Doc. Math. Extra Vol.* (2003), 479–538, Kazuya Kato’s fiftieth birthday. MR 2005c:11087 Zbl 1092.14504
- [JSS] U. JANNSEN, S. SAITO, and K. SATO, Étale duality for constructible sheaves on arithmetic schemes, preprint. Available at www.mathematik.uni-regensburg.de/Jannsen
- [Kah93] B. KAHN, Résultats de “pureté” pour les variétés lisses sur un corps fini, in *Algebraic K-Theory and Algebraic Topology* (Lake Louise, AB, 1991), *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.* **407**, Kluwer Acad. Publ., Dordrecht, 1993, pp. 57–62, appendix to the article “On the reciprocity sequence in the higher class field theory of function fields” by J.-L. Colliot-Thélène [in *Algebraic K-Theory and Algebraic Topology* (Lake Louise, AB, 1991), 35–55, Kluwer, Dordrecht, 1993; MR1367291 (97k:14019)]. MR 97k:14020
- [Kat86] K. KATO, A Hasse principle for two-dimensional global fields, *J. Reine Angew. Math.* **366** (1986), 142–183. MR 88b:11036 Zbl 0576.12012
- [KS83] K. KATO and S. SAITO, Unramified class field theory of arithmetical surfaces, *Ann. of Math.* **118** (1983), 241–275. MR 86c:14006 Zbl 0562.14011
- [KA79] S. L. KLEIMAN and A. B. ALTMAN, Bertini theorems for hypersurface sections containing a subscheme, *Comm. Algebra* **7** (1979), 775–790. MR 81i:14007 Zbl 0401.14002
- [Kol04] J. KOLLÁR, Specialization of zero cycles, *Publ. Res. Inst. Math. Sci.* **40** (2004), 689–708. MR 2005i:14007 Zbl 1081.14008
- [KS03] J. KOLLÁR and E. SZABÓ, Rationally connected varieties over finite fields, *Duke Math. J.* **120** (2003), 251–267. MR 2005h:14090 Zbl 1077.14068
- [Kun85] E. KUNZ, *Introduction to Commutative Algebra and Algebraic Geometry*, Birkhäuser, Boston, MA, 1985. MR 86e:14001 Zbl 0563.13001
- [Mad08] D. A. MADORE, Équivalence rationnelle sur les hypersurfaces cubiques de mauvaise réduction, *J. Number Theory* **128** (2008), 926–944. MR 2009c:14016 Zbl 1167.14007
- [Mat55] A. MATTUCK, Abelian varieties over p -adic ground fields, *Ann. of Math.* **62** (1955), 92–119. MR 17,87f Zbl 0066.02802
- [MS82] A. S. MERKUR’EV and A. A. SUSLIN, K -cohomology of Severi-Brauer varieties and the norm residue homomorphism, *Dokl. Akad. Nauk SSSR* **264** (1982), 555–559, In Russian; translated in *Math. USSR Izv.* **21** (1983), 307–340. MR 84h:14025
- [Mil80] J. S. MILNE, *Étale Cohomology*, *Princeton Math. Series* **33**, Princeton Univ. Press, Princeton, N.J., 1980. MR 81j:14002 Zbl 0433.14012
- [Mum68] D. MUMFORD, Rational equivalence of 0-cycles on surfaces, *J. Math. Kyoto Univ.* **9** (1968), 195–204. MR 40 #2673 Zbl 0184.46603

- [PS95] R. PARIMALA and V. SURESH, Zero-cycles on quadric fibrations: finiteness theorems and the cycle map, *Invent. Math.* **122** (1995), 83–117. MR 96i:14005 Zbl 0865.14002
- [Poo04] B. POONEN, Bertini theorems over finite fields, *Ann. of Math.* **160** (2004), 1099–1127. MR 2006a:14035 Zbl 1084.14026
- [Poo08] ———, Smooth hypersurface sections containing a given subscheme over a finite field, *Math. Res. Lett.* **15** (2008), 265–271. MR 2009c:14037 Zbl 05310630
- [RZ82] M. RAPOPORT and T. ZINK, Über die lokale Zetafunktion von Shimuravarietäten. Monodromiefiltration und verschwindende Zyklen in ungleicher Charakteristik, *Invent. Math.* **68** (1982), 21–101. MR 84i:14016 Zbl 0498.14010
- [Ras95] W. RASKIND, Higher l -adic Abel-Jacobi mappings and filtrations on Chow groups, *Duke Math. J.* **78** (1995), 33–57. MR 96d:14009 Zbl 0915.14003
- [RS00] W. RASKIND and M. SPIESS, Milnor K -groups and zero-cycles on products of curves over p -adic fields, *Compositio Math.* **121** (2000), 1–33. MR 2002b:14007 Zbl 0985.14003
- [SS95] S. SAITO and R. SUJATHA, A finiteness theorem for cohomology of surfaces over p -adic fields and an application to Witt groups, in *K-Theory and Algebraic Geometry: Connections with Quadratic Forms and Division Algebras* (Santa Barbara, CA, 1992), *Proc. Sympos. Pure Math.* **58**, Amer. Math. Soc., Providence, RI, 1995, pp. 403–415. MR 96d:19009 Zbl 0820.14013
- [Ser92] J.-P. SERRE, *Cohomologie Galoisienne*, 5^e éd., *Lecture Notes in Math.* **5**, Springer-Verlag, New York, 1992. Zbl 0812.12002

(Received May 8, 2006)

(Revised September 28, 2007)

E-mail address: sshuji@msb.biglobe.ne.jp

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO,
8-1 KOMABA 3-CHOME, MEGURO-KU, TOKYO 153-8914, JAPAN

E-mail address: kanetomo@math.nagoya-u.ac.jp

GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY, FUROCHO, CHIKUSA-KU,
NAGOYA 464-8602, JAPAN

E-mail address: uwe.jannsen@mathematik.uni-regensburg.de

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT REGENSBURG, UNIVERSITÄTSSTR. 31,
93040 REGENSBURG, GERMANY

This periodical is published bimonthly by the Department of Mathematics at Princeton University with the cooperation of the Institute for Advanced Study. *Annals* is typeset in \TeX by Sarah R. Warren and produced by Mathematical Sciences Publishers. The six numbers each year are divided into two volumes of three numbers each.

Editorial correspondence

Papers submitted for publication and editorial correspondence should be addressed to Maureen Schupsky, *Annals of Mathematics*, Fine Hall-Washington Road, Princeton University, Princeton, NJ, 08544-1000 U.S.A. The e-mail address is annals@math.princeton.edu.

Preparing and submitting papers

The *Annals* requests that all papers include an abstract of about 150 words which explains to the nonspecialist mathematician what the paper is about. It should not make any reference to the bibliography. Authors are encouraged to initially submit their papers electronically and in PDF format. Please send the file to: annals@math.princeton.edu or to the Mathematics e-print arXiv: front.math.ucdavis.edu/submissions. If a paper is submitted through the arXiv, then please e-mail us with the arXiv number of the paper.

Proofs

A PDF file of the galley proof will be sent to the corresponding author for correction. If requested, a paper copy will also be sent to the author.

Offprints

Authors of single-authored papers will receive 30 offprints. (Authors of papers with one co-author will receive 15 offprints, and authors of papers with two or more co-authors will receive 10 offprints.) Extra offprints may be purchased through the editorial office.

Subscriptions

The price for a print and online subscription, or an online-only subscription, is \$390 per year for institutions. In addition, there is a postage surcharge of \$40 for print subscriptions that are mailed to countries outside of the United States. Individuals interested in subscriptions for their own personal use should contact the publisher at the address below. Subscriptions and changes of address should be sent to Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840 (e-mail: contact@mathscipub.org; phone: 1-510-643-8638; fax: 1-510-295-2608). (Checks should be made payable to "Mathematical Sciences Publishers".)

Back issues and reprints

Orders for missing issues and back issues should be sent to Mathematical Sciences Publishers at the above address. Claims for missing issues must be made within 12 months of the publication date. Online versions of papers published five or more years ago are available through JSTOR (www.jstor.org).

Microfilm

Beginning with Volume 1, microfilm may be purchased from NA Publishing, Inc., 4750 Venture Drive, Suite 400, PO Box 998, Ann Arbor, MI 48106-0998; phone: 1-800-420-6272 or 734-302-6500; email: info@napubco.com, website: www.napubco.com/contact.html.

ALL RIGHTS RESERVED UNDER THE BERNE CONVENTION AND
THE UNIVERSAL COPYRIGHT CONVENTION

TABLE OF CONTENTS

SHIGERU MUKAI. Curves and symmetric spaces, II.....	1539–1558
MANJUL BHARGAVA. The density of discriminants of quintic rings and fields.....	1559–1591
SHUJI SAITO and KANETOMO SATO. A finiteness theorem for zero-cycles over p -adic fields.....	1593–1639
SYLVAIN CROVISIER. Birth of homoclinic intersections: a model for the central dynamics of partially hyperbolic systems.....	1641–1677
JOSEPH BERNSTEIN and ANDRE REZNIKOV. Subconvexity bounds for triple L -functions and representation theory.....	1679–1718
SÁNDOR J KOVÁCS and MAX LIEBLICH. Boundedness of families of canonically polarized manifolds: A higher dimensional analogue of Shafarevich’s conjecture.....	1719–1748
ANDREI TELEMAN. Instantons and curves on class VII surfaces.....	1749–1804
DANIJELA DAMJANOVIĆ and ANATOLE KATOK. Local rigidity of partially hyperbolic actions I. KAM method and \mathbb{Z}^k actions on the torus.....	1805–1858
JORGE LAURET. Einstein solvmanifolds are standard.....	1859–1877
PASCAL COLLIN and HAROLD ROSENBERG. Construction of harmonic diffeomorphisms and minimal graphs.....	1879–1906
JOHN LEWIS and KAJ NYSTRÖM. Boundary behavior and the Martin boundary problem for p harmonic functions in Lipschitz domains.....	1907–1948
JENS MARKLOF and ANDREAS STRÖMBERGSSON. The distribution of free path lengths in the periodic Lorentz gas and related lattice point problems.....	1949–2033
ÉTIENNE FOUVRY and JÜRGEN KLÜNERS. On the negative Pell equation.....	2035–2104
LEX G. OVERSTEEGEN and EDWARD D. TYMCHATYN. Extending isotopies of planar continua.....	2105–2133
JAN BRUINIER and KEN ONO. Heegner divisors, L -functions and harmonic weak Maass forms.....	2135–2181
FLORIAN POP. Henselian implies large.....	2183–2195
MIKHAIL BELOLIPETSKY, TSACHIK GELANDER, ALEXANDER LUBOTZKY and ANER SHALEV. Counting arithmetic lattices and surfaces.....	2197–2221
SASHA SODIN. The spectral edge of some random band matrices.....	2223–2251