

Energy quantization for Willmore surfaces and applications

By YANN BERNARD and TRISTAN RIVIÈRE

Abstract

We prove a bubble-neck decomposition together with an energy quantization result for sequences of Willmore surfaces into \mathbb{R}^m with uniformly bounded energy and nondegenerating conformal type. We deduce the strong compactness of Willmore closed surfaces of a given genus modulo the Möbius group action, below some energy threshold.

I. Introduction

Let $\vec{\Phi}$ be an immersion from a closed abstract two-dimensional manifold Σ into $\mathbb{R}^{m \geq 3}$. We denote by $g := \vec{\Phi}^* g_{\mathbb{R}^m}$ the pull back by $\vec{\Phi}$ of the flat canonical metric $g_{\mathbb{R}^m}$ of \mathbb{R}^m , also called the *first fundamental form of $\vec{\Phi}$* , and we let $d\text{vol}_g$ be its associated *volume form*. The *Gauss map* of the immersion $\vec{\Phi}$ is the map taking values in the Grassmannian of oriented $m - 2$ -planes in \mathbb{R}^m given by

$$\vec{n}_{\vec{\Phi}} := \star \frac{\partial_{x_1} \vec{\Phi} \wedge \partial_{x_2} \vec{\Phi}}{|\partial_{x_1} \vec{\Phi} \wedge \partial_{x_2} \vec{\Phi}|},$$

where \star is the usual Hodge star operator in the Euclidean metric. Denoting by $\pi_{\vec{n}_{\vec{\Phi}}}$ the orthonormal projection of vectors in \mathbb{R}^m onto the $m - 2$ -plane given by $\vec{n}_{\vec{\Phi}}$, the *second fundamental form* may be expressed as¹

$$\forall X, Y \in T_p \Sigma \quad \vec{\mathbb{I}}_p(X, Y) := \pi_{\vec{n}_{\vec{\Phi}}} d^2 \vec{\Phi}(X, Y).$$

The *mean curvature vector* of the immersion at p is

$$\vec{H}_{\vec{\Phi}} := \frac{1}{2} \text{tr}_g(\vec{\mathbb{I}}) = \frac{1}{2} \left[\vec{\mathbb{I}}(\varepsilon_1, \varepsilon_1) + \vec{\mathbb{I}}(\varepsilon_2, \varepsilon_2) \right],$$

where $(\varepsilon_1, \varepsilon_2)$ is an orthonormal basis of $T_p \Sigma$ for the metric $g_{\vec{\Phi}}$.

Y. B. was supported by the DFG SFB 71 fund (project B3).

© 2014 Department of Mathematics, Princeton University.

¹In order to define $d^2 \vec{\Phi}(X, Y)$ one has to extend locally around $T_p \Sigma$ the vector-fields X and Y . It is not difficult to check that $\pi_{\vec{n}_{\vec{\Phi}}} d^2 \vec{\Phi}(X, Y)$ is independent of this extension.

In the present paper, we study the Lagrangian given by the L^2 -norm of the second fundamental form:

$$E(\vec{\Phi}) := \int_{\Sigma} |\vec{\mathbb{I}}|_{g_{\vec{\Phi}}}^2 \, d\text{vol}_{g_{\vec{\Phi}}}.$$

An elementary computation gives

$$E(\vec{\Phi}) := \int_{\Sigma} |\vec{\mathbb{I}}|_{g_{\vec{\Phi}}}^2 \, d\text{vol}_{g_{\vec{\Phi}}} = \int_{\Sigma} |d\vec{n}_{\vec{\Phi}}|_{g_{\vec{\Phi}}}^2 \, d\text{vol}_{g_{\vec{\Phi}}}.$$

The energy E may accordingly be seen as the *Dirichlet Energy* of the Gauss map $\vec{n}_{\vec{\Phi}}$ with respect to the induced metric $g_{\vec{\Phi}}$. The Gauss-Bonnet theorem implies that

$$\begin{aligned} \text{(I.1)} \quad E(\vec{\Phi}) &:= \int_{\Sigma} |\vec{\mathbb{I}}|_{g_{\vec{\Phi}}}^2 \, d\text{vol}_{g_{\vec{\Phi}}} = 4 \int_{\Sigma} |\vec{H}_{\vec{\Phi}}|^2 \, d\text{vol}_{g_{\vec{\Phi}}} - 2 \int_{\Sigma} K_{\vec{\Phi}} \, d\text{vol}_{g_{\vec{\Phi}}} \\ &= 4 \int_{\Sigma} |\vec{H}_{\vec{\Phi}}|^2 \, d\text{vol}_g - 4\pi\chi(\Sigma), \end{aligned}$$

where $K_{\vec{\Phi}}$ is the Gauss curvature of the immersion and $\chi(\Sigma)$ is the *Euler characteristic* of the surface Σ . The energy

$$W(\vec{\Phi}) := \int_{\Sigma} |\vec{H}_{\vec{\Phi}}|^2 \, d\text{vol}_{g_{\vec{\Phi}}},$$

is called *Willmore energy*.

Critical points of the *Willmore energy*, comprising for example minimal surfaces,² are called *Willmore surfaces*. Although already known in the 19th century in the context of the elasticity theory of plates, it was first considered in conformal geometry by Blaschke in [Bla29], who sought to merge the *theory of minimal surfaces* and the *conformal invariance property*. This Lagrangian has indeed both desired features: its critical points contain minimal surfaces, and it is conformal invariant, owing to the following pointwise identity which holds for an arbitrary immersion $\vec{\Phi}$ of Σ into \mathbb{R}^m and at every point of Σ :

$$\begin{aligned} \forall \Xi \quad \text{conformal diffeo. of } \mathbb{R}^m \cup \{\infty\} \quad & [|\vec{H}_{\Xi \circ \vec{\Phi}}|^2 - K_{\Xi \circ \vec{\Phi}}] \, d\text{vol}_{g_{\Xi \circ \vec{\Phi}}} \\ &= [|\vec{H}_{\vec{\Phi}}|^2 - K_{\vec{\Phi}}] \, d\text{vol}_{g_{\vec{\Phi}}}. \end{aligned}$$

Using again the Gauss-Bonnet theorem, the latter implies the conformal invariance of W :

$$\forall \Xi \quad \text{conformal diffeo. from } \mathbb{R}^m \cup \{\infty\} \quad W(\Xi \circ \vec{\Phi}) = W(\vec{\Phi}).$$

This conformal invariance implies that the image of a *Willmore immersion* by a conformal transformation of \mathbb{R}^m is still a Willmore immersion. Starting for example from a minimal surface, one may then generate many new Willmore surfaces, simply by applying conformal transformations. (Naturally,

²Minimal surfaces satisfy $\vec{H} = 0$ and are hence absolute minimizers of W .

these surfaces need no longer be minimal.) In his time, Blaschke used the term *conformal minimal* for the critical points of W , seeking to insist on this idea of merging together the theory of minimal surface with conformal invariance.

An important task in the analysis of Willmore surfaces is to understand the closure of the space of *Willmore immersions* under a certain level of energy. Because of the noncompactness of the conformal group of transformation of \mathbb{R}^m , one cannot expect that the space of Willmore immersions is closed in the strong C^l -topology. However, locally, in isothermic coordinates,³ under some universal energy threshold, if the conformal factor of the induced metric $g_{\vec{\Phi}}$ is controlled in L^∞ , then the immersion is uniformly bounded in any C^l -norm. More precisely, there holds the following ε -regularity result.

THEOREM I.1 ([Riv08], [Riv13b]). *There exists $\varepsilon(m) > 0$ such that, for any Willmore conformal immersion $\vec{\Phi}$ from $B_1(0)$ into \mathbb{R}^m satisfying*

$$\int_{B_1(0)} |\nabla \vec{n}_{\vec{\Phi}}|^2 dx < \varepsilon(m),$$

then for any $l \in \mathbb{N}^$, we have*

$$\|e^{-\lambda} \nabla^l \vec{\Phi}\|_{L^\infty(B_{1/2})} \leq C_l(\|\nabla \lambda\|_1) \left[\int_{B_1(0)} |\nabla \vec{n}_{\vec{\Phi}}|^2 dx + 1 \right]^{1/2},$$

where C_l only depends on l and the L^1 -norm of $\nabla \lambda$, while λ denotes the conformal parameter of $\vec{\Phi}$. Namely, $\lambda = \|\log |\partial_{x_1} \vec{\Phi}|\|_{L^\infty(B_1)} = \|\log |\partial_{x_2} \vec{\Phi}|\|_{L^\infty(B_1)}$.

This theorem leads to the *concentration of compactness* “dialectic” developed by Sacks and Uhlenbeck in the context of harmonic maps from the two sphere into manifolds. In a conformal parametrization, assuming that the conformal factor is L^∞ -controlled in some subdomain of Σ , then a sequence of Willmore immersions can fail to converge strongly in C^l only at isolated points; namely, at those points where the $W^{1,2}$ -norm of the Gauss map concentrates. Assuming their induced metric generates a sequence of conformal classes that remains within a compact subdomain of the moduli space of Σ , the control of the conformal factor of a sequence of conformal immersions with uniformly bounded Willmore energy is also guaranteed, except at those isolated points of Σ where the $W^{1,2}$ -norm of the Gauss map concentrates. This fact is established in [Riv13c] (see also Proposition III.1 below), and it ultimately follows from the works of Toro [Tor95], of Müller and Šverák [MŠ95], and from the work of Hélein [Hél02] on immersions with totally bounded curvature.

³Analogously to other gauge-invariant problems, such as in Yang-Mills theory, isothermic coordinates or conformal parametrizations provide the optimal symmetry breaking method in the search of pertinent estimates. A detailed discussion on this topic is available in [Riv13b].

To fully understand the loss of strong compactness of a sequence of Willmore surfaces, the difficulty is of course to perform the necessary analysis at the “blow up points.” This is the aim of the present paper. Under the assumption that the sequence of conformal classes associated to the sequence of Willmore surfaces remains within a compact subdomain of the Moduli space, we show that, modulo extraction of a subsequence, the parametrizing abstract surface splits into three distinct regions: the main region where strong convergence holds, the concentrating parametrization of nontrivial Willmore spheres, and finally the bubble and neck regions connecting the two previous ones and in which we show the energy vanishes.

We now state our main result.

THEOREM I.2. *Let $\vec{\Phi}_k$ be a sequence of Willmore immersions of a closed surface Σ . Assume that*

$$\limsup_{k \rightarrow +\infty} W(\vec{\Phi}_k) < +\infty$$

and that the conformal class of $\vec{\Phi}_k^ g_{\mathbb{R}^m}$ remains within a compact subdomain of the moduli space of Σ . Then, modulo extraction of a subsequence, the following energy identity holds:*

$$(I.2) \quad \lim_{k \rightarrow +\infty} W(\vec{\Phi}_k) = W(\vec{\xi}_\infty) + \sum_{s=1}^p W(\vec{\eta}_s) + \sum_{t=1}^q [W(\vec{\zeta}_t) - 4\pi \theta_t],$$

where $\vec{\xi}_\infty$ is a possibly branched smooth immersion of Σ . The maps $\vec{\eta}_s$ and $\vec{\zeta}_t$ are smooth, possibly branched, immersions of S^2 , and θ_t is the integer density of the current $(\vec{\zeta}_t)_[S^2]$ at some point $p_t \in \vec{\zeta}_t(S^2)$; namely,*

$$\theta_t := \lim_{r \rightarrow 0} \frac{\mathcal{H}^2(B_r^m(p_t) \cap \vec{\zeta}_t(S^2))}{\pi r^2}.$$

Remark I.1. Due to the fact that the first residues are zeros at the blow up points are zeros (see the terminology in [BR13] and precisely identity (VIII.4) in Section VIII of the paper), the map $\vec{\xi}_\infty$ extends to a $C^{2,\alpha}$ map for any $\alpha < 1$ at the branched points

The second part of our main result describes how the maps $\vec{\xi}_\infty$, $\vec{\eta}^s$, and $\vec{\zeta}^t$ are obtained from the original sequence $\vec{\Phi}_k$.

THEOREM I.3. *With the same notation as in Theorem I.2, the immersion $\vec{\xi}_\infty$ is obtained as follows. There exist a sequence f_k of diffeomorphisms of Σ , a sequence $\vec{\Xi}_k$ of Möbius transformations of \mathbb{R}^m , and finitely many points $\{a^1, \dots, a^n\}$ such that $\vec{\Xi}_k \circ \vec{\xi}_k \circ f_k$ is conformal and*

$$(I.3) \quad \vec{\xi}_k := \vec{\Xi}_k \circ \vec{\xi}_k \circ f_k \longrightarrow \vec{\xi}_\infty \quad \text{in } C_{\text{loc}}^l(\Sigma \setminus \{a^1, \dots, a^n\}) \quad \forall l \in \mathbb{N}.$$

Furthermore, there holds

$$(I.4) \quad \lim_{k \rightarrow +\infty} W(\vec{\Phi}_k) = W(\vec{\xi}_\infty) \quad \iff \quad \vec{\xi}_k \longrightarrow \vec{\xi}_\infty \quad \text{in } C^l(\Sigma) \quad \forall l \in \mathbb{N}.$$

Finally there exists a sequence h_k of constant scalar curvature metric conformally equivalent to $\vec{\xi}_k = \Xi_k \circ \vec{\xi}_k \circ f_k$ and strongly converging in $C^l(\Sigma)$ such that for any $s \in \{1, \dots, p\}$ (resp. for any $t \in \{1, \dots, q\}$), there exist a sequence of points $x_k^s \in \Sigma$ converging to one of the a^i , a sequence of radius ρ_k^s converging to zero, and a sequence of Möbius transformations Ξ_k^s (resp. Ξ_k^t) for which (in converging h_k conformal coordinates φ_k around a^i), one has

$$\Xi_k^s \circ \vec{\xi}_k \circ \varphi_k(\rho_k^s y + \phi_k^{-1}(x_k^s)) \longrightarrow \vec{\eta}_s \circ \pi^{-1}(y) \quad \text{in } C_{\text{loc}}^l(\mathbb{C} \setminus \{a_s^1, \dots, a_s^{n_s}\})$$

and respectively

$$\Xi_k^t \circ \vec{\xi}_k \circ \varphi_k(\rho_k^t y + \phi_k^{-1}(x_k^t)) \longrightarrow \vec{\zeta}_t \circ \pi^{-1}(y) \quad \text{in } C_{\text{loc}}^l(\mathbb{C} \setminus \{a_t^1, \dots, a_t^{n_t}\})$$

for any $l \in \mathbb{N}$, where π denotes the stereographic projection from S^2 into \mathbb{C} and $\{a_s^1, \dots, a_s^{n_s}\}$ and $\{a_t^1, \dots, a_t^{n_t}\}$ are finite sets of points in the complex plane.

The maps Ξ_k^s are compositions of a dilation and isometries, while each map Ξ_k^t is a composition of one inversion, one dilation, and one isometry.

The results given in Theorems I.2 and I.3 are to be viewed in the context of other bubble-neck decomposition and energy quantization situations previously studied; in particular, for harmonic maps and other conformally invariant problems as those considered in [SU81], [Jos91], [Str85], [DT95], [Par96], [Riv02b], and in [Riv02a]. However, these problems are all of second-order elliptic or parabolic types. The novelty of the present work is an energy quantization result and bubble-neck decomposition for a fourth-order problem.⁴ Moreover, the analysis that we present here requires us to handle one additional delicate difficulty; namely, the need to first control the conformal parameter in “conformally degenerating” neck regions (annuli), prior to obtaining from PDE techniques the vanishing of the energy in these regions.

The method we develop to prove Theorems I.2 and I.3 takes partly its sources in techniques introduced by the second author in collaboration with F. H. Lin, in the context of the Ginzburg-Landau equations [LR01] and in the context of harmonic maps [LR02]. This technique relies on the properties of the Lorentz interpolation spaces. More precisely, in the present paper, we obtain estimates of the trace of the second fundamental form in the neck regions,

⁴Indeed, in a conformal parametrization $\vec{\xi}$, the Willmore functional may be recast as

$$W(\vec{\xi}) = \frac{1}{4} \int_{\Sigma} |\Delta_g \vec{\xi}|^2 \, d\text{vol}_g,$$

thereby giving rise to a fourth-order problem.

respectively in the weak Marcinkiewicz space $L^{2,\infty}$, and in its predual, the Lorentz space $L^{2,1}$. These estimates constitute the main achievement of our proof. Prior to deriving these estimates, it is first necessary to carefully study the conformal factor of the conformal Willmore immersions in the neck region. This is done through extracting a good Coulomb gauge. The existence of such a gauge is one of the novelties of the present work, for it requires only that the gradient of the Gauss map lie in L^2 and that it have a small $L^{2,\infty}$ -norm. Such a control on the $L^{2,\infty}$ -norm of the second fundamental form (i.e., of the gradient of the Gauss map) ultimately stems from the ε -regularity established in [Riv08]. The appropriate control in $L^{2,1}$ is then deduced using the standard methods of integrability by compensation and applying them to the conservation laws satisfied by Willmore immersions and originally derived in [Riv08].

It is legitimate to ask what happens if one removes the assumption on the control of the conformal structure. In this situation, the first important observation is provided by the examples constructed in [BP06], where inverted catenoids arise as limiting bubbles, which are thus not smooth Willmore immersion of S^2 . The second important observation is that harmonic mapping from a degenerating Riemann surface into a manifold are known in general not to satisfy the energy quantization property in the thin collar regions; cf. example constructed in [Par96] and the careful systematic study given in [Zhu10].

Several authors have recently undertaken the task of understanding the degeneracy of sequences in Moduli space. In [Sch13], it is shown that for a sequence $\Sigma_k \subset \mathbb{R}^{m \geq 3}$ of closed, orientable, embedded surfaces of fixed genus $g \geq 1$ with $\limsup_{k \rightarrow \infty} W(\Sigma_k) < 8\pi$, the convergence in Moduli space of the associated sequence of conformal classes $[\Sigma_k]$ is equivalent to a “no loss of topology” condition: any subsequence has a subsequence satisfying, after applying Möbius transformations, $\mathcal{H}^2 \llcorner \Sigma_k \rightarrow \mu$ as Radon measures, where $\text{spt}(\mu)$ is a closed, orientable, embedded topological surface with same genus g . Furthermore, the associated conformal class $[\Sigma_k]$ converges to $[\mu]$ in Moduli space. Alternatively, a sufficient energy condition ensuring the nondegeneracy of the conformal class has been given (in the cases $m = 3$ and $m = 4$) in [KS12], and for surfaces in any codimension, in [Riv13a] (and independently in [KL12]). Following the convention in [Sim93], we introduce

$$\beta_g^m := \inf \left\{ W(\vec{\Phi}) ; \vec{\Phi} \text{ is an immersion of the genus } g \text{ closed surface} \right\}.$$

and

$$\omega_g^m := \min \left\{ 4\pi + \sum_{i=1}^p (\beta_{g_i}^m - 4\pi) ; g = g_1 + \cdots + g_p, \quad 1 \leq g_i < g \right\}.$$

In [BK03], it is proved that for $g \geq 2$, there holds

$$(I.5) \quad \beta_g^m < \omega_g^m.$$

THEOREM I.4 ([Riv13a] [KL12]). *Let (Σ, c_k) be a sequence of closed Riemann surfaces of genus g , with degenerating conformal class $[c_k]$ diverging to the boundary of the Moduli space of Σ . Let $\vec{\Phi}_k$ be a sequence of conformal immersions from (Σ, c_k) into \mathbb{R}^m . Then*

$$(I.6) \quad \liminf_{k \rightarrow +\infty} \int_{\Sigma} |\vec{H}_{\vec{\Phi}_k}|^2 \, \text{dvol}_{\vec{\Phi}_k^* g_{\mathbb{R}^m}} \geq \min\{8\pi, \omega_g^m\}.$$

A consequence of our analysis below is the following result.

THEOREM I.5. *Let Σ be an arbitrary closed two-dimensional manifold. Modulo the action of the Möbius group of \mathbb{R}^m , the space of Willmore immersions into \mathbb{R}^m for $m = 3$ and $m = 4$, satisfying*

$$W(\Phi) < \min\{8\pi, \omega_g^m\} - \delta$$

is strongly compact in the C^l topology for any $l \in \mathbb{N}$ and any $\delta > 0$.

This result was obtained for $\Sigma = T^2$ and $m = 3$ in [KS04] and for $\Sigma = T^2$ and $m = 4$ in [Riv08]. They were based on the energy quantization for Willmore spheres in codimension 1 and 2 respectively due to R. Bryant [Bry84] and S. Montiel [Mon00]. Thanks to the recent proof of The Willmore conjecture by F. C. Marques and A. Neves, [MN14], one knows that $\omega_g^3 > 8\pi$; thus for $m = 3$, the assumption in Theorem I.5 is then reduced to

$$W(\Phi) < 8\pi - \delta.$$

The paper is organized as follows. In Section II, we recall the formalism introduced in [Riv13c] that enables us to appropriately “renormalize” a sequence of weak immersions uniformly bounded Willmore energy. In Section III, we outline a generic “energy tracking” procedure enabling us to detect bubbles and neck regions. Section IV, also general in nature, explains how to construct a Coulomb moving frame under the hypothesis that the Gauss map has small weak- L^2 -norm. In Section V, we develop uniform Harnack estimates giving control of the conformal factor of weak immersions in the neck regions. Willmore immersions first appear in Section VI, where we show how to control the $L^{2,1}$ -norm of the mean curvature vector in the neck regions. In Section VII, we derive an energy quantization result for a sequence of Gauss maps corresponding to a sequence of Willmore immersions. Finally, in Section VIII, all of the aforementioned estimates are seamed together, and the main Theorems I.2 and I.3 are proved.

Acknowledgements. This work was partly elaborated while the first author was visiting the *Forschungsinstitut für Mathematik* at the *ETH Zürich*. He would like to thank the institute for its hospitality and the excellent working conditions. The authors are also grateful to the referees for useful comments.

II. Preliminaries

In [Riv13c], the second author developed a suitable framework in which one can perform the calculus of variation of the Willmore Lagrangian of a surface. In this framework, a particularly useful space is that of Lipschitz immersions with L^2 -bounded second fundamental forms.

Let g_0 be a smooth “reference” metric on Σ . One defines the Sobolev spaces $W^{k,p}(\Sigma, \mathbb{R}^m)$ of measurable maps from Σ into \mathbb{R}^m as

$$W^{k,p}(\Sigma, \mathbb{R}^m) := \left\{ f \text{ meas. } \Sigma \rightarrow \mathbb{R}^m \text{ s.t. } \sum_{l=0}^k \int_{\Sigma} |\nabla^l f|_{g_0}^p \, d\text{vol}_{g_0} < +\infty \right\}.$$

As Σ is assumed to be compact, this space is independent of the chosen reference metric g_0 .

It is important to have a weak first fundamental form: we need $\vec{\Phi}^* g_{\mathbb{R}^m}$ to define an L^∞ metric with a bounded inverse. This is the case if we assume that $\vec{\Phi}$ lies in $W^{1,\infty}(\Sigma)$ and if $d\vec{\Phi}$ has maximal rank 2 at every point, with some uniform quantitative control of “how far” $d\vec{\Phi}$ is from being degenerate. Namely, there exists $c_0 > 0$ such that

$$(II.1) \quad |d\vec{\Phi} \wedge d\vec{\Phi}|_{g_0} \geq c_0 > 0,$$

where $d\vec{\Phi} \wedge d\vec{\Phi}$ is a 2-form on Σ with values in the space of 2-vectors of \mathbb{R}^m , given in local coordinates by $2 \partial_x \vec{\Phi} \wedge \partial_y \vec{\Phi} \, dx \wedge dy$. Note that the condition (II.1) is independent of the choice of the metric g_0 . To a Lipschitz immersion satisfying (II.1) we associate its Gauss map

$$\vec{n}_{\vec{\Phi}} := \star \frac{\partial_{x_1} \vec{\Phi} \wedge \partial_{x_2} \vec{\Phi}}{|\partial_{x_1} \vec{\Phi} \wedge \partial_{x_2} \vec{\Phi}|}.$$

It belongs to $L^\infty(\Sigma)$ and takes values in the Grassmannian of oriented $(m-2)$ -planes in \mathbb{R}^m .

We next introduce the space \mathcal{E}_Σ of Lipschitz immersions⁵ of Σ with square-integrable second fundamental form:

$$\mathcal{E}_\Sigma := \left\{ \begin{array}{l} \vec{\Phi} \in W^{1,\infty}(\Sigma, \mathbb{R}^m) \quad \text{s.t. } \vec{\Phi} \text{ satisfies (II.1) for some } c_0 \\ \text{and} \quad \int_{\Sigma} |d\vec{n}|_g^2 \, d\text{vol}_g < +\infty \end{array} \right\}.$$

As before, $g := \vec{\Phi}^* g_{\mathbb{R}^m}$ is the pullback by $\vec{\Phi}$ of the flat canonical metric $g_{\mathbb{R}^m}$ of \mathbb{R}^m , and $d\text{vol}_g$ is its corresponding volume form.

As in the case of the Yang-Mills functional, one of the main difficulties one encounters in the variational study of the Willmore functional is the size of the corresponding gauge group (i.e., the group of transformations through which

⁵Henceforth called *weak immersions*.

the energy is invariant). For the energy W , this gauge group is very large: it contains all diffeomorphisms of Σ . To “break” the gauge invariance, one of the first techniques considered in [Riv13c] consists in working with conformal immersions. Indeed, owing to the work of Müller and Šverák [MŠ95] and to the work of Hélein [Hél02], one gains an L^∞ -control of the conformal parameter under a small energy E assumption. It is then possible to show that the pullback metric of any element in \mathcal{E}_Σ defines a smooth conformal structure on Σ (cf. details in [Riv13b]). Thus, corresponding to any element $\vec{\Phi} \in \mathcal{E}_\Sigma$, there exists a Lipschitz diffeomorphism f of Σ such that $\vec{\Phi} \circ f$ is conformal. Having found a conformal structure c of Σ , one defines

$$\mathcal{E}_\Sigma^c := \left\{ \vec{\Phi} \in \mathcal{E}_\Sigma \quad \text{s.t. } \vec{\Phi} \text{ is conformal w.r.t. } c \right\}.$$

The existence of a minimizer in the class \mathcal{E}_Σ^c is proved in [Riv13c] (arbitrary codimension) and in [KS13] (codimensions 1 and 2). Except if the minimizer is globally isothermic, its smoothness is established in [Riv13c] away from finitely many branch points, and the minimizer satisfies the *conformal-constrained Willmore equation* (cf. [BR11]). Alternatively, under certain energy assumptions, or assuming there are no branch points, it is shown in [KS13] that the minimizer is smooth.

Naturally, in considering a sequence of elements in \mathcal{E}_Σ , the corresponding sequence of conformal classes it yields is not *a priori* controlled, and it might diverge in the moduli space of Σ , even when the Willmore energy of the immersions is uniformly bounded. This possible degeneracy of the conformal class is another source of trouble when trying to control a sequence of immersions with uniformly bounded Willmore energy. In the present work, we will assume that the aforementioned sequence of conformal classes does **not** degenerate. Sufficient energy conditions to avoid degeneracy of the conformal structure were given in [Riv13a] and independently in [KL12], while necessary and sufficient conditions were given in [Sch13] in topological terms.

Once the global gauge choice is made, the second difficulty one encounters is the invariance of the Willmore functional under conformal transformations of \mathbb{R}^m . Namely, any conformal diffeomorphism Ξ of \mathbb{R}^m (i.e., an element of the Möbius group of \mathbb{R}^m) leaves both \mathcal{E}_Σ^c and W (through composition). Since the Möbius group is noncompact, this invariance generates serious troubles in the study of a sequence of immersions with uniformly bounded Willmore energy. Indeed, such a sequence may *a priori* collapse to a point in the limit! To overcome this difficulty, various methods have been devised,⁶ such as Lemma 4.1 in [KS12], Lemma III.1 in [Riv13c], and Proposition 2.2 in [Sch13]. In the

⁶Analogously, in the “ambient approach” originally introduced by Leon Simon [Sim93], a renormalization of the image is performed.

present work, we extract a suitable gauge Ξ with the help of the “normalization lemma” A.4 of [Riv13c]. Doing so yields an “almost” weak closure result for sequences in \mathcal{E}_Σ^c modulo composition with suitable elements of the Möbius group. More precisely, the limit immersion is not necessarily a weak immersion of the whole Σ , but only a weak immersion away from possibly finitely many points. We are thus led to consider the space of weak branched immersions:

$$\mathcal{F}_\Sigma := \left\{ \begin{array}{l} \vec{\Phi} \in W^{1,\infty}(\Sigma, \mathbb{R}^m) \quad \text{s.t. } \exists \{a_1, \dots, a_N\} \in \Sigma \quad \text{with} \\ \quad \forall K \text{ compact subset of } \Sigma \setminus \{a_1, \dots, a_N\} \\ \exists c_K > 0 \quad \text{s.t. } \forall x \in K \quad |d\vec{\Phi} \wedge d\vec{\Phi}|_{g_0}(x) \geq c_K \\ \text{and} \quad \int_\Sigma |d\vec{n}|_g^2 \, d\text{vol}_g < +\infty \end{array} \right\}.$$

Finally, we introduce the space of elements in \mathcal{F}_Σ that are conformal with respect to some fixed conformal structure c :

$$\mathcal{F}_\Sigma^c := \left\{ \vec{\Phi} \in \mathcal{F}_\Sigma \quad \text{s.t.} \quad \vec{\Phi} \text{ is conformal w.r.t. } c \right\}.$$

The above discussion is summarized in the following result, which will be a central ingredient in the rest of this article.

LEMMA II.1 ([Riv13c]). *Let Σ be a closed two-dimensional manifold. Let $\vec{\Phi}_k$ be a sequence of elements in \mathcal{E}_Σ such that $W(\vec{\Phi}_k)$ is uniformly bounded. Assume that the conformal class of the conformal structure c_k (i.e., complex structure of Σ) defined by $\vec{\Phi}_k$ remains in a compact subspace of the Moduli space of Σ . Then, modulo extraction of a subsequence, the sequence c_k converges to a smooth limiting complex structure c_∞ , and there exist a sequence of Lipschitz diffeomorphisms f_k of Σ and a sequence of smooth conformal structures c_k of Σ such that $\vec{\Phi}_k \circ f_k$ is conformal from (Σ, c_k) into \mathbb{R}^m . Moreover, there exists a sequence Ξ_k of conformal diffeomorphisms of $\mathbb{R}^m \cup \{\infty\}$ and at most finitely many points $\{a^1, \dots, a^n\}$ such that*

$$(II.2) \quad \limsup_{k \rightarrow +\infty} \mathcal{H}(\Xi_k \circ \vec{\Phi}_k \circ f_k(\Sigma)) < +\infty, \quad \Xi_k \circ \vec{\Phi}_k \circ f_k(\Sigma) \subset B_R(0)$$

for some $R > 0$ independent of k , and

$$(II.3) \quad \vec{\xi}_k := \Xi_k \circ \vec{\Phi}_k \circ f_k \rightharpoonup \vec{\xi}_\infty \quad \text{weakly in } (W_{\text{loc}}^{2,2} \cap W_{\text{loc}}^{1,\infty})^*(\Sigma \setminus \{a^1, \dots, a^n\}).$$

The convergences are understood with respect to h_k , which is the constant scalar curvature metric of unit volume attached to the conformal structure c_k . Furthermore, there holds

$$(II.4) \quad \forall K \text{ compact subset of } \Sigma \setminus \{a^1, \dots, a^n\} \quad \limsup_{k \rightarrow +\infty} \|\log |d\vec{\xi}_k|_{h_k}\|_{L^\infty(K)} < +\infty.$$

Finally, $\vec{\xi}_\infty$ is a weak immersion of $\Sigma \setminus \{a^1, \dots, a^n\}$ and conformal from (Σ, c_∞) into \mathbb{R}^m .

III. The bubble-neck decomposition procedure

The goal of this section is to establish the following technical proposition. Although its statement might seem at first somewhat overwhelming, its proof involves no particularly challenging difficulties and is based on an iteration argument.

PROPOSITION III.1 (Bubble-neck decomposition). *Let Σ be a closed two-dimensional manifold. Let $\vec{\xi}_k$ be a sequence of weak Lipschitz immersions with L^2 -bounded second fundamental forms (i.e., $\vec{\xi}_k \in \mathcal{E}_\Sigma$) such that*

$$\int_{\Sigma} |d\vec{n}_{\vec{\xi}_k}|_{g_k}^2 \, d\text{vol}_{g_k} \leq \Lambda,$$

where $g_k := \vec{\xi}_k^* g_{\mathbb{R}^m}$.

We suppose that g_k is conformally equivalent to a constant scalar curvature metric h_k such that for all $l \in \mathbb{N}$,

$$h_k \longrightarrow h_\infty \quad \text{in } C^l(\Sigma),$$

where h_∞ is a constant scalar curvature on Σ .

Suppose there exist n points $\{a^1, \dots, a^n\} \subset \Sigma$ and a radius $R > 0$ such that the following holds:

(i)

$$(III.5) \quad \limsup_{k \rightarrow +\infty} \mathcal{H}(\vec{\xi}_k(\Sigma)) < +\infty,$$

(ii)

$$(III.6) \quad \vec{\xi}_k(\Sigma) \subset B_R(0),$$

(iii)

$$(III.7) \quad \vec{\xi}_k \rightharpoonup \vec{\xi}_\infty \quad \text{weakly in } (W_{\text{loc}}^{2,2} \cap W_{\text{loc}}^{1,\infty})^*(\Sigma \setminus \{a^1, \dots, a^n\}),$$

(iv)

$$(III.8) \quad \forall K \text{ compact subset of } \Sigma \setminus \{a^1, \dots, a^n\} \quad \limsup_{k \rightarrow +\infty} \|\log |d\vec{\xi}_k|_{h_k}\|_{L^\infty(K)} < +\infty.$$

Then for any $0 < \varepsilon_0 < 8\pi/3$, there exist a subsequence, still denoted $\vec{\xi}_k$, n integers $\{Q^1, \dots, Q^n\}$, n sequences of points $(x_k^{i,j})_{j=1, \dots, Q^i} \subset \Sigma$, and n sequences of radii $(\rho_k^{i,j})_{j=1, \dots, Q^i}$ satisfying

$$(III.9) \quad \forall i \in 1, \dots, n \quad \forall j \in \{1, \dots, Q^i\} \quad \lim_{k \rightarrow +\infty} x_k^{i,j} = a^i,$$

$$(III.10) \quad \forall i \in 1, \dots, n \quad \forall j \in \{1, \dots, Q^i\} \quad \lim_{k \rightarrow +\infty} \rho_k^{i,j} = 0,$$

and

(III.11)

$$\forall i \in \{1, \dots, n\} \quad \forall j \neq j' \in \{1, \dots, Q^i\} \quad \left\{ \begin{array}{l} \text{either} \quad \lim_{k \rightarrow +\infty} \frac{\rho_k^{i,j}}{\rho_k^{i,j'}} + \frac{\rho_k^{i,j'}}{\rho_k^{i,j}} = +\infty \\ \text{or} \quad \lim_{k \rightarrow +\infty} \frac{|x_k^{i,j} - x_k^{i,j'}|}{\rho_k^{i,j} + \rho_k^{i,j'}} = +\infty. \end{array} \right.$$

Moreover,

$$(III.12) \quad \forall i \in \{1, \dots, n\} \quad \forall j \in \{1, \dots, Q^i\} \quad \int_{B_{\rho_k^{i,j}}(x_k^{i,j})} |d\vec{n}_{\vec{\xi}_k}|_{g_k}^2 \, d\text{vol}_{g_k} > \varepsilon_0.$$

The set of balls $B_{\rho_k^{i,j}}$ are called “bubbles” associated to the sequence $\vec{\xi}_k$.

We have furthermore that for any $i \in \{1, \dots, n\}$ and for any $j \in \{1, \dots, Q^i\}$, the set of indices⁷

$$I^{i,j} := \left\{ j' \quad ; \quad x_k^{i,j'} \in B_{\rho_k^{i,j}}(x_k^{i,j}) \quad ; \quad \frac{\rho_k^{i,j}}{\rho_k^{i,j'}} \rightarrow +\infty \right\}$$

is independent of k . It is called the set of “bubbles contained in the bubble $B_{\rho_k^{i,j}}(x_k^{i,j})$ ”.

For any $\alpha < 1$, we denote

$$\forall i \in \{1, \dots, n\} \quad \Omega_k^i(\alpha) := B_\alpha(a^i) \setminus \bigcup_{j=1}^{Q^i} B_{\alpha^{-1}\rho_k^{i,j}}(x_k^{i,j})$$

and

$$\forall i \in \{1, \dots, n\} \quad \forall j \in \{1, \dots, Q^i\} \quad \Omega_k^{i,j}(\alpha) := \bigcup_{j' \in I^{i,j}} \left(B_{\alpha\rho_k^{i,j'}}(x_k^{i,j'}) \setminus \bigcup_{j'' \in I^{i,j}} B_{\alpha^{-1}\rho_k^{i,j''}}(x_k^{i,j''}) \right).$$

The sets $\Omega_k^i(\alpha)$ and the $\Omega_k^{i,j}(\alpha)$ are the “ α -neck regions” of $\vec{\xi}_k$. Let

$$\Omega_k(\alpha) := \bigcup_{i=1}^n \left(\Omega_k^i(\alpha) \cup \bigcup_{j=1}^{Q^i} \Omega_k^{i,j}(\alpha) \right).$$

Then there exists $0 < \alpha < 1$ independent of k such that

(III.13)

$$\forall i \in \{1, \dots, n\} \quad \forall j \in \{1, \dots, Q^i\} \quad \text{and} \quad \forall \rho > 0 \quad \text{with} \quad B_{2\rho}(x_k^{i,j}) \setminus B_\rho(x) \subset \Omega_k(\alpha)$$

$$\text{then} \quad \int_{B_{2\rho}(x_k^{i,j}) \setminus B_\rho(x)} |d\vec{n}_{\vec{\xi}_k}|_{g_k}^2 \, d\text{vol}_{g_k} < \varepsilon_0$$

⁷This set might be empty.

and such that

$$(III.14) \quad \forall i \in \{1, \dots, n\} \quad \forall j \in \{1, \dots, Q^i\}$$

$$\text{then} \quad \int_{B_{\rho_k^{i,j}}(x_k^{i,j}) \setminus \bigcup_{j' \in I^{i,j}} B_{\alpha \rho_k^{i,j}}(x_k^{i,j'})} |d\vec{n}_{\xi_k}|_{g_k}^2 \, d\text{vol}_{g_k} > \varepsilon_0.$$

Given $0 < \alpha < 1$, we set

$$\forall i \in \{1, \dots, n\} \quad \forall j \in \{1, \dots, Q^i\}$$

$$r_k^{i,j} := \inf \left\{ 0 < r < \rho_k^{i,j} ; \begin{array}{l} B_\rho(x) \subset B_{\alpha^{-1} \rho_k^{i,j}}(x_k^{i,j}) \setminus \bigcup_{j' \in I^{i,j}} B_{\alpha \rho_k^{i,j}}(x_k^{i,j'}) \\ \int_{B_r(x)} |d\vec{n}_{\xi_k}|_{g_k}^2 \, d\text{vol}_{g_k} = \varepsilon_0 \end{array} \right\}.$$

Then there holds

$$(III.15) \quad \forall i \in \{1, \dots, n\} \quad \forall j \in \{1, \dots, Q^i\} \quad \liminf_{k \rightarrow +\infty} \frac{r_k^{i,j}}{\rho_k^{i,j}} > 0.$$

We prove this proposition inductively by repeating finitely many iterations of the following lemma.

LEMMA III.2 (Energy tracking procedure). *Let $\vec{\xi}_k$ be a sequence conformal weak immersions from the two-dimensional annulus $\Omega_k := B_{R_k}(0) \setminus B_{r_k}(0) \subset \mathbb{C}$ into \mathbb{R}^m , with $R_k/r_k \rightarrow +\infty$. We assume that*

$$(III.16) \quad \int_{B_{R_k}(0) \setminus B_{r_k}(0)} |\nabla \vec{n}_{\xi_k}|^2 \, dx \leq \Lambda < +\infty.$$

Then for any $\varepsilon_0 > 0$, one can extract a subsequence, still denoted $\vec{\xi}_k$, such that there exist a fixed integer $N \in \mathbb{N}$ satisfying

$$(III.17) \quad N \leq 4 \left(\frac{\Lambda}{\varepsilon_0} + 1 \right)$$

and a sequence of decreasing families of $(N+1)$ radii

$$R_k^0 = R_k > R_k^1 > \dots > R_k^N = r_k$$

for which

$$(III.18) \quad \forall i \in \{0, \dots, N-1\} \quad \int_{\partial B_{R_k^i}} |\nabla \vec{n}_{\xi_k}| \, d\mathcal{H}^1 < \sqrt{\frac{5\pi}{2}} \varepsilon_0.$$

Moreover, the set $I = \{0, \dots, N-1\}$ can be decomposed into two disjoint subsets $I := I_0 \cup I_1$ with

$$(III.19) \quad \forall i \in I_0 \quad \log_2 \frac{4}{3} < \tau_i := \lim_{k \rightarrow +\infty} \log \frac{R_k^i}{R_k^{i+1}} < +\infty$$

and

(III.20)

$$\forall i \in I_1 \quad \int_{B_{R_k^i} \setminus B_{R_k^{i+1}}} |\nabla \vec{n}_{\xi_k}|^2 dx < \varepsilon \quad \text{and} \quad \lim_{k \rightarrow +\infty} \log \frac{R_k^i}{R_k^{i+1}} = +\infty.$$

Proof of Lemma III.2. We begin by constructing a decreasing sequence of radii $\{\rho_k^0 = R_k > \rho_k^1 > \dots > \rho_k^{Q-1} > \rho_k^Q = r_k\}$ such that

$$(III.21) \quad \forall j \in \{0, \dots, Q\} \quad \int_{\partial B_{\rho_k^j}(0)} |\nabla \vec{n}_{\xi_k}| d\mathcal{H}^1 < \sqrt{\frac{5\pi}{2}} \varepsilon_0$$

and

$$(III.22) \quad \forall j \in \{0, \dots, Q\} \quad \log_2 \frac{\rho_k^{j-1}}{\rho_k^j} > \log_2 \frac{4}{3}.$$

Moreover, for any $j \in J = \{0, \dots, Q-1\}$, one of the following three possibilities holds:

(i)

$$(III.23) \quad \varepsilon_0/2 < \int_{B_{\rho_k^j} \setminus B_{\rho_k^{j+1}}} |\nabla \vec{n}_{\xi_k}|^2 dx < \varepsilon_0$$

(ii)

$$(III.24) \quad \int_{B_{\rho_k^j} \setminus B_{\rho_k^{j+1}}} |\nabla \vec{n}_{\xi_k}|^2 dx < \varepsilon_0/2$$

and either $\rho_k^{j+1} = \rho_k^Q = r_k$ or

$$(III.25) \quad \int_{B_{\rho_k^{j+1}} \setminus B_{\rho_k^{j+2}}} |\nabla \vec{n}_{\xi_k}|^2 dx > \varepsilon_0/2 \quad \text{and} \quad \log_2 \frac{\rho_k^{j+1}}{\rho_k^{j+2}} \leq C$$

(iii)

$$(III.26) \quad \int_{B_{\rho_k^j} \setminus B_{\rho_k^{j+1}}} |\nabla \vec{n}_{\xi_k}|^2 dx > \varepsilon_0/2 \quad \text{and} \quad \log_2 \frac{\rho_k^j}{\rho_k^{j+1}} \leq C,$$

where⁸ $C := [2\Lambda/\varepsilon_0] + 2$.

The construction of ρ_k^j is done by induction on j . We take first $\rho_k^0 := R_k$. Assume that we have found some ρ_k^j satisfying (III.21)–(III.22) and one of the

⁸ $[a]$ denotes the integer part of $a \in \mathbb{R}$, namely, the largest integer less than or equal to a .

alternatives (i)–(iii) for $l \leq j$, and such that (III.24) fails for j . To construct ρ_k^{j+1} , we proceed as follows. Let $\rho \geq r_k$ be such that

$$\int_{B_{\rho^j} \setminus B_{\rho}(0)} |\nabla \vec{n}_{\xi_k}|^2 dx = \varepsilon_0/2.$$

If no such ρ exists, we simply choose $\rho_k^{j+1} = r_k$.

For any $0 < \alpha < 1$ and $R_k > t > \alpha^{-1} r_k$, there holds

$$(III.27) \quad \int_{\alpha t}^t ds \int_{\partial B_s(0)} |\nabla \vec{n}_{\xi_k}| \leq t \sqrt{\pi(1-\alpha^2)} \sqrt{\int_{B_t \setminus B_{\alpha t}} |\nabla \vec{n}_{\xi_k}|^2}.$$

Suppose first that

$$\int_{B_{\rho} \setminus B_{\rho/2}} |\nabla \vec{n}_{\xi_k}|^2 \leq \varepsilon_0/2.$$

Applying (III.27) with $t = 3\rho/4$ and $\alpha = 2/3$, and using the mean value theorem, yields some $s \in [\rho/2, 3\rho/4]$ such that

$$\int_{\partial B_s(0)} |\nabla \vec{n}_{\xi_k}| d\mathcal{H}^1 < \sqrt{\frac{5\pi}{2}} \varepsilon_0.$$

Then $\rho_k^{j+1} := s$ satisfies (III.21), (III.22), and the alternative (III.23).

Suppose next that

$$\int_{B_{\rho} \setminus B_{\rho/2}} |\nabla \vec{n}_{\xi_k}|^2 > \varepsilon_0/2.$$

Then there must exist some $l \in \{1, \dots, [2\Lambda/\varepsilon_0] + 1\}$ for which

$$\int_{B_{2^{-l}\rho} \setminus B_{2^{-l-1}\rho}} |\nabla \vec{n}_{\xi_k}|^2 \leq \varepsilon_0/2.$$

Calling again upon (III.27) with $t = 2^{-l}\rho$ and $\alpha = 1/2$, and using the mean value theorem, we obtain some $s \in [2^{-l-1}\rho, 2^{-l}\rho]$ such that

$$(III.28) \quad \int_{\partial B_s(0)} |\nabla \vec{n}_{\xi_k}| d\mathcal{H}^1 < \sqrt{\frac{3\pi}{2}} \varepsilon_0.$$

If $\rho_k^j \leq 2\rho$, then we let $\rho_k^{j+1} := s$ and note that (III.21), (III.22), and the third alternative (III.26) with $C = [2\Lambda/\varepsilon_0] + 2$ are satisfied. If instead $\rho_k^j > 2\rho$, we pick some $\sigma \in [2\rho, \rho]$ such that

$$\int_{\partial B_{\sigma}(0)} |\nabla \vec{n}_{\xi_k}| d\mathcal{H}^1 < \sqrt{\frac{3\pi}{2}} \varepsilon_0$$

and set $\rho_k^{j+1} := \sigma$ and $\rho_k^{j+2} := s$. One verifies that (III.21), (III.22), and the second alternative (III.24)–(III.25) is satisfied.

The above inductive construction must stop after a number of steps smaller than $[4\Lambda/\varepsilon_0] + 2$. Indeed, at least every other index, an amount of $\varepsilon_0/2$ of the (finite) Dirichlet energy of \vec{n}_{ξ_k} is exhausted.

To complete the proof of the lemma, we first extract a subsequence $\vec{\xi}_{k'}$ such that

- Q is independent of k' ;
- for each $j \in \{0, \dots, Q-1\}$, we have

$$\tau_j := \lim_{k' \rightarrow +\infty} \log_2 \frac{\rho_{k'}^j}{\rho_{k'}^{j+1}} \in \mathbb{R} \cup \{+\infty\}.$$

We keep denoting this subsequence by $\vec{\xi}_k$. From the sequence ρ_k^j , we extract the desired decreasing sequence of radii R_k^j by merging successive annuli such that

$$\lim_{k' \rightarrow +\infty} \log_2 \frac{\rho_{k'}^j}{\rho_{k'}^{j+1}} = \tau_j \in \mathbb{R}$$

and dropping the intermediate ρ_k^j . One easily verifies that the obtained sequence R_k^j satisfies (III.17)–(III.20), thus completing the proof of Lemma III.2. \square

Proof of Proposition III.1. Let $0 < \varepsilon_0 < 8\pi/3$. Consider

$$\rho_k := \inf \left\{ \rho ; \int_{B_\rho(x)} |d\vec{n}_{\xi_k}|_{g_k}^2 \, \text{dvol}_{g_k} = \varepsilon_0 \quad \forall x \in \Sigma \right\},$$

where $B_\rho(x)$ denotes the geodesic ball of center x and radius ρ for the constant scalar curvature h_k . If

$$\limsup_{k \rightarrow +\infty} \rho_k > 0,$$

we can extract a subsequence such that ρ_k converges to a positive constant. Then there is no a_j . This would indeed require the concentration of at least $8\pi/3$ of energy, which is precluded by the result proved in [Riv13c]. There is no bubble and the procedure stops.

Alternatively, we consider the case when

$$\lim_{k \rightarrow +\infty} \rho_k = 0.$$

Let $x_k \in \Sigma$ be such that

$$\int_{B_{\rho_k}(x_k)} |d\vec{n}_{\xi_k}|_{g_k}^2 \, \text{dvol}_{g_k} = \varepsilon_0.$$

Modulo the extraction of a subsequence, we have that x_k converges to one of the points a^i . We set $x_k^{i,1} := x_k$ and $\rho_k^{i,1} := \rho_k$. By hypothesis, we may choose converging conformal coordinates around a^i . For notational convenience and clarity, since the converging metrics h_k are uniformly equivalent to the flat

metric, we shall work with flat geodesic balls. We can choose a subsequence $\vec{\xi}_k$ and a fixed radius $\alpha > 0$ such that

$$\lim_{k \rightarrow +\infty} \sup_{0 < r \leq \alpha} \left\{ r ; \int_{B_\alpha(0) \setminus B_r(0)} |\nabla \vec{n}_{\vec{\xi}_k}|^2 dx = \varepsilon_0 \right\} = 0,$$

and we apply Lemma III.2 to this subsequence, still denoted $\vec{\xi}_k$, and to $R_k := \alpha$ and $r_k := \rho_k^{i,1} = \rho_k$. This gives then the existence of this family of radii

$$R_k^0 = \alpha > R_k^1 > \dots > R_k^N = \rho_k^{i,0}$$

satisfying (III.18)–(III.20) and

$$(III.29) \quad \lim_{k \rightarrow +\infty} \log \frac{R_k^0}{R_k^1} = +\infty.$$

We merge successive radii R_k^i and R_k^{i+1} for $i \in I_1$ or for $i \in I_0$ such that

$$\int_{B_{R_k^i}(x_k^{i,1}) \setminus B_{R_k^{i+1}}(x_k^{i,1})} |\nabla \vec{n}_{\vec{\xi}}|^2 dx_1 dx_2 \leq \varepsilon_0.$$

We then get a subsequence of radii $R_k^0 = \alpha > R_k^{i_1} > \dots > R_k^{i_P} = \rho_k^{i,0}$ for $P \leq N$ such that either, for $i_l \in I_0$,

$$\lim_{k \rightarrow +\infty} \log \frac{R_k^{i_l}}{R_k^{i_l+1}} < +\infty \quad \text{and} \quad \int_{B_{R_k^{i_l}}(0) \setminus B_{R_k^{i_l+1}}(0)} |\nabla \vec{n}_{\vec{\xi}}|^2 dx_1 dx_2 \geq \varepsilon_0$$

or, when $i_l \in I_1$, we have

$$\lim_{k \rightarrow +\infty} \log \frac{R_k^{i_l}}{R_k^{i_l+1}} = +\infty$$

and

$$\forall \rho \in (R_k^{i_l+1}, R_k^{i_l}/2) \quad \int_{B_{2\rho}(x_k^{i,1}) \setminus B_\rho(x_k^{i,1})} |\nabla \vec{n}_{\vec{\xi}}|^2 dx_1 dx_2 < \varepsilon_0.$$

We consider the smallest annulus $\Omega_k^{i_l} := B_{R_k^{i_l}} \setminus B_{R_k^{i_l+1}}$ of the first type $i_l \in I_0$.

For such an i_l we introduce

$$\rho_k^{i_l} := \inf \left\{ \rho < R_k^{i_l+1} ; \int_{B_\rho(x)} |d\vec{n}_{\vec{\xi}_k}|_{g_k}^2 d\text{vol}_{g_k} = \varepsilon_0 \quad \text{where} \quad x \in \Omega_k^{i_l} \right\}$$

We now consider the following alternative. Either

$$\limsup_{k \rightarrow +\infty} \frac{\rho_k^{i_l}}{R_k^{i_l}} > 0,$$

we then extract a subsequence still denoted $\vec{\xi}_k$ such that

$$\lim_{k \rightarrow +\infty} \frac{\rho_k^{i_l}}{R_k^{i_l}} > 0,$$

and we pass to the next $\Omega_k^{i'}$, if there is any, where $i' \in I_0$, or we have

$$\lim_{k \rightarrow +\infty} \frac{\rho_k^{i_l}}{R_k^{i_l}} = 0.$$

In such a case we again apply the *energy tracking Lemma* III.2 on the annulus $B_{R_k^{i_l}/2}(x) \setminus B_{\rho_k^{i_l}}(x)$, where $x \in \Omega_k^{i_l}$ satisfies

$$\int_{B_{\rho_k^{i_l}}(x)} |d\vec{n}_{\vec{\xi}_k}|_{g_k}^2 \, d\text{vol}_{g_k} = \varepsilon_0,$$

and we set $x_k^{i,2} := x$ and $\rho_k^{i,2} := \rho_k^{i_l}$. We keep this procedure running until all annuli of the type I_0 have been explored and until the neighborhood of each a^i has been considered. This can be done in a finite number of steps, since each application of Lemma III.2 takes place around a "bubble" exhausting at least ε_0 of the total energy of the sequence $\vec{\xi}_k$. With moderate effort, one checks that the sequence of finite families of bubbles constructed at the end of the procedure fulfills the desired requirements of Proposition III.1. \square

IV. Construction of a Coulomb moving frame in each component of the neck region

Let $\vec{\xi}_k$ be a family of weakly converging Willmore immersions of Σ fulfilling the conclusions of Lemma II.1. For some ε_0 chosen smaller than the constant ε appearing in the ε -regularity Theorem I.5 in [Riv08], we apply Proposition III.1 so as to obtain a subsequence for which we can decompose the surface into "converging regions" (complements of small neighborhoods of the points a^i), bubbles, and neck regions. In each annulus $B_{R_k}(x_k^{i,j}) \setminus B_{r_k}(x_k^{i,j}) \subset \Omega_k(\alpha)$ of a part of the neck region $\Omega_k(\alpha)$ satellite to some bubble centered on $x_k^{i,j}$, the condition (III.13) states that

$$\sup_{r_k < \rho < R_k} \int_{B_{2\rho}(x_k^{i,j}) \setminus B_\rho(x_k^{i,j})} |\nabla \vec{n}_{\vec{\xi}_k}| \, dx < \varepsilon_0.$$

Combining this fact with the ε -regularity Theorem I.5 in [Riv08] gives that

$$(IV.1) \quad \forall x \in B_{R_k/2}(x_k^{i,j}) \setminus B_{2r_k}(x_k^{i,j}) \quad |\nabla \vec{n}_{\vec{\xi}_k}| \leq C_m \frac{\sqrt{\varepsilon_0}}{|x - x_k^{i,j}|}.$$

This pointwise estimate translates into a control of the L^2 -weak norm of $\nabla \vec{n}_{\vec{\xi}}$ in the annulus, namely:

$$(IV.2) \quad \|\nabla \vec{n}_{\vec{\xi}_k}\|_{L^{2,\infty}(B_{R_k/2}(x_k^{i,j}) \setminus B_{2r_k}(x_k^{i,j}))} \leq C_m \sqrt{\varepsilon_0}.$$

This L^2 -weak estimate for the gradient of the Gauss map in neck regions already appears in previous works [LR01], [LR02], and [Riv02b], where it plays a decisive role. It is also of utmost importance in the present paper, as it will

enable us in Lemma IV.3 to construct a $W^{1,2}$ Coulomb gauge without having to resort to the stronger L^2 -control of the gradient of the Gauss map commonly required in similar situations. Our results are generic in nature, and they should be helpful to the study of other gauge invariant problems.

We first need an extension lemma.

LEMMA IV.1. *Let $\vec{n} \in W^{1,2}(B_{2\rho}(0) \setminus B_\rho(0), G_{m-2}(\mathbb{R}^m))$, where $G_{m-2}(\mathbb{R}^m)$ denotes the Grassmannian manifold of oriented $(m-2)$ -planes in \mathbb{R}^m . There exists $\delta(m) > 0$ depending only on m such that if*

$$(IV.3) \quad \int_{\partial B_\rho(0)} |\nabla \vec{n}| \, dl_{\partial B_\rho(0)} \leq \delta(m),$$

then we can extend \vec{n} to $\hat{\vec{n}} \in W^{1,2}(B_{2\rho}(0), G_{m-2}(\mathbb{R}^m))$ with $\hat{\vec{n}} = \vec{n}$ on the annulus $B_{2\rho}(0) \setminus B_\rho(0)$. Furthermore,

$$(IV.4) \quad \|\nabla \hat{\vec{n}}\|_{L^2(B_{2\rho}(0))} \leq C_m \|\nabla \vec{n}\|_{L^2(B_{2\rho}(0) \setminus B_\rho(0))}$$

and

$$(IV.5) \quad \|\nabla \hat{\vec{n}}\|_{L^{2,\infty}(B_\rho(0))} \leq C_m \delta(m),$$

where the constant $C_m > 0$ depends only on m .

Proof of Lemma IV.1. Let

$$\vec{n}_0 := \int_{\partial B_\rho(0)} \vec{n} \, dl_{\partial B_\rho(0)}.$$

Then

$$\|\vec{n} - \vec{n}_0\|_{L^\infty(\partial B_\rho(0))} \leq C \int_{\partial B_\rho(0)} |\nabla \vec{n}| \, dl_{\partial B_\rho(0)} \leq C \delta(m)$$

for some constant C independent of \vec{n} and ρ .

Thus, if $\delta(m) > 0$ is chosen small enough and independently of \vec{n}_0 , the plane $\vec{n}(p)$ is contained in a small geodesic ball $B_\delta(\vec{n}_0) \subset G_{m-2}(\mathbb{R}^m)$ on which coordinates $y = (y_1, \dots, y_N)$ exist. We may further choose these coordinates in such a way that $B_\delta(\vec{n}_0)$ corresponds to the ball $B_\delta^N(0)$ of radius δ and centered on $0 \in \mathbb{R}^N$. Owing to the compactness of $G_{m-2}(\mathbb{R}^m)$, we may always arrange for $\|\nabla y^{-1}\|_{L^\infty(B_\delta^N(0))}$ to depend only on m . We have⁹

$$(IV.6) \quad \|\vec{n} - \vec{n}_0\|_{H^{1/2}(\partial B_\rho(0))} \simeq \sum_{l=1}^N \|n_l - (n_l)_0\|_{H^{1/2}(\partial B_\rho(0))},$$

⁹On the left-hand side of (IV.6), we use the isometric embedding of $G_{m-2}(\mathbb{R}^m)$ into some \mathbb{R}^{K_m} , and \vec{n} is viewed as an \mathbb{R}^{K_m} -valued map.

where n_l are the coordinates of \vec{n} in the system $\{y_l\}_{l=1,\dots,N}$, and

$$(n_l)_0 := \int_{\partial B_\rho(0)} n_l.$$

Let \hat{n}_l be the harmonic extension of n_l into $B_\rho(0)$. Since the \mathbb{R}^N -valued map (n_l) is included in the ball $B_\delta^N(0)$, the maximum principle yields that $(\hat{n}_l)_{l=1,\dots,N} \in B_\delta^N(0)$. Moreover, standard estimates on harmonic extensions and Sobolev embedding estimates give

$$\begin{aligned} \int_{B_\rho(0)} |\nabla \hat{n}_l|^2 &\leq C \|n_l - (n_l)_0\|_{H^{1/2}(\partial B_\rho(0))}^2 \lesssim C \|\vec{n} - \vec{n}_0\|_{H^{1/2}(\partial B_\rho(0))}^2 \\ &\leq C \|\nabla \vec{n}\|_{L^1(\partial B_\rho(0))}^2 \leq C \delta^2, \end{aligned}$$

where the constants C and C_0 depend only on m . Note that we have again used the characterization of \vec{n} as an \mathbb{R}^{K_m} -valued map. Using the general fact that $\|\cdot\|_{L^{2,\infty}(U)} \leq 2\|\cdot\|_{L^2(U)}$ for all U , the latter yields

$$(IV.7) \quad \|\nabla \hat{n}_l\|_{L^{2,\infty}(B_\rho(0))} \leq C_1 \delta(m)$$

for some constant C_1 depending only on m . In addition, using a trace theorem, we obtain

$$(IV.8) \quad \begin{aligned} \int_{B_\rho(0)} |\nabla \hat{n}_l|^2 &\leq C \|n_l - (n_l)_0\|_{H^{1/2}(\partial B_\rho(0))}^2 \lesssim C \|\vec{n} - \vec{n}_0\|_{H^{1/2}(\partial B_\rho(0))}^2 \\ &\leq C_2 \|\nabla \vec{n}\|_{L^2(B_{2\rho}(0) \setminus B_\rho(0))}^2, \end{aligned}$$

where the constant C_2 depends only on m .

We next set $\hat{\vec{n}} := y^{-1}((\hat{n}_l))$. Since $\|\nabla y^{-1}\|_{L^\infty(B_\delta(0))}$ only depends on m , (IV.7) and (IV.8) imply the desired (IV.5) and (IV.4), thereby proving Lemma IV.1. \square

The main result of this section also requires an integrability-by-compensation lemma for second-order equations in divergence form. It was originally proved by Y. Ge in [Ge99] through a different method from the one used here.

LEMMA IV.2 (Wente estimates in Lorentz $L^{p,q}$ spaces). *Consider the divergence-form problem*

$$(IV.9) \quad \begin{cases} \Delta \varphi = \nabla^\perp a \cdot \nabla b & \text{in } B_1(0), \\ \varphi = 0 & \text{on } \partial B_1(0), \end{cases}$$

where $\nabla a \in L^{2,\infty}(B_1(0))$, and $\nabla b \in L^{p,q}(B_1(0))$ for some $p \in (1, \infty)$ and $q \in [1, \infty]$. Then there holds

$$(IV.10) \quad \|\nabla \varphi\|_{L^{p,q}(B_1(0))} \leq C_{p,q} \|\nabla a\|_{L^{2,\infty}(B_1(0))} \|\nabla b\|_{L^{p,q}(B_1(0))}$$

for some constant $C_{p,q} > 0$.

Proof. Owing to the interpolative nature of the Lorentz space $L^{p,q}$, it suffices to show that (IV.10) holds for $p = q \in (1, \infty)$. Consider first the case when $p > 2$. According to the Hölder inequality for Lorentz spaces, there holds

$$(IV.11) \quad L^{2,\infty} \cdot L^p \subset L^{\frac{2p}{2+p},p}.$$

Since $p > 2$, we call upon the usual Calderon-Zygmund theorem and the Sobolev embedding theorem to obtain that $\nabla\varphi \in W^{1,\frac{2p}{2+p}} \subset L^p$ with the estimate (IV.10), as claimed.

Next, when $1 < p \leq 2$, we use the divergence-form structure of the equation. Note that

$$(IV.12) \quad \Delta\varphi = \operatorname{div}(b \nabla^\perp a),$$

and that

$$(IV.13) \quad b \nabla^\perp a \in W^{1,p} \cdot L^{2,\infty} \subset L^{\frac{2p}{2-p},p} \cdot L^{2,\infty} \subset L^p,$$

where we have again used the Sobolev embedding theorem and the Hölder inequality for Lorentz spaces. The desired (IV.10) follows immediately from (IV.12) and (IV.13), and this concludes the proof of Lemma IV.2. \square

We next state and prove the main result of this section.

LEMMA IV.3 ($W^{1,2}$ -controlled Coulomb frame from small $L^{2,\infty}$ -control of the second fundamental form). *Let $\vec{n} \in W^{1,2}(B_1(0) \setminus B_r(0), G_{m-2}(\mathbb{R}^m))$, where $0 \leq r < 1/2$. There exists $\eta(m) > 0$ depending only on m (but not on r) such that if*

$$(IV.14) \quad \|\nabla\vec{n}\|_{L^{2,\infty}(B_1(0) \setminus B_r(0))} \leq \eta(m)$$

and

$$(IV.15) \quad \int_{\partial B_r(0)} |\nabla\vec{n}| \, dl_{\partial B_r(0)} \leq \eta(m),$$

then one can construct an associated Coulomb frame $\{\vec{e}_1, \vec{e}_2\}$ satisfying

$$\star \vec{n} = \vec{e}_1 \wedge \vec{e}_2 \quad \text{and} \quad \operatorname{div}(\vec{e}_1 \cdot \nabla \vec{e}_2) = 0.$$

Furthermore,

$$(IV.16) \quad \int_{B_1(0) \setminus B_r(0)} |\nabla \vec{e}_1|^2 + |\nabla \vec{e}_2|^2 \leq C \int_{B_1(0) \setminus B_r(0)} |\nabla \vec{n}|^2$$

and

$$(IV.17) \quad \sum_{i=1}^2 \|\nabla \vec{e}_i\|_{L^{2,\infty}(B_1(0) \setminus B_r(0))} \leq C \|\nabla \vec{n}\|_{L^{2,\infty}(B_1(0) \setminus B_r(0))}$$

for some constant C depending on m , but neither on r nor on \vec{n} . \square

Proof. Owing to Lemma IV.1, we can always reduce to the case when $r = 0$. Indeed, we have seen that the harmonic extension $\hat{\vec{n}}$ in $B_r(0)$ of the restriction of \vec{n} to $B_{2r}(0) \setminus B_r(0)$ satisfies $\hat{\vec{n}} = \vec{n}$ on $B_1(0) \setminus B_r(0)$, and we have the estimate

$$\|\nabla \hat{\vec{n}}\|_{L^2, \infty(B_1(0))} \leq \|\nabla \vec{n}\|_{L^2, \infty(B_1(0) \setminus B_r(0))} + \|\nabla \hat{\vec{n}}\|_{L^2, \infty(B_r(0))} \leq \eta(m) + C_m \eta(m).$$

In particular, if Lemma IV.3 holds for $r = 0$ with the constant $\eta(m) > 0$, then it also holds for an arbitrary $0 < r < 1/2$ with the constant $\eta(m)/(C_m + 1)$ playing the role of $\eta(m)$ in the statement. We shall thus focus on the case $r = 0$.

We first suppose that \vec{n} is smooth. According to [Hél02, Th. 5.2.1], there exists a smooth moving orthonormal two-frame $\{\vec{f}_1, \vec{f}_2\}$ in $W^{1,2}(B_1(0), \mathbb{R}^m)$ with $\star \vec{n} = \vec{f}_1 \wedge \vec{f}_2$. Its energy is however not controlled since we do not assume that $\|\nabla \vec{n}\|_{L^2(B_1(0))}$ is smaller than the required threshold $8\pi/3$.

For each $r \in [0, 1]$, let $\{\vec{f}_{1,r}(x), \vec{f}_{2,r}(x)\} := \{\vec{f}_1(rx), \vec{f}_2(rx)\}$. We minimize

$$(IV.18) \quad F_r(\theta_r) := \int_{B_r(0)} \left| \nabla \theta_r + (\vec{f}_{1,r} \cdot \nabla \vec{f}_{2,r}) \right|^2$$

over all rotations $\theta_r \in W^{1,2}(B_1(0), \mathbb{R})$. As explained in Theorem 4.1.1 from [Hél02], for each r , the minimum of F_r is attained at some frame $\{\vec{e}_{1,r}, \vec{e}_{2,r}\}$ satisfying

$$(IV.19) \quad \vec{e}_{1,r} + i \vec{e}_{2,r} = e^{i\theta_r} (\vec{f}_{1,r} + i \vec{f}_{2,r})$$

and

$$(IV.20) \quad \begin{cases} \operatorname{div}(\vec{e}_{1,r} \cdot \nabla \vec{e}_{2,r}) = 0 & \text{on } B_r(0), \\ \vec{e}_{1,r} \cdot \partial_\nu \vec{e}_{2,r} = 0 & \text{on } \partial B_r(0). \end{cases}$$

Thus, there exists λ_r equal to zero on $\partial B_r(0)$ and solving

$$\nabla \lambda_r = -\vec{e}_{1,r} \cdot \nabla^\perp \vec{e}_{2,r}.$$

In particular, it satisfies

$$(IV.21) \quad \begin{cases} \Delta \lambda_r = \nabla^\perp \vec{e}_{1,r} \cdot \nabla \vec{e}_{2,r} & \text{in } B_r(0), \\ \lambda_r = 0 & \text{on } \partial B_r(0). \end{cases}$$

Using the Wente inequality Lemma IV.2 with $(p, q) = (2, \infty)$ yields

$$(IV.22) \quad \|\nabla \lambda_r\|_{L^2, \infty(B_r(0))} \leq C_0 \|\nabla \vec{e}_{1,r}\|_{L^2, \infty(B_r(0))} \|\nabla \vec{e}_{2,r}\|_{L^2, \infty(B_r(0))}$$

for some constant C_0 independent of r . Moreover, as shown in the proof of Lemma 5.1.4 in [Hél02], an elementary computation gives

$$(IV.23) \quad |\nabla \vec{e}_{1,r}|^2 + |\nabla \vec{e}_{2,r}|^2 = 2 |\nabla \lambda_r|^2 + |\nabla \vec{n}|^2.$$

Combining altogether (IV.22) and (IV.23) gives the inequality

$$(IV.24) \quad \|\nabla\lambda_r\|_{L^{2,\infty}(B_r(0))}^2 + \frac{1}{4C_0} (C_0\|\nabla\vec{n}\|_{L^{2,\infty}} - 1)\|\nabla\lambda_r\|_{L^{2,\infty}(B_r(0))} \\ + C_0\|\nabla\vec{n}\|_{L^{2,\infty}(B_r(0))} \geq 0.$$

Accordingly, there exists some small enough threshold $\eta(C_0) \equiv \eta(m)$ for which the hypothesis that $\|\nabla\vec{n}\|_{L^{2,\infty}(B_1(0))} < \eta(m)$ guarantees that the range of $\|\nabla\lambda_r\|_{L^{2,\infty}(B_r(0))}$ is of the form $[0, \alpha(\eta)] \cup [\beta(\eta), \infty)$ for some $0 < \alpha(\eta) < \beta(\eta)$. One easily checks from (IV.24) that $\alpha(\eta) \leq C_1\|\nabla\vec{n}\|_{L^{2,\infty}(B_1(0))}$ with the constant C_1 depending only on C_0 , and thus only on m .

As $\vec{f}_{j,r}(x) = f_j(rx)$ and \vec{f}_j belongs to $W^{1,2}(B_1(0))$, it is clear that $\vec{f}_{j,r}$ is continuous in the parameter r . From this and the definition of θ_r , it follows that θ_r is likewise continuous in r , and hence from (IV.19) and (IV.21), that $\vec{e}_{j,r}$ and λ_r are continuous in r . We see that $\|\nabla\lambda_r\|_{L^{2,\infty}(B_r(0))}$ takes the value 0 at $r = 0$. From the above discussion, we deduce that at $r = 1$,

$$(IV.25) \quad \|\nabla\lambda\|_{L^{2,\infty}(B_1(0))} \equiv \|\nabla\lambda_1\|_{L^{2,\infty}(B_1(0))} \leq \alpha(\eta) \leq C_1\|\nabla\vec{n}\|_{L^{2,\infty}(B_1(0))}.$$

Let now $\{\vec{e}_1, \vec{e}_2\} := \{\vec{e}_{1,1}, \vec{e}_{2,1}\}$. By definition, this Coulomb frame satisfies

$$(IV.26) \quad \star \vec{n} = \vec{e}_1 \wedge \vec{e}_2, \quad \operatorname{div}(\vec{e}_1 \cdot \nabla \vec{e}_2) = 0, \quad \nabla^\perp \lambda = \vec{e}_1 \cdot \nabla \vec{e}_2 = -\vec{e}_2 \cdot \nabla \vec{e}_1.$$

Note next that

$$(IV.27) \quad |\nabla \vec{e}_j|^2 = |\vec{e}_{j'} \cdot \nabla \vec{e}_j|^2 + |\pi_{\vec{n}} \nabla \vec{e}_j|^2 \leq |\nabla \lambda|^2 + |\nabla \vec{n}|^2 \quad \text{for } (j, j') \in \{(1, 2), (2, 1)\}.$$

In particular, using (IV.25), we find

$$(IV.28) \quad \|\nabla \vec{e}_j\|_{L^{2,\infty}(B_1(0))} \leq \|\nabla \lambda\|_{L^{2,\infty}(B_1(0))} + \|\nabla \vec{n}\|_{L^{2,\infty}(B_1(0))} \\ \leq (C_1 + 1)\|\nabla \vec{n}\|_{L^{2,\infty}(B_1(0))},$$

thereby giving the required (IV.17).

Calling again upon the Wente inequality of Lemma IV.2 with $(p, q) = (2, 2)$, we obtain from (IV.21) with $r = 1$ that

$$\|\nabla \lambda\|_{L^2(B_1(0))} \leq C_2 \|\nabla \vec{e}_1\|_{L^{2,\infty}(B_1(0))} \|\nabla \vec{e}_2\|_{L^2(B_1(0))} \\ \leq C_2 (C_1 + 1) \eta(m) \left(\|\nabla \lambda\|_{L^2(B_1(0))} + \|\nabla \vec{n}\|_{L^2(B_1(0))} \right),$$

where we have used (IV.27), (IV.28), and the hypothesis (IV.14). Here C_2 is a constant that depends only on m . The latter shows that $\|\nabla \lambda\|_{L^2(B_1(0))} \leq C_3 \eta(m) \|\nabla \vec{n}\|_{L^2(B_1(0))}$ which, once introduced into (IV.27), gives the announced

$$(IV.29) \quad \int_{B_1(0)} |\nabla \vec{e}_1|^2 + |\nabla \vec{e}_2|^2 \leq C \int_{B_1(0)} |\nabla \vec{n}|^2,$$

where the constant C depends only on m .

In the general case when \vec{n} is not smooth, we use the limiting process outlined in step 6 of the proof of Lemma 5.1.4 from [Hél02]. Applying the known fact [SU83] that $C^\infty(B_1(0), G_{m-2}(\mathbb{R}^m))$ is dense in $W^{1,2}(B_1(0), G_{m-2}(\mathbb{R}^m))$, we have completed the proof of Lemma IV.3. \square

V. General estimates in the neck regions of weakly converging weak conformal immersions

LEMMA V.1 (Estimate of the “Euler characteristic production” in neck regions). *There exists a constant $\eta(m) > 0$ with the following property. Let $\eta < \eta(m)$, let $R > 4r > 0$, and let $\vec{\xi}$ be a conformal weak immersion from $B_R(0) \setminus B_r(0)$ into \mathbb{R}^m with L^2 -bounded second fundamental form satisfying*

$$(V.1) \quad \|\nabla \vec{n}_{\vec{\xi}}\|_{L^2, \infty(B_R \setminus B_r)} \leq \eta,$$

$$(V.2) \quad \int_{\partial B_r(0)} |\nabla \vec{n}_{\vec{\xi}}| \, dl_{\partial B_r(0)} \leq \eta,$$

and

$$(V.3) \quad \int_{B_R \setminus B_{R/2} \cup B_{2r} \setminus B_r} |\nabla \vec{n}_{\vec{\xi}}|^2 \, dx \leq \eta.$$

Then there exists a positive constant C_m depending only on m , but not on the conformal type of the annulus $\log \frac{R}{r}$, such that

$$(V.4) \quad \left| \int_{B_R(0) \setminus B_r(0)} K_{\vec{\xi}} \, d\text{vol}_{g_{\vec{\xi}}} \right| \leq C_m \eta,$$

where $K_{\vec{\xi}}$ denotes the Gaussian curvature associated with the immersion $\vec{\xi}$.

Proof. Let $\Omega = B_R(0) \setminus B_r(0)$. According to Lemma IV.3 (applicable here owing to the hypotheses (V.1) and (V.2)), there exists a Coulomb moving frame

$$\{\vec{e}_1, \vec{e}_2\} \in (W^{1,2}(\Omega, S^{m-1}))^2$$

satisfying

$$\star \vec{n}_{\vec{\xi}} = \vec{e}_1 \wedge \vec{e}_2$$

and

$$(V.5) \quad \sum_{i=1}^2 \|\nabla \vec{e}_i\|_{L^2, \infty(\Omega)} \leq C_1 \|\nabla \vec{n}\|_{L^2, \infty(\Omega)} \leq C_2 \eta,$$

where the constants C_j depend only on m . The mean value theorem yields the existence of some $\alpha \in (1/2, 1)$ with

$$\begin{aligned}
(V.6) \quad \sum_{i=1}^2 \int_{\partial(B_{\alpha R} \setminus B_{\alpha^{-1}r})} |\nabla \vec{e}_i| &\leq C_3 r^{-1} \sum_{i=1}^2 \|\nabla \vec{e}_i\|_{L^1(B_{2r} \setminus B_r)} \\
&\quad + C_4 R^{-1} \sum_{i=1}^2 \|\nabla \vec{e}_i\|_{L^1(B_R \setminus B_{R/2})} \\
&\leq C_5 \sum_{i=1}^2 \|\nabla \vec{e}_i\|_{L^{2,\infty}(\Omega)} \leq C_5 C_2 \eta,
\end{aligned}$$

where C_5 is a universal constant.

The Liouville equation states that

$$(V.7) \quad \Delta \lambda = -e^{2\lambda} K_{\vec{\xi}},$$

where λ is the conformal parameter of the conformal immersion $\vec{\xi}$. From the way the Coulomb frame $\{\vec{e}_1, \vec{e}_2\}$ was constructed, the following also holds on the annulus $B_R(0) \setminus B_r(0)$:

$$(V.8) \quad \Delta \lambda = -\nabla \vec{e}_1 \cdot \nabla^\perp \vec{e}_2 = -\operatorname{div}(\vec{e}_1 \cdot \nabla^\perp \vec{e}_2).$$

Hence, using (V.6) we find that

$$\begin{aligned}
(V.9) \quad \left| \int_{B_{\alpha R}(0) \setminus B_{\alpha^{-1}r}(0)} K_{\vec{\xi}} e^{2\lambda} dx \right| &= \left| \int_{B_{\alpha R}(0) \setminus B_{\alpha^{-1}r}(0)} \nabla \vec{e}_1 \cdot \nabla^\perp \vec{e}_2 dx \right| \\
&\leq \int_{\partial B_{\alpha R}(0)} |e_1 \cdot \partial_\tau e_2| dl + \int_{\partial B_{\alpha^{-1}r}(0)} |e_1 \cdot \partial_\tau e_2| dl \leq C \eta,
\end{aligned}$$

where τ denotes the unit tangent vector to the circles $\partial B_\rho(0)$, namely $\tau = \rho^{-1} \partial_\theta$. Moreover, owing to the hypothesis (V.3) and the fact that $\alpha \in (1/2, 1)$, we have that

$$(V.10) \quad \int_{B_{\alpha^{-1}r}(0) \setminus B_r(0)} |K_{\vec{\xi}}| e^{2\lambda} dx \leq 2^{-1} \int_{B_{\alpha^{-1}r}(0) \setminus B_r(0)} |\nabla \vec{n}_{\vec{\xi}}|^2 dx \leq 2^{-1} \eta,$$

and similarly

$$(V.11) \quad \int_{B_R(0) \setminus B_{\alpha R}(0)} |K_{\vec{\xi}}| e^{2\lambda} dx \leq 2^{-1} \int_{B_R(0) \setminus B_{\alpha R}(0)} |\nabla \vec{n}_{\vec{\xi}}|^2 dx \leq 2^{-1} \eta.$$

Altogether (V.9), (V.10), and (V.11) yield the announced (V.4). \square

LEMMA V.2 (Uniform Harnack estimates for the conformal factor in neck regions). *There exists a constant $\varepsilon(m) > 0$ with the following property. Let $4 < R < \infty$. If $\vec{\xi}$ is a Lipschitz conformal immersion from $\Omega_R := B_R(0) \setminus B_{R^{-1}}(0)$ into \mathbb{R}^m satisfying*

$$(V.12) \quad \sup_{r \in (R^{-1}, R/2)} \int_{B_{2r}(0) \setminus B_r(0)} |\nabla \vec{n}_{\vec{\xi}}|^2 dx \leq \varepsilon(m)$$

and¹⁰

$$(V.13) \quad \|\nabla\lambda\|_{L^2,\infty(\Omega_R)} = A < +\infty,$$

then for any $4R^{-2} < \alpha < 1$ there exists $C_\alpha(A) > 0$, depending only on α and A but not on R , such that

$$(V.14) \quad \sup_{r \in (4R^{-1}, \alpha R)} \frac{\sup_{x \in B_{\alpha^{-1}r}(0) \setminus B_r(0)} e^{\lambda(x)}}{\inf_{x \in B_{\alpha^{-1}r}(0) \setminus B_r(0)} e^{\lambda(x)}} \leq C_\alpha.$$

Proof. Let $r \in (4R^{-1}, R/4)$. From (V.12), we have

$$(V.15) \quad \int_{B_{3r}(0) \setminus B_{r/4}(0)} |\nabla \vec{n}_\xi|^2 dx \leq 4\varepsilon(m).$$

With the help of the Fubini and mean value theorems, we deduce from (V.12) the existence of $\rho \in (r/4, r/2)$ such that

$$(V.16) \quad \int_{\partial B_\rho(0)} |\nabla \vec{n}_\xi|^2 \leq \frac{4}{r} \varepsilon(m) \leq \frac{2}{\rho} \varepsilon(m),$$

and hence

$$(V.17) \quad \int_{\partial B_\rho(0)} |\nabla \vec{n}_\xi| dx \leq \sqrt{4\pi \varepsilon(m)}.$$

Following Lemma IV.1, we choose $\sqrt{4\pi \varepsilon(m)} < \delta(m)$ so as to obtain an extension \hat{n} of \vec{n} in $B_\rho(0)$ with

$$(V.18) \quad \int_{B_{4r}(0)} |\nabla \hat{n}|^2 dx \leq 4C_m \varepsilon(m),$$

where C_m is the constant depending only on m in the statement of Lemma IV.1. Insuring further that $12C_m \varepsilon_m \leq 8\pi$, we can apply Lemma 5.1.4 of [Hél02] in order to find a framing $\{\vec{e}_1, \vec{e}_2\} \in (W^{1,2}(B_{4r}(0)))^2$ satisfying $\star \hat{n} = \vec{e}_1 \wedge \vec{e}_2$ and

$$(V.19) \quad \int_{B_{4r}(0)} \sum_{i=1}^2 |\nabla \vec{e}_i|^2 dx \leq C \int_{B_{4r}(0)} |\nabla \hat{n}|^2 dx \leq 4C C_m \varepsilon(m).$$

The conformal parameter satisfies

$$(V.20) \quad \Delta\lambda = \nabla^\perp \vec{e}_1 \cdot \nabla \vec{e}_2 \quad \text{in } B_{4r}(0) \setminus B_{r/2}(0).$$

Let μ be the solution of

$$(V.21) \quad \begin{cases} \Delta\mu = \nabla^\perp \vec{e}_1 \cdot \nabla \vec{e}_2 & \text{in } B_{4r}(0) \setminus B_{r/2}(0), \\ \mu = 0 & \text{on } \partial(B_{4r}(0) \setminus B_{r/2}(0)). \end{cases}$$

A standard Wente estimate (see Theorem 3.3.8 in [Hél02]) yields on one hand

$$(V.22) \quad \|\mu\|_{L^\infty(B_{4r}(0) \setminus B_{r/2}(0))} \leq (2\pi)^{-1} \|\nabla \vec{e}_1\|_{L^2} \|\nabla \vec{e}_2\|_{L^2} \leq C \varepsilon(m)$$

¹⁰As before, λ is the conformal parameter of $\vec{\xi}$. Namely, $|\partial_{x_1} \vec{\xi}| = e^\lambda = |\partial_{x_2} \vec{\xi}|$.

and on the other hand

$$(V.23) \quad \|\nabla\mu\|_{L^2(B_{4r}(0)\setminus B_{r/2}(0))} \leq \sqrt{3/32\pi} \|\nabla\vec{e}_1\|_{L^2} \|\nabla\vec{e}_2\|_{L^2} \leq C \varepsilon(m).$$

Let $\nu := \lambda - \mu$ be the harmonic function in $B_{4r}(0) \setminus B_{r/2}(0)$ equal to λ on the boundary. Owing to (V.13) and (V.23), there holds

$$(V.24) \quad \|\nabla\nu\|_{L^{2,\infty}(B_{4r}(0)\setminus B_{r/2}(0))} \leq C \varepsilon(m) + A.$$

Let $\bar{\nu}$ be the average of ν on $B_{4r}(0) \setminus B_{r/2}(0)$. The Sobolev-Poincaré inequality gives

$$(V.25) \quad r^{-2} \int_{B_{4r}(0)\setminus B_{r/2}(0)} |\nu - \bar{\nu}| dx \leq C_1 r^{-1} \int_{B_{4r}(0)\setminus B_{r/2}(0)} |\nabla\nu| dx \\ \leq C_2 \|\nabla\nu\|_{L^{2,\infty}(B_{4r}(0)\setminus B_{r/2}(0))} \leq C_3 \varepsilon(m) + C_2 A,$$

where C_j are universal constants. Since $\nu - \bar{\nu}$ is harmonic, we deduce that on a smaller annulus the L^∞ -norm of $(\nu - \bar{\nu})$ is controlled by $\varepsilon(m)$ and A :

$$(V.26) \quad \|\nu - \bar{\nu}\|_{L^\infty(B_{2r}(0)\setminus B_r(0))} \leq C r^{-2} \int_{B_{4r}(0)\setminus B_{r/2}(0)} |\nu - \bar{\nu}| dx_1 dx_2 \leq C_4(\varepsilon(m) + A),$$

where C_4 is a universal constant. We set $\lambda_- := \inf\{\lambda(x) ; x \in B_{2r}(0) \setminus B_r(0)\}$, $\lambda_+ := \max\{\lambda(x) ; x \in B_{2r}(0) \setminus B_r(0)\}$, $\nu_- := \inf\{\nu(x) ; x \in B_{2r}(0) \setminus B_r(0)\}$, and $\nu_+ := \max\{\nu(x) ; x \in B_{2r}(0) \setminus B_r(0)\}$. One has the estimates

$$(V.27) \quad \frac{e^{\lambda_+}}{e^{\lambda_-}} \leq \frac{e^{\nu_+ + \|\mu\|_\infty}}{e^{\nu_- - \|\mu\|_\infty}} \leq e^{2\|\mu\|_\infty} e^{\nu_+ - \nu_-} \leq e^{2C\varepsilon(m)} e^{2C_4(\varepsilon(m) + A)},$$

where we have used respectively (V.22) and (V.26).

We have thus proved (V.14) for $\alpha = 1/2$. Let now $\alpha = 2^{-j}$ for some $j \in \mathbb{N}^*$. The idea is to apply (V.14) successively j times, with $\alpha = 1/2$ and with $2^j r$ in place of r . For $i = 1, \dots, j$, we let $\lambda_-^i := \inf\{\lambda(x) ; x \in B_{2^i r}(0) \setminus B_{2^{i-1} r}(0)\}$ and $\lambda_+^i := \max\{\lambda(x) ; x \in B_{2^i r}(0) \setminus B_{2^{i-1} r}(0)\}$. Using the fact that

$$\lambda_-^i \leq \lambda_+^{i\pm 1}$$

gives

$$\frac{e^{\lambda_+^i}}{e^{\lambda_-^{i\pm 1}}} \leq \frac{e^{\lambda_+^i}}{e^{\lambda_-^i}} \frac{e^{\lambda_-^i}}{e^{\lambda_+^{i\pm 1}}} \frac{e^{\lambda_+^{i\pm 1}}}{e^{\lambda_-^{i\pm 1}}} \leq \frac{e^{\lambda_+^i}}{e^{\lambda_-^i}} \frac{e^{\lambda_+^{i\pm 1}}}{e^{\lambda_-^{i\pm 1}}}.$$

The latter enables an iteration and eventually yields that (V.14) holds with $\alpha = 2^{-j}$ for an arbitrary positive integer j satisfying $4R^{-2} < 2^{-j}$, thereby concluding the proof of Lemma V.2. \square

LEMMA V.3 (Pointwise control of the conformal factor in the neck region). *There exists a constant $\eta(m) > 0$ with the following property. Let $0 < \eta < \eta(m)$ and $0 < 4r < R < +\infty$. If $\vec{\xi}$ is any (weak) conformal immersion of*

$\Omega := B_R(0) \setminus B_r(0)$ into \mathbb{R}^m with L^2 -bounded second fundamental form, and satisfying

$$(V.28) \quad \|\nabla \vec{n}_{\vec{\xi}}\|_{L^2, \infty(\Omega)} \leq \sqrt{\eta},$$

then there exist $\alpha \in (1/2, 1)$ and $A \in \mathbb{R}$ (depending on R, r, m , and $\vec{\xi}$) such that

$$(V.29) \quad \|\lambda(x) - d \log |x| - A\|_{L^\infty(B_{\alpha R}(0) \setminus B_{\alpha^{-1}r}(0))} \leq C_m \left(\|\nabla \lambda\|_{L^2, \infty(\Omega)} + \int_{\Omega} |\nabla \vec{n}_{\vec{\xi}}|^2 dx \right),$$

where d satisfies

$$(V.30) \quad \left| 2\pi d - \int_{\partial B_r} \partial_r \lambda \, dl_{\partial B_r} \right| \leq C_m \left[\int_{B_{2r}(0) \setminus B_r(0)} |\nabla \vec{n}_{\vec{\xi}}|^2 dx + \frac{1}{\log R/r} \left(\|\nabla \lambda\|_{L^2, \infty(\Omega)} + \int_{\Omega} |\nabla \vec{n}_{\vec{\xi}}|^2 dx \right) \right],$$

and C_m depends only on m , and λ is as in Lemma V.2.

Proof. It is convenient to introduce the constant

$$\Lambda := \|\nabla \lambda\|_{L^2, \infty(\Omega)} + \int_{\Omega} |\nabla \vec{n}_{\vec{\xi}}|^2 dx.$$

We choose $\eta(m)$ smaller than the constant $\eta(m)$ given in Lemma IV.3 so that there exists a framing

$$\{\vec{e}_1, \vec{e}_2\} \in (W^{1,2}(\Omega, S^{m-1}))^2$$

with

$$(V.31) \quad \star \vec{n}_{\vec{\xi}} = \vec{e}_1 \wedge \vec{e}_2 \quad \text{and} \quad \sum_{i=1}^2 \int_{\Omega} |\nabla \vec{e}_i|^2 dx \leq C \int_{\Omega} |\nabla \vec{n}_{\vec{\xi}}|^2 dx.$$

The conformal parameter λ satisfies

$$(V.32) \quad \Delta \lambda = \nabla^\perp \vec{e}_1 \cdot \nabla \vec{e}_2 \quad \text{in } \Omega.$$

Let μ be the solution of

$$(V.33) \quad \begin{cases} \Delta \mu = \nabla^\perp \vec{e}_1 \cdot \nabla \vec{e}_2 & \text{in } \Omega, \\ \mu = 0 & \text{on } \partial\Omega. \end{cases}$$

As in the proof of Lemma V.2, Wente estimates give

$$(V.34) \quad \|\mu\|_{L^\infty(\Omega)} \leq (2\pi)^{-1} \|\nabla \vec{e}_1\|_{L^2} \|\nabla \vec{e}_2\|_{L^2} \leq C \Lambda$$

and

$$(V.35) \quad \|\nabla \mu\|_{L^2(\Omega)} \leq \sqrt{3/32\pi} \|\nabla \vec{e}_1\|_{L^2} \|\nabla \vec{e}_2\|_{L^2} \leq C \Lambda.$$

Let $\nu := \lambda - \mu$ be the harmonic function in Ω equal to λ on the boundary $\partial\Omega$. Analogously to (V.6) in the proof of Lemma V.1, there exists $\alpha \in (1/2, 1)$ with

$$(V.36) \quad \int_{\partial(B_{\alpha R} \setminus B_{\alpha^{-1}r})} |\nabla \nu| \leq r^{-1} \|\nabla \nu\|_{L^1(B_{2r} \setminus B_r)} + 2R^{-1} \|\nabla \nu\|_{L^1(B_R \setminus B_{R/2})} \\ \leq C_4 \|\nabla \nu\|_{L^{2,\infty}(\Omega)} \leq C_4 \left[\|\nabla \lambda\|_{L^{2,\infty}(\Omega)} + \|\nabla \mu\|_{L^2(\Omega)} \right] \leq C_4 (1 + C) \Lambda,$$

where C_4 is a universal constant. From (V.36) we deduce the existence of two constants a and b depending on $\vec{\xi}$ such that

$$(V.37) \quad \forall x \in \partial B_{\alpha^{-1}r}(0) \quad -C_m \Lambda \leq \nu(x) - a \leq C_m \Lambda$$

and

$$(V.38) \quad \forall x \in \partial B_{\alpha R}(0) \quad -C_m \Lambda \leq \nu(x) - b \leq C_m \Lambda,$$

where $C_m > 0$ only depends on m . Since the functions

$$\tau^\pm(x) := \frac{b-a}{\log R/r} \log \frac{|x|}{r} + a \pm C_m \Lambda$$

are harmonic, and since (V.37) and (V.38) imply

$$\tau^- \leq \nu \leq \tau^+ \quad \text{on } \partial(B_{\alpha R} \setminus B_{\alpha^{-1}r}),$$

the maximum principle yields

$$(V.39) \quad \forall x \in B_{\alpha R} \setminus B_{\alpha^{-1}r} \quad -C_m \Lambda \leq \nu(x) - \left[\frac{b-a}{\log R/r} \log \frac{|x|}{r} + a \right] \leq C_m \Lambda.$$

Let $\rho \in (r, R)$, and define $2\pi d := \int_{\partial B_\rho} \partial_\rho \nu$. Integrating by parts gives

$$0 = \int_{B_{\alpha R} \setminus B_{\alpha^{-1}r}} \Delta \nu \log |x| dx \\ = 2\pi d \log \frac{R}{r} + \int_0^{2\pi} \nu(\alpha^{-1}r, \theta) d\theta - \int_0^{2\pi} \nu(\alpha R, \theta) d\theta.$$

Hence

$$(V.40) \quad \left| d - \frac{b-a}{\log R/r} \right| \leq \frac{C_m \Lambda}{\log R/r}.$$

Combining (V.39) and (V.40) yields

$$(V.41) \quad \forall x \in B_{\alpha R} \setminus B_{\alpha^{-1}r} \quad -C_m \Lambda \leq \nu(x) - \frac{1}{2\pi} \left[\int_{\partial B_r(0)} \partial_r \nu \right] \log \frac{|x|}{r} - a \leq C_m \Lambda.$$

We next estimate the difference $\int_{\partial B_r(0)} \partial_r \mu = \int_{\partial B_r(0)} \partial_r(\nu - \lambda)$. There holds

$$(V.42) \quad \begin{aligned} 0 &= \int_{\Omega} \mu \Delta \log(|x|/R) dx = - \int_{\Omega} \nabla \mu \cdot \nabla \log(|x|/R) dx \\ &= -\log(r/R) \int_{\partial B_r} \partial_r \mu + \int_{\Omega} (\nabla^\perp \vec{e}_1 \cdot \nabla \vec{e}_2) \log(|x|/R) dx. \end{aligned}$$

As seen in the proof of Lemma IV.3, the construction of the moving frame $\{\vec{e}_1, \vec{e}_2\}$ requires the construction of an extension \hat{n} of $\vec{n}_{\vec{\xi}}$ into $B_r(0)$. Since $\{\vec{e}_1, \vec{e}_2\}$ is also a moving frame for \hat{n} in $B_R(0)$, Lemma IV.1 shows that

$$\sum_{i=1}^2 \int_{B_R(0)} |\nabla \vec{e}_i|^2 dx \leq C \int_{\Omega} |\nabla \vec{n}_{\vec{\xi}}|^2 dx \leq \Lambda.$$

Using this extension, we may also recast (V.42) in form

$$(V.43) \quad \begin{aligned} \log(R/r) \int_{\partial B_r} \partial_r \mu &= \int_{B_R} (\nabla^\perp \vec{e}_1 \cdot \nabla \vec{e}_2) \log(|x|/R) dx \\ &\quad - \int_{B_r} (\nabla^\perp \vec{e}_1 \cdot \nabla \vec{e}_2) \log(|x|/R) dx. \end{aligned}$$

Let Ψ be the solution of

$$\begin{cases} \Delta \Psi = \nabla^\perp \vec{e}_1 \cdot \nabla \vec{e}_2 & \text{in } B_R(0), \\ \Psi = 0 & \text{on } \partial B_R(0). \end{cases}$$

As in the proof of Lemma V.2, Wente estimates give

$$(V.44) \quad \|\nabla \Psi\|_{L^{2,1}(B_R(0))} \leq C_0 \|\nabla \vec{e}_1\|_{L^2(B_R(0))} \|\nabla \vec{e}_2\|_{L^2(B_R(0))} \leq C_m \Lambda$$

for some universal constant C_0 independent of R . Hence,

$$(V.45) \quad \begin{aligned} \left| \int_{B_R} (\nabla^\perp \vec{e}_1 \cdot \nabla \vec{e}_2) \log(|x|/R) dx \right| &= \left| \int_{B_R} \nabla \Psi \nabla \log(|x|/R) dx \right| \\ &\leq \|\nabla \Psi\|_{L^{2,1}(B_R(0))} \|\nabla \log(|x|/R)\|_{L^{2,\infty}(B_R(0))} \leq C_m \Lambda. \end{aligned}$$

On the other hand, Lemmas IV.3 and IV.1 yield a frame

$$\{\vec{f}_1, \vec{f}_2\} \in (W^{1,2}(B_r(0), S^{m-1}))^2$$

satisfying

$$\star \hat{n} = \vec{f}_1 \wedge \vec{f}_2$$

and

$$(V.46) \quad \sum_{i=1}^2 \int_{B_r(0)} |\nabla \vec{f}_i|^2 dx \leq C \int_{B_r(0)} |\nabla \hat{n}|^2 dx \leq C_m \int_{B_{2r}(0) \setminus B_r(0)} |\nabla \vec{n}_{\vec{\xi}}|^2 dx.$$

Clearly, there exists $\theta \in W^{1,2}(B_r, \mathbb{R})$ such that

$$e^{i\theta}(\vec{e}_1 + i\vec{e}_2) = \vec{f}_1 + i\vec{f}_2,$$

and thus

$$\vec{e}_1 \cdot \nabla \vec{e}_2 + \nabla \theta = \vec{f}_1 \cdot \nabla \vec{f}_2 \quad \text{in } B_r(0),$$

whence there holds

$$(V.47) \quad \nabla^\perp \vec{e}_1 \cdot \nabla \vec{e}_2 = \nabla^\perp \vec{f}_1 \cdot \nabla \vec{f}_2 \quad \text{in } B_r(0).$$

Let ψ be the solution of

$$\begin{cases} \Delta \psi = \nabla^\perp \vec{f}_1 \cdot \nabla \vec{f}_2 - |B_r|^{-1} \int_{B_r} \nabla^\perp \vec{f}_1 \cdot \nabla \vec{f}_2 & \text{in } B_r(0), \\ \partial_r \psi = 0 & \text{on } \partial B_r(0). \end{cases}$$

Calling again upon Wente estimates (this time with Neuman boundary data) gives the following control of the $L^{2,1}$ -norm of $\nabla \psi$:

$$(V.48) \quad \begin{aligned} \|\nabla \psi\|_{L^{2,1}(B_R(0))} &\leq C_0 \|\nabla \vec{f}_1\|_{L^2(B_R(0))} \|\nabla \vec{f}_2\|_{L^2(B_R(0))} \\ &\leq C_m \int_{B_{2r}(0) \setminus B_r(0)} |\nabla \vec{n}_\xi|^2 dx. \end{aligned}$$

Consequently, we derive

$$(V.49) \quad \begin{aligned} &\left| \int_{B_r} (\nabla^\perp \vec{e}_1 \cdot \nabla \vec{e}_2) \log(|x|/R) dx \right| \\ &\leq \log(R/r) \left| \int_{B_r} \nabla^\perp \vec{f}_1 \cdot \nabla \vec{f}_2 \right| + \left| \int_{B_r} \nabla \psi \nabla \log(|x|/R) dx \right| \\ &\leq C_m \log(R/r) \int_{B_{2r}(0) \setminus B_r(0)} |\nabla \vec{n}_\xi|^2 dx \\ &\quad + \|\nabla \psi\|_{L^{2,1}(B_r(0))} \|\nabla \log(|x|/R)\|_{L^{2,\infty}(B_R(0))} \\ &\leq C_m \log(R/r) \int_{B_{2r}(0) \setminus B_r(0)} |\nabla \vec{n}_\xi|^2 dx. \end{aligned}$$

Finally, combining (V.43), (V.45), and (V.49) gives the announced

$$(V.50) \quad \log(R/r) \left| \int_{\partial B_r} \partial_r \mu \right| \leq C_m \left[\Lambda + \log(R/r) \int_{B_{2r}(0) \setminus B_r(0)} |\nabla \vec{n}_\xi|^2 dx \right]. \quad \square$$

VI. Uniform $L^{2,1}$ -control of the mean curvature vector of a Willmore immersion in neck regions

The goal of this section is to prove the following result.

LEMMA VI.1. *Let $m \geq 3$. There exists a constant $\varepsilon(m) > 0$ with the following property. Let $0 < 8r < R$, and let $\vec{\xi}$ be any conformal Willmore immersion of $B_R(0)$ into \mathbb{R}^m such that*

$$(VI.1) \quad \sup_{r < s < R/2} \int_{B_{2s}(0) \setminus B_s(0)} |\nabla \vec{n}_\xi|^2 dx \leq \varepsilon(m).$$

We set

$$\Lambda := \|\nabla\lambda\|_{L^{2,\infty}(B_R \setminus B_r)} + \int_{B_R(0)} |\nabla\vec{n}_\xi|^2 dx,$$

where λ denotes as before the conformal parameter of $\vec{\xi}$. Then there holds

$$(VI.2) \quad \|e^\lambda \vec{H}_\xi\|_{L^{2,1}(B_{R/2} \setminus B_{2r})} \leq C(m, \Lambda),$$

where \vec{H}_ξ is the mean curvature vector of the immersion $\vec{\xi}$ and $C(m, \Lambda)$ is a positive constant depending¹¹ only on m and Λ .

Proof of Lemma VI.1. According to the work in [Riv08], an immersion $\vec{\xi}$ is conformal and Willmore on $B_R(0)$ if and only if there exists an \mathbb{R}^m -valued map \vec{L} (uniquely defined up to an additive constant vector) satisfying

$$(VI.3) \quad \nabla\vec{L} = 2\nabla^\perp \vec{H} - 3\nabla^\perp(\pi_{\vec{n}})(\vec{H}) + \star(\nabla\vec{n}_\xi \wedge \vec{H}),$$

where \star is the canonical Hodge operation on multivectors of \mathbb{R}^m and $\pi_{\vec{n}}$ is the projection operator onto the normal space to $\vec{\xi}_*T\Sigma$. It is given by two consecutive applications of the contraction operator¹² \lrcorner with \vec{n} (see (VI.87) in [Riv13b]):

$$\forall \vec{w} \in \mathbb{R}^m \quad \pi_{\vec{n}}(\vec{w}) := (-1)^{m-1} \vec{n} \lrcorner (\vec{n} \lrcorner \vec{w}).$$

Hence the gradient of $\pi_{\vec{n}}$ in (VI.3) is to be understood as

$$(VI.4) \quad \forall \vec{w} \in \mathbb{R}^m \quad \nabla(\pi_{\vec{n}})(\vec{w}) := (-1)^{m-1} (\nabla\vec{n}) \lrcorner (\vec{n} \lrcorner \vec{w}) + (-1)^{m-1} \vec{n} \lrcorner ((\nabla\vec{n}) \lrcorner \vec{w}).$$

The first condition to impose on the constant $\varepsilon(m)$ is that it should be smaller than the one in Lemma V.2 so that the *uniform Harnack estimate* (V.14) holds on the annulus $B_{\alpha R} \setminus B_{\alpha^{-1}r}$ for every $0 < \alpha < 1$. If, in addition, we ensure that $\varepsilon(m)$ be smaller than the threshold constant appearing the ϵ -regularity Theorem I.5 in [Riv08], then the uniform Harnack estimate enables¹³ us to conclude that for any $x \in B_{4R/5}(0) \setminus B_{5r/4}(0)$, there holds

$$(VI.5) \quad |\nabla\vec{n}_\xi(x)|^2 \leq C|x|^{-2} \int_{B_{2|x|}(0) \setminus B_{|x|/2}(0)} |\nabla\vec{n}_\xi|^2 dx \leq C\varepsilon(m)|x|^{-2},$$

¹¹The constant $C(m, \Lambda)$ is independent of the conformal type of the annulus.

¹²The contraction operator \lrcorner is the linear operation defined as follows. For every α, β , and γ , respectively p, q , and $(p - q)$ vectors in \mathbb{R}^m , one has

$$\langle \alpha \lrcorner \beta, \gamma \rangle = \langle \alpha, \beta \wedge \gamma \rangle.$$

¹³Indeed, the proof of Theorem I.5 in [Riv08] requires that the conformal parameter λ be “roughly” constant on balls around the origin. We are instead working on annuli, so the estimate (V.14) is exactly what is needed.

where C is some constant independent of the data of the problem. This implies, in particular, that

$$(VI.6) \quad \|\nabla \vec{n}_{\xi'}\|_{L^{2,\infty}(B_{4R/5}(0) \setminus B_{5r/4}(0))} \leq \sqrt{\frac{15\pi}{16}} C \varepsilon(m) = C' \sqrt{\varepsilon(m)}.$$

Finally, we ask that $\varepsilon(m) > 0$ satisfy $C' \sqrt{\varepsilon(m)} \leq \sqrt{\eta(m)}$, where $\eta(m)$ is the constant given in Lemma V.3. Accordingly, there exist a constant A (which depends *a priori* on all parameters r, R, Λ , and ξ) and a constant d such that

$$(VI.7) \quad |d| \leq C_m \Lambda$$

and

$$(VI.8) \quad \|\lambda(x) - d \log|x| - A\|_{L^\infty(B_{R/2} \setminus B_{2r})} \leq C_m \Lambda.$$

We introduce for $0 < s < R/2$ the function

$$\delta(s) := \left(s^{-2} \int_{B_{2s}(0) \setminus B_{s/2}(0)} |\nabla \vec{n}_{\xi'}|^2 dx \right)^{1/2}.$$

Clearly,

$$(VI.9) \quad s \delta(s) \leq \sqrt{\Lambda} \quad \forall s \in (0, R/2).$$

Inequality (VI.5) gives

$$(VI.10) \quad e^{\lambda(x)} |\vec{H}_{\xi'}(x)| \leq |\nabla \vec{n}_{\xi'}(x)| \leq C \delta(|x|) \quad \forall x \in B_{R/2}(0) \setminus B_{2r}(0).$$

Clearly, since $8r < R$, there holds

$$(VI.11) \quad \int_r^{R/2} s \delta^2(s) ds \leq \log(4) \int_{B_R(0) \setminus B_{r/2}(0)} |\nabla \vec{n}_{\xi'}|^2 dx \leq 2\Lambda.$$

The identity (VI.3) shows that on any dyadic annulus $B_{2s} \setminus B_s$ included in $B_R(0) \setminus B_r(0)$, the mean curvature vector satisfies

$$\Delta \vec{H} = \frac{1}{2} \operatorname{div} \left(3\nabla(\pi_{\vec{n}})(\vec{H}) + \star(\nabla^\perp \vec{n}_{\xi'} \wedge \vec{H}) \right).$$

From (VI.10), a standard argument from the theory of second-order uniformly elliptic equations in divergence form then yields the existence of a constant $C_1 > 0$ such that

$$(VI.12) \quad e^{\lambda(x)} |\nabla \vec{H}_{\xi'}(x)| \leq C_1 \frac{\delta(|x|)}{|x|} \quad \forall x \in B_{R/2}(0) \setminus B_{2r}(0).$$

For any $t \in (r, R)$, we denote

$$\vec{L}_t := \frac{1}{|\partial B_t(0)|} \int_{\partial B_t(0)} \vec{L},$$

where the function \vec{L} is as (VI.10). Using (VI.3), (VI.8), and (VI.12), there holds

$$(VI.13) \quad |\vec{L}(x) - \vec{L}_{|x|}| \leq \int_{\partial B_{|x|}} |\nabla \vec{L}| \leq 2\pi C_1 e^{2C_m \Lambda} e^{-\lambda(x)} \delta(|x|) \quad \forall x \in B_{R/2}(0) \setminus B_{2r}(0).$$

This implies, in particular, that

$$(VI.14) \quad \int_{B_{R/2}(0) \setminus B_{2r}(0)} e^{2\lambda(x)} |\vec{L}(x) - \vec{L}_{|x|}|^2 dx \leq C_2 \Lambda$$

for some constant C_2 . Note that we have used (VI.11). We next have

$$(VI.15) \quad \frac{d\vec{L}_t}{dt} = \frac{1}{2\pi} \int_0^{2\pi} \frac{\partial \vec{L}}{\partial t}(t, \theta) d\theta = \frac{3}{2\pi} \int_0^{2\pi} \frac{1}{r} \frac{\partial \pi_{\vec{n}}}{\partial \theta}(\vec{H}) d\theta + \frac{1}{2\pi} \int_0^{2\pi} \star \left(\frac{\partial \vec{n}_{\xi}}{\partial \nu} \wedge \vec{H} \right) d\theta.$$

Setting $a(t) := |\vec{L}_t|$ and putting (VI.10) into the latter gives the estimate

$$(VI.16) \quad |\dot{a}(t)| = \left| \frac{d|\vec{L}_t|}{dt} \right| = \left| \frac{d\vec{L}_t}{dt} \right| \leq C^2 e^{-\lambda} \delta^2(t)$$

whence, using (VI.11), we reach

$$(VI.17) \quad \int_{2r}^{R/2} s e^{\lambda} |\dot{a}(s)| ds \leq 2C^2 \Lambda.$$

The Harnack estimate (VI.8) yields

$$(VI.18) \quad e^{A-C_m \Lambda} |x|^d \leq e^{\lambda(x)} \leq e^{A+C_m \Lambda} |x|^d \quad \forall x \in B_{R/2} \setminus B_{2r}(0),$$

thereby showing that (VI.17) may be recast in the form

$$(VI.19) \quad \int_{2r}^{R/2} s^{1+d} |\dot{a}(s)| ds \leq 2C^2 e^{C_m \Lambda - A} \Lambda.$$

An elementary integration by parts gives, for any $r \leq \tau < T \leq R$,

$$\int_{\tau}^T s^{1+d} \dot{a}(s) ds = T^{1+d} a(T) - \tau^{1+d} a(\tau) - (1+d) \int_{\tau}^T s^d a(s) ds.$$

Hence, since $a \geq 0$, we have

$$(VI.20) \quad \begin{cases} \forall d \leq -1 \quad \forall 2r < t < R/2 \quad t^{1+d} a(t) \leq (2r)^{1+d} a(2r) + \int_{2r}^{R/2} s^{1+d} |\dot{a}(s)| ds, \\ \forall d \geq -1 \quad \forall 2r < t < R/2 \quad t^{1+d} a(t) \leq (R/2)^{1+d} a(R/2) + \int_{2r}^{R/2} s^{1+d} |\dot{a}(s)| ds. \end{cases}$$

Recall that we still have the freedom to adjust the vector \vec{L} by an additive constant vector. It will be convenient to us to choose \vec{L} in such a way that

$$\text{If } d \leq -1, \text{ we take } \int_{\partial B_{2r}} \vec{L} = 0 \quad \text{whereas if } d \geq -1, \text{ we take } \int_{\partial B_{R/2}} \vec{L} = 0.$$

This particular choice implies

$$(VI.21) \quad t e^\lambda |\vec{L}_t| \equiv t e^\lambda a(t) \leq 2 C^2 e^{2C_m \Lambda} \Lambda \quad \forall 2r < t < R/2,$$

where we have used (VI.20), (VI.18), and (VI.19). With the help of an elementary computation, the latter then implies that

$$(VI.22) \quad \|e^{\lambda(x)} |\vec{L}_{|x|}|\|_{L^{2,\infty}(B_{R/2} \setminus B_{2r})} \leq C_0(m, \Lambda)$$

for some positive constant $C_0(m, \Lambda)$. Combining this inequality with (VI.14) gives

$$(VI.23) \quad \|e^{\lambda(x)} \vec{L}(x)\|_{L^{2,\infty}(B_{R/2} \setminus B_{2r})} \leq C(m, \Lambda),$$

where $C(m, \Lambda)$ depends only on m and Λ . Moreover, the estimates (VI.13) and (VI.21) altogether yield

$$(VI.24) \quad \begin{aligned} e^{\lambda(x)} |\vec{L}(x)| &\leq C(\Lambda, C_m) [1 + |x|\delta(x)] |x|^{-1} \\ &\leq C'(m, \Lambda) |x|^{-1} \quad \forall x \in B_{R/2}(0) \setminus B_{2r}(0). \end{aligned}$$

Note that we have also used (VI.9).

As is done in [Riv08] (see also Section VI.7.2 of [Riv13b]), we introduce on $B_R(0)$ the smooth function S and the smooth $\wedge^2(\mathbb{R}^m)$ -valued map \vec{R} via

$$(VI.25) \quad \begin{cases} \nabla S = \vec{L} \cdot \nabla \vec{\xi}, \\ \nabla \vec{R} = \vec{L} \wedge \nabla \vec{\xi} + 2 (\star(\vec{n}_\xi \lrcorner \vec{H})) \lrcorner \nabla \vec{\xi}. \end{cases}$$

The existence of S and \vec{R} have been proved in [Ber13] to be consequences of applications of Noether theorem.

Since $|\nabla \vec{\xi}|^2 = 2e^{2\lambda}$, the estimates (VI.6) and (VI.23) give the bound

$$(VI.26) \quad \|\nabla S\|_{L^{2,\infty}(B_{R/2} \setminus B_{2r})} + \|\nabla \vec{R}\|_{L^{2,\infty}(B_{R/2} \setminus B_{2r})} \leq C(m, \Lambda).$$

Furthermore, using (VI.10), (VI.24), and (VI.9), we obtain the pointwise bound

$$(VI.27) \quad |\nabla S(x)| + |\nabla \vec{R}(x)| \leq C(m, \Lambda) |x|^{-1} \quad \forall x \in B_{R/2} \setminus B_{2r}.$$

One verifies (cf. [Riv08] and Theorem VI.15 of [Riv13b]) that the following equations¹⁴ hold:

$$(VI.28) \quad \begin{cases} \nabla S = -(\star \vec{n}) \cdot \nabla^\perp \vec{R}, \\ \nabla \vec{R} = (-1)^m \star (\vec{n} \bullet \nabla^\perp \vec{R}) + (\star \vec{n}) \nabla^\perp S. \end{cases}$$

For any $t \in (2r, R/2)$, we let

$$S_t := \frac{1}{|\partial B_t(0)|} \int_{\partial B_t(0)} S, \quad \vec{R}_t := \frac{1}{|\partial B_t(0)|} \int_{\partial B_t(0)} \vec{R}, \quad \vec{n}_t := \frac{1}{|\partial B_t(0)|} \int_{\partial B_t(0)} \vec{n}.$$

The equations (VI.28) then yield

$$(VI.29) \quad \begin{cases} \frac{dS_t}{dt} = \frac{1}{2\pi} \int_0^{2\pi} \frac{\partial S}{\partial t}(t, \theta) d\theta = \frac{1}{2\pi} \int_0^{2\pi} \star (\vec{n} - \vec{n}_t) \cdot \frac{1}{t} \frac{\partial \vec{R}}{\partial \theta} d\theta, \\ \frac{d\vec{R}_t}{dt} = \frac{1}{2\pi} \int_0^{2\pi} \frac{\partial \vec{R}}{\partial t}(t, \theta) d\theta \\ \quad = (-1)^{m-1} \frac{1}{2\pi} \int_0^{2\pi} (\vec{n} - \vec{n}_t) \bullet \frac{1}{t} \frac{\partial \vec{R}}{\partial \theta} d\theta - \frac{1}{2\pi} \int_0^{2\pi} \star (\vec{n} - \vec{n}_t) \frac{1}{t} \frac{\partial S}{\partial \theta} d\theta. \end{cases}$$

Note that (VI.10) gives

$$(VI.30) \quad |\vec{n}(x) - \vec{n}_{|x|}| \leq C|x|\delta(|x|) \quad \forall x \in B_{R/2}(0) \setminus B_{2r}(0).$$

Thus, (VI.27), (VI.29), and (VI.30) altogether yield

$$(VI.31) \quad \left| \frac{dS_t}{dt} \right| + \left| \frac{d\vec{R}_t}{dt} \right| \leq C(m, \Lambda) \delta(t).$$

Hence, with the help of (VI.11), we deduce

$$(VI.32) \quad \int_{2r}^{R/2} \left[\left| \frac{dS_t}{dt} \right|^2 + \left| \frac{d\vec{R}_t}{dt} \right|^2 \right] t dt \leq C(m, \Lambda).$$

Our study requires the following result, proved in [LR].

¹⁴The linear operator \bullet is the contraction that to a pair of p and q -vectors of \mathbb{R}^m assigns a $(p+q-2)$ -vector of \mathbb{R}^m in such a way that

$$\forall \vec{a} \in \wedge^p \mathbb{R}^m \quad \forall \vec{b} \in \wedge^1 \mathbb{R}^m \quad \vec{a} \bullet \vec{b} := \vec{a} \lrcorner \vec{b}$$

and

$$\begin{aligned} \forall \vec{a} \in \wedge^p \mathbb{R}^m \quad \forall \vec{b} \in \wedge^r \mathbb{R}^m \quad \forall \vec{c} \in \wedge^s \mathbb{R}^m \\ \vec{a} \bullet (\vec{b} \wedge \vec{c}) := (\vec{a} \bullet \vec{b}) \wedge \vec{c} + (-1)^{r \cdot s} (\vec{a} \bullet \vec{c}) \wedge \vec{b}. \end{aligned}$$

LEMMA VI.2 ([LR]). *Let a and b be two functions on $B_1(0)$ such that $\nabla a \in L^{2,\infty}$ and $\nabla b \in L^2$. Let $0 < \varepsilon < 1/4$ and ϕ satisfy*

$$-\Delta\phi = \partial_{x_1}a \partial_{x_2}b - \partial_{x_1}b \partial_{x_2}a \quad \text{in } B_1(0) \setminus B_\varepsilon(0).$$

For $\varepsilon \leq r \leq 1$, we set $\phi_0(r) := (2\pi r)^{-1} \int_{\partial B_r(0)} \phi$, and we assume that

$$(VI.33) \quad \int_\varepsilon^1 |\dot{\phi}_0|^2 r \, dr < +\infty.$$

Then $\nabla\phi \in L^2(B_{1/2}(0) \setminus B_{2\varepsilon}(0))$ and there exists a positive constant C independent of ε and ϕ such that

$$(VI.34) \quad \begin{aligned} \|\nabla\phi\|_{L^2(B_{1/2}(0) \setminus B_{2\varepsilon}(0))} &\leq C \|\nabla a\|_{L^{2,\infty}(B_1(0))} \|\nabla b\|_{L^2(B_1(0))} \\ &\quad + C \|\nabla\phi_0\|_{L^2(B_1(0) \setminus B_\varepsilon(0))} + C \|\nabla\phi\|_{L^{2,\infty}(B_1(0) \setminus B_\varepsilon(0))}. \end{aligned}$$

Applying divergence to both sides of equations (VI.28) gives the *conservative conformal Willmore system*:

$$(VI.35) \quad \begin{cases} \Delta S = -\star \nabla \vec{n} \cdot \nabla^\perp \vec{R}, \\ \Delta \vec{R} = (-1)^m \star (\nabla \vec{n} \bullet \nabla^\perp \vec{R}) + \star \nabla \vec{n} \cdot \nabla^\perp S. \end{cases}$$

Owing to (VI.1), (VI.26), and (VI.32), we may apply Lemma VI.2 to the system (VI.35) and find

$$(VI.36) \quad \|\nabla S\|_{L^2(B_{R/4}(0) \setminus B_{4r}(0))} + \|\nabla \vec{R}\|_{L^2(B_{R/4}(0) \setminus B_{4r}(0))} \leq C(m, \Lambda).$$

Since the L^2 -norm of $|x|^{-1}$ on annuli of the form $B_{as}(0) \setminus B_{bs}(0)$ is independent of s , we can use the pointwise estimate (VI.27) to get an upper bound independent of R and r for

$$\|\nabla S\|_{L^2(B_{R/2}(0) \setminus B_{R/4}(0) \cup B_{4r}(0) \setminus B_{2r}(0))} + \|\nabla \vec{R}\|_{L^2(B_{R/2}(0) \setminus B_{R/4}(0) \cup B_{4r}(0) \setminus B_{2r}(0))}.$$

Combining it to (VI.36) then yields

$$(VI.37) \quad \|\nabla S\|_{L^2(B_{R/2}(0) \setminus B_{2r}(0))} + \|\nabla \vec{R}\|_{L^2(B_{R/2}(0) \setminus B_{2r}(0))} \leq C(m, \Lambda).$$

This new information, along with (VI.11), is injected into (VI.29) so as to produce

$$(VI.38) \quad \begin{aligned} \int_{2r}^{R/2} \left| \frac{dS_t}{dt} \right| + \left| \frac{d\vec{R}_t}{dt} \right| dt &\leq C \int_{2r}^{R/2} \delta(t) \int_{\partial B_t} |\nabla S| + |\nabla \vec{R}| \, dt \\ &\leq C \left(\int_{2r}^{R/2} \delta^2(t) t \, dt \right)^{1/2} \left(\int_{B_{R/2} \setminus B_{2r}} |\nabla S|^2 + |\nabla \vec{R}|^2 \, dx \right)^{1/2} \leq C(m, \Lambda). \end{aligned}$$

Since the functions S and \vec{R} are both defined up to an additive ‘‘constant,’’ we have the freedom to impose the conditions $S_{2r} = 0$ and $\vec{R}_{2r} = \vec{0}$. From (VI.31)

we deduce, in particular, that

$$(VI.39) \quad |S_t| + |\vec{R}_t| \leq C(m, \Lambda) \quad \forall t \in (2r, R/2).$$

Paired to the pointwise estimate (VI.27) on the gradient of S and \vec{R} , the latter implies

$$(VI.40) \quad \|S\|_{L^\infty(B_{R/2}(0) \setminus B_{2r}(0))} + \|\vec{R}\|_{L^\infty(B_{R/2}(0) \setminus B_{2r}(0))} \leq C(m, \Lambda).$$

We are sufficiently geared to apply the following general result, whose proof may be found in [LR].

LEMMA VI.3 ([LR]). *Let a and b be two functions on $B_1(0)$ such that $\nabla a \in L^2$ and $\nabla b \in L^2$. Let $0 < \epsilon < 1/4$ and ϕ satisfy*

$$-\Delta \phi = \partial_{x_1} a \partial_{x_2} b - \partial_{x_1} b \partial_{x_2} a \quad \text{in } B_1(0) \setminus B_\epsilon(0).$$

Assume that

$$(VI.41) \quad \|\phi\|_\infty < +\infty.$$

Then $\nabla \phi \in L^{2,1}(B_{1/2}(0) \setminus B_{2\epsilon}(0))$, and there exists a positive constant C independent of ϵ and ϕ such that

$$(VI.42) \quad \|\nabla \phi\|_{L^{2,1}(B_{1/2}(0) \setminus B_{2\epsilon}(0))} \leq C \|\nabla a\|_2 \|\nabla b\|_2 + C \|\phi\|_{L^\infty(B_1(0) \setminus B_\epsilon(0))} \\ + C \|\nabla \phi\|_{L^2(B_1(0) \setminus B_\epsilon(0))}.$$

Just as we did above, owing to (VI.1), (VI.26), and (VI.40), we apply Lemma VI.3 to the system (VI.35) and find

$$(VI.43) \quad \|\nabla S\|_{L^{2,1}(B_{R/4}(0) \setminus B_{4r}(0))} + \|\nabla \vec{R}\|_{L^{2,1}(B_{R/4}(0) \setminus B_{4r}(0))} \leq C(m, \Lambda).$$

Since the $L^{2,1}$ -norm of $|x|^{-1}$ on annuli of the form $B_{as}(0) \setminus B_{bs}(0)$ is independent of s , we use (VI.27) to get an upper bound independent of R and r for

$$\|\nabla S\|_{L^{2,1}(B_{R/2}(0) \setminus B_{R/4}(0) \cup B_{4r}(0) \setminus B_{2r}(0))} + \|\nabla \vec{R}\|_{L^{2,1}(B_{R/2}(0) \setminus B_{R/4}(0) \cup B_{4r}(0) \setminus B_{2r}(0))}.$$

Combining it to (VI.43) then yields

$$(VI.44) \quad \|\nabla S\|_{L^{2,1}(B_{R/2}(0) \setminus B_{2r}(0))} + \|\nabla \vec{R}\|_{L^{2,1}(B_{R/2}(0) \setminus B_{2r}(0))} \leq C(m, \Lambda).$$

It is shown in [Riv08] that

$$4e^{2\lambda} \vec{H}_{\vec{\xi}} \equiv 2\Delta \vec{\xi} = \nabla^\perp S \cdot \nabla \vec{\xi} - \nabla \vec{R} \lrcorner \nabla^\perp \vec{\xi}.$$

In particular, since $|\nabla \vec{\xi}|^2 = 2e^\lambda$, calling upon (VI.44) in the latter yields the announced (VI.2), thereby completing the proof. \square

VII. L^2 -weak energy quantization for the Gauss map of a Willmore immersion in neck regions

LEMMA VII.1 (L^2 -weak energy quantization for the Gauss map in neck regions). *There exists a constant $\varepsilon(m) > 0$ with the following property. Let $\vec{\xi}_k$ be a sequence of conformal Willmore immersions from $B_{R_k}(0)$ into \mathbb{R}^m , with $R_k \rightarrow +\infty$. Whenever $r_k \rightarrow 0$ satisfies*

$$(VII.1) \quad \sup_{r \in (r_k, R_k/2)} \int_{B_{2r}(0) \setminus B_r(0)} |\nabla \vec{n}_{\vec{\xi}_k}|^2 dx \leq \varepsilon(m)$$

and¹⁵

$$(VII.2) \quad \|\nabla \lambda_k\|_{L^2, \infty(\Omega_k)} + \int_{\Omega_k} |\nabla \vec{n}_{\vec{\xi}_k}|^2 dx \leq \Lambda < +\infty \quad \forall k,$$

where $\Lambda > 0$ is independent of k and $\Omega_k = B_{R_k}(0) \setminus B_{r_k}(0)$, then

$$(VII.3) \quad \forall \varepsilon \in (0, \varepsilon(m)) \exists \alpha \in (0, 1) \text{ s.t. } \limsup_{k \rightarrow +\infty} \||x| \nabla \vec{n}_{\vec{\xi}_k}(x)\|_{L^\infty(B_{\alpha R_k/2}(0) \setminus B_{\alpha^{-1} r_k}(0))} \leq \varepsilon.$$

Hence, in particular,

$$(VII.4) \quad \forall \varepsilon \in (0, \varepsilon(m)) \exists \alpha \in (0, 1) \text{ s.t. } \limsup_{k \rightarrow +\infty} \|\nabla \vec{n}_{\vec{\xi}_k}\|_{L^2, \infty(B_{\alpha R_k/2}(0) \setminus B_{\alpha^{-1} r_k}(0))} \leq \varepsilon.$$

Proof. Analogously to the argument given in the paragraph following (VI.4) in the proof of Lemma VI.1, the constant $\varepsilon(m)$ may be chosen so as to ensure that there holds

$$(VII.5) \quad \begin{aligned} |\nabla \vec{n}_{\vec{\xi}_k}(x)|^2 &\leq C |x|^{-2} \int_{B_{2|x|}(0) \setminus B_{|x|/2}(0)} |\nabla \vec{n}_{\vec{\xi}_k}|^2 dx \\ &\leq C \varepsilon(m) |x|^{-2} \quad \forall x \in B_{R_k/2}(0) \setminus B_{2r_k}(0), \end{aligned}$$

where C is some constant independent of the data of the problem. This implies, in particular, that

$$(VII.6) \quad \|\nabla \vec{n}_{\vec{\xi}_k}\|_{L^2, \infty(B_{R_k/2}(0) \setminus B_{2r_k}(0))} \leq C' \sqrt{\varepsilon(m)}.$$

We will argue by contradiction, by assuming that there exists a sequence of conformal Willmore immersions on $B_{R_k}(0)$ satisfying (VII.1) and (VII.2), but for which there exist $\varepsilon_1 > 0$ and $x_k \in \Omega_k$ with

$$(VII.7) \quad \log \left| \frac{|x_k|}{r_k} \right| \rightarrow +\infty, \quad \log \left| \frac{|x_k|}{R_k} \right| \rightarrow -\infty$$

and with

$$(VII.8) \quad |x_k| |\nabla \vec{n}_{\vec{\xi}_k}(x_k)| \geq \varepsilon_1 > 0.$$

¹⁵As usual, λ_k is the conformal parameter of $\vec{\xi}_k$.

From (VII.5) we deduce, in particular, that

$$(VII.9) \quad \int_{B_{2|x_k|}(0) \setminus B_{|x_k|/2}(0)} |\nabla \vec{n}_{\vec{\xi}_k}|^2 dx \geq \frac{\varepsilon_1^2}{C} > 0.$$

We next demand that $\varepsilon(m)$ be smaller than its counterpart from Lemma VI.1. Since all hypotheses of this lemma are fulfilled, we have the existence of \vec{L}_k in $C^\infty(B_{R_k}(0), \mathbb{R}^m)$, S_k in $C^\infty(B_{R_k}(0), \mathbb{R})$, and \vec{R}_k in $C^\infty(B_{R_k}(0), \wedge^2 \mathbb{R}^m)$ such that

$$(VII.10) \quad \begin{cases} \nabla^\perp \vec{L}_k := \nabla \vec{H}_k - 3\pi \vec{n}_{\vec{\xi}_k} (\nabla \vec{H}_k) + \star (\nabla^\perp \vec{n}_{\vec{\xi}_k} \wedge \vec{H}_k), \\ \nabla S_k := \vec{L}_k \cdot \nabla \vec{\xi}_k, \\ \nabla \vec{R}_k := \vec{L}_k \wedge \nabla \vec{\xi}_k + 2\vec{H}_k \wedge \nabla^\perp \vec{\xi}_k. \end{cases}$$

In the course of the proof of Lemma VI.1, we have seen that

$$(VII.11) \quad \|e^{\lambda_k} \vec{L}_{\vec{\xi}_k}\|_{L^{2,\infty}(B_{R_k/2} \setminus B_{2r_k})} + \|\nabla S_k\|_{L^{2,1}(B_{R_k/2} \setminus B_{2r_k})} + \|\nabla \vec{R}_k\|_{L^{2,1}(B_{R_k/2} \setminus B_{2r_k})} \leq C(m, \Lambda).$$

Moreover, we have the conservative conformal Willmore system

$$(VII.12) \quad \begin{cases} \Delta S_k = -(\star \nabla \vec{n}_{\vec{\xi}_k}) \cdot \nabla^\perp \vec{R}_k, \\ \Delta \vec{R}_k = (-1)^m \star (\nabla \vec{n}_{\vec{\xi}_k} \bullet \nabla^\perp \vec{R}_k) + (\star \nabla \vec{n}_{\vec{\xi}_k}) \cdot \nabla^\perp S_k, \\ 2\Delta \vec{\xi}_k = \nabla^\perp S_k \cdot \nabla \vec{\xi}_k - \nabla \vec{R}_k \lrcorner \nabla^\perp \vec{\xi}_k. \end{cases}$$

Consider the conformal mapping

$$\vec{\xi}_k(y) := e^{-\lambda_k(x_k)} (\vec{\xi}_k(|x_k|y) - \vec{\xi}_k(x_k)).$$

An elementary computation shows that

$$\nabla_y \vec{n}_{\vec{\xi}_k}(y) = |x_k| \nabla_x \vec{n}_{\vec{\xi}_k}(|x_k|y)$$

and

$$\nabla_y \vec{H}_k(y) = |x_k| e^{-\lambda_k(x_k)} \nabla_x \vec{H}_k(|x_k|y),$$

where \vec{H}_k is the mean curvature vector of the immersion $\vec{\xi}_k$. The corresponding \mathbb{R}^m -valued map \vec{L}_k satisfies

$$\nabla_y \vec{L}_k(y) = |x_k| e^{-\lambda_k(x_k)} \nabla_x \vec{L}_k(|x_k|y),$$

so that

$$\vec{L}_k(y) = e^{-\lambda_k(x_k)} \vec{L}_k(|x_k|y).$$

Clearly, the conformal factor of the rescaled immersion satisfies

$$\tilde{\lambda}_k(y) = \lambda_k(|x_k|y) - \lambda_k(x_k).$$

This implies, in particular, that the corresponding \tilde{S}_k and \tilde{R}_k are given by

$$\tilde{S}_k(y) = S_k(|x_k|y) \quad \text{and} \quad \tilde{R}_k(y) = \vec{R}_k(|x_k|y).$$

We thus see that the maps S_k and \vec{R}_k behave under rescaling like the Gauss map $\vec{n}_{\vec{\xi}}$. This remarkable fact was first observed in [BR13], where it played a decisive role. From the latter and (VII.11), it follows that

$$(VII.13) \quad \limsup_{k \rightarrow +\infty} \|\nabla \tilde{S}_k\|_{L^{2,1}(K)} + \|\nabla \tilde{R}_k\|_{L^{2,1}(K)} \leq C(m, \Lambda)$$

for any compact subdomain K of $\mathbb{C} \setminus \{0\}$.

The pointwise control of the conformal factor in neck regions provided by Lemma V.3 shows that the sequence of Willmore immersions $\tilde{\xi}_k$ satisfies

$$(VII.14) \quad \limsup_{k \rightarrow +\infty} \|\nabla \vec{n}_{\tilde{\xi}_k}\|_{L_{\text{loc}}^\infty(\mathbb{C} \setminus \{0\})} < +\infty$$

and

$$(VII.15) \quad \limsup_{k \rightarrow +\infty} \|\log |\nabla \tilde{\xi}_k|\|_{L_{\text{loc}}^\infty(\mathbb{C} \setminus \{0\})} < +\infty.$$

From the work in [Riv08], we conclude that for all $l \in \mathbb{N}$, the sequence $\tilde{\xi}_k$ converges in $C_{\text{loc}}^l(\mathbb{C} \setminus \{0\})$ to a Willmore immersion $\tilde{\xi}_\infty$ of $\mathbb{C} \setminus \{0\}$. This strong convergence implies that condition (VII.9) and the system (VII.12) pass to the limit. Hence,

$$(VII.16) \quad \int_{B_2(0) \setminus B_{1/2}(0)} |\nabla \vec{n}_{\tilde{\xi}_\infty}|^2 dx \geq \frac{\varepsilon_1^2}{C} > 0$$

and

$$(VII.17) \quad \begin{cases} \Delta \tilde{S}_\infty = -(\star \nabla \vec{n}_{\tilde{\xi}_\infty}) \cdot \nabla^\perp \tilde{R}_\infty, \\ \Delta \tilde{R}_\infty = (-1)^m \star (\nabla \vec{n}_{\tilde{\xi}_\infty} \bullet \nabla^\perp \tilde{R}_\infty) + (\star \nabla \vec{n}_{\tilde{\xi}_\infty}) \cdot \nabla^\perp \tilde{S}_\infty, \\ 4e^{2\tilde{\lambda}_\infty} \vec{H}_{\tilde{\xi}_\infty} = 2\Delta \tilde{\xi}_\infty = \nabla^\perp \tilde{S}_\infty \cdot \nabla \tilde{\xi}_\infty - \nabla \tilde{R}_\infty \lrcorner \nabla^\perp \tilde{\xi}_\infty. \end{cases}$$

On one hand, we obtain from (VII.6) that

$$(VII.18) \quad \|\nabla \vec{n}_{\tilde{\xi}_\infty}\|_{L^{2,\infty}(\mathbb{C})} \leq C \sqrt{\varepsilon(m)}.$$

On the other hand, it follows from (VII.13) that

$$(VII.19) \quad \|\nabla \tilde{S}_\infty\|_{L^{2,1}(\mathbb{C})} + \|\nabla \tilde{R}_\infty\|_{L^{2,1}(\mathbb{C})} \leq C(m, \Lambda) < +\infty.$$

Applying to the system (VII.17) the Wente inequality from Lemma IV.2 with $(p, q) = (2, 1)$ now yields

$$(VII.20) \quad \begin{aligned} & \|\nabla \tilde{S}_\infty\|_{L^{2,1}(\mathbb{C})} + \|\nabla \tilde{R}_\infty\|_{L^{2,1}(\mathbb{C})} \\ & \leq C \|\nabla \vec{n}_{\tilde{\xi}_\infty}\|_{L^{2,\infty}(\mathbb{C})} \left[\|\nabla \tilde{S}_\infty\|_{L^{2,1}(\mathbb{C})} + \|\nabla \tilde{R}_\infty\|_{L^{2,1}(\mathbb{C})} \right] \\ & \leq C_m \sqrt{\varepsilon(m)} \left[\|\nabla \tilde{S}_\infty\|_{L^{2,1}(\mathbb{C})} + \|\nabla \tilde{R}_\infty\|_{L^{2,1}(\mathbb{C})} \right], \end{aligned}$$

where we have used (VII.18). Accordingly, when $C_m \sqrt{\varepsilon(m)} < 1$, we deduce that $\nabla \tilde{S}_\infty = 0$ and $\nabla \tilde{R}_\infty = 0$. Injecting this information in the last identity of (VII.17) gives

$$\vec{H}_{\tilde{\xi}_\infty} \equiv \vec{0}.$$

In particular, since the (weighted) mean curvature vector is the trace part of the gradient of the Gauss map, it follows easily that

$$(VII.21) \quad |\nabla \vec{n}_{\tilde{\xi}_\infty}|^2 = -2 e^{2\tilde{\lambda}_\infty} K_{\tilde{\xi}_\infty} \quad \text{on } \mathbb{C},$$

where $K_{\tilde{\xi}_\infty}$ is the Gauss curvature of the limit immersion $\tilde{\xi}_\infty$.

Let us next choose the constant $\varepsilon(m)$ small enough that we can apply Lemma IV.3 with condition (VII.18). Namely, on any ball $B_\rho(0)$, we construct a moving frame $\{\vec{e}_1^\rho, \vec{e}_2^\rho\}$ satisfying

$$\star \hat{n}_{\tilde{\xi}_\infty} = \vec{e}_1^\rho \wedge \vec{e}_2^\rho, \quad \text{div}(\vec{e}_1^\rho \cdot \nabla \vec{e}_2^\rho) = 0$$

and

$$(VII.22) \quad \int_{B_R(0)} |\nabla \vec{e}_1^\rho|^2 + |\nabla \vec{e}_2^\rho|^2 \leq C \int_{\mathbb{C}} |\nabla \vec{n}_{\tilde{\xi}_\infty}|^2.$$

We now consider a sequence of radii ρ_j converging to infinity. Extracting a subsequence gives the existence of a limiting frame $(\vec{e}_1^\infty, \vec{e}_2^\infty)$ on \mathbb{C} satisfying

$$\star \hat{n}_{\tilde{\xi}_\infty} = \vec{e}_1^\infty \wedge \vec{e}_2^\infty, \quad \text{div}(\vec{e}_1^\infty \cdot \nabla \vec{e}_2^\infty) = 0,$$

and

$$(VII.23) \quad \int_{\mathbb{C}} |\nabla \vec{e}_1^\infty|^2 + |\nabla \vec{e}_2^\infty|^2 \leq C \int_{\mathbb{C}} |\nabla \vec{n}_{\tilde{\xi}_\infty}|^2.$$

We have at every point of the *minimal* conformal immersion $\tilde{\xi}_\infty$ the identity

$$e^{2\tilde{\lambda}_\infty} K_{\tilde{\xi}_\infty} = -\nabla^\perp \vec{e}_1^\infty \cdot \nabla \vec{e}_2^\infty.$$

Integrating over \mathbb{C} then yields

$$\int_{\mathbb{C}} K_{\tilde{\xi}_\infty} e^{2\tilde{\lambda}_\infty} dx = \int_{\mathbb{C}} d\vec{e}_1^\infty \wedge d\vec{e}_2^\infty = 0.$$

Paired with (VII.21), the latter implies that $\nabla \vec{n}_{\tilde{\xi}_\infty} \equiv 0$, which contradicts (VII.16). This is the desired contradiction, and thus Lemma VII.1 is proved. \square

VIII. Proof of theorems

VIII.1. *Proof of Theorem I.2.* Let $\vec{\Phi}_k$ be a sequence of Willmore immersions of Σ with uniformly bounded energy

$$\limsup_{k \rightarrow +\infty} E(\vec{\Phi}_k) < +\infty$$

and such that the conformal class of the induced metric $g_k = \vec{\Phi}_k^* g_{\mathbb{R}^m}$ remains in a compact subdomain of the moduli space of Σ .

We consider a subsequence, still denoted $\vec{\Phi}_k$, as given by Lemma II.1. Let Ξ_k be the corresponding sequence of Möbius transformations, and let f_k be the sequence of Lipschitz diffeomorphisms of Σ , as given by Lemma II.1. We set $\vec{\xi}_k := \Xi_k \circ \vec{\Phi}_k \circ f_k$. The conclusion of Lemma II.1 states that $\vec{\xi}_k$ weakly converges in $W^{2,2}$ to a weak immersion $\vec{\xi}_\infty$, away from finitely many points a_i . This limiting immersion might be branched at the points a_i .

Let h_k be constant scalar curvature metric with respect to which $\vec{\xi}_k$ is conformal, and let u_k be the function satisfying

$$g_k = \vec{\xi}_k^* g_{\mathbb{R}^m} = e^{2u_k} h_k.$$

The classical Liouville equation reads

$$\Delta_{h_k} u_k + K_{g_k} e^{2u_k} = K_{h_k}.$$

Since h_k strongly converges to some limiting constant scalar curvature h_∞ , classical results from geometric analysis on manifolds (see, for instance, [Aub82]) show that the gradient of the Green function associated with the Laplace-Beltrami operator Δ_{h_k} is uniformly bounded in $L^{2,\infty}$. Hence,

$$\begin{aligned} \text{(VIII.1)} \quad \|\nabla u_k\|_{L^{2,\infty}(\Sigma)} &\leq C \left[\int_{\Sigma} |K_{g_k}| \, d\text{vol}_{g_k} + \int_{\Sigma} |K_{h_k}| \, d\text{vol}_{h_k} \right] \\ &\leq C \left[E(\vec{\Phi}_k) + 2\pi |\chi(\Sigma)| \right], \end{aligned}$$

where the $L^{2,\infty}$ -norm is to be understood with respect to a reference metric g_0 on Σ . As before, $\chi(\Sigma)$ denotes the Euler characteristic of Σ . We define the constant

$$\text{(VIII.2)} \quad \Lambda := \sup_{k \in \mathbb{N}} E(\vec{\Phi}_k) + \|\nabla u_k\|_{L^{2,\infty}(\Sigma)} < +\infty.$$

Owing to the ε -regularity Theorem I.5 in [Riv08], the sequence $\vec{\xi}_k$ actually converges strongly in $C_{\text{loc}}^l(\Sigma \setminus \{a_1, \dots, a_N\}, \mathbb{R}^m)$ for all $l \in \mathbb{N}$. Namely,

$$\text{(VIII.3)} \quad \vec{\xi}_k \longrightarrow \vec{\xi}_\infty \quad \text{in} \quad C_{\text{loc}}^l(\Sigma \setminus \{a_1, \dots, a_N\}, \mathbb{R}^m).$$

Accordingly, the limiting immersion $\vec{\xi}_\infty$ is Willmore, although *a priori* only away from the points $\{a_1, \dots, a_N\}$. Since $\vec{\xi}_k$ is Willmore, for any fixed small

radius $\rho > 0$ and any $i = 1, \dots, N$, one has

$$\int_{B_\rho(a_i)} \operatorname{div} \left(2 \nabla \vec{H}_{\xi_k}^- - 3 \nabla (\pi \vec{n}_{\xi_k}^-) (\vec{H}_{\xi_k}^-) - \star (\nabla^\perp \vec{n}_{\xi_k}^- \wedge \vec{H}_{\xi_k}^-) \right) dx = 0,$$

where $B_\rho(a_i)$ is the geodesic ball centered at a_i and of radius ρ for the flat metric, in some converging conformal coordinate system for $\vec{\xi}_k^* g_{\mathbb{R}^m}$. Whence,

$$\int_{\partial B_\rho(a_i)} 2 \partial_\nu \vec{H}_{\xi_k}^- - 3 \partial_\nu (\pi \vec{n}_{\xi_k}^-) (\vec{H}_{\xi_k}^-) + \star (\partial_\tau \vec{n}_{\xi_k}^- \wedge \vec{H}_{\xi_k}^-) \rho d\theta = 0,$$

where ν is the unit normal vector $x/|x|$ and τ the unit tangent vector $x^\perp/|x|$ to the boundary of $B_\rho(a_i)$. The strong convergence of $\vec{\xi}$ to $\vec{\xi}_\infty$ enables passing to the limit in the last identity, and we thus find

$$(VIII.4) \quad \int_{\partial B_\rho(a_i)} 2 \partial_\nu \vec{H}_{\xi_\infty}^- - 3 \partial_\nu (\pi \vec{n}_{\xi_\infty}^-) (\vec{H}_{\xi_\infty}^-) + \star (\partial_\tau \vec{n}_{\xi_\infty}^- \wedge \vec{H}_{\xi_\infty}^-) \rho d\theta = 0$$

for all fixed ρ .

This is precisely the condition required to apply the ‘‘point removability’’ theorem established in [Riv08]. In particular, $\vec{\xi}_\infty$ extends to a (possibly branched) smooth Willmore immersion in $B_\rho(a_i)$. More detailed information on the behavior of the immersion near the points a_i are given in [BR13].

We now choose a fixed $\varepsilon_0 > 0$ smaller than the constants $\varepsilon(m)$ of lemmas VI.1 and VII.1. For this particular choice of ε_0 , we apply the bubble-neck decomposition procedure outlined in Proposition III.1. Owing to (III.13), both Lemma VI.1 and Lemma VII.1 apply in each connected component of $\Omega_k(\alpha)$, which, recall, is a union of disjoint annuli. From Lemma VI.1, we thus have

$$(VIII.5) \quad \limsup_{k \rightarrow +\infty} \|e^\lambda \vec{H}_{\xi_k}^-\|_{L^{2,1}(\Omega_k(1/2))} \leq C(m, \Lambda).$$

As for Lemma VII.1, it yields

$$(VIII.6) \quad \forall \varepsilon > 0 \quad \exists \alpha > 0 \quad \text{s.t.} \quad \|\nabla \vec{n}_{\xi_k}^-\|_{L^{2,\infty}(\Omega_k(\alpha))} \leq \varepsilon.$$

Combining this last assertion with the result of Lemma V.1 implies that

$$(VIII.7) \quad \forall \varepsilon > 0 \quad \exists \alpha > 0 \quad \text{s.t.} \quad \left| \int_{\Omega_k(\alpha)} K_{\xi_k}^- \operatorname{dvol}_{g_k} \right| \leq \varepsilon.$$

Altogether now, (VIII.5) and (VIII.6) give

$$(VIII.8) \quad \forall \varepsilon > 0 \quad \exists \alpha > 0 \quad \exists k_0 \in \mathbb{N} \quad \text{s.t.} \quad \forall k \geq k_0 \\ \|e^\lambda \vec{H}_{\xi_k}^-\|_{L^1(\Omega_k(\alpha))} \leq \|e^\lambda \vec{H}_{\xi_k}^-\|_{L^{2,1}(\Omega_k(\alpha))}^{1/2} \|e^\lambda \vec{H}_{\xi_k}^-\|_{L^{2,\infty}(\Omega_k(\alpha))}^{1/2} \leq \sqrt{C(m, \Lambda)} \varepsilon,$$

where we used the fact that $L^{2,\infty}$ is the dual space of $L^{2,1}$.

Combining (VIII.8) and (VIII.7) shows that

$$(VIII.9) \quad \lim_{\alpha \rightarrow 0} \lim_{k \rightarrow +\infty} \int_{\Omega_k(\alpha)} |\nabla \vec{n}_{\xi_k}^-|^2 dx = 0.$$

This is the *no-neck energy property* that we aimed at establishing. To finish the proof of Theorem I.2, we still need to identify the energy of the bubbles as converging in the limit to that of Willmore spheres. This is what we do below.

Let $i \in \{1, \dots, N\}$, and let $j \in \{1, \dots, Q^i\}$. From (III.15), on the bubble domain

$$B(i, j, \alpha, k) := B_{\alpha^{-1}\rho_k^{i,j}}(x_k^{i,j}) \setminus \cup_{j' \in I^{i,j}} B_{\alpha\rho_k^{i,j}}(x_k^{i,j'}),$$

there is no concentration of energy. Thus, arguing as in the proof of Lemma V.2 (see also lemma III.1 in [Riv13c]), the conformal factor satisfies a uniform Harnack estimate:

(VIII.10)

$$\forall 0 < \alpha < 1 \quad \exists C_\alpha > 0 \quad \text{s.t.} \quad 1 \leq \frac{\sup_{\{x \in B(i,j,\alpha,k)\}} e^{\lambda_k(x)}}{\inf_{\{x \in B(i,j,\alpha,k)\}} e^{\lambda_k(x)}} \leq C_\alpha.$$

Choose an arbitrary point

$$z_k^{i,j} \in B(i, j, \alpha, k) \cap B_{\rho_k^{i,j}}(x_k^{i,j}) \setminus \left(\cup_{j' \in I^{i,j}} B_{M^{-1}\rho_k^{i,j}}(x_k^{i,j'}) \right),$$

where $M > \text{card}(I^{i,j})$, and set $\lambda(i, j, k, \alpha) := \lambda_k(x_k)$. We introduce the renormalized Willmore immersion

$$\vec{\xi}_k(y) := e^{-\lambda(i,j,\alpha,k)} \left[\vec{\xi}_k(\rho_k^{i,j} y + x_k^{i,j}) - \vec{\xi}_k(z_k^{i,j}) \right].$$

We can extract a subsequence in such a way that the following limits exist:

$$\forall j' \in I^{i,j} \quad \lim_{k \rightarrow +\infty} \frac{x_k^{i,j} - z_k^{i,j}}{\rho_k^{i,j}} = a_i^{j,j'} \in B_1(0) \setminus \{0\}.$$

As the energy does not concentrate on any $B(i, j, \alpha, k)$ for $0 < \alpha < 1$, we obtain, using once more the ε -regularity and the local control of the conformal factor ensured by (VIII.11), that

$$\vec{\xi}_k \longrightarrow \vec{\xi}_\infty \quad \text{in} \quad C_{\text{loc}}^l(\mathbb{C} \setminus \cup_{j' \in I^{i,j}} \{a_i^{j,j'}\}) \quad \forall l \in \mathbb{N},$$

where $\vec{\xi}_\infty$ is a Willmore immersion of $\mathbb{C} \setminus \cup_{j' \in I^{i,j}} \{a_i^{j,j'}\}$. Suppose that

$$\int_{\mathbb{C}} |\nabla \vec{\xi}_\infty|^2 dx < +\infty.$$

Then, using the stereographic projection of S^2 onto \mathbb{C} with respect to the north pole, $\vec{\xi}_\infty$ realizes an immersion of $S^2 \setminus \cup_{j' \in I^{i,j}} \{a_i^{j,j'}\} \cup \{\text{North}\}$ included in a bounded domain. Just as we explained above for the limit immersion $\vec{\xi}_\infty$, one can remove the singularities $\cup_{j' \in I^{i,j}} \{a_i^{j,j'}\} \cup \{\text{North}\}$, which possibly realize branch points of our Willmore S^2 . We have moreover

$$(VIII.11) \quad \lim_{\alpha \rightarrow 0} \lim_{k \rightarrow +\infty} \int_{B(i,j,\alpha,k)} e^{2\lambda_k} |\vec{H}_{\vec{\xi}_k}|^2 = W(\vec{\xi}_\infty).$$

On the other hand, suppose that¹⁶

$$\int_{\mathbb{C}} |\nabla \tilde{\xi}_{\infty}|^2 dx_1 dx_2 = +\infty.$$

Following [Riv13a], we can find a point $p_k \in \mathbb{R}$ such that

$$\forall k \in \mathbb{N} \quad e^{-\lambda(i,j,\alpha,k)} [\vec{\xi}_k(\Sigma) - \vec{\xi}_k(z_k^{i,j})] \cap B_1(p_k) = \emptyset \quad \text{and} \quad \limsup_{k \rightarrow +\infty} |p_k| < +\infty.$$

Let \mathcal{I}_k be the inversion with respect to p_k , and $\hat{\xi}_k := \mathcal{I}_k \circ \tilde{\xi}_k$. We may assume, modulo extraction of a subsequence if necessary, that the sequence p_k converges to a point p_{∞} . We let \mathcal{I}_{∞} be the inversion in \mathbb{R}^m with respect to this limit-point. There holds

$$\hat{\xi}_k \longrightarrow \hat{\xi}_{\infty} = \mathcal{I}_{\infty} \circ \tilde{\xi}_{\infty} \quad \text{in} \quad C^l_{\text{loc}}(\mathbb{C} \setminus \cup_{j' \in I^{i,j}} \{a_i^{j,j'}\}) \quad \forall l \in \mathbb{N},$$

where $\hat{\xi}_{\infty}$ is a Willmore immersion of $\mathbb{C} \setminus \cup_{j' \in I^{i,j}} \{a_i^{j,j'}\}$ satisfying

$$\int_{\mathbb{C}} |\nabla \hat{\xi}_{\infty}|^2 dx < +\infty.$$

Arguing as above, $\hat{\xi}_{\infty}$ extends to a smooth, possibly branched, immersion of S^2 . Using lemma A.4 in [Riv13a], we then obtain that

$$(VIII.12) \quad \lim_{\alpha \rightarrow 0} \lim_{k \rightarrow +\infty} \int_{B(i,j,\alpha,k)} e^{2\lambda_k} |\vec{H}_{\vec{\xi}_k}|^2 dx = W(\hat{\xi}_{\infty}) - 4\pi\theta_0,$$

where θ_0 is the integer density of $\vec{\xi}_{\infty}(\mathbb{C})$ at the image point $0 \in \mathbb{R}^m$.

Altogether, (VIII.3), (VIII.11), and (VIII.12) imply that there exists a subsequence (indexed as the original, for notational convenience) such that

$$(VIII.13) \quad \lim_{k \rightarrow +\infty} W(\vec{\Phi}_k) = W(\vec{\xi}_{\infty}) + \sum_{s=1}^p W(\tilde{\xi}_{\infty}^s) + \sum_{t=1}^q \left[W(\hat{\xi}_{\infty}^t) - 4\pi\theta_0^t \right].$$

The index s is used for the first alternative, in which the bubble has no infinite end, while the index t is used for the second alternative, in which the bubble has at least one infinite planar end, and it must thus be inverted to become compact. This is exactly the desired *energy identity* (I.2), thereby ending the proof of Theorem I.2. \square

VIII.2. *Proof of Theorem I.5.* Let $\vec{\Phi}_k$ be a sequence of Willmore immersions satisfying

$$W(\vec{\Phi}_k) < \min\{8\pi, \omega_g^m\} - \delta.$$

¹⁶This should correspond to the case when $\lambda(i, j, \alpha, k) \rightarrow -\infty$.

We want to show that it is possible extract a subsequence $\vec{\Phi}_{k'}$ and to find a sequence of diffeomorphisms $f_{k'}$ of Σ and Möbius transformation $\Xi_{k'}$, such that

$$(VIII.14) \quad \vec{\xi}_{k'} := \Xi_{k'} \circ \vec{\Phi}_{k'} \circ f_{k'} \longrightarrow \vec{\xi}_{\infty} \quad \text{strongly in } C^l(\Sigma) \quad \forall l \in \mathbb{N}.$$

From the work in [Riv13a] (cf. also [KL12]), the sequence of conformal classes of the metrics induced by $\vec{\Phi}_{k'}$ remains in a compact subdomain of the Moduli space. Theorems I.3 and I.4 thus apply.

If

$$\lim_{k' \rightarrow \infty} W(\vec{\Phi}_{k'}) = W(\vec{\xi}_{\infty}).$$

then from Theorem I.3, we get (VIII.14) directly. If not, then there must be a bubble. We take it to be the most concentrated one, centered at $x_{k'}$ and of radius $\rho_{k'}$, and converging to a point a^1 . As is explained at the end of the previous section, we can find a Möbius transformation $\Lambda_{k'}$ such that in local conformal coordinates there holds

$$\Lambda_{k'} \circ \vec{\xi}_{k'}(\rho_{k'}y + x_{k'}) \longrightarrow \hat{\xi}_{\infty},$$

where $\hat{\xi}_{\infty}$ is a Willmore sphere. Assume this sphere is homothetic to the round sphere. Then, if $\Lambda_{k'}$ contains no inversion and $\hat{\xi}_{\infty}$ is of the type η_s from Theorem I.2, we conclude that

$$\lim_{k' \rightarrow \infty} W(\vec{\Phi}_{k'}) \geq W(\vec{\xi}_{\infty}) + W(\hat{\xi}_{\infty}) \geq 4\pi + 4\pi = 8\pi,$$

which is a contradiction.

If now $\Lambda_{k'}$ contains an inversion, then modulo a dilation in the image, we can arrange for $\xi_{k'}(\rho_{k'}y + x_{k'})$ to converge strongly to a plane, and thus

$$\int_{B_{\rho_{k'}}(x_{k'})} |\nabla \vec{\xi}_{k'}| \rightarrow 0.$$

This contradicts the very definition of a bubble, which requires that

$$\int_{B_{\rho_{k'}}(x_{k'})} |\nabla \vec{\xi}_{k'}| \geq \varepsilon(m) > 0.$$

Accordingly, $\hat{\xi}_{\infty}$ cannot be homothetic to the round sphere. Calling upon the results from [Bry84] and [Mon00], it follows that

$$\begin{aligned} \lim_{k' \rightarrow \infty} W(\vec{\Phi}_{k'}) &= \lim_{k' \rightarrow \infty} W(\Lambda_{k'} \circ \vec{\xi}_{k'}) \\ &\geq \lim_{k' \rightarrow \infty} W(\Lambda_{k'} \circ \xi_{k'}(\rho_{k'}y + x_{k'})) \longrightarrow W(\hat{\xi}_{\infty}) \geq 8\pi, \end{aligned}$$

which is again a contradiction.

Therefore, we have shown that in all cases, the assertion (VIII.14) holds, thereby proving Theorem I.5.

References

- [Aub82] T. AUBIN, *Nonlinear Analysis on Manifolds. Monge-Ampère Equations*, *Grundl. Math. Wissen.* **252**, Springer-Verlag, New York, 1982. MR 0681859. Zbl 0512.53044. <http://dx.doi.org/10.1007/978-1-4612-5734-9>.
- [BK03] M. BAUER and E. KUWERT, Existence of minimizing Willmore surfaces of prescribed genus, *Int. Math. Res. Not.* **2003** (2003), 553–576. MR 1941840. Zbl 1029.53073. <http://dx.doi.org/10.1155/S1073792803208072>.
- [Ber13] Y. BERNARD, Noether’s theorem and the Willmore functional, 2013, preprint.
- [BR11] Y. BERNARD and T. RIVIÈRE, Local Palais-Smale sequences for the Willmore functional, *Comm. Anal. Geom.* **19** (2011), 563–599. MR 2843242. Zbl 1245.53051. <http://dx.doi.org/10.4310/CAG.2011.v19.n3.a5>.
- [BR13] Y. BERNARD and T. RIVIÈRE, Singularity removability at branch points for Willmore surfaces, *Pacific J. Math.* **265** (2013), 257–311. MR 3096502. Zbl 06242169. <http://dx.doi.org/10.2140/pjm.2013.265.257>.
- [Bla29] W. BLASCHKE, Vorlesungen über Differentialgeometrie und geometrische Grundlagen von Einsteins relativitätstheorie, III, in *Differentialgeometrie der Kreise und Kugeln*, *Grundlehren Math. Wissen.* **39**, Gerhard Thomsen, 1929. JFM 55.0422.01.
- [Bry84] R. L. BRYANT, A duality theorem for Willmore surfaces, *J. Differential Geom.* **20** (1984), 23–53. MR 0772125. Zbl 0555.53002. Available at <http://projecteuclid.org/euclid/jdg/1214438991>.
- [BP06] A. BUTSCHER and F. PACARD, Doubling constant mean curvature tori in S^3 , *Ann. Sc. Norm. Super. Pisa Cl. Sci.* **5** (2006), 611–638. MR 2297724. Zbl 1170.53303. Available at http://www.numdam.org/item?id=ASNSP_2006_5_5_4_611_0.
- [DT95] W. DING and G. TIAN, Energy identity for a class of approximate harmonic maps from surfaces, *Comm. Anal. Geom.* **3** (1995), 543–554. MR 1371209. Zbl 0855.58016.
- [Ge99] Y. GE, A remark on generalized harmonic maps into spheres, *Nonlinear Anal.* **36** (1999), 495–506. MR 1675268. Zbl 0932.35071. [http://dx.doi.org/10.1016/S0362-546X\(98\)00079-0](http://dx.doi.org/10.1016/S0362-546X(98)00079-0).
- [Hél02] F. HÉLEIN, *Harmonic Maps, Conservation Laws and Moving Frames*, second ed., *Cambridge Tracts in Math.* **150**, Cambridge Univ. Press, Cambridge, 2002. MR 1913803. Zbl 1010.58010. <http://dx.doi.org/10.1017/CBO9780511543036>.
- [Jos91] J. JOST, *Two-Dimensional Geometric Variational Problems*, *Pure Appl. Math. (NY)*, John Wiley & Sons Ltd., Chichester, 1991, A Wiley-Interscience Publication. MR 1100926. Zbl 0729.49001.
- [KL12] E. KUWERT and Y. LI, $W^{2,2}$ -conformal immersions of a closed Riemann surface into \mathbb{R}^n , *Comm. Anal. Geom.* **20** (2012), 313–340. MR 2928715. Zbl 1271.53010. <http://dx.doi.org/10.4310/CAG.2012.v20.n2.a4>.

- [KS04] E. KUWERT and R. SCHÄTZLE, Removability of point singularities of Willmore surfaces, *Ann. of Math.* **160** (2004), 315–357. MR 2119722. Zbl 1078.53007. <http://dx.doi.org/10.4007/annals.2004.160.315>.
- [KS12] E. KUWERT and R. SCHÄTZLE, Closed surfaces with bounds on their Willmore energy, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* **11** (2012), 605–634. MR 3059839. Zbl 1260.53027. Available at <http://annaliscienze.sns.it/index.php?page=Article&id=251>.
- [KS13] E. KUWERT and R. SCHÄTZLE, Minimizers of the Willmore functional under fixed conformal class, *J. Differential Geom.* **93** (2013), 471–530. MR 3024303. Zbl 06201325. Available at <http://projecteuclid.org/euclid.jdg/1361844942>.
- [LR] P. LAURAIN and T. RIVIÈRE, Angular energy quantization for linear elliptic systems with antisymmetric potentials and applications, to appear in *Anal. PDE*. arXiv 1109.3599.
- [LR01] F.-H. LIN and T. RIVIÈRE, A quantization property for moving line vortices, *Comm. Pure Appl. Math.* **54** (2001), 826–850. MR 1823421. Zbl 1029.35127. <http://dx.doi.org/10.1002/cpa.3003>.
- [LR02] F.-H. LIN and T. RIVIÈRE, Energy quantization for harmonic maps, *Duke Math. J.* **111** (2002), 177–193. MR 1876445. Zbl 1014.58008. <http://dx.doi.org/10.1215/S0012-7094-02-11116-8>.
- [MN14] F. C. MARQUES and A. NEVES, Min-max theory and the Willmore conjecture, *Ann. of Math.* **179** (2014), 683–782. <http://dx.doi.org/10.4007/annals.2014.179.2.6>.
- [Mon00] S. MONTIEL, Willmore two-spheres in the four-sphere, *Trans. Amer. Math. Soc.* **352** (2000), 4469–4486. MR 1695032. Zbl 0961.53035. <http://dx.doi.org/10.1090/S0002-9947-00-02571-X>.
- [MŠ95] S. MÜLLER and V. ŠVERÁK, On surfaces of finite total curvature, *J. Differential Geom.* **42** (1995), 229–258. MR 1366547. Zbl 0853.53003. Available at <http://projecteuclid.org/euclid.jdg/1214457233>.
- [Par96] T. H. PARKER, Bubble tree convergence for harmonic maps, *J. Differential Geom.* **44** (1996), 595–633. MR 1431008. Zbl 0874.58012. Available at <http://projecteuclid.org/euclid.jdg/1214459224>.
- [Riv02a] T. RIVIÈRE, Bubbling and regularity issues in geometric non-linear analysis, in *Proceedings of the International Congress of Mathematicians, Vol. III* (Beijing, 2002), Higher Ed. Press, Beijing, 2002, pp. 197–208. MR 1957532. Zbl 1136.35334.
- [Riv02b] T. RIVIÈRE, Interpolation spaces and energy quantization for Yang-Mills fields, *Comm. Anal. Geom.* **10** (2002), 683–708. MR 1925499. Zbl 1018.58006.
- [Riv08] T. RIVIÈRE, Analysis aspects of Willmore surfaces, *Invent. Math.* **174** (2008), 1–45. MR 2430975. Zbl 1155.53031. <http://dx.doi.org/10.1007/s00222-008-0129-7>.
- [Riv13a] T. RIVIÈRE, Lipschitz conformal immersions from degenerating Riemann surfaces with L^2 -bounded second fundamental forms, *Adv. Calc. Var.*

- 6** (2013), 1–31. MR 3008339. Zbl 06137860. <http://dx.doi.org/10.1515/acv-2012-0108>.
- [Riv13b] T. RIVIÈRE, Weak immersions of surfaces with L^2 bounded second fundamental form, 2013, IAS/Park City Mathematics Institute summer school.
- [Riv13c] T. RIVIÈRE, Variational principles for immersed surfaces with L^2 -bounded second fundamental form, *J. reine angew. Math.*, published online: 01/23/2013. <http://dx.doi.org/10.1515/crelle-2012-0106>.
- [SU81] J. SACKS and K. UHLENBECK, The existence of minimal immersions of 2-spheres, *Ann. of Math.* **113** (1981), 1–24. MR 0604040. Zbl 0462.58014. <http://dx.doi.org/10.2307/1971131>.
- [Sch13] R. M. SCHÄTZLE, Estimation of the conformal factor under bounded Willmore energy, *Math. Z.* **274** (2013), 1341–1383. MR 3078270. Zbl 1275.53011. <http://dx.doi.org/10.1007/s00209-012-1119-4>.
- [SU83] R. SCHOEN and K. UHLENBECK, Boundary regularity and the Dirichlet problem for harmonic maps, *J. Differential Geom.* **18** (1983), 253–268. MR 0710054. Zbl 0547.58020. Available at <http://projecteuclid.org/euclid.jdg/1214437663>.
- [Sim93] L. SIMON, Existence of surfaces minimizing the Willmore functional, *Comm. Anal. Geom.* **1** (1993), 281–326. MR 1243525. Zbl 0848.58012.
- [Str85] M. STRUWE, On the evolution of harmonic mappings of Riemannian surfaces, *Comment. Math. Helv.* **60** (1985), 558–581. MR 0826871. Zbl 0595.58013. <http://dx.doi.org/10.1007/BF02567432>.
- [Tor95] T. TORO, Geometric conditions and existence of bi-Lipschitz parameterizations, *Duke Math. J.* **77** (1995), 193–227. MR 1317632. Zbl 0847.42011. <http://dx.doi.org/10.1215/S0012-7094-95-07708-4>.
- [Zhu10] M. ZHU, Harmonic maps from degenerating Riemann surfaces, *Math. Z.* **264** (2010), 63–85. MR 2564932. Zbl 1213.53086. <http://dx.doi.org/10.1007/s00209-008-0452-0>.

(Received: July 1, 2011)

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT REGENSBURG, REGENSBURG, GERMANY
E-mail: yann.bernard@mathematik.uni-regensburg.de

ETH ZENTRUM, ZÜRICH, SWITZERLAND
E-mail: tristan.riviere@fim.math.ethz.ch