

Chern slopes of simply connected complex surfaces of general type are dense in $[2,3]$

By XAVIER ROULLEAU and GIANCARLO URZÚA

Dedicated to Igor Dolgachev on the occasion of his 70th birthday

Abstract

We prove that for any number $r \in [2,3]$, there are spin (resp. non-spin and minimal) simply connected complex surfaces of general type X with $c_1^2(X)/c_2(X)$ arbitrarily close to r . In particular, this shows the existence of simply connected surfaces of general type arbitrarily close to the Bogomolov-Miyaoka-Yau line. In addition, we prove that for any $r \in [1,3]$ and any integer $g \geq 0$, there are minimal complex surfaces of general type X with $c_1^2(X)/c_2(X)$ arbitrarily close to r and $\pi_1(X)$ isomorphic to the fundamental group of a compact Riemann surface of genus g .

1. Introduction

Let X be a minimal nonsingular projective surface of general type over \mathbb{C} . Its Chern numbers $c_1^2(X), c_2(X)$ satisfy $c_1^2(X) > 0$, $c_2(X) > 0$, the Noether inequality $\frac{1}{5}c_2(X) - \frac{36}{5} \leq c_1^2(X)$, and the Bogomolov-Miyaoka-Yau inequality [Bog78], [Miy77], [Yau77]

$$c_1^2(X) \leq 3c_2(X),$$

where $c_1^2(X) = 3c_2(X)$ if and only if the universal cover of X is the complex two dimensional ball [Yau77], [Miy83]. The geography problem [Per87] asks: for which pair of integers (a, b) does there exist X such that $c_1^2(X) = a$ and $c_2(X) = b$? We refer to [BHPVdV04, VII §8] for general existence results. One of them, due to Sommese [Som84], states that every rational point in $[1/5, 3]$ occurs as the slope $c_1^2(X)/c_2(X)$ of some (irregular) surface X .

Since the early stages of the geography problem, the question was naturally sharpened by imposing the constraint of simply connectedness. Because of the above result of Yau and Miyaoka, a surface X (of general type) satisfying $c_1^2(X) = 3c_2(X)$ is not simply connected. *Is this the only additional restriction?* If we divide surfaces according to their index $\frac{1}{3}(c_1^2(X) - 2c_2(X))$, then

there are satisfactory results for surfaces with negative index due to Persson [Per81]. For instance, all rational slopes between $1/5$ and 2 are realized as Chern slopes of simply connected surfaces.

Simply connected surfaces with nonnegative index are more difficult to find. In the late 70s, it was actually conjectured that any simply connected surface of general type has negative index (the Bogomolov Watershed Conjecture; see [VdV78], [Per87]). In 1984, Moishezon and Teicher [MT87] gave the first examples with positive index. After this, there were several attempts to fill out this positive index region. One would at least want to fill out the Chern slope interval $[2, 3]$ as much as possible. In particular, *are there simply connected surfaces of general type with Chern slope arbitrarily close to 3?* This is a well-known open problem; see, for example, [Hol78], [Hol79a], [Hol79b], [Hol81], [Per87], [Che87], [PP96, §4], [PPX96], [BHPVdV04, VII§8B], [Urz10], [Urz11]. The construction of this type of surfaces becomes harder as we approach 3 . The highest known slope was given in [Urz10], proving the existence of such surfaces with c_1^2/c_2 arbitrarily close to $\frac{71}{26} \approx 2.730769$. Our main theorem is

THEOREM 1.1. *For any number $r \in [2, 3]$, there are spin (resp. nonspin and minimal) simply connected surfaces of general type X with $c_1^2(X)/c_2(X)$ arbitrarily close to r .*

In particular, this shows that 3 is indeed the sharp upper bound for slopes of both spin and nonspin simply connected surfaces. Let us recall that a simply connected surface is called *spin* if its canonical class is 2 -divisible. For the topological interest in the existence of spin surfaces we refer to [PP96, §4], [PPX96].

A key ingredient in the construction of the surfaces X is the existence of a family of special arrangements of elliptic curves \mathcal{H}'_n in the blow-up H of \mathbb{P}^2 at the 12 triple points of the dual Hesse arrangement of lines. These arrangements are related to special arrangements of elliptic curves in $\mathbb{C}/\mathbb{Z}[\zeta] \times \mathbb{C}/\mathbb{Z}[\zeta]$, where $\zeta = e^{2\pi i/6}$. The latter were discovered by Hirzebruch in [Hir84]. They are connected to open ball quotients. (See [KH05] for more on this type of special arrangements.) The arrangements \mathcal{H}'_n in H have many simple 4 -fold points as singularities (we later call them 4 -points) and a particular divisibility property in $\text{Pic}(H)$. This allows us to consider certain cyclic covers [EV92] which are branched along the arrangement, together with other curves, but “avoid” the exceptional divisors from all the 4 -points. We make use of the more detailed description of cyclic covers developed in [Urz08], [Urz10] which, in particular, makes explicit the use of log invariants, classical Dedekind sums, and lengths of Hirzebruch-Jung continued fractions, to estimate the Chern slope. To demonstrate that these surfaces are actually simply connected, we

show that certain blow-ups of the surfaces X admit a fibration to \mathbb{P}^1 , which has sections and a simply connected fiber; we then use a method from [Xia91]. The spin issue is more delicate; among other things it involves a choice of specific multiplicities for the curves in the branch locus.

The construction has many degrees of freedom due to the properties of the key arrangements of elliptic curves. For example, the surfaces we construct in the previous theorem admit many deformations. For this, we add a general arrangement of d lines to the branch locus, where d is essentially independent of the parameters used in the construction. In addition, the construction allows us the use of a specific base change, over the fibration to \mathbb{P}^1 mentioned above, to prove the following.

COROLLARY 1.2. *Let $q > 0$ be an integer. Then, for any number $r \in [1, 3]$, there are minimal surfaces of general type X with $c_1^2(X)/c_2(X)$ arbitrarily close to r and $\pi_1(X)$ isomorphic to the fundamental group of a compact Riemann surface of genus q .*

The following is an overall outline of the construction of the surfaces X in the above theorem, in connection to the sections of the paper. Let us fix $r \in [2, 3]$ and $\varepsilon > 0$. We construct the surfaces $X (= X_n)$ via morphisms $f: X_n \rightarrow Y_n$, such that there exists a function $\lambda: \mathbb{Q}_{>0} \rightarrow \mathbb{Q}$ and a rational number x satisfying $|\lambda(x) - r| < \varepsilon$ and $\lim_{n \rightarrow \infty} c_1^2(X_n)/c_2(X_n) = \lambda(x)$. To achieve this, it is crucial to have a special branch divisor $A \subset Y_n$ for the morphisms f . The surfaces Y_n are built through blow-ups at specific points of the surface H explained above. The key part of A comes from the arrangements of elliptic curves \mathcal{H}'_n in H . Roughly speaking, these arrangements are essentially defined as images of the Hirzebruch's arrangements \mathcal{H}_n in [Hir84], by means of a cyclic quotient of a blow-up of $\mathbb{C}/\mathbb{Z}[\zeta] \times \mathbb{C}/\mathbb{Z}[\zeta]$. In Section 2 we review the construction of the arrangements \mathcal{H}_n . In Section 3 we give the quotient construction of H , and we define the arrangements \mathcal{H}'_n . In Section 4 we compute their log Chern numbers. These log invariants are central for our asymptotic results. In Section 5 we prove our main result for spin surfaces. There, we also discuss all the morphisms involved, the branch divisors A , various divisibilities in Picard groups, formulas for Chern invariants, and arithmetic numbers which are relevant in these formulas. Using particular curves in A , we also prove in Section 5 that X is simply connected. In Section 6, by applying the previous work in a similar set up, we prove our main result for nonspin minimal surfaces. We point out that the functions λ have as image either $]1.375, 3[$ for spin surfaces or $]1, 3[$ for nonspin surfaces, and so we indeed obtain density for the closure of these two intervals. At the end of Section 6, we use the nonspin surfaces to prove the above corollary via particular base changes.

Conventions. For a nonsingular surface S , K_S denotes a canonical divisor of S . An arrangement of curves is a collection of curves $\{C_1, \dots, C_r\}$ on a surface. We loosely consider it as either a set, a divisor $C_1 + \dots + C_r$, or a curve $\bigcup_{i=1}^r C_i$. A k -point of an arrangement is a point of it locally of the form $(0, 0) \in \{(x - a_1 y) \cdots (x - a_k y) = 0\} \subset \mathbb{C}_{x,y}^2$ for some $a_i \neq a_j$. For an invertible sheaf \mathcal{L} , we write $\mathcal{L}^r := \mathcal{L}^{\otimes r}$. The genus of a compact Riemann surface C is denoted by $g(C)$. The (topological) fundamental group of Z is denoted by $\pi_1(Z)$.

Acknowledgements. We would like to thank Igor Dolgachev for helpful discussions and Fabrizio Catanese for pointing out that Proposition 3.1 is a classical fact due to A. Comessatti (see [CC93, §2]). The main results of the present article were found while the first author was visiting Pontificia Universidad Católica de Chile under an invitation from the second. Xavier Roulleau benefited from a sabbatical semester *délegation* CNRS. Giancarlo Urzúa was supported by the FONDECYT Inicio grant 11110047 funded by the Chilean Government.

2. Hirzebruch elliptic arrangements

Let $\zeta = e^{2\pi i/6}$, and let T be the elliptic curve

$$T = \mathbb{C}/\mathbb{Z}[\zeta],$$

where $\mathbb{Z}[\zeta] = \{a + \zeta b \mid a, b \in \mathbb{Z}\}$ are the Eisenstein integers. In [Hir84], Hirzebruch constructs special arrangements of elliptic curves in $T \times T$. We recall his construction, trying not to deviate much from his notation. Write (z, w) for points in $T \times T$. For each $n \in \mathbb{Z}[\zeta]$, let us denote by U_n the group of n -torsion points

$$U_n = \{(z, w) \in T \times T \mid nz = nw = 0\}.$$

The group U_n has order $(n\bar{n})^2$, where \bar{n} denotes the complex conjugation of n . (In [Hir84], n is an integer.)

Let us consider the following four elliptic curves:

$$T_0 = \{z = 0\}, \quad T_1 = \{z = w\}, \quad T_\infty = \{w = 0\}, \quad T_\zeta = \{z = \zeta w\}.$$

These four curves pass through $(0, 0)$ and do *not* intersect anywhere else. The group U_n acts on $T \times T$ by translation. The stabilizer of each of the four curves T_i in U_n has order $n\bar{n}$. Therefore, the orbit $D_i = U_n(T_i)$ of T_i ($i = 0, 1, \zeta, \infty$) consists of $n\bar{n}$ disjoint nonsingular elliptic curves, all being fibers of the fibrations $\pi_i: T \times T \rightarrow T$ defined as

$$\pi_0(z, w) = z, \quad \pi_1(z, w) = z - w, \quad \pi_\infty(z, w) = w, \quad \pi_\zeta(z, w) = z - \zeta w.$$

The arrangement

$$\mathcal{H}_n := D_0 + D_1 + D_\zeta + D_\infty$$

has $4n\bar{n}$ elliptic curves. The singularities of \mathcal{H}_n are precisely $(n\bar{n})^2$ 4-points.

For example, let us take $U_{1-\zeta^2}$, which is the group of smallest order among all nontrivial groups U_n with $n \in \mathbb{Z}[\zeta]$. The arrangement $\mathcal{H}_{1-\zeta^2}$ consists of twelve elliptic curves, each containing three points of $U_{1-\zeta^2}$, and nine 4-points; it gives a Hesse type configuration $(12_3, 9_4)$.

For $n \in \mathbb{Z}$, Holzapfel proves in [Hol86] that the complement in the blow-up at the n^4 4-points of the strict transform of the \mathcal{H}_n is a ball quotient by an arithmetic group. Holzapfel also considered other similar special elliptic arrangements, for example using the Gaussian integers; cf. [KH05]. They would also work in our upcoming constructions.

3. Dual Hesse and elliptic arrangements

The endomorphism $[\zeta^2]$ of $T \times T$ given by

$$[\zeta^2](z, w) = (\zeta^2 z, \zeta^2 w)$$

(where $\zeta = e^{2i\pi/6}$) is an order 3 automorphism of $T \times T$. Its fixed point set is the group of nine elements $U_{1-\zeta^2} \subset U_3$.

Let $\sigma: B \rightarrow T \times T$ be the blow-up at the nine points in $U_{1-\zeta^2}$. The strict transform of the T_i is denoted again by T_i . Thus $T_i^2 = -3$ in B . The automorphism $[\zeta^2]$ acts on B , and it is the identity on each of the nine exceptional curves of σ . Let $H = B/\langle[\zeta^2]\rangle$, and let $\pi: B \rightarrow H$ be the quotient map. For each i , we have the commutative diagram

$$\begin{array}{ccc} B & \xrightarrow{\pi} & H \\ \downarrow \pi_i \circ \sigma & & \downarrow \pi'_i \\ T & \longrightarrow & \mathbb{P}^1, \end{array}$$

where the horizontal arrows are quotients by $\mathbb{Z}/3\mathbb{Z}$ and the π'_i is the induced elliptic fibration on H .

PROPOSITION 3.1. *The surface H is the blow-up $\tau: H \rightarrow \mathbb{P}^2$ at the twelve 3-points of the dual Hesse arrangement*

$$(x^3 - y^3)(x^3 - z^3)(y^3 - z^3) = 0.$$

The image of the strict transform of $\mathcal{H}_{1-\zeta^2}$ by σ is the exceptional divisor of τ . The image of the nine (-1) -curves in B is the strict transform under τ of the nine lines in the dual Hesse arrangement.

Proof. For any i , the elliptic fibration $\pi'_i: H \rightarrow \mathbb{P}^1$ has three trees of four \mathbb{P}^1 's and nine sections from the image of the curves in $\mathcal{H}_{1-\zeta^2}$. This implies that H is simply connected; cf. [Xia91]. Moreover, if C is the exceptional divisor of σ , then $\pi^*(K_H) + 2C = K_B = C$, and so $\pi^*(K_H) = -C$ and $K_H^2 = -3$. If $e(X)$ is the topological Euler characteristic of X , then we also have $e(H) = \frac{1}{3}e(B) + \frac{2}{3}e(C) = 15$. The image of C under π are nine disjoint

(-3) -curves, and the image of the twelve elliptic curves in the strict transform of $\mathcal{H}_{1-\zeta^2}$ is a divisor in H formed by twelve disjoint (-1) -curves. After we blow down these 12 curves, we obtain \mathbb{P}^2 , and the nine (-3) -curves become an arrangement of nine lines with twelve 3-points. It is well known that, up to a projective equivalence, this is the dual Hesse arrangement; for example, see [Urz10, Thm.7.2]. \square

We now define the arrangements of elliptic curves in H which are going to be key in the next sections. Let us first to come back to $T \times T$ and the automorphism $[\zeta^2]$. For each $i \in \{0, 1, \zeta, \infty\}$, the curve T_i is stabilized by $[\zeta^2]$ and contains three fixed points including 0. These points are in $U_{1-\zeta^2}$, therefore the orbit $U_{1-\zeta^2}(T_i)$ consists of three elliptic curves. For $n \in \mathbb{Z}[\zeta]$, we recall that D_i is a union of $n\bar{n}$ disjoint elliptic curves which are fibers of π_i . Let us denote again by D_i the strict transform in B of D_i . We have two types of elliptic arrangements in H according to the $(1 - \zeta^2)$ -divisibility of n .

Suppose that $1 - \zeta^2$ divides n . In this paper we consider only this case. For each $i \in \{0, 1, \zeta, \infty\}$, it is immediate to verify that $[\zeta^2]$ acts on the divisor D_i , stabilizing the three curves in the orbit $U_{1-\zeta^2}(T_i)$ and permuting the other $n\bar{n} - 3$ curves.

Let us consider the situation in B . Each curve in $U_{1-\zeta^2}(T_i)$ cuts three (-1) -curves; the other components of D_i are disjoint from (-1) -curves. As we have said, the (-1) -curves are pointwise fixed by $[\zeta^2]$. The image of $U_{1-\zeta^2}(T_i)$ under π is formed by three (-1) -curves. The other $(n\bar{n} - 3)/3$ disjoint elliptic curves in the image of D_i are fibers of π'_i . We denote them by \mathcal{E}_i . Thus we define in H the following arrangement:

$$\mathcal{H}'_n := \mathcal{E}_0 + \mathcal{E}_1 + \mathcal{E}_\infty + \mathcal{E}_\zeta.$$

On $T \times T$, the arrangement \mathcal{H}_n has $(n\bar{n})^2$ 4-points, which are the n -torsion points. Since $U_{1-\zeta^2} \cap U_n = U_{1-\zeta^2}$, the arrangement in B contains $((n\bar{n})^2 - 9)$ 4-points, none of them lying on the ramification locus of π . Therefore, \mathcal{H}'_n consists of $4(n\bar{n} - 3)/3$ elliptic curves, and its singularities are precisely $4(n\bar{n} - 3)$ 3-points and $(n\bar{n} - 3)(n\bar{n} - 9)/3$ 4-points. Notice that for any $i \neq j$, if Γ_i and Γ_j are elliptic curves in \mathcal{E}_i and \mathcal{E}_j respectively, then they intersect transversally at three points.

4. Log Chern numbers

In this section we recall some basic definitions and facts around log differentials; cf. [EV92], [Urz10, §2]. We then compute the log Chern invariants for the arrangements \mathcal{H}'_n and for certain modifications $\overline{\mathcal{H}}_n$. This computation shows the importance of $\overline{\mathcal{H}}_n$ for the upcoming constructions.

Let Y be a nonsingular projective surface. Let A be a simple normal crossing divisor in Y . This means A is an arrangement of nonsingular curves

in Y and its singularities are only nodes (i.e., 2-points). The sheaf of log differentials along A , denoted by $\Omega_Y^1(\log A)$, is the \mathcal{O}_Y -submodule of $\Omega_Y^1 \otimes \mathcal{O}_Y(A)$ satisfying

- (1) $\Omega_Y^1(\log A)|_{Y \setminus A} = \Omega_{Y \setminus A}^1$.
- (2) At any point P in A , we have $\omega_P \in \Omega_Y^1(\log A)_P$ if and only if $\omega_P = \sum_{i=1}^m a_i \frac{dy_i}{y_i} + \sum_{j=m+1}^2 b_j dy_j$, where (y_1, y_2) is a local system around P for Y , and $\{y_1 \cdots y_m = 0\}$ defines A around P . (And so it is either $\{y_1 = 0\}$ or $\{y_1 y_2 = 0\}$.)

Hence, $\Omega_Y^1(\log A)$ is a locally free sheaf of rank two. Notice that

$$\bigwedge^2 \Omega_Y^1(\log A) \simeq \mathcal{O}_Y(K_Y + A).$$

In analogy to the Chern invariants of a nonsingular projective surface, the log Chern invariants of the pair (Y, A) are defined as $\bar{c}_i(Y, A) := c_i(\Omega_Y^1(\log A)^\vee)$ for $i = 1, 2$, where $\Omega_Y^1(\log A)^\vee$ is the dual sheaf of $\Omega_Y^1(\log A)$.

The log Chern numbers of (Y, A) are

$$\bar{c}_1^2(Y, A) := c_1(\Omega_Y^1(\log A)^\vee)^2 \quad \text{and} \quad \bar{c}_2(Y, A) := c_2(\Omega_Y^1(\log A)^\vee).$$

Hence for $A = 0$, we recover the Chern numbers of Y , which are denoted by $c_1^2(Y)$ and $c_2(Y)$. Sometimes we may drop Y or (Y, A) if the context is understood.

An arrangement of nonsingular curves \mathcal{A} in a nonsingular projective surface Z is called *simple crossings* if it has only k -points as singularities. The model example are line arrangements in \mathbb{P}^2 . The log Chern numbers of the pair (Z, \mathcal{A}) are defined as the log Chern numbers of the pair (Y, A) , where $Y \rightarrow Z$ is the blow-up of Z at all k -points of \mathcal{A} with $k > 2$ and A is the (reduced) total transform of \mathcal{A} .

PROPOSITION 4.1 ([Urz10, Prop.4.6]). *Let $\mathcal{A} = \{C_1, \dots, C_d\}$ be a simple crossing arrangement in Z . Let t_k be the number of k -points in \mathcal{A} . Then,*

$$\bar{c}_1^2(Z, \mathcal{A}) = c_1^2(Z) - \sum_{i=1}^d C_i^2 + \sum_{k \geq 2} (3k - 4)t_k + 4 \sum_{i=1}^d (g(C_i) - 1)$$

and $\bar{c}_2(Z, \mathcal{A}) = c_2(Z) + \sum_{k \geq 2} (k - 1)t_k + 2 \sum_{i=1}^d (g(C_i) - 1)$.

Let $\varphi_n: Z_n \rightarrow H$ be the blow-up at all the 4-points of \mathcal{H}'_n . Let $\bar{\mathcal{H}}_n$ be the strict transform of \mathcal{H}'_n under φ_n . Using the formulas in Proposition 4.1, we have

$$\bar{c}_1^2(H, \mathcal{H}'_n) = \frac{8}{3}(n\bar{n})^2 - 12n\bar{n} + 9, \quad \bar{c}_2(H, \mathcal{H}'_n) = (n\bar{n})^2 - 4n\bar{n} + 18$$

and

$$\bar{c}_1^2(Z_n, \bar{\mathcal{H}}_n) = (n\bar{n})^2 + 8n\bar{n} - 36, \quad \bar{c}_2(Z_n, \bar{\mathcal{H}}_n) = \frac{1}{3}n\bar{n}(n\bar{n} + 12).$$

At this point we highlight that the elimination of all 4-points from \mathcal{H}'_n make a big asymptotical difference with respect to log invariants.

5. Density for spin simply connected surfaces

Recall that for each $i \in \{0, 1, \infty, \zeta\}$, we have the commutative diagram

$$\begin{array}{ccccc} T \times T & \xleftarrow{\sigma} & B & \xrightarrow{\pi} & H & \xleftarrow{\varphi_n} & Z_n \\ & & & \nearrow \pi'_i & \downarrow \tau & & \\ & & T & \longrightarrow & \mathbb{P}^1 & & \mathbb{P}^2. \end{array}$$

The three singular fibers of π'_i are denoted by $F_{i,1}, F_{i,2}, F_{i,3}$. Each $F_{i,j}$ consists of four \mathbb{P}^1 's: one central curve $N_{i,j}$ with multiplicity 3, and three reduced curves transversal to $N_{i,j}$ at one point each. We write $N_i = N_{i,1} + N_{i,2} + N_{i,3}$. Let M be the nine \mathbb{P}^1 's from the lines of the dual Hesse arrangement, and let N be the twelve exceptional \mathbb{P}^1 's from its twelve 3-points. We have $N = \sum_{i=0,1,\zeta,\infty} N_i$ and

$$F_{i,1} + F_{i,2} + F_{i,3} = M + 3N_i.$$

Let $\alpha > 0, \beta > 0$ be integers. Let us take $n = 12\alpha p$, where $p \geq 5$ is a prime number. Let \mathcal{E}'_i be $8\beta^2 p^2$ general fibers of π'_i . Let $d > 0$ be an integer, and let $\mathcal{A}_{8d} = \sum_{i=1}^{8d} L_i$ be the pre-image under τ of an arrangement of $8d$ general lines in \mathbb{P}^2 (so it does not contain any singularity of $\mathcal{H}'_n + \sum_{i=0,1,\zeta,\infty} \mathcal{E}'_i$, and it is transversal to all of its curves).

For each i , we have that $\mathcal{E}_i \sim \frac{n^2-3}{3} F_i$, where F_i is the class of a fiber of π'_i , and so

$$3\mathcal{E}_i + F_{i,1} + F_{i,2} + F_{i,3} \sim n^2 F_i.$$

We define $a_0 = a_1 = b_i = 1$ for $1 \leq i \leq 4d$ and $a_\infty = a_\zeta = b_i = 2p - 1$ for $4d + 1 \leq i \leq 8d$. Let L be the pull-back of the class of a line. Then

$$\mathcal{O}_H \left(\sum_{i=0,1,\zeta,\infty} 3a_i \mathcal{E}_i + \sum_{i=0,1,\zeta,\infty} 3a_i \mathcal{E}'_i + \sum_{i=0,1,\zeta,\infty} a_i (F_{i,1} + F_{i,2} + F_{i,3}) + \sum_{i=1}^{8d} 3b_i L_i \right)$$

is isomorphic to \mathcal{L}_0^{4p} where

$$\mathcal{L}_0 := \mathcal{O}_H \left(6p(6\alpha^2 + \beta^2) \left(\sum_{i=0,1,\zeta,\infty} a_i F_i \right) + 6dL \right).$$

For each i , we denote by the same symbol the strict transform in Z_n of the divisors $\mathcal{E}_i, \mathcal{E}'_i, L_j, F_{i,j}$. Then

$$\mathcal{O}_{Z_n} \left(\sum_{i=0,1,\zeta,\infty} 3a_i \mathcal{E}_i + \sum_{i=0,1,\zeta,\infty} 3a_i \mathcal{E}'_i + \sum_{i=0,1,\zeta,\infty} a_i (F_{i,1} + F_{i,2} + F_{i,3}) + \sum_{i=1}^{8d} 3b_i L_i \right)$$

is \mathcal{L}_1^{4p} where $\mathcal{L}_1 := \varphi_n^*(\mathcal{L}_0) \otimes \mathcal{O}_{Z_n}(-3E)$, and E is the exceptional divisor of φ_n .

Consider the blow-up $\sigma_n: Y_n \rightarrow Z_n$ at the $4(n^2 - 3)$ 3-points of $\overline{\mathcal{H}}_n$. The exceptional divisor of σ_n is denoted by G . We denote again the strict transform of $\mathcal{E}_i, \mathcal{E}'_i, L_j, F_{i,j}, M, N_i, N$ in Y_n by the same symbol. Then we have

$$\mathcal{O}_{Y_n} \left(\sum_{i=0,1,\zeta,\infty} 3a_i \mathcal{E}_i + \sum_{i=0,1,\zeta,\infty} 3a_i \mathcal{E}'_i + \sum_{i=0,1,\zeta,\infty} 3a_i N_i + \sum_{i=1}^{8d} 3b_i L_i \right) \simeq \mathcal{L}^{4p},$$

where $\mathcal{L} := \sigma_n^*(\mathcal{L}_1) \otimes \mathcal{O}_{Y_n}(-M - 3G)$.

As in [EV92], this is the data to construct a $4p$ -th root cover of Y_n branch along the nodal curve

$$\sum_{i=0,1,\zeta,\infty} \mathcal{E}_i + \sum_{i=0,1,\zeta,\infty} \mathcal{E}'_i + \sum_{i=0,1,\zeta,\infty} N_i + \sum_{i=1}^{8d} L_i.$$

We now follow the description in [Urz08, Urz10]. Notice that the branch multiplicities are either 3 or $3(2p - 1)$, both coprime to $4p$.

We write

$$\sum_j \nu_j A_j := \sum_{i=0,1,\zeta,\infty} 3a_i \mathcal{E}_i + \sum_{i=0,1,\zeta,\infty} 3a_i \mathcal{E}'_i + \sum_{i=0,1,\zeta,\infty} 3a_i N_i + \sum_{i=1}^{8d} 3b_i L_i,$$

where the A_j are distinct irreducible curves and ν_j are the corresponding multiplicities. Set $A = \sum_j A_j$. The construction follows three steps:

$$X_n \xrightarrow{f_3} \text{Spec}_{Y_n} \left(\bigoplus_{i=0}^{4p-1} \mathcal{L}^{(i)-1} \right) \xrightarrow{f_2} \text{Spec}_{Y_n} \left(\bigoplus_{i=0}^{4p-1} \mathcal{L}^{-i} \right) \xrightarrow{f_1} Y_n,$$

where f_1 is the $4p$ -th root cover defined by the \mathcal{O}_{Y_n} -algebra $\bigoplus_{i=0}^{4p-1} \mathcal{L}^{-i}$ induced by $\mathcal{L}^{4p} \hookrightarrow \mathcal{O}_{Y_n}$, the morphism f_2 is the normalization, where

$$\mathcal{L}^{(i)} := \mathcal{L}^i \otimes \mathcal{O}_{Y_n} \left(- \sum_j \left[\frac{\nu_j i}{4p} \right] A_j \right)$$

for $i \in \{0, 1, \dots, 4p - 1\}$ and $[x]$ is the integral part of x , and f_3 is the minimal resolution of the singularities of $\text{Spec}_{Y_n} \left(\bigoplus_{i=0}^{4p-1} \mathcal{L}^{(i)-1} \right)$. The surface X_n is a nonsingular irreducible projective surface; cf. [Urz10, §1].

Let $f: X_n \rightarrow Y_n$ be the composition of the above morphisms. We have the \mathbb{Q} -numerical equivalence

$$K_{X_n} \equiv f^* \left(K_{Y_n} + \frac{(4p - 1)}{4p} A \right) + \Delta,$$

where Δ is a \mathbb{Q} -divisor supported on the exceptional divisor of f_3 .

Remark 5.1. In this construction, we can always arrange the multiplicities ν_i so that $0 < \nu_i < 4p$. At the end the corresponding surfaces, X_n are isomorphic over Y_n ; cf. [Urz10, §1]. For the sake of simplicity, in this paper we will not adjust multiplicities.

The following are key numbers to compute the Chern invariants.

Definition 5.2. Let q, m be coprime integers such that $0 < q < m$.

- (1) The associated *Hirzebruch-Jung continued fraction* is

$$\frac{m}{q} = e_1 - \frac{1}{e_2 - \frac{1}{\ddots - \frac{1}{e_l}}}$$

We denote its *length* as $l(q, m) := l$.

- (2) The *Dedekind sum* associated to the pair (q, m) is defined as

$$s(q, m) := \sum_{i=1}^{m-1} \left(\left(\frac{i}{m} \right) \right) \left(\left(\frac{iq}{m} \right) \right),$$

where $((x)) = x - [x] - \frac{1}{2}$ for any rational number x .

These numbers play a central role for the defect caused by the nodes of A in the Chern numbers of X_n . General connections between Dedekind sums and geometry can be found in [HZ74].

Remark 5.3. Each of the singularities in $\text{Spec}_{Y_n} \left(\bigoplus_{i=0}^{4p-1} \mathcal{L}^{(i)-1} \right)$ is over a node of A . Let $P \in A$ be a node in $A_i \cap A_j$. Then the singularity over P is locally isomorphic to the normalization of

$$(0, 0, 0) \in \{z^{4p} = x^{\nu_i} y^{\nu_j}\} \subset \mathbb{C}_{x,y,z}^3,$$

which is the cyclic quotient singularity $\frac{1}{4p}(1, q)$, where $0 < q < 4p$ with $\nu_i + q\nu_j \equiv 0 \pmod{4p}$. For details on these singularities, we refer to [BHPVdV04, III §5]. Let $R = R_1 + \dots + R_{l(q,4p)}$ be the prime decomposition of the exceptional (reduced) divisor over P . Let \tilde{A}_i, \tilde{A}_j be the strict pre-images of A_i, A_j in X_n locally over P . Then we have

$$f^*(A_i) = 4p\tilde{A}_i + \sum_{k=1}^{l(q,4p)} c_k R_k \quad \text{and} \quad f^*(A_j) = 4p\tilde{A}_j + \sum_{k=1}^{l(q,4p)} d_k R_k,$$

for certain integers $0 < c_k, d_k < 4p$, such that the discrepancy of R_k is (see, for example, [Urz08, §4.3.2], [Urz10] for precise numbers)

$$\text{disc}(R_k) = -1 + \frac{c_k}{4p} + \frac{d_k}{4p}.$$

We recall that $\text{disc}(R_k) \in]-1, 0] \cap \mathbb{Q}$ is the coefficient of R_k in Δ .

In our case, we have two types of singularities. We have singularities of type $\frac{1}{4p}(1, 4p - 1)$ over the nodes in a $A_i \cap A_j$ with $\nu_i = \nu_j$ and of type $\frac{1}{4p}(1, 2p + 1)$ over the other nodes.

PROPOSITION 5.4. *The surface X_n is simply connected.*

Proof. This is the same proof as in [Urz10, §8]; we recall it. Take a general point in $P \in L_1 \subset Y_n$. Through that point we have the trivial pencil of lines from \mathbb{P}^2 . Let $\psi: Y'_n \rightarrow Y_n$ be the blow-up of P . We now take the pulled-back $4p$ -th root cover $f': X'_n \rightarrow Y'_n$, coming from the $f: X_n \rightarrow Y_n$, so that we have the commutative diagram

$$\begin{array}{ccc} X'_n & \xrightarrow{f'} & Y'_n \\ \downarrow \psi' & & \downarrow \psi \\ X_n & \xrightarrow{f} & Y_n, \end{array}$$

where $\psi': X'_n \rightarrow X_n$ is the blow-down of a chain of $4p$ \mathbb{P}^1 's, starting with a (-1) -curve followed by $4p - 1$ (-2) -curves. Notice that X'_n has a fibration $h: X'_n \rightarrow \mathbb{P}^1$ with connected fibers, sections, and at least one simply connected fiber which comes from the line L_1 . (Singularities coming from points in L_1 are resolved through chains of smooth rational curves.) Then we apply [Xia91] to conclude that X'_n is simply connected, and so is X_n . For more details, see [Urz10, Prop.8.3]. \square

PROPOSITION 5.5. *The surface X_n is spin. In particular, it is minimal.*

Proof. By Proposition 5.4, the surface X_n is simply connected. Thus we only need to prove that K_{X_n} is numerically 2-divisible.

Let $\mathcal{E} := \sum_{i=0,1,\zeta,\infty} \mathcal{E}_i$ and $\mathcal{E}' := \sum_{i=0,1,\zeta,\infty} \mathcal{E}'_i$. We have

$$K_{X_n} \equiv f^* \left(-3L + N + E + 2G + \frac{(4p-1)}{4p} (\mathcal{E} + \mathcal{E}' + N + \mathcal{A}_{8d}) \right) + \Delta,$$

where Δ is a \mathbb{Q} -divisor supported on the exceptional locus of the singularities $\frac{1}{4p}(1, 2p+1)$ over the nodes of A , which are locally of the type $\{x^3y^{3(2p-1)}=0\}$ in $\sum_j \nu_j A_j$. For the other nodes, we have rational double points of type $\frac{1}{4p}(1, 4p-1)$, and so they do not contribute to Δ .

Notice that

$$\begin{aligned} & \mathcal{L}^{4p} - 2 \left(\mathcal{E}_0 + N_0 + \mathcal{E}'_0 + \mathcal{E}_1 + \mathcal{E}'_1 + N_1 + \sum_{i=1}^{4d} L_i \right) \\ & - (6p-4) \left(\mathcal{E}_\infty + N_\infty + \mathcal{E}'_\infty + \mathcal{E}_\zeta + N_\zeta + \mathcal{E}'_\zeta + \sum_{i=4d+1}^{8d} L_i \right) \sim \mathcal{E} + \mathcal{E}' + N + \mathcal{A}_{8d}. \end{aligned}$$

Then $f^* \left(\frac{(4p-1)}{4p} (\mathcal{E} + \mathcal{E}' + N + \mathcal{A}_{8d}) \right)$ is numerically equivalent to

$$\begin{aligned} & f^* \left((4p-1)\mathcal{L} \right) - 2(4p-1) \left(\bar{\mathcal{E}}_0 + \bar{\mathcal{E}}'_0 + \bar{N}_0 + \bar{\mathcal{E}}_1 + \bar{\mathcal{E}}'_1 + \bar{N}_1 + \sum_{i=1}^{4d} \bar{L}_i \right) \\ & - (6p-4)(4p-1) \left(\bar{\mathcal{E}}_\infty + \bar{\mathcal{E}}'_\infty + \bar{N}_\infty + \bar{\mathcal{E}}_\zeta + \bar{\mathcal{E}}'_\zeta + \bar{N}_\zeta + \sum_{i=4d+1}^{8d} \bar{L}_i \right) - \bar{\Delta}, \end{aligned}$$

where $\bar{\Gamma}$ denotes the strict (reduced) transform of a curve Γ under f , and $\bar{\Delta}$ is \mathbb{Q} -numerically effective and supported in the exceptional divisors from $\frac{1}{4p}(1, 4p - 1)$, and $\frac{1}{4p}(1, 2p + 1)$.

Over one $\frac{1}{4p}(1, 4p - 1)$, we have that $\bar{\Delta}$ is either $(6p - 4)(4p - 1)R$ or $2(4p - 1)R$ according to the multiplicities $\nu_i = \nu_j$ in the corresponding node, where R is the chain of $4p - 1$ (-2) -curves which resolves $\frac{1}{4p}(1, 4p - 1)$. In this case $\Delta = 0$, so $-\bar{\Delta} + \Delta$ is even.

The singularity $\frac{1}{4p}(1, 2p + 1)$ is resolved by a chain of three \mathbb{P}^1 's, say R_1, R_2, R_3 , such that $R_1^2 = R_3^2 = -2$ and $R_2^2 = -(p + 1)$. One can compute that over one $\frac{1}{4p}(1, 2p + 1)$, we have that

$$\bar{\Delta} = \frac{4p - 1}{4p} \left((10p - 2)R_1 + (12p - 4)R_2 + (12p^2 - 2p - 2)R_3 \right)$$

and $-\Delta = \frac{p-1}{2p}R_1 + \frac{p-1}{p}R_2 + \frac{p-1}{2p}R_3$. Therefore, $-\bar{\Delta} + \Delta \in \mathbb{Z}$ is even.

Thus we only need to prove that $f^*(-3L + N + E + 2G + (4p - 1)\mathcal{L})$ is divisible by 2. But

$$\mathcal{L} \sim 6p(6\alpha^2 + \beta^2)(a_0F_0 + a_1F_1 + a_\infty F_\infty + a_\zeta F_\zeta) - 3E - 3G - M + 6dL,$$

so we only need $f^*(-3L + N + G + M)$ to be divisible by 2. But $M + 3N + 3G \sim 9L$, since this is the pull-back of the dual Hesse arrangement. \square

The arrangement A has only 2-points, and its number is

$$t_2 = 13824\beta^2\alpha^2p^4 + 1152\beta^4p^4 + 4608d\alpha^2p^2 + 768d\beta^2p^2 + 32d^2 - 100d.$$

By Proposition 4.1, its log Chern numbers are

$$\bar{c}_1^2 = n^4 + 2t_2 - 40d - 48 \quad \text{and} \quad \bar{c}_2 = \frac{n^4}{3} + t_2 - 16d - 12.$$

For the Chern numbers, we have

$$c_1^2(X_n) = 4p\bar{c}_1^2 - 2\left(t_2 + 2\sum_j (g(A_j) - 1)\right) + \frac{1}{4p}\sum_j A_j^2 - \sum_{i < j} c(q_{i,j}, 4p)A_i \cdot A_j$$

and

$$c_2(X_n) = 4p\bar{c}_2 - \left(t_2 + 2\sum_j (g(A_j) - 1)\right) + \sum_{i < j} l(q_{i,j}, 4p)A_i \cdot A_j,$$

where $0 < q_{i,j} < 4p$ with $\nu_i + q_{i,j}\nu_j \equiv 0 \pmod{4p}$, and

$$c(q_{i,j}, 4p) := 12s(q_{i,j}, 4p) + l(q_{i,j}, 4p).$$

See [Urz08, §4.3] for proofs which work whenever the multiplicities of the branch divisor (in our case 3 and $2p - 3$ after reduction modulo $4p$) are coprime to the order of the cyclic cover (in our case $4p$).

We also have that $c(4p - 1, 4p) = \frac{4p-1}{2p}$ and $c(2p + 1, 4p) = \frac{2p^2+1}{2p}$ [Urz10, p. 345], and $l(4p - 1, 4p) = 4p - 1$ and $l(2p + 1, 4p) = 3$. Therefore,

$$\sum_{i < j} c(q_{i,j}, 4p) A_i \cdot A_j = \frac{(4p - 1)}{2p} t_{2,1} + \frac{(2p^2 + 1)}{2p} t_{2,2}$$

and

$$\sum_{i < j} l(q_{i,j}, 4p) A_i \cdot A_j = (4p - 1)t_{2,1} + 3t_{2,2},$$

where $t_{2,1}$ and $t_{2,2}$ are the number of 2-points corresponding to the singularities $\frac{1}{4p}(1, 4p - 1)$ and $\frac{1}{4p}(1, 2p + 1)$ respectively. Hence

$$t_{2,1} = 384\beta^4 p^4 + 4608\alpha^2 \beta^2 p^4 + 2304d\alpha^2 p^2 - 52d + 384d\beta^2 p^2 + 16d^2$$

and

$$t_{2,2} = 768\beta^4 p^4 + 9216\alpha^2 \beta^2 p^4 + 2304d\alpha^2 p^2 - 48d + 384d\beta^2 p^2 + 16d^2.$$

By plugging in the formulas above, we obtain that

$$\lim_{p \rightarrow \infty} \frac{c_1^2}{c_2} = \frac{108x^4 + 132x^2 + 11}{36x^4 + 96x^2 + 8} =: \lambda(x),$$

where $x = \alpha/\beta$. Notice that $[2, 3] \subset \lambda([0, \infty^+]) = [1.375, 3]$.

THEOREM 5.6. *For any number $r \in [1.375, 3]$, there are spin simply connected minimal surfaces of general type X with $c_1^2(X)/c_2(X)$ arbitrarily close to r .*

Proof. Let $\varepsilon > 0$. Then there exists a positive rational number α/β such that $|\lambda(\alpha/\beta) - r| < \varepsilon$. We take those α, β for the construction above. The surfaces X_n are spin by Proposition 5.5 and simply connected by Proposition 5.4. Using Enriques' classification of surfaces, it is clear that the surfaces X_n are of general type for $p \gg 0$, since $c_1^2(X_n)$ and $c_2(X_n)$ approach $+\infty$ as p tends to $+\infty$. □

Example 5.7. For parameters $\alpha = 1, \beta = 0, d = 1$, and $p = 5$, we obtain a spin simply connected surface with $c_1^2 = 2^6 \cdot 3^2 \cdot 455393$ and $c_2 = 2^5 \cdot 3^3 \cdot 103997$, thus with Chern slope around 2.919. For $p = 100003$, we get a Chern slope close to 3 up to $2 \cdot 10^{-10}$.

6. Nonspin surfaces and other fundamental groups

For nonspin surfaces, we consider the same set up as in Section 5.5 but for a p -th root cover. We keep same notation as in the previous section up to some explicitly redefined numbers and symbols.

Let $p \geq 5$ be a prime number, and let $\alpha > 0, \beta > 0$ be integers. Let $n = 3\alpha p$. As before, we consider the arrangement \mathcal{H}'_n in H . Let \mathcal{E}'_i be $\beta^2 p^2$

general fibers of π'_i , and let $\mathcal{A}_{2d} = L_1 + \cdots + L_{2d}$ be the strict transform of an arrangement of $2d$ general lines in \mathbb{P}^2 , where $3 \leq 2d \leq p$. We define $a_0 = a_1 = b_i = 1$ for $1 \leq i \leq d$ and $a_\infty = a_\zeta = b_i = p - 1$ for $d + 1 \leq i \leq 2d$. Then

$$\mathcal{O}_H \left(\sum_{i=0,1,\zeta,\infty} 3a_i \mathcal{E}_i + \sum_{i=0,1,\zeta,\infty} 3a_i \mathcal{E}'_i + \sum_{i=0,1,\zeta,\infty} a_i (F_{i,1} + F_{i,2} + F_{i,3}) + \sum_{i=1}^{2d} 3b_i L_i \right)$$

is isomorphic to \mathcal{L}_0^p , where

$$\mathcal{L}_0 := \mathcal{O}_H \left(3p(3\alpha^2 + \beta^2) \left(\sum_{i=0,1,\zeta,\infty} a_i F_i \right) + 3dL \right).$$

For each i , we denote the strict transform of $\mathcal{E}_i, \mathcal{E}'_i, L_j, F_{i,j}$ in Z_n by the same symbol. Then

$$\mathcal{O}_{Z_n} \left(\sum_{i=0,1,\zeta,\infty} 3a_i \mathcal{E}_i + \sum_{i=0,1,\zeta,\infty} 3a_i \mathcal{E}'_i + \sum_{i=0,1,\zeta,\infty} a_i (F_{i,1} + F_{i,2} + F_{i,3}) + \sum_{i=1}^{2d} 3b_i L_i \right)$$

is \mathcal{L}_1^p , where $\mathcal{L}_1 := \varphi_n^*(\mathcal{L}_0) \otimes \mathcal{O}_{Z_n}(-6E)$, and E is the exceptional divisor of φ_n . Again, we denote the strict transform of $\mathcal{E}_i, \mathcal{E}'_i, L_j, F_{i,j}, M, N_i, N$ in Y_n by the same symbol. Then we have

$$\mathcal{O}_{Y_n} \left(\sum_{i=0,1,\zeta,\infty} 3a_i \mathcal{E}_i + \sum_{i=0,1,\zeta,\infty} 3a_i \mathcal{E}'_i + \sum_{i=0,1,\zeta,\infty} 3a_i N_i + \sum_{i=1}^{2d} 3b_i L_i \right) \simeq \mathcal{L}^p,$$

where $\mathcal{L} := \sigma_n^*(\mathcal{L}_1) \otimes \mathcal{O}_{Y_n}(-2M - 6G)$. With this data, we construct a p -th root cover of Y_n branch along

$$A := \sum_{i=0,1,\zeta,\infty} \mathcal{E}_i + \sum_{i=0,1,\zeta,\infty} \mathcal{E}'_i + \sum_{i=0,1,\zeta,\infty} N_i + \sum_{i=1}^{2d} L_i.$$

Let $f: X_p \rightarrow Y_n$ be the corresponding morphism for the p -th root cover, as in the previous section.

PROPOSITION 6.1. *The surfaces X_p are simply connected.*

Proof. This is the same as Proposition 5.4 using \mathcal{A}_{2d} . □

Let us write

$$\sum_j \nu_j A_j = \sum_{i=0,1,\zeta,\infty} 3a_i \mathcal{E}_i + \sum_{i=0,1,\zeta,\infty} 3a_i \mathcal{E}'_i + \sum_{i=0,1,\zeta,\infty} 3a_i N_i + \sum_{i=1}^{2d} 3b_i L_i,$$

where A_j are the irreducible curves in A . Hence ν_j is equal to either $3a_i$ or $3b_k$ for some i, k . The absence of \mathbb{P}^1 's coming from k -points in the branch divisor allows us to prove the following [Urz08, IV §4.3.2].

PROPOSITION 6.2. *The surfaces X_p are minimal.*

Proof. The argument will be two parts. First we show that K_{X_p} is \mathbb{Q} -numerically equal to an effective divisor supported on the pre-image of $A + E + G$ and the exceptional divisor of f_3 . (See construction of f in the previous section.) The second part is to show that no curve in the pre-image of $A + E + G$ is a (-1) -curve.

We know that $K_{X_p} \equiv f^*\left(K_{Y_n} + \frac{(p-1)}{p}A\right) + \Delta$, where Δ is supported on the exceptional divisor of f_3 . We have that $K_{Y_n} \sim -3L + N + E + 2G$ and

$$-\frac{1}{4}(F_0 + F_1 + F_\infty + F_\zeta) \equiv -3L + \frac{3}{4}N + \frac{3}{4}G.$$

If $u = \frac{p-1}{p}$ and $v = \beta^2 p^2$, then K_{X_p} is numerically \mathbb{Q} -equivalent to

$$f^*\left(\sum_{i=0,1,\zeta,\infty} u\mathcal{E}_i + \sum_{i=0,1,\zeta,\infty} \left(u - \frac{1}{4v}\right)\mathcal{E}'_i + \left(u + \frac{1}{4}\right)N + u\mathcal{A}_{2d} + E + \frac{5}{4}G\right) + \Delta.$$

On the other hand, for a node $P \in A_i \cap A_j$, we locally have

$$f^*(A_i + A_j) = p\tilde{A}_i + p\tilde{A}_j + \sum_{k=1}^l (c_k + d_k)R_k$$

as in Remark 5.3, where $R_1 + \dots + R_l$ is the exceptional divisor above P , and $\text{discr}(R_k) = -1 + \frac{c_k}{p} + \frac{d_k}{p}$. We clearly have

$$xc_k + yd_k - 1 + \frac{c_k}{p} + \frac{d_k}{p} > 0,$$

where $x, y \in \{u, u - \frac{1}{4v}, u + \frac{1}{4}\}$. Therefore, K_{X_p} is \mathbb{Q} -numerically effective with support equal to the pre-image of $A + E + G$ and the exceptional divisor of f_3 .

Notice that in this support the only \mathbb{P}^1 's are the ones coming from N , lines in \mathcal{A}_{2d} , and Δ . The ones in Δ are not (-1) -curves. For the ones in N , we have \mathbb{P}^1 's intersecting $\sum_{i=0,1,\zeta,\infty} \mathcal{E}'_i$ at $3v$ points. Similarly with the \mathbb{P}^1 's coming from lines L_j , we have intersection at $4(n^2 - 3) + 12v$ points with $\sum_{i=0,1,\zeta,\infty} \mathcal{E}_i + \sum_{i=0,1,\zeta,\infty} \mathcal{E}'_i$. Using the notation from Remark 5.3, for the strict pre-image \tilde{A}_j of A_j under f , we have (see [Urz08, IV §4.3.2])

$$\tilde{A}_j^2 = \frac{1}{p}\left(A_j^2 - \sum_{i \neq j} q_{i,j}A_i \cdot A_j\right) < -1.$$

Therefore, we have no (-1) -curves in the support. □

The arrangement A has only 2-points, and its number is

$$t_2 = 108\alpha^2\beta^2p^4 + 18\beta^4p^4 + 72d\alpha^2p^2 - 25d + 24d\beta^2p^2 + 2d^2.$$

By Proposition 4.1, its log Chern numbers are

$$\bar{c}_1^2 = n^4 + 2t_2 - 10d - 48 \quad \text{and} \quad \bar{c}_2 = \frac{n^4}{3} + t_2 - 4d - 12.$$

We also have that $c(p - 1, p) = \frac{2p-2}{p}$ and $c(1, p) = \frac{p^2-2p+2}{p}$ [Urz10, p.345], and $l(p - 1, p) = p - 1$ and $l(1, p) = 1$. Therefore,

$$\sum_{i < j} c(q_{i,j}, 4p) A_i \cdot A_j = \frac{(2p - 2)}{p} t_{2,1} + \frac{(p^2 - 2p + 2)}{p} t_{2,2}$$

and

$$\sum_{i < j} l(q_{i,j}, 4p) A_i \cdot A_j = (p - 1) t_{2,1} + t_{2,2},$$

where $t_{2,1}$ and $t_{2,2}$ are the number of 2-points corresponding to the singularities $\frac{1}{p}(1, p - 1)$ and $\frac{1}{p}(1, 1)$ respectively. Hence

$$t_{2,1} = 6\beta^4 p^4 + 36\alpha^2 \beta^2 p^4 + 36d\alpha^2 p^2 - 13d + 12d\beta^2 p^2 + d^2$$

and

$$t_{2,2} = 12\beta^4 p^4 + 72\alpha^2 \beta^2 p^4 + 36d\alpha^2 p^2 - 12d + 12d\beta^2 p^2 + d^2.$$

By plugging in the formulas for Chern numbers in Section 5.5, we obtain that

$$\lim_{p \rightarrow \infty} \frac{c_1^2}{c_2} = \frac{27x^4 + 48x^2 + 8}{9x^4 + 48x^2 + 8} =: \lambda(x),$$

where $x = \alpha/\beta$. Notice that $[2, 3] \subset \lambda([0, \infty^+]) = [1, 3]$.

THEOREM 6.3. *For any number $r \in [1, 3]$, there are nonspin simply connected minimal surfaces of general type X with $c_1^2(X)/c_2(X)$ arbitrarily close to r .*

Proof. Let $\varepsilon > 0$. Then there exists a positive rational number α/β such that $|\lambda(\alpha/\beta) - r| < \varepsilon$. We take those α, β for the construction above. The surfaces X_p are minimal by Proposition 6.2, simply connected by Proposition 6.1, and nonspin because of the degree of f . The surfaces X_p are of general type for $p \gg 0$. □

Let $q > 0$ be an integer, and let $\rho: C_q \rightarrow \mathbb{P}^1$ be a $(q + 1)$ -cyclic cover completely branch at four general points. Hence the genus of C_q is q .

COROLLARY 6.4. *For any number $r \in [1, 3]$, there are minimal surfaces of general type X with $c_1^2(X)/c_2(X)$ arbitrarily close to r , and $\pi_1(X) \simeq \pi_1(C_q)$.*

Proof. Consider the X_p in Theorem 6.3. Take a general point in $P \in L_1 \subset Y_n$. Through that point we have the trivial pencil of lines from \mathbb{P}^2 . As we did in Proposition 5.4, let $\psi: Y'_n \rightarrow Y_n$ be the blow-up of P . We now take the pulled-back p -th root cover $f': X'_p \rightarrow Y'_n$, coming from the $f: X_p \rightarrow Y_n$, so

that we have the commutative diagram

$$\begin{array}{ccc} X'_p & \xrightarrow{f'} & Y'_n \\ \downarrow \psi' & & \downarrow \psi \\ X_p & \xrightarrow{f} & Y_n, \end{array}$$

where $\psi' : X'_p \rightarrow X_p$ is the blow-down of a chain of p \mathbb{P}^1 's, starting with a (-1) -curve followed by $p - 1$ (-2) -curves. Notice that X'_p has a fibration $h : X'_p \rightarrow \mathbb{P}^1$ with connected fibers, sections, and at least one simply connected fiber which comes from the line L_1 . The genus g of the general fiber of h satisfies the Riemann-Hurwitz formula $2g - 2 = p(-2) + (p - 1)(4(n^2 - 3) + 12\beta^2 p^2 + 2d)$. Consider the base change fibration

$$\begin{array}{ccc} X & \xrightarrow{\rho'} & X'_p \\ \downarrow h' & & \downarrow h \\ C_q & \xrightarrow{\rho} & \mathbb{P}^1, \end{array}$$

where the surface X is nonsingular projective of general type such that

$$\frac{c_1^2(X)}{c_2(X)} = \frac{(q + 1)(c_1^2(X_p) - p) + 16q(g - 1)}{(q + 1)(c_2(X_p) + p) + 8q(g - 1)},$$

and so, since the highest power of p in $g - 1$ is p^3 , $\frac{c_1^2(X)}{c_2(X)}$ approaches $\lambda(\alpha/\beta)$ as p tends to infinity. For minimality, we notice that

$$K_X \equiv \rho'^* \left(\psi'^*(K_{X_p}) + \sum_{i=1}^p iS_i + \frac{4q}{q + 1} F \right),$$

where $\sum_{i=1}^p S_i$ is the chain of \mathbb{P}^1 explained above, with $S_p^2 = -1$, and F is a general fiber of h . We know that K_{X_p} is \mathbb{Q} -effective. Therefore, K_X is \mathbb{Q} -effective. One can verify that its support does not contain (-1) -curves and so X is minimal. Also, notice that h' has sections, for instance, the pre-image of S_p , and it has some simply connected fibers from the pre-image of the one in h . Hence, by [Xia91], we have that $\pi_1(X) \simeq \pi_1(C_q)$. \square

Remark 6.5. The arrangement of either $8d$ or $2d$ general lines in the constructions above gives not only simply connectedness but also shows that the constructed surfaces have unbounded deformations. It also exhibits some freedom for the constructions, where we can add branching curves in such a way that they either do not depend on p or do depend on p but affecting Chern numbers in a suitable power of p .

References

- [BHPVdV04] W. P. BARTH, K. HULEK, C. A. M. PETERS, and A. VAN DE VEN, *Compact Complex Surfaces*, second ed., *Ergeb. Math. Grenzgeb.* **4**, Springer-Verlag, New York, 2004. MR 2030225. Zbl 1036.14016.
- [Bog78] F. A. BOGOMOLOV, Holomorphic tensors and vector bundles on projective manifolds, *Izv. Akad. Nauk SSSR Ser. Mat.* **42** (1978), 1227–1287, 1439, in Russian; translated in *Math. USSR-Izv.* **13** (1979), 499–555. MR 0522939. Zbl 0439.14002. <http://dx.doi.org/10.1070/IM1979v013n03ABEH002076>.
- [CC93] F. CATANESE and C. CILIBERTO, On the irregularity of cyclic coverings of algebraic surfaces, in *Geometry of Complex Projective Varieties* (Cetraro, 1990), *Sem. Conf.* **9**, Mediterranean, Rende, 1993, pp. 89–115. MR 1225590. Zbl 0937.14007.
- [Che87] Z. J. CHEN, On the geography of surfaces. Simply connected minimal surfaces with positive index, *Math. Ann.* **277** (1987), 141–164. MR 0884652. Zbl 0595.14027. <http://dx.doi.org/10.1007/BF01457284>.
- [EV92] H. ESNAULT and E. VIEHWEG, *Lectures on Vanishing Theorems, DMV Seminar* **20**, Birkhäuser, Boston, 1992. MR 1193913. Zbl 0779.14003. <http://dx.doi.org/10.1007/978-3-0348-8600-0>.
- [Hir84] F. HIRZEBRUCH, Chern numbers of algebraic surfaces: an example, *Math. Ann.* **266** (1984), 351–356. MR 0730175. Zbl 0504.14030. <http://dx.doi.org/10.1007/BF01475584>.
- [HZ74] F. HIRZEBRUCH and D. ZAGIER, *The Atiyah-Singer Theorem and Elementary Number Theory, Math. Lecture Series* **3**, Publish or Perish, Boston, 1974. MR 0650832. Zbl 0288.10001.
- [Hol86] R.-P. HOLZAPFEL, Chern numbers of algebraic surfaces—Hirzebruch’s examples are Picard modular surfaces, *Math. Nachr.* **126** (1986), 255–273. MR 0846579. Zbl 0625.14018. <http://dx.doi.org/10.1002/mana.19861260117>.
- [Hol78] R.-P. HOLZAPFEL, *Arithmetische Kugelquotientenflächen, Seminarberichte [Seminar Reports]* **14**, Humboldt Universität, Sektion Mathematik, Berlin, 1978, II. Klassifikation von Kugelspitzen- und Kugelquotientensingularitäten. [II. Classification of conic double-point and quotient singularities of spheres], with English and Russian summaries. MR 0521083. Zbl 0417.14038.
- [Hol79a] R.-P. HOLZAPFEL, *Arithmetische Kugelquotientenflächen, Seminarberichte [Seminar Reports]* **20**, Humboldt Universität, Sektion Mathematik, Berlin, 1979, IV. Dimensionsformeln für Räume von Spitzenformen. [IV. Dimension formulas for spaces of cusp forms], with English and Russian summaries. MR 0538624. Zbl 0417.14039.
- [Hol79b] R.-P. HOLZAPFEL, *Arithmetische Kugelquotientenflächen, Seminarberichte [Seminar Reports]* **21**, Humboldt Universität, Sektion Mathematik, Berlin, 1979, VI. Kugelquotientenflächen vom allgemeinen Typ.

- [VI. Spherical quotient surfaces of general type], with English and Russian summaries. MR 0550140. Zbl 0477.14021.
- [Hol81] R.-P. HOLZAPFEL, Arithmetic surfaces with great K^2 , in *Proceedings of the Week of Algebraic Geometry* (Bucharest, 1980), *Teubner-Texte Math.* **40**, Teubner, Leipzig, 1981, pp. 80–91. MR 0712517. Zbl 0485.14008.
- [KH05] A. KASPARIAN and R.-P. HOLZAPFEL, Arithmetic proportional elliptic configurations with comparatively large number of irreducible components, in *Geometry, Integrability and Quantization*, Softex, Sofia, 2005, pp. 252–261. MR 2161772. Zbl 1158.14311.
- [Miy77] Y. MIYAOKA, On the Chern numbers of surfaces of general type, *Invent. Math.* **42** (1977), 225–237. MR 0460343. Zbl 0374.14007. <http://dx.doi.org/10.1007/BF01389789>.
- [Miy83] Y. MIYAOKA, Algebraic surfaces with positive indices, in *Classification of Algebraic and Analytic Manifolds* (Katata, 1982), *Progr. Math.* **39**, Birkhäuser, Boston, 1983, pp. 281–301. MR 0728611. Zbl 0584.14022.
- [MT87] B. MOISHEZON and M. TEICHER, Simply-connected algebraic surfaces of positive index, *Invent. Math.* **89** (1987), 601–643. MR 0903386. Zbl 0627.14019. <http://dx.doi.org/10.1007/BF01388987>.
- [PP96] U. PERSSON and C. PETERS, Some aspects of the topology of algebraic surfaces, in *Proceedings of the Hirzebruch 65 Conference on Algebraic Geometry* (Ramat Gan, 1993), *Israel Math. Conf. Proc.* **9**, Bar-Ilan Univ., Ramat Gan, 1996, pp. 377–392. MR 1360515. Zbl 0862.14011.
- [Per81] U. PERSSON, Chern invariants of surfaces of general type, *Compositio Math.* **43** (1981), 3–58. MR 0631426. Zbl 0479.14018. Available at http://www.numdam.org/item?id=CM_1981__43_1_3_0.
- [Per87] U. PERSSON, An introduction to the geography of surfaces of general type, in *Algebraic Geometry, Bowdoin, 1985* (Brunswick, Maine, 1985), *Proc. Sympos. Pure Math.* **46**, Amer. Math. Soc., Providence, RI, 1987, pp. 195–218. MR 0927957. Zbl 0656.14020.
- [PPX96] U. PERSSON, C. PETERS, and G. XIAO, Geography of spin surfaces, *Topology* **35** (1996), 845–862. MR 1404912. Zbl 0874.14031. [http://dx.doi.org/10.1016/0040-9383\(95\)00046-1](http://dx.doi.org/10.1016/0040-9383(95)00046-1).
- [Som84] A. J. SOMMESE, On the density of ratios of Chern numbers of algebraic surfaces, *Math. Ann.* **268** (1984), 207–221. MR 0744608. Zbl 0521.14016. <http://dx.doi.org/10.1007/BF01456086>.
- [Urz08] G. A. URZUA, Arrangements of curves and algebraic surfaces, 2008, ProQuest LLC, Ann Arbor, MI, Thesis (Ph.D.)—University of Michigan. MR 2712257. Available at http://gateway.proquest.com/openurl?url_ver=Z39.88-2004&rft_val_fmt=info:ofi/fmt:kev:mtx:dissertation&res_dat=xri:pqdiss&rft_dat=xri:pqdiss:3328979.
- [Urz10] G. A. URZÚA, Arrangements of curves and algebraic surfaces, *J. Algebraic Geom.* **19** (2010), 335–365. MR 2580678. Zbl 1192.14033. <http://dx.doi.org/10.1090/S1056-3911-09-00520-7>.

- [Urz11] G. A. URZÚA, Arrangements of rational sections over curves and the varieties they define, *Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei Mat. Appl.* **22** (2011), 453–486. MR 2904994. Zbl 06007192.
- [VdV78] A. VAN DE VEN, Some recent results on surfaces of general type, in *Séminaire Bourbaki, 29e année (1976/77), Lecture Notes in Math.* **677**, Springer-Verlag, New York, 1978, pp. Exp. No. 500, pp. 155–166. MR 0521766. Zbl 0432.14023.
- [Xia91] G. XIAO, π_1 of elliptic and hyperelliptic surfaces, *Internat. J. Math.* **2** (1991), 599–615. MR 1124285. Zbl 0762.14013. <http://dx.doi.org/10.1142/S0129167X91000338>.
- [Yau77] S. T. YAU, Calabi’s conjecture and some new results in algebraic geometry, *Proc. Nat. Acad. Sci. U.S.A.* **74** (1977), 1798–1799. MR 0451180. Zbl 0355.32028. Available at <http://www.pnas.org/content/74/5/1798.full.pdf+html>.

(Received: February 26, 2014)

(Revised: October 24, 2014)

UNIVERSITÉ DE POITIERS, POITIERS, FRANCE
E-mail: Xavier.Roulleau@math.univ-poitiers.fr

PONTIFICIA UNIVERSIDAD CATÓLICA DE CHILE, SANTIAGO, CHILE
E-mail: urzua@mat.puc.cl