

Universal hierarchical structure of quasiperiodic eigenfunctions

By SVETLANA JITOMIRSKAYA and WENCAI LIU

Abstract

We determine exact exponential asymptotics of eigenfunctions and of corresponding transfer matrices of the almost Mathieu operators for all frequencies in the localization regime. This uncovers a universal structure in their behavior, governed by the continued fraction expansion of the frequency, explaining some predictions in physics literature. In addition it proves the arithmetic version of the frequency transition conjecture. Finally, it leads to an explicit description of several non-regularity phenomena in the corresponding non-uniformly hyperbolic cocycles, which is also of interest as both the first natural example of some of those phenomena and, more generally, the first non-artificial model where non-regularity can be explicitly studied.

1. Introduction

A very captivating question and a longstanding theoretical challenge in solid state physics is to determine/understand the hierarchical structure of spectral features of operators describing 2D Bloch electrons in perpendicular magnetic fields, as related to the continued fraction expansion of the magnetic flux. Such structure was first predicted in the work of Azbel in 1964 [Azb64]. It was numerically confirmed through the famous butterfly plot and further argued for by Hofstadter in [Hof76] for the spectrum of the almost Mathieu operator. This was even before this model was linked to the integer quantum Hall effect [Tho84] and other important phenomena. Mathematically, it is known that the spectrum is a Cantor set for all irrational fluxes [AJ09] and, moreover, even all gaps predicted by the gap labeling are open in the non-critical case [AJ10], [AYZ]. Both were very important challenges in themselves, even though these results, while strongly indicated, do not describe or explain the hierarchical structure, and the problem of its description/explanation remains

Keywords: Anderson localization, spectral transition, universal hierarchical structure

AMS Classification: Primary: 47B36; Secondary: 37C55, 82B26.

© 2018 Department of Mathematics, Princeton University.

open, even in physics. As for the understanding the hierarchical behavior of the eigenfunctions, despite significant numerical studies and even a discovery of Bethe Ansatz solutions [WZ94], [ATW98] it has also remained an important open challenge even at the physics level. Certain results indicating the hierarchical structure in the corresponding semi-classical/perturbative regimes were previously obtained in the works of Sinai, Helffer-Sjöstrand, and Buslaev-Fedotov. (See [HS89], [Fed13], [Sin87], and also [Zhi91] for a different model.)

In this paper we address the latter problem by describing the universal self-similar exponential structure of eigenfunctions throughout the entire localization regime. We determine explicit universal functions $f(k)$ and $g(k)$, depending only on the Lyapunov exponent and the position of k in the hierarchy defined by the denominators q_n of the continued fraction approximants of the flux α , that completely define the exponential behavior of, correspondingly, eigenfunctions and norms of the transfer matrices of the almost Mathieu operators for all eigenvalues corresponding to almost every phase; see Theorem 2.1.¹ Our result holds for *all* frequency and coupling pairs in the localization regime. Since the behavior is fully determined by the frequency and does not depend on the phase, it is the same, eventually, around any starting point, so it is also seen unfolding at different scales when magnified around local eigenfunction maxima, thus describing the exponential universality in the hierarchical structure; see, for example, Theorems 2.2 and 2.4.

While the almost Mathieu family is precisely the one of main interest in physics literature, it also presents the simplest case of an analytic quasiperiodic operator, so a natural question is which features discovered for the almost Mathieu would hold for this more general class. Not all do; in particular, the ones that exploit the self-dual nature of the family $H_{\lambda,\alpha,\theta}$ often cannot be expected to hold in general. In case of Theorems 2.1 and 2.2, we conjecture that they should in fact hold for general analytic (or even more general) potentials, for almost every phase and with $\ln|\lambda|$ replaced by the Lyapunov exponent $L(E)$ (see footnote 4), but with otherwise the same or very similar statements. The hierarchical structure theorems 2.2 and 2.4 are also expected to hold universally for most (albeit not all, as in the present paper) appropriate local maxima. Some of our qualitative corollaries may hold in even higher generality. Establishing this fully would require certain new ideas since so far even an arithmetic version of localization for the Diophantine case has not been established for the general analytic family, the current state-of-the-art result by Bourgain-Goldstein [BG00] being measure theoretic in α . However, some

¹This paper supplants our earlier preprint entitled “Asymptotics of quasiperiodic eigenfunctions.” The latter preprint is not intended for publication.

ideas of our method can already be transferred to general trigonometric polynomials [JL]. Moreover, our method was used recently in [HJY] to show that the same f and g govern the asymptotics of eigenfunctions and universality around the local maxima throughout the a.e. localization regime in another popular object, the Maryland model.

Since we are interested in exponential growth/decay, the behavior of f and g becomes most interesting in the case of frequencies with exponential rate of approximation by the rationals. In general, localization for quasiperiodic operators is a classical case of a small denominator problem and has been traditionally approached in a perturbative way: through KAM-type schemes for large couplings [Sin87], [FSW90], [Eli97] (which, being KAM-type schemes, all required Diophantine conditions on frequencies) or through perturbation of periodic operators (Liouville frequency). Unlike the random case, where, in dimension one, localization holds for all couplings, a distinctive feature of quasiperiodic operators is the presence of metal-insulator transitions as couplings increase. Even when non-perturbative methods, for the almost Mathieu and then for general analytic potentials, were developed in the 1990s [Jit99], [BG00], allowing to obtain localization for almost every frequency throughout the regime of positive Lyapunov exponents, they still required Diophantine conditions, and exponentially approximated frequencies that are neither far from nor close enough to rationals remained a challenge, as for them there was nothing left to perturb about or to remove. Moreover, it has become clear that for all frequencies, the true localization threshold should be arithmetically determined and happen precisely where the exponential growth provided by the Lyapunov exponent beats the exponential strength of the small denominators. Thus the most interesting regime — the neighborhood of the transition — required dealing with the exponential frequencies not amenable to perturbations/parameter removals, adding a strong number theoretic flavor to the problem. The precise second transition conjecture was stated for the almost Mathieu operator [Jit95]. Our analysis provides also a (constructive) solution to the full arithmetic version of the transition in frequency and explains the role of frequency resonances in the phenomenon of localization in a sharp way.

The almost Mathieu operator (AMO) is the quasiperiodic Schrödinger operator on $\ell^2(\mathbb{Z})$:

$$(H_{\lambda,\alpha,\theta}u)(n) = u(n+1) + u(n-1) + 2\lambda \cos 2\pi(\theta + n\alpha)u(n),$$

where λ is the coupling, α is the frequency, and θ is the phase.

It is the central quasiperiodic model due to coming from physics and attracting continued interest there. First appearing in Peierls [Pei33], it arises as related, in two different ways, to a two-dimensional electron subject to a perpendicular magnetic field and plays a central role in the Thouless et al.

theory of the integer quantum Hall effect. For further background, history, and surveys of results, see [MJ17], [Dam17], [Jit07], [Las05] and references therein.

The Almost Mathieu operator has a transition from zero to positive Lyapunov exponents on the spectrum at $|\lambda| = 1$ (the critical coupling) leading to the conjecture, dating back to [AA80], that it induces a transition from absolutely continuous to pure point spectrum. For Diophantine α , this was proved in [Jit99]. The result was extended to all α, θ for $|\lambda| < 1$ (the subcritical regime) in [Avi08], solving one of the Simon's problems [Sim00]. For the supercritical regime ($|\lambda| > 1$), it is known however that the nature of the spectrum should depend on the arithmetic properties of α [Gor76], [AS82].

Set

$$(1) \quad \beta = \beta(\alpha) = \limsup_{n \rightarrow \infty} \frac{\ln q_{n+1}}{q_n},$$

where $\frac{p_n}{q_n}$ are the continued fraction approximants of α .

For any irrational number α , we say that the phase $\theta \in (0, 1)$ is Diophantine with respect to α if there exist $\kappa > 0$ and $\nu > 0$ such that

$$(2) \quad \||2\theta + k\alpha\|_{\mathbb{R}/\mathbb{Z}} > \frac{\kappa}{|k|^\nu}$$

for any $k \in \mathbb{Z} \setminus \{0\}$, where $\|x\|_{\mathbb{R}/\mathbb{Z}} = \text{dist}(x, \mathbb{Z})$. Clearly, for any irrational number α , the set of phases that are Diophantine with respect to α is of full Lebesgue measure. The conjecture in [Jit95] states that for α -Diophantine (thus almost every) θ , $H_{\lambda, \alpha, \theta}$ satisfies Anderson localization (i.e., has only pure point spectrum with exponentially decaying eigenfunctions) if $|\lambda| > e^\beta$ and has, for all θ , purely singular continuous spectrum for $1 < |\lambda| < e^\beta$.²

For $\beta = 0$, this follows from [Jit99]. A progress towards the localization side of the above conjecture was made in [AJ09] (localization for $|\lambda| > e^{\frac{16}{9}\beta}$, as a step in solving the Ten Martini problem) and in [YZ13] (in a limited sense for $|\lambda| > Ce^\beta$). The method developed in [AJ09] that allowed one to approach exponentially small denominators on the localization side was brought to its technical limits in [LY15a], where the result for $|\lambda| > e^{\frac{3}{2}\beta}$ was obtained.

Lately, with the development of Avila's global theory and the proof of the almost reducibility conjecture [Avi10], it has become possible to obtain non-perturbative reducibility directly ([AFK11], [HY12], [YZ13]), allowing one to potentially argue localization for the dual model by duality, as was first done in a perturbative regime in [BLT83], avoiding the localization method

² The original conjecture is slightly stronger in that it allows for not just polynomial, but any subexponential approximation of 2θ by $k\alpha$. The same goes for our proof, with obvious modifications. We choose to present the result, and thus also present the conjecture, for a slightly stronger Diophantine case in order to slightly simplify the argument.

completely. This was done recently by Avila-You-Zhou [AYZ17] who proved the full singular continuous part of the conjecture and a measure-theoretic (i.e., almost all θ) version of the pure point part. (See also [JK16], where a simple alternative way to argue completeness in the duality argument was presented.) The measure-theoretic (in phase) nature of the pure point result of [AYZ17] is, in fact, inherent in the duality argument. In contrast, our analysis provides a direct constructive proof for an arithmetically defined set of α -Diophantine θ , thus proving the full arithmetic version of the conjecture.

Our method can be used to also obtain *precise* asymptotics of *arbitrary* solutions of $H_{\lambda,\alpha,\theta}\varphi = E\varphi$, where E is an eigenvalue. Combined with the arguments of Last-Simon [LS99], this allows us to find precise asymptotics of the norms of the transfer-matrices, providing the first example of this sort for non-uniformly hyperbolic dynamics. Since those norms sometimes differ significantly from the reciprocals of the eigenfunctions, this leads to further interesting and unusual consequences; for example, exponential tangencies between contracted and expanded directions at the resonant sites.

From this point of view, our analysis also provides, as far as we know, the first study of the dynamics of Lyapunov-Perron non-regular points, in a natural setting. An artificial example of irregular dynamics can be found in [BP07, p. 23], however it is not even a cocycle over an ergodic transformation, and we are not aware of other such, even artificial, ergodic examples where the dynamics has been studied. Loosely, for a cocycle A over a transformation f acting on a space X (Lyapunov-Perron), non-regular points $x \in X$ are the ones at which Oseledets multiplicative ergodic theorem does not hold coherently in both directions. They therefore form a measure zero set with respect to any invariant measure on X .³ Yet, it is precisely the non-regular points that are of interest in the study of Schrödinger cocycles in the non-uniformly hyperbolic (positive Lyapunov exponent) regime, since spectral measures, for every fixed phase, are always supported on energies where there exists a solution polynomially bounded in both directions, so the (hyperbolic) cocycle defined at such energies is always non-regular at precisely the relevant phases. Thus the non-regular points capture the entire action from the point of view of spectral theory and so become the most important ones to study. One can also discuss stronger non-regularity notions: absence of forward regularity and, even stronger, non-exactness of the Lyapunov exponent [BP07]. While it is not difficult to see that energies in the support of singular continuous spectral measure in the non-uniformly hyperbolic regime always provide examples of non-exactness, our analysis gives the first non-trivial example of non-exactness

³Although in the uniformly hyperbolic situations this set can be of full Hausdorff dimension [BS00].

with a non-zero upper limit (Corollary 2.13). Finally, as we understand, this work also provides the first natural example of an even stronger manifestation of the lack of regularity, the exponential tangencies (Corollary 2.14). Tangencies between contracted and expanded directions are a characteristic feature of non-uniform hyperbolicity (and, in particular, always happen at the maxima of the eigenfunctions). They complicate proofs of positivity of the Lyapunov exponents and are viewed as a difficulty to avoid through, e.g., the parameter exclusion [BC91], [You97], [Bje15]. However, when the tangencies are only subexponentially deep, they do not in themselves lead to non-exactness. Here we observe the first natural example of *exponentially* strong tangencies (with the rate determined by the arithmetics of α and the positions precisely along the sequence of resonances).

The localization-for-the-exponential-regime method of [AJ09] consists of different arguments for non-resonant (meaning sufficiently far from jq_n on the corresponding scale) sites and for the resonant ones (the rest). It is the resonant sites that lead to dealing with the smallest denominators and that necessitate the $|\lambda| > e^{\frac{16}{9}\beta}$ requirement in [AJ09]. Here we start with the same basic setup and only technically modify the non-resonant statement of [AJ09]. However we develop a completely new bootstrap technique to handle the resonant sites, allowing us to get to the transition and obtain the fine estimates. The estimates from below (which coincide with our estimates from above) are also new. In general, the statements that are technically similar to the ones in the existing literature are collected in the appendices, while all the results/proofs in the body of the paper are, in their pivotal parts, not like anything that has appeared before.

The key elements of the technique developed in this paper are robust and have made it possible to approach other scenarios. As such, in the upcoming work we prove the sharp phase transition for Diophantine α and all θ and establish sharp exponential asymptotics of eigenfunctions and transfer matrices in the corresponding pure point regime [JL18]. Moreover, our analysis reveals there a universal *reflective-hierarchical* structure in the entire regime of phase-induced resonances, a phenomenon not even previously discovered in physics literature. Thus while in this paper we develop a complete understanding of frequency induced resonances, in [JL18] we develop new methods motivated by the ideas of this manuscript to obtain a complete understanding of phase induced resonances. In other follow-up works we determine the exact exponent of the exponential decay rate in expectation for the Diophantine case [JKL] and study delicate properties of the singular continuous regime, obtaining upper bounds on fractal dimensions of the spectral measure and quantum dynamics for the almost Mathieu operator [JLT], as well as potentials defined by general trigonometric polynomials [JL].

Except for a few standard, general (e.g., uniform upper semicontinuity) or very simple to verify statements, this paper is entirely self contained. The only technically involved fact that we use without proving it in the paper is Lemma 3.1 [BJ02], but this is not even necessary if we replace $\ln |\lambda|$ by the Lyapunov exponent $L(E)$ throughout the manuscript.⁴

2. Main results

Let

$$(3) \quad A_k(\theta) = \prod_{j=k-1}^0 A(\theta + j\alpha) = A(\theta + (k-1)\alpha)A(\theta + (k-2)\alpha) \cdots A(\theta)$$

and

$$(4) \quad A_{-k}(\theta) = A_k^{-1}(\theta - k\alpha)$$

for $k \geq 1$, where $A(\theta) = \begin{pmatrix} E-2\lambda \cos 2\pi\theta & -1 \\ 1 & 0 \end{pmatrix}$. The (k -step) transfer matrix is called A_k . As is clear from the definition, it also depends on θ and E , but since those parameters will be usually fixed, we omit this from the notation.

Given $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, we define functions $f, g : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$ in the following way. Let $\frac{p_n}{q_n}$ be the continued fraction approximants to α . For any $\frac{q_n}{2} \leq k < \frac{q_{n+1}}{2}$, define $f(k), g(k)$ as follows.

Case 1: $q_{n+1}^{\frac{8}{9}} \geq \frac{q_n}{2}$ or $k \geq q_n$. If $\ell q_n \leq k < (\ell + 1)q_n$ with $\ell \geq 1$, set

$$(5) \quad f(k) = e^{-|k-\ell q_n| \ln |\lambda| \bar{r}_\ell^n} + e^{-|k-(\ell+1)q_n| \ln |\lambda| \bar{r}_{\ell+1}^n}$$

and

$$(6) \quad g(k) = e^{-|k-\ell q_n| \ln |\lambda| \frac{q_{n+1}}{\bar{r}_\ell^n}} + e^{-|k-(\ell+1)q_n| \ln |\lambda| \frac{q_{n+1}}{\bar{r}_{\ell+1}^n}},$$

where for $\ell \geq 1$,

$$\bar{r}_\ell^n = e^{-(\ln |\lambda| - \frac{\ln q_{n+1}}{q_n} + \frac{\ln \ell}{q_n}) \ell q_n}.$$

Also set $\bar{r}_0^n = 1$ for convenience.

If $\frac{q_n}{2} \leq k < q_n$, set

$$(7) \quad f(k) = e^{-k \ln |\lambda|} + e^{-|k-q_n| \ln |\lambda| \bar{r}_1^n}$$

and

$$(8) \quad g(k) = e^{k \ln |\lambda|}.$$

⁴ In fact, $\ln |\lambda|$ is being used in this paper as a shortcut for $L(E)$.

Case 2: $q_{n+1}^{8/9} < \frac{q_n}{2}$ and $\frac{q_n}{2} \leq k \leq \min\{q_n, \frac{q_{n+1}}{2}\}$. Set

$$(9) \quad f(k) = e^{-k \ln |\lambda|}$$

and

$$(10) \quad g(k) = e^{k \ln |\lambda|}.$$

Notice that f, g only depend on α and λ but not on θ or E . Here $f(k)$ decays and $g(k)$ grows exponentially, globally, at varying rates that depend on the position of k in the hierarchy defined by the continued fraction expansion of α ; see Figures 1 and 2.




Figure 1.




Figure 2.

We say that ϕ is a generalized eigenfunction of H with generalized eigenvalue E if

$$(11) \quad H\phi = E\phi \text{ and } |\phi(k)| \leq \hat{C}(1 + |k|).$$

Our first main result is that in the entire regime $|\lambda| > e^\beta$, the exponential asymptotics of the generalized eigenfunctions and norms of transfer matrices at the generalized eigenvalues are completely determined by $f(k), g(k)$.

THEOREM 2.1. *Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ be such that $|\lambda| > e^{\beta(\alpha)}$. Suppose θ is Diophantine with respect to α , E is a generalized eigenvalue of $H_{\lambda, \alpha, \theta}$ and ϕ is the generalized eigenfunction. Let $U(k) = \begin{pmatrix} \phi(k) \\ \phi(k-1) \end{pmatrix}$. Then for any $\varepsilon > 0$, there exists K (depending on $\lambda, \alpha, \hat{C}, \varepsilon$ and Diophantine constants κ, ν) such that for any $|k| \geq K$, $U(k)$ and A_k satisfy*

$$(12) \quad f(|k|)e^{-\varepsilon|k|} \leq \|U(k)\| \leq f(|k|)e^{\varepsilon|k|}$$

and

$$(13) \quad g(|k|)e^{-\varepsilon|k|} \leq \|A_k\| \leq g(|k|)e^{\varepsilon|k|}.$$

Certainly, there is nothing special about $k = 0$, so the behavior described in Theorem 2.1 happens around an arbitrary point $k = k_0$. This implies the self-similar nature of the eigenfunctions: $U(k)$ behave as described at scale q_n but when looked at in windows of size $q_k, q_k < q_{n-1}$ will demonstrate the same universal behavior around appropriate local maxima/minima.

To make the above precise, let ϕ be an eigenfunction, and let $U(k) = \begin{pmatrix} \phi(k) \\ \phi(k-1) \end{pmatrix}$. Let $I_{\varsigma_1, \varsigma_2}^j = [-\varsigma_1 q_j, \varsigma_2 q_j]$ for some $0 < \varsigma_1, \varsigma_2 \leq 1$. We will say k_0 is a local j -maximum of ϕ if $\|U(k_0)\| \geq \|U(k)\|$ for $k - k_0 \in I_{\varsigma_1, \varsigma_2}^j$. Occasionally, we will also use the terminology (j, ς) -maximum for a local j -maximum on an interval $I_{\varsigma, \varsigma}^j$.

We will say a local j -maximum k_0 is *non-resonant* if

$$\|2\theta + (2k_0 + k)\alpha\|_{\mathbb{R}/\mathbb{Z}} > \frac{\kappa}{q_{j-1}^\nu}$$

for all $|k| \leq 2q_{j-1}$ and

$$(14) \quad \|2\theta + (2k_0 + k)\alpha\|_{\mathbb{R}/\mathbb{Z}} > \frac{\kappa}{|k|^\nu}$$

for all $2q_{j-1} < |k| \leq 2q_j$.

We will say a local j -maximum is *strongly non-resonant* if

$$(15) \quad \|2\theta + (2k_0 + k)\alpha\|_{\mathbb{R}/\mathbb{Z}} > \frac{\kappa}{|k|^\nu}$$

for all $0 < |k| \leq 2q_j$.

An immediate corollary of Theorem 2.1 is the universality of behavior at all (strongly) non-resonant local maxima.

THEOREM 2.2. *Given $\varepsilon > 0$, there exists $j(\varepsilon) < \infty$ such that if k_0 is a local j -maximum for $j > j(\varepsilon)$, then the following two statements hold:*

If k_0 is non-resonant, then

$$(16) \quad f(|s|)e^{-\varepsilon|s|} \leq \frac{\|U(k_0 + s)\|}{\|U(k_0)\|} \leq f(|s|)e^{\varepsilon|s|}$$

for all $2s \in I_{s_1, s_2}^j$, $|s| > \frac{q_{j-1}}{2}$.

If k_0 is strongly non-resonant, then

$$(17) \quad f(|s|)e^{-\varepsilon|s|} \leq \frac{\|U(k_0 + s)\|}{\|U(k_0)\|} \leq f(|s|)e^{\varepsilon|s|}$$

for all $2s \in I_{s_1, s_2}^j$.

Remark 2.3.

- (1) For the neighborhood of a local j -maximum described in Theorem 2.2, only the behavior of $f(s)$ for $q_{j-1}/2 < |s| \leq q_j/2$ is relevant. Thus f implicitly depends on j but through the scale-independent mechanism described in (5), (7) and (9).
- (2) Actually, a modification in our proof us allows to formulate (16) in Theorem 2.2 with the non-resonant condition (14) only required for $2q_{j-1} < |k| \leq q_j$ rather than for $2q_{j-1} < |k| \leq 2q_j$.

In case $\beta(\alpha) > 0$, Theorem 2.1 also guarantees an abundance (and a hierarchical structure) of local maxima of each eigenfunction. Let k_0 be a global maximum.⁵

We first describe the hierarchical structure of local maxima informally. We will say that a scale n_{j_0} is exponential if $\ln q_{n_{j_0}+1} > cq_{n_{j_0}}$. Then there is a constant scale \hat{n}_0 and thus a constant $C := q_{\hat{n}_0+1}$ such that for any exponential scale n_j and any eigenfunction, there are local n_j -maxima within distance C of $k_0 + sq_{n_{j_0}}$ for each $0 < |s| < e^{cq_{n_{j_0}}}$. Moreover, these are all the local n_{j_0} -maxima in $[k_0 - e^{cq_{n_{j_0}}}, k_0 + e^{cq_{n_{j_0}}}]$. The exponential behavior of the eigenfunction in the local neighborhood (of size $\sim q_{n_{j_0}}$) of each such local maximum, normalized by the value at the local maximum is given by f . Note that only exponential behavior at the corresponding scale is determined by f and fluctuations of much smaller size are invisible. Now, let $n_{j_1} < n_{j_0}$ be another exponential scale. Denoting “depth 1” local maximum located near $k_0 + a_{n_{j_0}} q_{n_{j_0}}$ by $b_{a_{n_{j_0}}}$, we then have a similar picture around $b_{a_{n_{j_0}}}$; there are local n_{j_1} -maxima in the vicinity of $b_{a_{n_{j_0}}} + sq_{n_{j_1}}$ for each $0 < |s| < e^{cq_{n_{j_1}}}$. Again, this describes all the local $q_{n_{j_1}}$ -maxima within an exponentially large interval. And again, the exponential (for the n_{j_1} scale) behavior in the local

⁵If there are several, what follows is true for each.




Figure 3. Hierarchical structure of an eigenfunction.

neighborhood (of size $\sim q_{n_{j_1}}$) of each such local maximum, normalized by the value at the local maximum, is given by f . Denoting those “depth 2” local maxima located near $b_{a_{n_{j_0}}} + a_{n_{j_1}} q_{n_{j_1}}$ by $b_{a_{n_{j_0}}, a_{n_{j_1}}}$, we then get the same picture taking the magnifying glass another level deeper and so on. At the end we obtain a complete hierarchical structure of local maxima, which we denote by $b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}}$ with each “depth $s + 1$ ” local maximum $b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}}$ being in the corresponding vicinity of the “depth s ” local maximum $b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_{s-1}}}}$ and with universal behavior at the corresponding scale around each. The quality of the approximation of the position of the next maximum gets lower with each level of depth, yet the depth of the hierarchy that can be so achieved is at least $j/2 - C$; see Corollary 2.7. Figure 3 schematically illustrates the structure of local maxima of depth one and two, and Figure 4 illustrates that the neighborhood of a local maximum appropriately magnified looks like a picture of the global maximum.

We now describe the hierarchical structure precisely. Suppose

$$(18) \quad \|2(\theta + k_0\alpha) + k\alpha\|_{\mathbb{R}/\mathbb{Z}} > \frac{\kappa}{|k|^\nu}$$

for any $k \in \mathbb{Z} \setminus \{0\}$. Fix $0 < \varsigma, \epsilon$ with $\varsigma + 2\epsilon < 1$. Let $n_j \rightarrow \infty$ be such that $\ln q_{n_{j+1}} \geq (\varsigma + 2\epsilon) \ln |\lambda| q_{n_j}$. Let $\mathfrak{c}_j = (\ln q_{n_{j+1}} - \ln |a_{n_j}|) / \ln |\lambda| q_{n_j} - \epsilon$. We have $\mathfrak{c}_j > \epsilon$ for $0 < a_{n_j} < e^{\varsigma \ln |\lambda| q_{n_j}}$. Then we have




Figure 4. Window I.

THEOREM 2.4. *There exists $\hat{n}_0(\alpha, \lambda, \kappa, \nu, \epsilon) < \infty$ such that for any $j_0 > j_1 > \dots > j_k$, $n_{j_k} \geq \hat{n}_0 + k$, and $0 < a_{n_{j_i}} < e^{\epsilon \ln |\lambda| q_{n_{j_i}}}$, $i = 0, 1, \dots, k$, for all $0 \leq s \leq k$, there exists a local n_{j_s} -maximum $b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}}$ on the interval $b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}} + I_{c_{j_s}, 1}^{n_{j_s}}$ for all $0 \leq s \leq k$ such that the following hold:*

- (I) $|b_{a_{n_{j_0}}} - (k_0 + a_{n_{j_0}} q_{n_{j_0}})| \leq q_{\hat{n}_0+1}$;
- (II) for any $1 \leq s \leq k$, $|b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}} - (b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_{s-1}}}} + a_{n_{j_s}} q_{n_{j_s}})| \leq q_{\hat{n}_0+s+1}$;
- (III) if $2(x - b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_k}}}) \in I_{c_{j_k}, 1}^{n_{j_k}}$ and $|x - b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_k}}}| \geq q_{\hat{n}_0+k}$, then for each $s = 0, 1, \dots, k$,

$$(19) \quad f(x_s) e^{-\epsilon |x_s|} \leq \frac{\|U(x)\|}{\|U(b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}})\|} \leq f(x_s) e^{\epsilon |x_s|},$$

where $x_s = |x - b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}}|$ is large enough.

Moreover, every local n_{j_s} -maximum on the interval

$$b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_{s-1}}}} + [-e^{\epsilon \ln \lambda q_{n_{j_s}}}, e^{\epsilon \ln \lambda q_{n_{j_s}}}]$$

is of the form $b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}}$ for some $a_{n_{j_s}}$.

Remark 2.5. By (I) of Theorem 2.4, the local maximum can be determined up to a constant $K_0 = q_{\hat{n}_0+1}$. Actually, if k_0 is only a local $n_j + 1$ -maximum, we can still make sure that (I), (II) and (III) of Theorem 2.4 hold. This is the local version of Theorem 2.4; see Theorem 7.3.

Remark 2.6. Here $q_{\hat{n}_0+1}$ is the last scale at which *phase* resonances of $\theta + k_0\alpha$ still can appear. Notably, it determines the precision of pinpointing local n_{j_0} -maxima in a (exponentially large in $q_{n_{j_0}}$) neighborhood of k_0 for any j_0 . When we go down the hierarchy, the precision decreases, but note that except for the very last scale, it stays at least iterated logarithmically⁶ small in the corresponding scale $q_{n_{j_s}}$

Thus for $x \in b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}} + [-\frac{c_{j_s}}{2}q_{n_{j_s}}, \frac{1}{2}q_{n_{j_s}}]$, the behavior of $\phi(x)$ is described by the same universal f in each $q_{n_{j_s}}$ -window around the corresponding local maximum $b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}}$, $s = 0, 1, \dots, k$. We call such a structure *hierarchical*, and we will say that a local j -maximum is k -hierarchical if the complete hierarchy goes down at least k levels; for a precise definition, see Section 7. We then have an immediate corollary.

COROLLARY 2.7. *There exists $C = C(\alpha, \lambda, \kappa, \nu, \epsilon)$ such that every local n_j -maximum in $[k_0 - e^{\zeta \ln |\lambda|^{q_{n_j}}}, k_0 + e^{\zeta \ln |\lambda|^{q_{n_j}}}]$ is at least $(j/2 - C)$ -hierarchical.*

Remark 2.8. The estimate on the depth of the hierarchy in the corollary assumes the worst case scenario when all scales after \hat{n}_0 are Liouville. Otherwise the hierarchical structure will go much deeper. Note that a local n_j -maximum that is not an n_{j+1} -maximum cannot be k -hierarchical for $k > j$.

Another interesting corollary of Theorem 2.1 is

THEOREM 2.9. *Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ be such that $|\lambda| > e^{\beta(\alpha)}$ and θ is Diophantine with respect to α . Then $H_{\lambda, \alpha, \theta}$ has Anderson localization, with eigenfunctions decaying at the rate $\ln |\lambda| - \beta$.*

This solves the arithmetic version of the second transition conjecture in that it establishes localization throughout the entire regime of (α, λ) where localization may hold for any θ (see the discussion in the introduction) for an arithmetically defined full measure set of θ .

We note that Theorem 2.9 cannot be upgraded to *all* θ in the regime $|\lambda| > e^\beta$ [JS94], so exclusion of a certain arithmetically defined set where the spectrum must be singular continuous is necessary. There is a conjecture about where in this regime the transition in θ happens [Jit95], but we do not explore it in this work. The sharp transition in θ for Diophantine α will be established in the follow-up work [JL18]. Also, it could be added that, for all θ , $H_{\lambda, \alpha, \theta}$ has no localization (i.e., no exponentially decaying eigenfunctions) if $|\lambda| = e^\beta$; see Appendix A.1.

⁶for most scales even much less

Remark 2.10. Theorems 2.1 and 2.9 cover the optimal range of (α, λ) for almost every θ . For Theorem 2.9, even though some θ have to be excluded [JS94], we do not claim the Diophantine condition on θ is optimal. At the same time, exponentially strong θ -resonances (exponentially small lower bound in (2) instead of a polynomial) will make Theorem 2.1 false as stated, no matter how small the exponent, and they would require differently defined f and g . In [JL18] we obtain f' and g' that govern the exponential behavior of eigenfunctions and transfermatrices for *all* θ throughout the entire pure point regime corresponding to Diophantine α .

Let $\psi(k)$ denote any solution to $H_{\lambda, \alpha, \theta} \psi = E\psi$ that is linearly independent with respect to $\phi(k)$. Let $\tilde{U}(k) = \begin{pmatrix} \psi(k) \\ \psi(k-1) \end{pmatrix}$. An immediate counterpart of (13) is the following

COROLLARY 2.11. *Under the conditions of Theorem 2.1 for large k , vectors $\tilde{U}(k)$ satisfy*

$$(20) \quad g(|k|)e^{-\varepsilon|k|} \leq \|\tilde{U}(k)\| \leq g(|k|)e^{\varepsilon|k|}.$$

Thus every solution is expanding at the rate $g(k)$ except for one that is exponentially decaying at the rate $f(k)$.

It is well known that for E in the spectrum, the dynamics of the transfer-matrix cocycle A_k is non-uniformly hyperbolic. Moreover, E being a generalized eigenvalue of $H_{\lambda, \alpha, \theta}$ already implies that the behavior of A_k is non-regular. Theorem 2.1 provides precise information on how the non-regular behavior unfolds in this case. Previously, a study of some features of the non-regular behavior for the almost Mathieu operator was made in [FK08]. We are not aware though of other non-artificially constructed examples of non-uniformly hyperbolic systems where non-regular behavior can be described with such precision as in the present work.

The information provided by Theorem 2.1 leads to many interesting corollaries, which will be explored elsewhere. Here we only want to list a few immediate sharp consequences.

COROLLARY 2.12. *Under the condition of Theorem 2.1, we have*

- (i) $\limsup_{k \rightarrow \infty} \frac{\ln \|A_k\|}{k} = \limsup_{k \rightarrow \infty} \frac{\ln \|\tilde{U}(k)\|}{k} = \ln |\lambda|;$
- (ii) $\liminf_{k \rightarrow \infty} \frac{\ln \|A_k\|}{k} = \liminf_{k \rightarrow \infty} \frac{\ln \|\tilde{U}(k)\|}{k} = \ln |\lambda| - \beta;$
- (iii) *outside an explicit sequence of lower density zero,*⁷

$$\lim_{k \rightarrow \infty} \frac{\ln \|A_k\|}{k} = \lim_{k \rightarrow \infty} \frac{\ln \|\tilde{U}(k)\|}{k} = \ln |\lambda|.$$

⁷It will be clear from the proof that the sequence with convergence to the Lyapunov exponent contains $q_n, n = 1, \dots$.

Therefore the Lyapunov behavior for the norm fails to hold only along a sequence of density zero. It is interesting that the situation is different for the eigenfunctions. While, just like the overall growth of $\|A_k\|$ is $\ln |\lambda| - \beta$, the overall rate of decay of the eigenfunctions is also $\ln |\lambda| - \beta$, they however decay at the Lyapunov rate only outside a sequence of positive upper density. That is,

COROLLARY 2.13. *Under the condition of Theorem 2.1, we have*

- (i) $\limsup_{k \rightarrow \infty} \frac{-\ln \|U(k)\|}{k} = \ln |\lambda|$;
- (ii) $\liminf_{k \rightarrow \infty} \frac{-\ln \|\tilde{U}(k)\|}{k} = \ln |\lambda| - \beta$;
- (iii) *there is an explicit sequence of upper density $1 - \frac{1}{2} \frac{\beta}{\ln |\lambda|}$,⁸ along which*

$$\lim_{k \rightarrow \infty} \frac{-\ln \|U(k)\|}{k} = \ln |\lambda|;$$

- (iv) *there is an explicit sequence of upper density $\frac{1}{2} \frac{\beta}{\ln |\lambda|}$,⁹ along which*

$$\limsup_{k \rightarrow \infty} \frac{-\ln \|U(k)\|}{k} < \ln |\lambda|.$$

The fact that g is not always the reciprocal of f also leads to another interesting phenomenon.

Let $0 \leq \delta_k \leq \frac{\pi}{2}$ be the angle between vectors $U(k)$ and $\tilde{U}(k)$.

COROLLARY 2.14. *We have*

$$(21) \quad \limsup_{k \rightarrow \infty} \frac{\ln \delta_k}{k} = 0$$

and

$$(22) \quad \liminf_{k \rightarrow \infty} \frac{\ln \delta_k}{k} = -\beta.$$

As becomes clear from the proof, neighborhoods of resonances q_n are the places of exponential tangencies between contracted and expanded directions, with the rate approaching $-\beta$ along a subsequence.¹⁰ Exponential tangencies also happen around points of the form jq_n but at lower strength. This means, in particular, that A_k with $k \sim q_n$ is exponentially close to a matrix with the trace $e^{(\ln |\lambda| - \beta)k}$.

The rest of this paper is organized in the following way. We list the definitions and standard preliminaries in Section 3. We also include there the

⁸It will be clear from the proof that the sequence contains $\lfloor \frac{q_n}{2} \rfloor, n = 1, \dots$

⁹As will be clear from the proof, this sequence can have lower density ranging from 0 to $\frac{1}{2} \frac{\beta}{\ln |\lambda|}$ depending on finer continued fraction properties of α .

¹⁰In fact the rate is close to $-\frac{\ln q_{n+1}}{q_n}$ for any large n .

non-resonant regularity statement. While similar to the corresponding statements in [AJ09], [LY15a], [LY15b], it differs in enough technical details that a proof is needed for completeness. We present this proof in Appendix B. Section 4 is devoted to the bootstrap localization argument, establishing sharp upper bounds for the resonant case. Section 5 is devoted to the lower bounds. In Section 6 we prove the statements about eigenfunctions: (12) of Theorem 2.1, and Theorems 2.2 and 2.9. In Section 7, we will prove the hierarchical structure, Theorem 2.4 and Corollary 2.7. In Section 8, we study the growth of transfer matrices and prove (13) of Theorem 2.1. The remaining corollaries are proved in Section 9.

3. Preliminaries

Fix $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ such that $\beta(\alpha) < \infty$. Unless stated otherwise, we always assume $\lambda > e^\beta$ (for $\lambda < -e^\beta$, notice that $H_{\lambda,\alpha,\theta} = H_{-\lambda,\alpha,\theta+\frac{1}{2}}$), θ is Diophantine with respect to α and E is a generalized eigenvalue. We also assume ϕ is the corresponding generalized eigenfunction of $H_{\lambda,\alpha,\theta}$. Without loss of generality assume $|\phi(0)|^2 + |\phi(-1)|^2 = 1$. Let ψ be any solution to $H_{\lambda,\alpha,\theta}\psi = E\psi$ linear independent with respect to ϕ , i.e., $|\psi(0)|^2 + |\psi(-1)|^2 = 1$ and

$$(23) \quad \phi(-1)\psi(0) - \phi(0)\psi(-1) = c,$$

where $c \neq 0$.

Then by the constancy of the Wronskian, one has

$$(24) \quad \phi(k+1)\psi(k) - \phi(k)\psi(k+1) = c.$$

We also will denote by φ an *arbitrary* solution, so either ψ or ϕ . Thus for any k, m , one has

$$(25) \quad \begin{pmatrix} \varphi(k+m) \\ \varphi(k+m-1) \end{pmatrix} = A_k(\theta + m\alpha) \begin{pmatrix} \varphi(m) \\ \varphi(m-1) \end{pmatrix}.$$

The Lyapunov exponent is given by

$$(26) \quad L(E) = \lim_{k \rightarrow \infty} \frac{1}{k} \int_{\mathbb{R}/\mathbb{Z}} \ln \|A_k(\theta)\| d\theta.$$

The Lyapunov exponent can be computed precisely for E in the spectrum of $H_{\lambda,\alpha,\theta}$. We denote the spectrum by $\Sigma_{\lambda,\alpha}$. (It does not depend on θ .)

LEMMA 3.1. [BJ02] *For $E \in \Sigma_{\lambda,\alpha}$ and $\lambda > 1$, we have $L(E) = \ln \lambda$.*

Recall that we always assume $E \in \Sigma_{\lambda,\alpha}$ so by upper semicontinuity and unique ergodicity (e.g., [Fur97]), one has

$$(27) \quad \ln \lambda = \lim_{k \rightarrow \infty} \sup_{\theta \in \mathbb{R}/\mathbb{Z}} \frac{1}{k} \ln \|A_k(\theta)\|;$$

that is, the convergence in (27) is uniform with respect to $\theta \in \mathbb{R}$. Precisely, for all $\varepsilon > 0$,

$$(28) \quad \|A_k(\theta)\| \leq e^{(\ln \lambda + \varepsilon)k} \text{ for } k \text{ large enough.}$$

We start with the basic setup going back to [Jit99]. Let us denote

$$P_k(\theta) = \det(R_{[0,k-1]}(H_{\lambda,\alpha,\theta} - E)R_{[0,k-1]}).$$

It is easy to check that

$$(29) \quad A_k(\theta) = \begin{pmatrix} P_k(\theta) & -P_{k-1}(\theta + \alpha) \\ P_{k-1}(\theta) & -P_{k-2}(\theta + \alpha) \end{pmatrix}.$$

By Cramer’s rule for given x_1 and $x_2 = x_1 + k - 1$, with $y \in I = [x_1, x_2] \subset \mathbb{Z}$, one has

$$(30) \quad |G_I(x_1, y)| = \left| \frac{P_{x_2-y}(\theta + (y + 1)\alpha)}{P_k(\theta + x_1\alpha)} \right|,$$

$$(31) \quad |G_I(y, x_2)| = \left| \frac{P_{y-x_1}(\theta + x_1\alpha)}{P_k(\theta + x_1\alpha)} \right|.$$

By (28) and (29), the numerators in (30) and (31) can be bounded uniformly with respect to θ . Namely, for any $\varepsilon > 0$,

$$(32) \quad |P_k(\theta)| \leq e^{(\ln \lambda + \varepsilon)k}$$

for k large enough.

Definition 3.2. Fix $\tau > 0$, $0 < \delta < 1/2$. A point $y \in \mathbb{Z}$ will be called (τ, k) regular with δ if there exists an interval $[x_1, x_2]$ containing y , where $x_2 = x_1 + k - 1$, such that

$$|G_{[x_1, x_2]}(y, x_i)| < e^{-\tau|y-x_i|} \text{ and } |y - x_i| \geq \delta k \text{ for } i = 1, 2.$$

It is easy to check that

$$(33) \quad \varphi(x) = -G_{[x_1, x_2]}(x_1, x)\varphi(x_1 - 1) - G_{[x_1, x_2]}(x, x_2)\varphi(x_2 + 1),$$

where $x \in I = [x_1, x_2] \subset \mathbb{Z}$.

Definition 3.3. We say that the set $\{\theta_1, \dots, \theta_{k+1}\}$ is ε -uniform if

$$(34) \quad \max_{x \in [-1, 1]} \max_{i=1, \dots, k+1} \prod_{j=1, j \neq i}^{k+1} \frac{|x - \cos 2\pi\theta_j|}{|\cos 2\pi\theta_i - \cos 2\pi\theta_j|} < e^{k\varepsilon}.$$

Let $A_{k,r} = \{\theta \in \mathbb{R} \mid |P_k(\cos 2\pi(\theta - \frac{1}{2}(k - 1)\alpha))| \leq e^{(k+1)r}\}$ with $k \in \mathbb{N}$ and $r > 0$. We have the following lemma.

LEMMA 3.4 ([AJ09, Lemma 9.3]). *Suppose $\{\theta_1, \dots, \theta_{k+1}\}$ is ε_1 -uniform. Then there exists some θ_i in the set $\{\theta_1, \dots, \theta_{k+1}\}$ such that $\theta_i \notin A_{k, \ln \lambda - \varepsilon}$ if $\varepsilon > \varepsilon_1$ and k is sufficiently large.*

Proof. Straightforward calculation. □

We say θ is *n-Diophantine* with respect to α if, for some $\kappa > 0, \nu > 1$, the following hold:

$$(35) \quad \|2\theta + k\alpha\|_{\mathbb{R}/\mathbb{Z}} > \frac{\kappa}{q_n^\nu}$$

for all $|k| \leq 2q_n$ and

$$(36) \quad \|2\theta + k\alpha\|_{\mathbb{R}/\mathbb{Z}} > \frac{\kappa}{|k|^\nu}$$

for all $2q_n < |k| \leq 2q_{n+1}$.

Define $b_n = q_n^t$ with $\frac{8}{9} \leq t < 1$; t will be defined later. For any $k > 0$, we will distinguish two cases with respect to n :

- (i) $|k - \ell q_n| \leq b_n$ for some $\ell \geq 1$, called *n-resonance*;
- (ii) $|k - \ell q_n| > b_n$ for all $\ell \geq 0$, called *n-non-resonance*.

For the *n-non-resonant* y , let n_0 be the least positive integer such that $4q_{n-n_0} \leq \text{dist}(y, q_n\mathbb{Z})$. Let s be the largest positive integer such that $4sq_{n-n_0} \leq \text{dist}(y, q_n\mathbb{Z})$. Notice that $n_0 \leq C(\alpha)$.

The following theorem is similar to a statement appearing in[AJ09] with modifications in [LY15a], [LY15b]. We present a proof in Appendix B.

THEOREM 3.5. *Assume $\lambda > e^{\beta(\alpha)}$. Suppose either*

- (i) $b_n \leq |y| < Cb_{n+1}$, where $C > 1$ is a fixed constant and θ is *n-Diophantine with respect to α* ; or
- (ii) $0 \leq |y| < q_n$, and θ satisfies (35).

*Then for any $\varepsilon > 0$ and n large enough, if y is *n-non-resonant*, we have y is $(\ln \lambda + 8 \ln(sq_{n-n_0}/q_{n-n_0+1})/q_{n-n_0} - \varepsilon, 4sq_{n-n_0} - 1)$ regular with $\delta = \frac{1}{4}$.*

Remark 3.6. If θ is *n - 1-Diophantine* with respect to α , then (35) holds.

Remark 3.7. In the non-resonant case, for any $\varepsilon > 0, \frac{8}{9} \leq t < 1$, one has $\ln \lambda + 8 \ln(sq_{n-n_0}/q_{n-n_0+1})/q_{n-n_0} \geq \ln \lambda - 8(1-t)\beta - \varepsilon > 0$. In addition, we have $\ln \lambda + 8 \ln(sq_{n-n_0}/q_{n-n_0+1})/q_{n-n_0} \geq \ln \lambda - 2\varepsilon$ if t is close to 1.

Remark 3.8. In the present paper, we only use Theorem 3.5 with $C = 50C_*$, where C_* is given by (37); see the next section.

4. Bootstrap resonant localization

In this section we assume θ is *n-Diophantine* with respect to α . Clearly, it is enough to consider $k > 0$. In this section we study the resonant case. Suppose there exists some $k \in [b_n, b_{n+1}]$ such that k is *n-resonant*. Then we have $b_{n+1} \geq \frac{q_n}{2}$. For any $\varepsilon > 0$, choose $\eta = \frac{\varepsilon}{C}$, where C is a large constant (depending on λ, α).

Let

$$(37) \quad C_* = 2 \left(1 + \left\lfloor \frac{\ln \lambda}{\ln \lambda - \beta} \right\rfloor \right),$$

where $\lfloor m \rfloor$ denotes the smallest integer not exceeding m .

For an arbitrary solution φ satisfying $H\varphi = E\varphi$, let

$$r_j^{n,\varphi} = \sup_{|r| \leq 10\eta} |\varphi(jq_n + rq_n)|,$$

where $|j| \leq 50C_* \frac{b_{n+1}}{q_n}$.

Fix ψ satisfying (23), and write

$$R_j^n = r_j^{n,\psi}$$

and

$$r_j^n = r_j^{n,\phi}.$$

Since we keep n fixed in this section, we omit the dependence on n from the notation and write r_j^φ, R_j , and r_j .

Note that below we always assume n is large enough.¹¹ In the next lemma and its variant, Lemma 4.2, we establish exponential decay of the eigenfunctions at non-resonant points, at the nearly Lyapunov rate, with respect to the distance to the resonances.

LEMMA 4.1. *Let $k \in [jq_n, (j + 1)q_n]$ with $\text{dist}(k, q_n\mathbb{Z}) \geq 10\eta q_n$. Suppose either*

- (i) $|j| \leq 48C_* \frac{b_{n+1}}{q_n}$ and $b_{n+1} \geq \frac{q_n}{2}$, or
- (ii) $j = 0$;

then for sufficiently large n ,

$$(38) \quad |\varphi(k)| \leq \max \left\{ r_j^\varphi \exp\{-(\ln \lambda - 2\eta)(d_j - 3\eta q_n)\}, \right. \\ \left. r_{j+1}^\varphi \exp\{-(\ln \lambda - 2\eta)(d_{j+1} - 3\eta q_n)\} \right\},$$

where $d_j = |k - jq_n|$ and $d_{j+1} = |k - jq_n - q_n|$.

Proof. The proof builds on the ideas used in the proof of Lemma 9.11 in [AJ09] and Lemma 3.2 in [LY15a]. However it requires a more careful approach.

We first prove case (i). For any $y \in [jq_n + \eta q_n, (j + 1)q_n - \eta q_n]$, apply (i) of Theorem 3.5 with $C = 50C_*$. Note that in this case, we have

$$\ln \lambda + 8 \ln(sq_{n-n_0}/q_{n-n_0+1})/q_{n-n_0} - \eta \geq \ln \lambda - 2\eta.$$

¹¹ The required largeness of n will depend on α, θ, \hat{C} in (11) and ε whenever ε is (implicitly) present in the statement.

Thus y is regular with $\tau = \ln \lambda - 2\eta$. Therefore there exists an interval $I(y) = [x_1, x_2] \subset [jq_n, (j+1)q_n]$ such that $y \in I(y)$ and

$$(39) \quad \text{dist}(y, \partial I(y)) \geq \frac{1}{4}|I(y)| \geq q_{n-n_0}$$

and

$$(40) \quad |G_{I(y)}(y, x_i)| \leq e^{-(\ln \lambda - 2\eta)|y - x_i|}, \quad i = 1, 2,$$

where $\partial I(y)$ is the boundary of the interval $I(y)$, i.e., $\{x_1, x_2\}$, and $|I(y)|$ is the size of $I(y) \cap \mathbb{Z}$, i.e., $|I(y)| = x_2 - x_1 + 1$. For $z \in \partial I(y)$, let z' be the neighbor of z , (i.e., $|z - z'| = 1$) not belonging to $I(y)$.

If $x_2 + 1 \leq (j+1)q_n - \eta q_n$ or $x_1 - 1 \geq jq_n + \eta q_n$, we can expand $\varphi(x_2 + 1)$ or $\varphi(x_1 - 1)$ using (33). We can continue this process until we arrive to z such that $z + 1 > (j+1)q_n - \eta q_n$ or $z - 1 < jq_n + \eta q_n$, or the iterating number reaches $\lfloor \frac{2q_n}{q_{n-n_0}} \rfloor$. Thus, by (33),

$$(41) \quad \varphi(k) = \sum_{s; z_{i+1} \in \partial I(z'_i)} G_{I(k)}(k, z_1) G_{I(z'_1)}(z'_1, z_2) \cdots G_{I(z'_s)}(z'_s, z_{s+1}) \varphi(z'_{s+1}),$$

where in each term of the summation one has $jq_n + \eta q_n + 1 \leq z_i \leq (j+1)q_n - \eta q_n - 1$, $i = 1, \dots, s$, and either $z_{s+1} \notin [jq_n + \eta q_n + 1, (j+1)q_n - \eta q_n - 1]$, $s+1 < \lfloor \frac{2q_n}{q_{n-n_0}} \rfloor$, or $s+1 = \lfloor \frac{2q_n}{q_{n-n_0}} \rfloor$. We should mention that $z_{s+1} \in [jq_n, (j+1)q_n]$.

If $z_{s+1} \in [jq_n, jq_n + \eta q_n]$, $s+1 < \lfloor \frac{2q_n}{q_{n-n_0}} \rfloor$, this implies

$$|\varphi(z'_{s+1})| \leq r_j^\varphi.$$

By (40), we have

$$(42) \quad \begin{aligned} & |G_{I(k)}(k, z_1) G_{I(z'_1)}(z'_1, z_2) \cdots G_{I(z'_s)}(z'_s, z_{s+1}) \varphi(z'_{s+1})| \\ & \leq r_j^\varphi e^{-(\ln \lambda - 2\eta)(|k - z_1| + \sum_{i=1}^s |z'_i - z_{i+1}|)} \\ & \leq r_j^\varphi e^{-(\ln \lambda - 2\eta)(|k - z_{s+1}| - (s+1))} \\ & \leq r_j^\varphi e^{-(\ln \lambda - 2\eta)(d_j - 2\eta q_n - 4 - \frac{2q_n}{q_{n-n_0}})}. \end{aligned}$$

If $z_{s+1} \in [(j+1)q_n - \eta q_n, (j+1)q_n]$, $s+1 < \lfloor \frac{2q_n}{q_{n-n_0}} \rfloor$, by the same arguments, we have

$$(43) \quad \begin{aligned} & |G_{I(k)}(k, z_1) G_{I(z'_1)}(z'_1, z_2) \cdots G_{I(z'_s)}(z'_s, z_{s+1}) \varphi(z'_{s+1})| \\ & \leq r_{j+1}^\varphi e^{-(\ln \lambda - 2\eta)(d_{j+1} - 2\eta q_n - 4 - \frac{2q_n}{q_{n-n_0}})}. \end{aligned}$$

If $s+1 = \lfloor \frac{2q_n}{q_{n-n_0}} \rfloor$, using (39) and (40), we obtain

$$(44) \quad \begin{aligned} & |G_{I(k)}(k, z_1) G_{I(z'_1)}(z'_1, z_2) \cdots G_{I(z'_s)}(z'_s, z_{s+1}) \varphi(z'_{s+1})| \\ & \leq e^{-(\ln \lambda - 2\eta)q_{n-n_0} \lfloor \frac{2q_n}{q_{n-n_0}} \rfloor} |\varphi(z'_{s+1})|. \end{aligned}$$

Notice that the total number of terms in (41) is at most $2^{\lfloor \frac{2q_n}{q_n - n_0} \rfloor}$ and $d_j, d_{j+1} \geq 10\eta q_n$. By (42), (43) and (44), we have

$$(45) \quad |\varphi(k)| \leq \max \left\{ r_j^\varphi e^{-(\ln \lambda - 2\eta)(d_j - 3\eta q_n)}, r_{j+1}^\varphi e^{-(\ln \lambda - 2\eta)(d_{j+1} - 3\eta q_n)}, \right. \\ \left. \max_{p \in [jq_n, (j+1)q_n]} \{ e^{-(\ln \lambda - 2\eta)q_n} |\varphi(p)| \} \right\}.$$

Now we will show that for any $p \in [jq_n, (j+1)q_n]$, one has $|\varphi(p)| \leq \max\{r_j^\varphi, r_{j+1}^\varphi\}$. Then (45) implies case (i) of Lemma 4.1. Otherwise, by the definition of r_j^φ , if $|\varphi(p')|$ is the largest one of $|\varphi(z)|$, $z \in [jq_n + 10\eta q_n + 1, (j+1)q_n - 10\eta q_n - 1]$, then $|\varphi(p')| > \max\{r_j^\varphi, r_{j+1}^\varphi\}$. Applying (45) to $\varphi(p')$ and noticing that $\text{dist}(p', q_n\mathbb{Z}) \geq 10\eta q_n$, we get

$$|\varphi(p')| \leq e^{-7(\ln \lambda - 2\eta)\eta q_n} \max\{r_j^\varphi, r_{j+1}^\varphi, |\varphi(p')|\}.$$

This is impossible because $|\varphi(p')| > \max\{r_j^\varphi, r_{j+1}^\varphi\}$.

Now we turn to the proof of case (ii). Notice that in proving case (i) of Lemma 4.1, we only used case (i) of Theorem 3.5. Using case (ii) of Theorem 3.5 instead, we can prove case (ii) of Lemma 4.1 by the same reasoning. In order to avoid repetition, we omit the details. \square

Lemma 4.1 is sufficient for our current purposes, but for the purposes of Section 7 we will need a similar statement that allows for shifts and reflections. For $B \in \mathbb{Z}$, let $r_{j,\pm}^{n,\varphi}(B) = \sup_{|r| \leq 10\eta} |\varphi(B \pm (jq_n + rq_n))|$. For $y \in [B \pm jq_n \pm \eta q_n, B \pm (j+1)q_n \mp \eta q_n]$, let n_0 be the least positive integer such that $4q_{n-n_0} \leq \text{dist}(y - B, q_n\mathbb{Z})$, and let s be the largest positive integer such that $4sq_{n-n_0} \leq \text{dist}(y - B, q_n\mathbb{Z})$. Since we only used the appropriate regularity of the non-resonant y , the proof of Lemma 4.1 also establishes the following lemma.

LEMMA 4.2. *Suppose that for any $y \in [B \pm jq_n \pm \eta q_n, B \pm (j+1)q_n \mp \eta q_n]$, y is $(\ln \lambda + 8 \ln(sq_{n-n_0}/q_{n-n_0+1})/q_{n-n_0} - \varepsilon, 4sq_{n-n_0} - 1)$ regular with $\delta = \frac{1}{4}$. Let $k - B \in \pm[jq_n, (j+1)q_n]$ with $\text{dist}(k - B, q_n\mathbb{Z}) \geq 10\eta q_n$. Suppose either*

- (i) $|j| \leq 48C_* \frac{b_{n+1}}{q_n}$ and $b_{n+1} \geq \frac{q_n}{2}$, or
- (ii) $j = 0$.

Then for sufficiently large n , we have

$$(46) \quad |\varphi(k)| \leq \max \left\{ r_{j,\pm}^\varphi(B) \exp\{-(\ln \lambda - 2\eta)(d_j - 3\eta q_n)\}, \right. \\ \left. r_{j+1,\pm}^\varphi(B) \exp\{-(\ln \lambda - 2\eta)(d_{j+1} - 3\eta q_n)\} \right\},$$

where $d_j = |k - B \mp jq_n|$ and $d_{j+1} = |k - B \mp (j+1)q_n|$.

By Theorem 3.5, Lemma 4.1 is a particular case of Lemma 4.2, when $B = 0$ and the sign is a $+$. Going back to this case, we will prove

LEMMA 4.3. For $1 \leq j \leq 46C_* \frac{b_{n+1}}{q_n}$ with $b_{n+1} \geq \frac{q_n}{2}$, the following holds:

$$(47) \quad r_j^\varphi \leq \max\{r_{j\pm 1}^\varphi \frac{q_{n+1}}{j} \exp\{-(\ln \lambda - C\eta)q_n\}\}.$$

Proof. Fix j with $1 \leq j \leq 46C_* \frac{b_{n+1}}{q_n}$ and $|r| \leq 10\eta q_n$. Set $I_1, I_2 \subset \mathbb{Z}$ as follows:

$$I_1 = [-\lfloor \frac{1}{2}q_n \rfloor, q_n - \lfloor \frac{1}{2}q_n \rfloor - 1],$$

$$I_2 = [jq_n - \lfloor \frac{1}{2}q_n \rfloor, (j+1)q_n - \lfloor \frac{1}{2}q_n \rfloor - 1].$$

Let $\theta_m = \theta + m\alpha$ for $m \in I_1 \cup I_2$. The set $\{\theta_m\}_{m \in I_1 \cup I_2}$ consists of $2q_n$ elements.

By arguments similar to those in Lemma 9.13 in [AJ09] or Theorem 3.1 in [LY15a], one has $\{\theta_m\}$ is $\frac{\ln q_{n+1} - \ln j}{2q_n} + \varepsilon$ uniform for any $\varepsilon > 0$. Since our case is slightly different, we prove it as Theorem B.5 in Appendix B. Combining with Lemma 3.4, there exists some j_0 with $j_0 \in I_1 \cup I_2$ such that $\theta_{j_0} \notin A_{2q_n-1, \ln \lambda - \frac{\ln q_{n+1} - \ln j}{2q_n} - \eta}$.

First, we assume $j_0 \in I_2$. Set $I = [j_0 - q_n + 1, j_0 + q_n - 1] = [x_1, x_2]$. In (32), let $\varepsilon = \eta$. Combining with (30) and (31), it is easy to verify

$$|G_I(jq_n + r, x_i)| \leq e^{(\ln \lambda + \eta)(2q_n - 1 - |jq_n + r - x_i|) - (2q_n - 1)(\ln \lambda - \frac{\ln q_{n+1} - \ln j}{2q_n} - \eta)}.$$

Using (33), we obtain

$$(48) \quad |\varphi(jq_n + r)| \leq \sum_{i=1,2} \frac{q_{n+1}}{j} e^{5\eta q_n} |\varphi(x'_i)| e^{-|jq_n + r - x_i| \ln \lambda},$$

where $x'_1 = x_1 - 1$ and $x'_2 = x_2 + 1$.

Let $d_j^i = |x_i - jq_n|$, $i = 1, 2$. It is easy to check that

$$(49) \quad |jq_n + r - x_i| + d_j^i, |jq_n + r - x_i| + d_{j\pm 1}^i \geq q_n - |r|$$

and

$$(50) \quad |jq_n + r - x_i| + d_{j\pm 2}^i \geq 2q_n - |r|.$$

If $\text{dist}(x_i, q_n\mathbb{Z}) \geq 10\eta q_n$, then we bound $\varphi(x_i)$ in (48) using (38). If $\text{dist}(x_i, q_n\mathbb{Z}) \leq 10\eta q_n$, then we bound $\varphi(x_i)$ in (48) by some proper r_j . Combining with (49) and (50), we have

$$r_j^\varphi \leq \max\left\{r_{j\pm 1}^\varphi \frac{q_{n+1}}{j} \exp\{-(\ln \lambda - C\eta)q_n\},\right. \\ \left. r_j^\varphi \frac{q_{n+1}}{j} \exp\{-(\ln \lambda - C\eta)q_n\}, r_{j\pm 2}^\varphi \frac{q_{n+1}}{j} \exp\{-2(\ln \lambda - C\eta)q_n\}\right\}.$$

However,

$$\begin{aligned} r_j^\varphi &\leq r_j^\varphi \frac{q_{n+1}}{j} \exp\{-(\ln \lambda - C\eta)q_n\} \\ &\leq r_j^\varphi \exp\{-(\ln \lambda - \beta - C\eta)q_n\} \end{aligned}$$

cannot happen, so we must have

$$(51) \quad r_j^\varphi \leq \max\{r_{j\pm 1}^\varphi \frac{q_{n+1}}{j} \exp\{-(\ln \lambda - C\eta)q_n\}, r_{j\pm 2}^\varphi \frac{q_{n+1}}{j} \exp\{-2(\ln \lambda - C\eta)q_n\}\}.$$

In particular,

$$(52) \quad r_j^\varphi \leq \exp\{-(\ln \lambda - \beta - C\eta)q_n\} \max\{r_{j\pm 1}^\varphi r_{j\pm 2}^\varphi\}.$$

If $j_0 \in I_1$, then (52) holds for $j = 0$. Let $\varphi = \phi$ in (52). We get

$$|\phi(0)|, |\phi(-1)| \leq \exp\{-(\ln \lambda - \beta - C\eta)q_n\};$$

this is in contradiction to $|\phi(0)|^2 + |\phi(-1)|^2 = 1$. Therefore $j_0 \in I_2$, so (51) holds for any φ .

By (25) and (28), we have

$$(53) \quad \left\| \begin{pmatrix} \varphi(k_1) \\ \varphi(k_1 - 1) \end{pmatrix} \right\| \geq C e^{-(\ln \lambda + \varepsilon)|k_1 - k_2|} \left\| \begin{pmatrix} \varphi(k_2) \\ \varphi(k_2 - 1) \end{pmatrix} \right\|.$$

This implies

$$r_{j\pm 2}^\varphi \leq r_{j\pm 1}^\varphi \exp\{(\ln \lambda + C\eta)q_n\},$$

and thus (51) becomes

$$(54) \quad r_j^\varphi \leq \max\{r_{j\pm 1}^\varphi \frac{q_{n+1}}{j} \exp\{-(\ln \lambda - C\eta)q_n\}\}$$

for any $1 \leq j \leq 46C_* \frac{b_{n+1}}{q_n}$. □

For solutions ϕ and ψ , we can also get a more subtle estimate.

THEOREM 4.4. *For $1 \leq j \leq 10 \frac{b_{n+1}}{q_n}$ with $b_{n+1} \geq \frac{q_n}{2}$, the following holds:*

$$(55) \quad r_j \leq r_{j-1} \exp\{-(\ln \lambda - C\eta)q_n\} \frac{q_{n+1}}{j}.$$

Proof. Let $\varphi = \phi$ in Lemma 4.3. We must have

$$(56) \quad r_j \leq \max\{r_{j\pm 1} \frac{q_{n+1}}{j} \exp\{-(\ln \lambda - C\eta)q_n\}\}$$

for any $1 \leq j \leq 46C_* \frac{b_{n+1}}{q_n}$.

Suppose that for some $1 \leq j \leq 10 \frac{b_{n+1}}{q_n}$, the following holds:

$$(57) \quad r_j \leq r_{j+1} \frac{q_{n+1}}{j} \exp\{-(\ln \lambda - C\eta)q_n\} \leq r_{j+1} \exp\{-(\ln \lambda - \beta - C\eta)q_n\}.$$

Applying (56) to $j + 1$, we obtain

$$(58) \quad r_{j+1} \leq \max\{r_j, r_{j+2}\} \frac{q_{n+1}}{j+1} \exp\{-(\ln \lambda - C\eta)q_n\}.$$

Combining with (57), we must have

$$(59) \quad r_{j+1} \leq r_{j+2} \exp\{-(\ln \lambda - \beta - C\eta)q_n\}.$$

Generally, for any $0 < p \leq (C_* + 1)j - 1$, we obtain

$$(60) \quad r_{j+p} \leq r_{j+p+1} \exp\{-(\ln \lambda - \beta - C\eta)q_n\}.$$

Thus

$$(61) \quad r_{(C_*+1)j} \geq r_j \exp\{(\ln \lambda - \beta - C\eta)C_*j q_n\}.$$

Clearly, by (53), one has

$$r_j \geq \exp\{-(\ln \lambda + C\eta)j q_n\}.$$

Then

$$(62) \quad r_{(C_*+1)j} \geq \exp\{((C_* - 1) \ln \lambda - C_*\beta - C\eta)j q_n\}.$$

By the definition of C_* , one has

$$(C_* - 1) \ln \lambda - C_*\beta > 0.$$

Thus (62) is in contradiction to the fact that $|\phi(k)| \leq 1 + |k|$.

Now that (57) cannot happen, from (56), we must have

$$(63) \quad r_j \leq r_{j-1} \frac{q_{n+1}}{j} \exp\{-(\ln \lambda - C\eta)q_n\}. \quad \square$$

THEOREM 4.5. For $0 \leq j \leq 8 \frac{b_{n+1}}{q_n}$ with $b_{n+1} \geq \frac{q_n}{2}$, the following holds:

$$(64) \quad R_j \leq R_{j+1} \exp\{-(\ln \lambda - C\eta)q_n\} \frac{q_{n+1}}{j+1}.$$

Proof. If $j = 0$, then (64) holds directly by (55) (applying it with $j = 1$) and (24). Now we consider $j \geq 1$. Let $\varphi = \psi$ in Lemma 4.3. Then (54) also holds for R_j with $j \geq 1$; that is,

$$(65) \quad R_j \leq \max\{R_{j\pm 1} \frac{q_{n+1}}{j+1} \exp\{-(\ln \lambda - C\eta)q_n\}\}.$$

Suppose that for some $j \geq 1$,

$$(66) \quad R_j \leq R_{j-1} \frac{q_{n+1}}{j+1} \exp\{-(\ln \lambda - C\eta)q_n\}.$$

Applying (65) to $j - 1$ and taking into account (66), one has

$$(67) \quad R_{j-1} \leq R_{j-2} \frac{q_{n+1}}{j} \exp\{-(\ln \lambda - C\eta)q_n\}.$$

Iterating j times, we must have

$$(68) \quad R_j \leq R_0 \frac{q_{n+1}^j}{(j+1)!} \exp\{-(\ln \lambda - C\eta)jq_n\} \leq R_0 \exp\{-(\ln \lambda - \beta - C\eta)jq_n\}.$$

Similarly, iterating (55) j times, we have

$$(69) \quad r_j \leq r_0 \exp\{-(\ln \lambda - \beta - C\eta)jq_n\}.$$

Equations (68) and (69) contradict (24). This implies (66) cannot happen. Thus we must have (64). \square

5. Lower bounds on decaying solution in the resonant case

In this section we assume θ is n -Diophantine with respect to α . We will study the lower bound on ϕ for the resonant sites. Recall that $b_{n+1} \geq \frac{q_n}{2}$ in this case.

THEOREM 5.1. *Let $\tilde{r}_j = \left\| \begin{pmatrix} \phi(jq_n) \\ \phi(jq_{n-1}) \end{pmatrix} \right\|$. Suppose $1 \leq j \leq 8\frac{b_{n+1}}{q_n}$ with $b_{n+1} \geq \frac{q_n}{2}$. Then we must have*

$$(70) \quad \tilde{r}_j \geq \frac{q_{n+1}}{j} e^{-(\ln \lambda + \varepsilon)q_n} \tilde{r}_{j-1}.$$

We first list two standard facts.

LEMMA 5.2 ([Sim85]). *Let A^1, A^2, \dots, A^n and B^1, B^2, \dots, B^n be 2×2 matrices with $\| \prod_{m=0}^{\ell-1} A^{j+m} \| \leq Ce^{d\ell}$ for some constant C and d . Then*

$$\| (A^n + B^n) \cdots (A^1 + B^1) - A^n \cdots A^1 \| \leq Ce^{dn} \left(\prod_{j=1}^n (1 + Ce^{-d} \|B^j\|) - 1 \right).$$

LEMMA 5.3. *For any $\varepsilon > 0$ and large n , the following hold:*

$$(71) \quad \|A_{q_n}(\theta + q_n\alpha) - A_{q_n}(\theta)\| \leq \frac{1}{q_{n+1}} e^{(\ln \lambda + \varepsilon)q_n}$$

and

$$(72) \quad \|A_{q_n}^{-1}(\theta + q_n\alpha) - A_{q_n}^{-1}(\theta)\| \leq \frac{1}{q_{n+1}} e^{(\ln \lambda + \varepsilon)q_n}.$$

Proof. We only prove (71) for simplicity. By the DC approximation (see (150) in Appendix B), we have

$$\|q_n\alpha\|_{\mathbb{R}/\mathbb{Z}} \leq \frac{1}{q_{n+1}}.$$

This implies

$$\|A(\theta + q_n\alpha) - A(\theta)\| \leq \frac{C}{q_{n+1}}.$$

Applying Lemma 5.2 and (28), one has

$$(73) \quad \|A_{q_n}(\theta + q_n\alpha) - A_{q_n}(\theta)\| \leq e^{(\ln \lambda + \varepsilon)q_n} \left(\left(1 + \frac{C}{q_{n+1}}\right)^{q_n} - 1 \right).$$

Using the fact $|e^y - 1| \leq ye^y$ for $y > 0$, we obtain

$$\begin{aligned} \left(1 + \frac{C}{q_{n+1}}\right)^{q_n} - 1 &\leq q_n \left(1 + \frac{C}{q_{n+1}}\right)^{q_n} \ln \left(1 + \frac{C}{q_{n+1}}\right) \\ &\leq C \frac{q_n}{q_{n+1}}. \end{aligned}$$

Combining this with (73) completes the proof. □

LEMMA 5.4. *For any $0 \leq j \leq 8 \frac{b_{n+1}}{q_n} - 1$, one of the following two estimates must hold:*

$$(74) \quad \tilde{r}_{j+1} \geq \frac{q_{n+1}}{j+1} e^{-(\ln \lambda + \varepsilon)q_n} \tilde{r}_j$$

or

$$(75) \quad \tilde{r}_{j+1} \tilde{r}_{j-1} \geq \left(1 - \frac{1}{10(j+1)}\right)^2 \left(1 - \frac{1}{10(j+1)^2}\right) \tilde{r}_j^2.$$

Proof. Suppose

$$(76) \quad \tilde{r}_{j+1} \leq \frac{q_{n+1}}{j+1} e^{-(\ln \lambda + \varepsilon)q_n} \tilde{r}_j.$$

Let $U_j = \begin{pmatrix} \phi(jq_n) \\ \phi(jq_{n-1}) \end{pmatrix}$. Then for $n > 0$, one has

$$U_j = A_{q_n}(\theta + (j-1)q_n\alpha)U_{j-1}.$$

Denote $B = A_{q_n}(\theta + jq_n\alpha)$. Notice that $\det B = 1$. We have

$$(77) \quad B^2 + (\text{Tr } B)B + I = 0.$$

Case 1: $\text{Tr } B \leq \frac{\tilde{r}_j}{\gamma \tilde{r}_{j+1}}$, where

$$1 - \frac{1}{\gamma} = \frac{1}{10(j+1)}.$$

Applying (77) to U_j , one has

$$(78) \quad B^2U_j + (\text{Tr } B)BU_j + U_j = 0.$$

Notice that $U_{j+1} = BU_j$; thus

$$\|(\text{Tr } B)BU_j\| \leq \frac{1}{\gamma} \tilde{r}_j.$$

Therefore we have

$$(79) \quad \|B^2U_j\| \geq \left(1 - \frac{1}{\gamma}\right) \tilde{r}_j = \frac{1}{10(j+1)} \tilde{r}_j.$$

This is impossible. Indeed, from the following estimate,

$$\begin{aligned} \|U_{j+2} - B^2U_j\| &\leq \|A_{q_n}(\theta + (j + 1)q_n\alpha) - A_{q_n}(\theta + jq_n\alpha)\| \|U_{j+1}\| \\ &\leq e^{(\ln \lambda + \frac{1}{2}\varepsilon)q_n} \frac{1}{q_{n+1}} \tilde{r}_{j+1} \\ &\leq \frac{1}{100(j + 1)} \tilde{r}_j, \end{aligned}$$

where the second inequality holds by (71) and the third inequality holds by assumption (76), combining with (79), one has

$$(80) \quad \|U_{j+2}\| = \tilde{r}_{j+2} \geq \frac{9}{100(j + 1)} \tilde{r}_j.$$

However, by (55) and (53),

$$\tilde{r}_{j+2} \leq \frac{q_{n+1}^2}{(j + 1)(j + 2)} e^{-2(\ln \lambda - C\eta)q_n} \tilde{r}_j.$$

This is in contradiction to (80).

Case 2: It remains to consider

$$(81) \quad \text{Tr } B \geq \frac{\tilde{r}_j}{\gamma \tilde{r}_{j+1}}.$$

From (77),

$$(82) \quad BU_j + (\text{Tr } B)U_j + B^{-1}U_j = 0.$$

First by assumption (76), one has

$$\tilde{r}_{j+1} \leq \frac{1}{10(j + 1)} \tilde{r}_j.$$

Then

$$\begin{aligned} \|BU_j\| = \tilde{r}_{j+1} &\leq \frac{\tilde{r}_j}{\gamma \tilde{r}_{j+1}} \frac{\tilde{r}_j}{10(j + 1)^2} \\ &\leq \|(\text{Tr } B)U_j\| \frac{1}{10(j + 1)^2}. \end{aligned}$$

Thus by (82), we have

$$\begin{aligned} \|B^{-1}U_j\| &\geq \left(1 - \frac{1}{10(j + 1)^2}\right) \|(\text{Tr } B)U_j\| \\ (83) \quad &\geq \left(1 - \frac{1}{10(j + 1)^2}\right) \frac{\tilde{r}_j^2}{\gamma \tilde{r}_{j+1}} \\ (84) \quad &\geq \left(1 - \frac{1}{10(j + 1)^2}\right) \frac{1}{\gamma} \frac{j + 1}{q_{n+1}} e^{(\ln \lambda + \varepsilon)q_n} \tilde{r}_j, \end{aligned}$$

where the second inequality holds by (81) and the third inequality holds by (76).

By (72), the following holds:

$$\begin{aligned}
 \|U_{j-1} - B^{-1}U_j\| &\leq \|A_{q_n}^{-1}(\theta + (j-1)q_n\alpha) - A_{q_n}^{-1}(\theta + jq_n\alpha)\| \|U_j\| \\
 &\leq e^{(\ln \lambda + \frac{1}{2}\varepsilon)q_n} \frac{1}{q_{n+1}} \tilde{r}_j \\
 (85) \qquad &\leq \frac{1}{10(j+1)} \|B^{-1}U_j\|,
 \end{aligned}$$

where the third inequality holds by (84).

Putting (83) and (85) together, we have

$$\begin{aligned}
 \tilde{r}_{j-1} = \|U_{j-1}\| &\geq (1 - \frac{1}{10(j+1)}) \|B^{-1}U_j\| \\
 &\geq (1 - \frac{1}{10(j+1)})^2 (1 - \frac{1}{10(j+1)^2}) \frac{\tilde{r}_j^2}{\tilde{r}_{j+1}}.
 \end{aligned}$$

This implies (75). □

Proof of Theorem 5.1. We can proceed by induction.

Set $j = 0$ in Lemma 5.4. By (55), the second case (75) cannot happen; thus Theorem 5.1 holds for $j = 1$.

Suppose (70) holds for $p = j - 1$; that is,

$$(86) \qquad \tilde{r}_{j-1} \geq \frac{q_{n+1}}{j-1} e^{-(\ln \lambda + \varepsilon)q_n} \tilde{r}_{j-2}.$$

We will show that (70) holds for $p = j$. Let us apply Lemma 5.4 to $p = j - 1$.

If (74) holds for $p = j - 1$, the result follows. Otherwise by (75), we have

$$\begin{aligned}
 \tilde{r}_j &\geq (1 - \frac{1}{10j})^2 (1 - \frac{1}{10j^2}) \tilde{r}_{j-1} \frac{\tilde{r}_{j-1}}{\tilde{r}_{j-2}} \\
 &\geq (1 - \frac{1}{10j})^2 (1 - \frac{1}{10j^2}) \tilde{r}_{j-1} \frac{q_{n+1}}{j-1} e^{-(\ln \lambda + \varepsilon)q_n} \\
 &\geq \tilde{r}_{j-1} \frac{q_{n+1}}{j} e^{-(\ln \lambda + \varepsilon)q_n},
 \end{aligned}$$

where the second inequality holds by (86). □

6. Decaying solutions. Proof of (12), Theorems 22, 24 and 29

In this section the dependence on n will play a role, so we go back to the r_j^n, \tilde{r}_j^n notation. We first give a series of auxiliary facts. Recall footnote 11.

THEOREM 6.1. *Assume θ is n -Diophantine with respect to α . For any $1 \leq j \leq 10 \frac{b_{n+1}}{q_n}$ with $b_{n+1} \geq \frac{q_n}{2}$, we have*

$$\bar{r}_j^n e^{-\varepsilon j q_n} \leq r_j^n \leq \bar{r}_j^n e^{\varepsilon j q_n}$$

and

$$\bar{r}_j^n e^{-\varepsilon j q_n} \leq \tilde{r}_j^n \leq \bar{r}_j^n e^{\varepsilon j q_n}.$$

Proof. For any $\varepsilon > 0$, we choose η small enough. Using (55) j times, we have

$$r_j^n \leq \frac{q_{n+1}^j}{j!} \exp\{-(\ln \lambda - \varepsilon)jq_n\}.$$

Similarly, using (70) j times, we have

$$\tilde{r}_j^n \geq \frac{q_{n+1}^j}{j!} \exp\{-(\ln \lambda + \varepsilon)jq_n\}.$$

By the Stirling formula and (53), we obtain the theorem. □

THEOREM 6.2. *Assume θ is n -Diophantine with respect to α . Assume $jq_n \leq k < (j + 1)q_n$ with $0 \leq j \leq 8\frac{b_{n+1}}{q_n}$, $b_{n+1} \geq \frac{q_n}{2}$ and $k \geq \frac{q_n}{4}$. We have*

$$(87) \quad \|U(k)\| \leq \max\{e^{-|k-jq_n| \ln \lambda \tilde{r}_j^n}, e^{-|k-(j+1)q_n| \ln \lambda \tilde{r}_{j+1}^n}\} e^{\varepsilon q_n},$$

$$(88) \quad \|U(k)\| \geq \max\{e^{-|k-jq_n| \ln \lambda \tilde{r}_j^n}, e^{-|k-(j+1)q_n| \ln \lambda \tilde{r}_{j+1}^n}\} e^{-\varepsilon q_n}.$$

In particular, we have

$$(89) \quad \|U(k)\| \leq \max\{e^{-|k-jq_n| \ln \lambda \tilde{r}_j^n}, e^{-|k-(j+1)q_n| \ln \lambda \tilde{r}_{j+1}^n}\} e^{\varepsilon |k|},$$

$$(90) \quad \|U(k)\| \geq \max\{e^{-|k-jq_n| \ln \lambda \tilde{r}_j^n}, e^{-|k-(j+1)q_n| \ln \lambda \tilde{r}_{j+1}^n}\} e^{-\varepsilon |k|}.$$

Proof. (89) and (90) just follows from (87), (88) and Theorem 6.1. Thus it suffices to prove (87) and (88). Clearly, by (53), one has

$$\|U(k)\| \geq \max\{e^{-|k-jq_n| \ln \lambda \tilde{r}_j^n}, e^{-|k-(j+1)q_n| \ln \lambda \tilde{r}_{j+1}^n}\} e^{-\varepsilon q_n}.$$

This implies (90) by Theorem 6.1.

We now turn to (89). If $|k - jq_n| \leq 10\eta q_n$ or $|k - (j + 1)q_n| \leq 10\eta q_n$, the result follows from Theorem 6.1 and (53). If $|k - jq_n| \geq 10\eta q_n$ and $|k - (j + 1)q_n| \geq 10\eta q_n$, it follows from Lemma 4.1, Theorem 6.1 and (53). □

THEOREM 6.3. *For $q_n^{\frac{8}{9}} \leq k \leq \frac{q_n}{2}$, let n_0 be the smallest positive integer such that $q_{n-n_0} \leq k < q_{n-n_0+1}$. Suppose $jq_{n-n_0} \leq k < (j+1)q_{n-n_0}$ with $j \geq 1$. If θ is k -Diophantine with respect to α for $k = n - n_0$ and $k = n - 1$, then*

$$(91) \quad \|U(k)\| \leq \max\{e^{-|k-jq_{n-n_0}| \ln \lambda \tilde{r}_j^{n-n_0}}, e^{-|k-(j+1)q_{n-n_0}| \ln \lambda \tilde{r}_{j+1}^{n-n_0}}\} e^{\varepsilon k}$$

and

$$(92) \quad \|U(k)\| \geq \max\{e^{-|k-jq_{n-n_0}| \ln \lambda \tilde{r}_j^{n-n_0}}, e^{-|k-(j+1)q_{n-n_0}| \ln \lambda \tilde{r}_{j+1}^{n-n_0}}\} e^{-\varepsilon k}.$$

Proof. Set $t_0 = 1 - \frac{\varepsilon}{8\beta}$. Let $t = t_0$ in the definition of resonance, i.e., $b_n = q_n^{t_0}$.

Case 1: $k \leq q_{n-n_0+1}^{t_0}$. In this case, one has

$$q_{n-n_0} \leq k \leq q_{n-n_0+1}^{t_0}.$$

The result holds by Theorem 6.2.

Case 2: $k \geq q_{n-n_0+1}^{t_0}$. Then

$$\begin{aligned} \bar{r}_j^{n-n_0} &\leq \exp\left\{-\left(\ln \lambda - \frac{\ln q_{n-n_0+1}}{q_{n-n_0}} + \frac{\ln q_{n-n_0+1}^{t_0}}{q_{n-n_0}} - \varepsilon\right)jq_{n-n_0}\right\} \\ &\leq \exp\left\{-\left(\ln \lambda - (1-t_0)\beta - \varepsilon\right)jq_{n-n_0}\right\} \\ &\leq \exp\left\{-\left(\ln \lambda - 2\varepsilon\right)jq_{n-n_0}\right\}, \end{aligned}$$

where the third inequality holds by the definition of t_0 . Noting that $k \leq q_{n-n_0+1}$, one has

$$\bar{r}_j^{n-n_0} \geq \exp\{-jq_{n-n_0}(\ln \lambda + \varepsilon)\}.$$

Similarly,

$$\exp\{-(j+1)q_{n-n_0}(\ln \lambda + \varepsilon)\} \leq \bar{r}_{j+1}^{n-n_0} \leq \exp\{-(j+1)q_{n-n_0}(\ln \lambda - \varepsilon)\}.$$

Thus in order to prove Case 2, it suffices to show

$$(93) \quad e^{-(\ln \lambda + \varepsilon)k} \leq \|U(k)\| \leq e^{-(\ln \lambda - \varepsilon)k}.$$

The left inequality holds by (53).

We start to prove the right inequality. For any $y \in [\varepsilon k, k]$ or $y \in [q_n - k, q_n - \varepsilon k]$, let n'_0 be the least positive integer such that $4q_{n-n'_0} \leq \text{dist}(y, q_n\mathbb{Z})$, and thus $n'_0 \geq n_0$. Let s be the largest positive integer such that $4sq_{n-n'_0} \leq \text{dist}(y, q_n\mathbb{Z})$.

Set $I_1, I_2 \subset \mathbb{Z}$ as follows:

$$\begin{aligned} I_1 &= [-sq_{n-n'_0}, sq_{n-n'_0} - 1], \\ I_2 &= [y - sq_{n-n'_0}, y + sq_{n-n'_0} - 1], \end{aligned}$$

and let $\theta_j = \theta + j\alpha$ for $j \in I_1 \cup I_2$. The set $\{\theta_j\}_{j \in I_1 \cup I_2}$ consists of $4sq_{n-n'_0}$ elements. By case (ii) of Theorem 3.5, y is $(\ln \lambda + 8 \ln(sq_{n-n'_0}/q_{n-n'_0+1})/q_{n-n'_0} - \varepsilon, 4sq_{n-n'_0} - 1)$ regular with $\delta = \frac{1}{4}$. Notice that

$$(s+1)q_{n-n'_0} \geq \varepsilon q_{n-n_0+1}^{t_0} \geq \varepsilon q_{n-n'_0+1}^{t_0}.$$

Thus we have

$$\begin{aligned} \ln \lambda + 8 \ln(sq_{n-n'_0}/q_{n-n'_0+1})/q_{n-n'_0} &\geq \ln \lambda - 8(1-t_0)\beta - \varepsilon \\ &\geq \ln \lambda - 2\varepsilon. \end{aligned}$$

This implies that for any $y \in [\varepsilon k, k]$ or $y \in [q_n - k, q_n - \varepsilon k]$, there exists an interval $I(y) = [x_1, x_2] \subset [0, q_n]$ with $y \in I(y)$ such that

$$(94) \quad \text{dist}(y, \partial I(y)) \geq \frac{1}{2}q_{n-n'_0}$$

and

$$(95) \quad |G_{I(y)}(y, x_i)| \leq e^{-(\ln \lambda - \varepsilon)|y-x_i|}, \quad i = 1, 2.$$

For any $y \in (k, q_n - k)$, let s be the largest positive integer such that $sq_{n-n_0} \leq \text{dist}(y, q_n\mathbb{Z})$, and set $I_1, I_2 \subset \mathbb{Z}$ as follows:

$$I_1 = \left[-\left\lfloor \frac{sq_{n-n_0}}{2} \right\rfloor, sq_{n-n_0} - \left\lfloor \frac{sq_{n-n_0}}{2} \right\rfloor - 1 \right],$$

$$I_2 = \left[y - \left\lfloor \frac{sq_{n-n_0}}{2} \right\rfloor, y + sq_{n-n_0} - \left\lfloor \frac{sq_{n-n_0}}{2} \right\rfloor - 1 \right].$$

By the same reason, (94) and (95) also hold for $n'_0 = n_0$.

Arguing exactly as in the proof of Lemma 4.1, with (39) replaced with (94) and (40) with (95), we obtain

(96)

$$\|U(k)\| \leq \max\{\hat{r}_0 \exp\{-(\ln \lambda - 2\varepsilon)(k - 3\varepsilon k)\}, \hat{r}_1^n \exp\{-(\ln \lambda - 2\varepsilon)(q_n - k - 3\varepsilon k)\}\},$$

where $\hat{r}_j = \max_{|r| \leq 10\varepsilon} \|U(jq_n + rk)\|$ with $j = 0, 1$. Using that $k \leq \frac{q_n}{2}$, one has

$$\|U(k)\| \leq e^{-(\ln \lambda - \varepsilon)k}.$$

This implies (93) and thus the theorem. □

Remark 6.4. The assumption that θ is $n - n_0$ Diophantine with respect to α is sufficient for the proof of Case 1. The assumption that θ satisfies (35) is sufficient for the proof of case (ii) of Theorem 3.5, and therefore for the proof of Case 2. Then by Remark 3.6, the assumption that θ is $n - 1$ Diophantine with respect to α is sufficient for the proof of Case 2.

Remark 6.5. Suppose we only consider $q_n^{\frac{8}{9}} \leq k \leq cq_n$ with $c \leq \frac{1}{2}$ in Theorem 6.3. Then Theorem 6.3 still holds if we only have (11) for function $\phi(k)$ on $[-cq_n, 2cq_n]$ for some $c > 0$.

In order to prove (12), it suffices to prove the following theorem, which is a stronger local version of (12).

THEOREM 6.6. *Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ be such that $|\lambda| > e^{\beta(\alpha)}$. Suppose E is a generalized eigenvalue of $H_{\lambda, \alpha, \theta}$ and ϕ is the generalized eigenfunction. Let $U(k) = \begin{pmatrix} \phi(k) \\ \phi(k-1) \end{pmatrix}$. Then for any $\varepsilon > 0, \kappa > 0, \nu > 1$, there exists \hat{n}_0 (depending on $\alpha, E, \kappa, \nu, \varepsilon$ ¹²) such that, if θ is n -Diophantine with respect to α with Diophantine constants κ, ν for some $n \geq \hat{n}_0$, we have that $U(k)$ satisfies*

(97)
$$f(|k|)e^{-\varepsilon|k|} \leq \|U(k)\| \leq f(|k|)e^{\varepsilon|k|}$$

for $\frac{q_n}{2} \leq |k| \leq \frac{q_{n+1}}{2}$.

Proof. It remains to collect several already proved statements that cover different scenarios.

¹²The dependence on E is through the constant \hat{C} in (11).

Case (i): $\frac{q_n}{2} \leq q_{n+1}^{\frac{8}{9}}$. For $\frac{q_n}{2} \leq k \leq 4q_{n+1}^{\frac{8}{9}}$ the result follows from Theorem 6.2. For $4q_{n+1}^{\frac{8}{9}} \leq k \leq \frac{q_{n+1}}{2}$, (97) follows from Theorem 6.3. (Notice that now $k \geq 2q_n$, so $n_0 = 1$.)

Case (ii): $q_{n+1}^{\frac{8}{9}} \leq \frac{q_n}{2}$.

Case (ii.1): $\frac{q_n}{2} \leq k \leq \min\{q_n, \frac{q_{n+1}}{2}\}$. If $q_n = q_{n-1} + q_{n-2}$, then $q_{n-1} \geq \frac{q_n}{2}$. By the proof of Case 2 in Theorem 6.3 (by Remark 6.4, the assumption that θ is n -Diophantine is enough), for any $q_{n-1} \leq k \leq \min\{q_n, \frac{q_{n+1}}{2}\}$, one has

$$|\phi(k)| \leq e^{-(\ln \lambda - \varepsilon)k}.$$

This leads to

$$|\phi(k)| \leq e^{-(\ln \lambda - \varepsilon)k}$$

for $\frac{q_n}{2} \leq k \leq \min\{q_n, \frac{q_{n+1}}{2}\}$. This also implies (12).

If $q_n = jq_{n-1} + q_{n-2}$ with $j \geq 2$, we have $\frac{q_n}{2} \geq q_{n-1}$. By Case 2 in Theorem 6.3 (by Remark 6.4, the assumption that θ is n -Diophantine is enough) again (with $n + 1 - n_0 = n - 1$), we obtain (97).

Case (ii.2): $q_n \leq k \leq \frac{q_{n+1}}{2}$. In this case (97) follows directly from Theorem 6.3 (with $n + 1 - n_0 = n$), because $n_0 = 1$, so that the fact that θ is n Diophantine can guarantee both Cases 1 and 2 of Theorem 6.3. \square

Proof of Theorem 2.2. The proof follows that of (97) by shifting by k_0 units and Remark 6.5. \square

Proof of Theorem 2.9. Assume θ is Diophantine with respect to α . First by the definition of $\beta(\alpha)$, for any large n and any ℓ , one has

$$\bar{r}_\ell^n \leq e^{-(\ln \lambda - \beta - \varepsilon)\ell q_n}.$$

Combining with the definition of $f(k)$ and (12), we have

$$(98) \quad |\phi(k)| \leq e^{-(\ln \lambda - \beta - \varepsilon)k}.$$

We therefore established that every generalized eigenfunction decays exponentially, which by Schnol’s Theorem [Ber68] implies the localization statement.

By the definition of $\beta(\alpha)$ again, there exists a subsequence q_{n_k} of q_n such that

$$(99) \quad q_{n_k+1} \geq e^{(\beta - \varepsilon)q_{n_k}}.$$

By Theorem 2.1 (or Theorem 6.1) and the definition of \bar{r}_j^n , for any $k > 0$, we have

$$(100) \quad \|U(q_{n_k})\| \geq e^{-(\ln \lambda - \beta + \varepsilon)q_{n_k}}.$$

Thus (98) and (100) imply that the decay rate is just $\ln \lambda - \beta$. \square

7. Hierarchical structure

As we have already established Theorem 2.9 we know that each generalized eigenfunction decays exponentially so has a global maximum. Assume its global maximum (see footnote 2) is at 0 and ϕ is normalized by $\|\phi\|_\infty = 1$. Note that then \hat{C} in (11) is equal to 1 and so all dependence of the largeness of n on E (see footnote 12) disappears. Theorem 2.1 provides, for sufficiently large n , plenty of local n -maxima in the vicinity of aq_n , but determined only with εq_n precision. In the next theorem we show that this precision can be improved all the way to an n -independent constant. We have

THEOREM 7.1. *Fix κ, ν, ϵ . Then for sufficiently small ε , there exists $\hat{n}_0(\kappa, \nu, \lambda, \alpha, \epsilon, \varepsilon)$ such that if θ is k -Diophantine for all $\hat{n}_0 \leq k \leq n$ with Diophantine constants κ, ν and $\frac{\ln q_{n+1} - \ln j}{q_n} > \epsilon \ln \lambda$ with $\epsilon > 0$. Then*

$$(101) \quad \sup_{k \in [jq_n - \epsilon q_n + \varepsilon q_n, jq_n]} \|U(k)\| = \sup_{k \in [jq_n - K_0, jq_n]} \|U(k)\|$$

and

$$(102) \quad \sup_{k \in [jq_n, jq_n + \varepsilon q_n - \epsilon q_n]} \|U(k)\| = \sup_{k \in [jq_n, jq_n + K_0]} \|U(k)\|,$$

where $K_0 = q_{\hat{n}_0+1}$.

Proof. We first give the proof of (101). Let $k_0 \in [jq_n - \varepsilon q_n, jq_n]$ be such that

$$\|U(k_0)\| = \sup_{k \in [jq_n - \varepsilon q_n, jq_n]} \|U(k)\|.$$

By (55), (70), (87) and (88), one has

$$\|U(k_0)\| = \sup_{k \in [jq_n - \epsilon q_n + \varepsilon q_n, jq_n]} \|U(k)\|.$$

Suppose (101) does not hold, i.e., $k_0 \in [jq_n - \varepsilon q_n, jq_n - K_0]$.

Now we will reflect the elements in $[jq_n - \varepsilon q_n, jq_n]$ at $\frac{j}{2}q_n$. That is, for any element $k \in [jq_n - \varepsilon q_n, jq_n]$, let $k' = jq_n - k$. Then $k' \in [0, \varepsilon q_n]$.

Choose n' such that $b_{n'} \leq k'_0 < b_{n'+1}$ (where $k'_0 = jq_n - k_0$). Then $n' \geq \hat{n}_0$.

Case 1: k'_0 is n' -non-resonant, i.e., $\text{dist}(k'_0, q_{n'}\mathbb{Z}) \geq b_{n'}$. Let n'_0 be the least positive integer such that $4q_{n'-n'_0} \leq \text{dist}(k'_0, q_{n'}\mathbb{Z})$. Let s be the largest positive integer such that $4sq_{n'-n'_0} \leq \text{dist}(k'_0, q_{n'}\mathbb{Z})$. Set $I_1, I_2, I'_2 \subset \mathbb{Z}$ as follows:

$$\begin{aligned} I_1 &= [-sq_{n'-n'_0}, sq_{n'-n'_0} - 1], \\ I_2 &= [k_0 - sq_{n'-n'_0} + 1, k_0 + sq_{n'-n'_0}], \\ I'_2 &= [k'_0 - sq_{n'-n'_0}, k'_0 + sq_{n'-n'_0} - 1]. \end{aligned}$$

Notice that I'_2 and I_2 are reflections of each other about $\frac{j}{2}q_n$.

By the Diophantine condition on θ with respect to α , for any $k_2 \in I_2(k'_2 \in I'_2)$ and $k_1 \in I_1$, we have

$$\begin{aligned}
 (103) \quad & \|2\theta + (k_1 + k_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} = \|2\theta - k'_2\alpha + jq_n\alpha + k_1\alpha\|_{\mathbb{R}/\mathbb{Z}} \\
 & \geq \|2\theta + (k_1 - k'_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} - \|jq_n\alpha\|_{\mathbb{R}/\mathbb{Z}} \\
 & \geq \|2\theta + (k_1 - k'_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} - \frac{j}{q_{n+1}} \\
 & \geq \|2\theta + (k_1 - k'_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} - e^{-\epsilon \ln \lambda q_n} \\
 & \geq \frac{1}{2} \|2\theta + (k_1 - k'_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{C}{q_{n'}},
 \end{aligned}$$

where the last inequality holds by the fact $|k_1|, |k'_2| \leq C_* b_{n'+1}$ so that we can apply Lemma B.2.

For any $k_2 \in I_2$ and $k_1 \in I_1$ ($k_1 + k'_2 \neq 0$ by the construction of I_1, I_2), we also have

$$\begin{aligned}
 (104) \quad & \|(k_2 - k_1)\alpha\|_{\mathbb{R}/\mathbb{Z}} = \|-k'_2\alpha + jq_n\alpha - k_1\|_{\mathbb{R}/\mathbb{Z}} \\
 & \geq \|(-k_1 - k'_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} - \|jq_n\alpha\|_{\mathbb{R}/\mathbb{Z}} \\
 & \geq \|(k_1 + k'_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} - e^{-\epsilon \ln \lambda q_n} \\
 & \geq \frac{1}{2} \|(k_1 + k'_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{C}{q_{n'}},
 \end{aligned}$$

where the last inequality holds by the fact $|k_1|, |k'_2| \leq C_* b_{n'+1}$ and $k_1 - k'_2 \neq q_{n'}\mathbb{Z}$ so that we can apply Lemma B.3.

By Theorem B.4, (103) and (104), we have that k_0 is $(\hat{k}_0, \ln \lambda - \beta - \epsilon)$ regular, where $\hat{k}_0 = 4sq_{n'-n'_0} - 1$. Let $I_2 = [x_1, x_2] \subset [jq_n - 2\epsilon q_n, jq_n]$.

By (33), we have

$$|\phi(k_0)| \leq e^{-(\ln \lambda - \beta - \epsilon) \frac{\hat{k}_0}{10}} (|\phi(x_1)| + |\phi(x_0)|) \leq e^{-(\ln \lambda - \beta - \epsilon) \frac{\hat{k}_0}{10}} \|U(k_0)\|.$$

Similarly,

$$|\phi(k_0 - 1)| \leq e^{-(\ln \lambda - \beta - \epsilon) \frac{\hat{k}_0}{10}} \|U(k_0)\|.$$

The last two inequalities imply that

$$\|U(k_0)\| \leq e^{-(\ln \lambda - \beta - \epsilon) \frac{\hat{k}_0}{10}} \|U(k_0)\|.$$

This is impossible.

Case 2: k'_0 is n -resonant, i.e., $|k'_0 - \ell q_{n'}| \leq b_{n'}$ for some ℓ . From (103) and (104), we know that the small divisor condition does not change under reflection at $\frac{j}{2}q_n$. Following the proof of (54) and replacing Lemma 4.1 with a combination of Lemma 4.2 and Theorem B.4, we have

$$\mathbf{r}_{\ell}^{n', \phi} \leq \exp\{-(\ln \lambda - \beta - \epsilon)q_{n'}\} \max\{\mathbf{r}_{\ell \pm 1}^{n', \phi}\},$$

where

$$r_\ell^{n',\phi} = \sup_{|r| \leq 10\epsilon} |\phi(jq_n - (\ell q_{n'} + r q_{n'}))|.$$

This contradicts the fact that k_0 is the maximal point because $|k'_0 - \ell q_{n'}| \leq b_{n'}$. This completes the proof of (101).

Now we turn to the proof of (102). Let $k_0^r \in [jq_n, jq_n + \epsilon q_n]$ be such that

$$||U(k_0^r)|| = \sup_{k \in [jq_n, jq_n + \epsilon q_n]} ||U(k)||.$$

Suppose the theorem does not hold, i.e., $k_0^r \in [jq_n + K_0, jq_n + \epsilon q_n]$. In this case we shift the elements in $[jq_n, jq_n + \epsilon q_n]$ by $-jq_n$. That is for any element $k \in [jq_n, jq_n + \epsilon q_n]$, let $k^{r'} = k - jq_n$. Then $k^{r'} \in [0, \epsilon q_n]$. Then (102) holds by the same proof, only replacing all k' with $k^{r'}$. \square

We restate the result of Theorem 7.1 as a more convenient

THEOREM 7.2. *Fix κ, ν, ϵ . Then for sufficiently small ϵ , there exists $\hat{n}_0(\kappa, \nu, \lambda, \alpha, \epsilon, \epsilon)$ such that if k_0 is a local $(n+1)$ -maximum, θ is k -Diophantine for all $\hat{n}_0 \leq k \leq n$ with Diophantine constants κ, ν , and*

$$\frac{\ln q_{n+1} - \ln j}{q_n} > \epsilon \ln \lambda.$$

Then

$$(105) \quad \sup_{k \in [k_0 + jq_n - \epsilon q_n + \epsilon q_n, k_0 + (j+1)q_n]} ||U(k)|| = \sup_{k \in [k_0 + jq_n - K_0, k_0 + jq_n + K_0]} ||U(k)||,$$

where $K_0 = q_{\hat{n}_0+1}$.

Proof. By shifting the operator by k_0 units, we can assume $k_0 = 0$. Theorem 7.1 still holds if 0 is a local $(n+1)$ -maximum by Remark 6.5. \square

We will now formulate a local version of the hierarchical structure Theorem 2.4. Fix $0 < \varsigma, \epsilon$ with $\varsigma + 2\epsilon < 1$. Let $n_j \rightarrow \infty$ be such that $\ln q_{n_j+1} \geq (\varsigma + 2\epsilon) \ln |\lambda| q_{n_j}$. Let $\mathbf{c}_j = (\ln q_{n_j+1} - \ln |a_{n_j}|) / \ln |\lambda| q_{n_j} - \epsilon$. $\mathbf{c}_j > \epsilon$ for $0 < a_{n_j} < e^{\varsigma \ln |\lambda| q_{n_j}}$.

THEOREM 7.3. *Suppose k_0 is a local $(n_{j_0} + 1)$ -maximum. Suppose $\theta + k_0 \alpha$ is Diophantine with respect to α (with Diophantine constants κ, ν). Then there exists $\hat{n}_0(\alpha, \lambda, \kappa, \nu, \epsilon) < \infty$ such that for any $j_0 > j_1 > \dots > j_k$, $n_{j_k} \geq \hat{n}_0 + k$, and $0 < a_{n_{j_i}} < e^{\varsigma \ln |\lambda| q_{n_{j_i}}}$, $i = 0, 1, \dots, k$ for all $0 \leq s \leq k$, there exists a local n_{j_s} -maximum $b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}}$ on the interval $b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}} + I_{\mathbf{c}_{j_s}, 1}^{n_{j_s}}$ for all $0 \leq s \leq k$ such that the following holds:*

- (I) $|b_{a_{n_{j_0}}} - (k_0 + a_{n_{j_0}} q_{n_{j_0}})| \leq q_{\hat{n}_0+1}$;
 - (II) for any $1 \leq s \leq k$,
- $$|b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}} - (b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_{s-1}}} + a_{n_{j_s}} q_{n_{j_s}})| \leq q_{\hat{n}_0+s+1};$$

(III) if $2(x - b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_k}}}) \in I_{c_{j_k}, 1}^{n_{j_k}}$, then for each $s = 0, 1, \dots, k$,

$$(106) \quad f(x_s)e^{-\varepsilon|x_s|} \leq \frac{\|U(x)\|}{\|U(b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}})\|} \leq f(x_s)e^{\varepsilon|x_s|},$$

where $x_s = |x - b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}}|$ is large enough.

Moreover, every local n_{j_s} -maximum on the interval

$$b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_{s-1}}}} + [-e^{\varepsilon \ln \lambda q_{n_{j_s}}}, e^{\varepsilon \ln \lambda q_{n_{j_s}}}]$$

is of the form $b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}}$ for some $a_{n_{j_s}}$.

Proof. Let $\hat{n}_0 = \hat{n}_0(\kappa, 3\nu, \lambda, \alpha, \varepsilon, \varepsilon/10)$ be given by Theorem 7.2.¹³ As long as

$$(107) \quad (\ln q_{n+1} - \ln |a_n|)/q_n \geq 2\varepsilon \ln |\lambda|$$

with $0 < \varsigma, \varepsilon < 1$, where $0 < |a_n| \leq \frac{q_{n+1}}{2q_n}$, Theorem 7.2 (upon shifting by k_0 units) implies that there exists a local n -maximum b_{a_n} on the interval $b_{a_n} + I_{\varepsilon, 1}^n$ such that

$$(108) \quad |b_{a_n} - (a_n q_n + k_0)| \leq K_0 = q_{\hat{n}_0+1}.$$

Now let n_i be such that $\ln q_{n_i+1} \geq (\varsigma + 2\varepsilon) \ln |\lambda| q_{n_i}$, $i = t_0, t_0 + 1, \dots, j$ for some $0 < \varsigma, \varepsilon < 1$. By (108), one has that there exists a local n_{j_0} maximum $b_{a_{n_{j_0}}} = a_{n_{j_0}} q_{n_{j_0}} + k_0 + \hat{K}_{n_{j_0}}$ with $|\hat{K}_{n_{j_0}}| \leq K_0$.

Now we will prove that for $0 \leq s \leq k$, there exists

$$|b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}} - a_{n_{j_s}} q_{n_{j_s}} - b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_{s-1}}}}| \leq K_s = q_{\hat{n}_0+s+1}.$$

Notice that $\sum_{i=0}^k K_i \leq 4K_k$. We will now prove Theorem 2.4 by induction on s . By the assumption, one has

$$(109) \quad \||2\theta + 2k_0 + k\alpha\|_{\mathbb{R}/\mathbb{Z}} > \frac{\kappa}{|k|^\nu}$$

for any $k \in \mathbb{Z} \setminus \{0\}$.

¹³Here 3ν can be easily relaxed to $(1 + \varepsilon)\nu$.

First we prove the case $s = 1$. By the Diophantine condition on θ (109), we have that for $|\ell| \leq q_{n_{j_1}+1}$, the following holds:

$$\begin{aligned}
 (110) \quad & \|2\theta + (2b_{a_{n_{j_0}}} + \ell)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \|2\theta + (2k_0 + \ell + 2K_{n_{j_0}})\alpha\|_{\mathbb{R}/\mathbb{Z}} - \|2a_{n_{j_0}}q_{n_{j_0}}\alpha\|_{\mathbb{R}/\mathbb{Z}} \\
 & \geq \frac{\kappa}{(2K_0 + |\ell|)^\nu} - \frac{2a_{n_{j_0}}}{q_{n_{j_0}+1}} \\
 & \geq \frac{\kappa}{|\max\{K_0, \ell\}|^{2\nu}} - e^{-\epsilon \ln \lambda q_{n_{j_0}}} \\
 & \geq \frac{\kappa}{|\max\{K_0, \ell\}|^{3\nu}}.
 \end{aligned}$$

Therefore $\theta + b_{a_{n_{j_0}}}\alpha$ is $\hat{n}_0 + 1$ -Diophantine with respect to α with parameters $3\nu, \kappa$, and by Theorem 7.2 again, there exists a local n_{j_1} -maximum such that $b_{a_{n_{j_0}}, a_{n_{j_1}}} = a_{n_{j_1}}q_{n_{j_1}} + b_{a_{n_{j_0}}} + \hat{K}_{n_{j_1}}$ with $|\hat{K}_{n_{j_1}}| \leq K_1 = q_{\hat{n}_0+2}$. This completes the first step.

Assume the theorem holds for $s = k - 1$. It suffices to show that it holds for $s = k$. By the Diophantine condition on θ (109) again, we have for $|\ell| \leq q_{n_{j_k}+1}$, the following holds:

$$\begin{aligned}
 & \|2\theta + (2b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_{k-1}}}} + \ell)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \|2\theta + (2k_0 + \ell + 2\sum_{s=0}^{k-1} K_s)\alpha\|_{\mathbb{R}/\mathbb{Z}} \\
 & \quad - \sum_{s=0}^{k-1} \|2a_{n_{j_s}}q_{n_{j_s}}\alpha\|_{\mathbb{R}/\mathbb{Z}} \\
 & \geq \frac{\kappa}{(8K_{k-1} + |\ell|)^\nu} - \sum_{s=0}^{k-1} \|2a_{n_{j_s}}q_{n_{j_s}}\alpha\|_{\mathbb{R}/\mathbb{Z}} \\
 & \geq \frac{\kappa}{|\max\{K_{k-1}, \ell\}|^{2\nu}} - \sum_{s=0}^{k-1} e^{-\epsilon q_{n_{j_s}}} \\
 & \geq \frac{\kappa}{|\max\{K_{k-1}, \ell\}|^{3\nu}}.
 \end{aligned}$$

Thus $\theta + b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_{k-1}}}}\alpha$ is $\hat{n}_0 + k$ -Diophantine with respect to α with parameters $3\nu, \kappa$, and by Theorem 7.2 again, there exists a local n_{j_k} -maximum such that $b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_k}}} = a_{n_{j_k}}q_{n_{j_k}} + b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_{k-1}}}} + \hat{K}_{n_{j_k}}$ with $|\hat{K}_{n_{j_k}}| \leq K_k = q_{\hat{n}_0+k+1}$. This implies (II) holds for $s = k$. Thus we complete the proof of (I) and (II).

(III), as well as the moreover part, follows from Theorem 2.2 directly. \square

Proof of Theorem 2.4. Since k_0 is a local $n_{j_0} + 1$ -maximum for every j , Theorem 2.4 follows from Theorem 7.3 directly. \square

Theorem 7.3 describes a hierarchical structure around every local $(n_{j_0} + 1)$ -maximum.

We will say that a local n_{j_0} -maximum is k -hierarchical if there exists $\epsilon > 0$, $j_0 > j_1 > \dots > j_k$ with $n_{j_{i+1}} > e^{\epsilon n_{j_i}}$ and, for each $s = 0, 1, \dots, k$, a collection of local n_{j_s} -maxima, $\{b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}}\}$ such that

(I) all local (n_{j_s}, ϵ) -maxima in

$$[b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}} - e^{\epsilon q n_{j_s}}, b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_{s-1}}} + e^{\epsilon q n_{j_s}}]$$

are given by $\{b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_{s-1}}}, a_{n_{j_s}}}\}$ with all possible choices of $a_{n_{j_s}}$;

(II) if $2(x - b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_k}}}) \in I_{\epsilon, \epsilon}^{n_{j_k}}$, then for each $s = 0, 1, \dots, k$,

$$(111) \quad f(x_s) e^{-\epsilon |x_s|} \leq \frac{\|U(x)\|}{\|U(b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}})\|} \leq f(x_s) e^{\epsilon |x_s|},$$

where $x_s = |x - b_{a_{n_{j_0}}, a_{n_{j_1}}, \dots, a_{n_{j_s}}}|$ is large enough.

Proof of Corollary 2.7. Clearly, $b_{a_{n_1}, \dots, a_{n_s}}$ of Theorem 7.3 form the collection required for the definition of k -hierarchy, so it remains to estimate the number of levels of the hierarchy; that is, find k such that $n_{j_k} \geq \hat{n}_0 + k$. Clearly, $k = j/2 - \lfloor \hat{n}_0/2 \rfloor$ works. □

8. Growth of transfer matrices. Proof of (13)

Assume that θ is Diophantine with respect to α in this and the following section.

THEOREM 8.1. *Let $A(j) = \|A_{jq_n}\|$. Assume $jq_n \leq k < (j + 1)q_n$ with $0 \leq j \leq 48C_* \frac{b_{n+1}}{q_n}$, $b_{n+1} \geq \frac{q_n}{2}$ and $k \geq \frac{q_n}{4}$. We have*

$$(112) \quad \|A_k\| \leq \max\{e^{-|k-jq_n| \ln \lambda} A(j), e^{-|k-(j+1)q_n| \ln \lambda} A(j+1)\} e^{\epsilon k},$$

$$(113) \quad \|A_k\| \geq \max\{e^{-|k-jq_n| \ln \lambda} A(j), e^{-|k-(j+1)q_n| \ln \lambda} A(j+1)\} e^{-\epsilon k}.$$

Proof. Let $\tilde{U}(k) = \begin{pmatrix} \psi(k) \\ \psi(k-1) \end{pmatrix}$. By Last-Simon's arguments ((8.6) in [LS99]), one has

$$(114) \quad \|A_k\| \geq \|A_k \tilde{U}(0)\| \geq c \|A_k\|.$$

Then (112) holds by (114), (53) and (38). Equation (113) holds directly by (28). □

THEOREM 8.2. *Assume $1 \leq j \leq 8 \frac{b_{n+1}}{q_n}$ and $b_{n+1} \geq \frac{q_n}{2}$. Then*

$$(115) \quad \frac{q_{n+1}}{\bar{r}_j^n} e^{-\epsilon j q_n} \leq A(j) \leq \frac{q_{n+1}}{\bar{r}_j^n} e^{\epsilon j q_n}.$$

Proof. We first show the left inequality. Clearly

$$(116) \quad \|A_k\| \geq \|U(k)\|^{-1}.$$

Thus by (89) and (112), for any $j q_n \leq k < (j + 1) q_n$ with $j \geq 0$ and $k \geq \frac{q_n}{4}$, we must have

$$(117) \quad \begin{aligned} & \max\{e^{-|k-jq_n|\ln \lambda} A(j), e^{-|k-(j+1)q_n|\ln \lambda} A(j+1)\} e^{\varepsilon k} \\ & \geq (\max\{e^{-|k-jq_n|\ln \lambda} \bar{r}_j^n, e^{-|k-(j+1)q_n|\ln \lambda} \bar{r}_{j+1}^n\})^{-1} e^{-\varepsilon k}. \end{aligned}$$

Let

$$k_0 = (j + 1)q_n - \frac{\ln q_{n+1} - \ln(j + 1)}{2 \ln \lambda}.$$

One has $k_0 \geq \frac{q_n}{4}$. Thus

$$\max\{e^{-|k_0-jq_n|\ln \lambda} \bar{r}_j^n, e^{-|k_0-(j+1)q_n|\ln \lambda} \bar{r}_{j+1}^n\} \leq \bar{r}_{j+1}^n \left(\frac{j+1}{q_{n+1}}\right)^{\frac{1}{2}} e^{\varepsilon k_0}.$$

Combining with (117), we have

$$(118) \quad \max\{e^{-|k_0-jq_n|\ln \lambda} A(j), e^{-|k_0-(j+1)q_n|\ln \lambda} A(j+1)\} \geq \frac{q_{n+1}^{\frac{1}{2}}}{(j+1)^{\frac{1}{2}}} (\bar{r}_{j+1}^n)^{-1} e^{-\varepsilon k_0}.$$

This implies that either

$$(119) \quad A(j) \geq e^{\ln \lambda q_n} (\bar{r}_{j+1}^n)^{-1} e^{-\varepsilon k_0},$$

or

$$(120) \quad A(j+1) \geq \frac{q_{n+1}}{j+1} (\bar{r}_{j+1}^n)^{-1} e^{-\varepsilon k_0}.$$

Notice that by (64) and (114), we have

$$(121) \quad A(j+1) \geq A(j) e^{(\ln \lambda - \varepsilon) q_n} \frac{j+1}{q_{n+1}}.$$

By (119), (120) and (121), we obtain the left inequality of (115).

Now we turn to the proof of the right inequality of (115). By (8.5) and (8.7) in [LS99], we have

$$(122) \quad \|A_k U(0)\|^2 \leq \|A_k\|^2 m(k)^2 + \|A_k\|^{-2},$$

where

$$(123) \quad m(k) \leq C \sum_{p=k}^{\infty} \frac{1}{\|A_p\|^2}.$$

If $k \geq C_* j q_n$ with $j \geq 1$, by (98) we have

$$\begin{aligned} \|A_k\| & \geq \|U(k)\|^{-1} \\ & \geq e^{(\ln \lambda - \beta - \varepsilon) k} \end{aligned}$$

and by (28) we have

$$A(j) \leq e^{(\ln \lambda + \varepsilon)jq_n}.$$

This implies

$$(124) \quad \|A_k\| \geq A(j)e^{\frac{\ln \lambda - \beta}{2}k}.$$

If $jq_n \leq k \leq C_*jq_n$ with $j \geq 1$, let $j_0q_n \leq k < (j_0+1)q_n$ with $j \leq j_0 \leq C_*j$. By (113) and (121), we have

$$(125) \quad \begin{aligned} \|A_k\| &\geq A(j_0) \max \left\{ e^{-|k-j_0q_n| \ln \lambda}, e^{-|k-(j_0+1)q_n| \ln \lambda} e^{q_n \ln \lambda \frac{j_0+1}{q_{n+1}}} \right\} e^{-\varepsilon j_0q_n} \\ &\geq \left(\frac{j_0+1}{q_{n+1}} \right)^{\frac{1}{2}} A(j_0) e^{-\varepsilon j_0q_n} \\ &\geq \left(\frac{j+1}{q_{n+1}} \right)^{\frac{1}{2}} A(j) e^{-\varepsilon jq_n}. \end{aligned}$$

Thus by (124) and (125), we have

$$(126) \quad m(jq_n) \leq \frac{q_{n+1}}{jA(j)^2} e^{\varepsilon jq_n}.$$

Let $k = jq_n$ in (122). One has

$$\tilde{r}_j^2 \leq \frac{q_{n+1}^2}{j^2 A(j)^2} e^{\varepsilon jq_n}.$$

Thus by (6.1), we obtain

$$(127) \quad A(j) \leq \frac{q_{n+1}}{j\tilde{r}_j^n} e^{\varepsilon jq_n}.$$

This implies the right inequality of (115). □

Theorems 8.1 and 8.2 imply the following theorem directly.

THEOREM 8.3. *Assume $jq_n \leq k < (j+1)q_n$ with $0 \leq j \leq 6\frac{b_{n+1}}{q_n}$, $b_{n+1} \geq \frac{q_n}{2}$. For $k \geq q_n$, we have*

$$(128) \quad \|A_k\| \leq \max \left\{ e^{-|k-jq_n| \ln \lambda \frac{q_{n+1}}{\tilde{r}_j^n}}, e^{-|k-(j+1)q_n| \ln \lambda \frac{q_{n+1}}{\tilde{r}_{j+1}^n}} \right\} e^{\varepsilon|k|}$$

and

$$(129) \quad \|A_k\| \geq \max \left\{ e^{-|k-jq_n| \ln \lambda \frac{q_{n+1}}{j\tilde{r}_j^n}}, e^{-|k-(j+1)q_n| \ln \lambda \frac{q_{n+1}}{(j+1)\tilde{r}_{j+1}^n}} \right\} e^{-\varepsilon|k|}.$$

and for $\frac{q_n}{4} \leq k < q_n$, we have

$$(130) \quad \|A_k\| \leq \max \left\{ e^{-|k| \ln \lambda}, e^{-|k-q_n| \ln \lambda \frac{q_{n+1}}{\tilde{r}_1^n}} \right\} e^{\varepsilon|k|}$$

and

$$(131) \quad \|A_k\| \geq \max \left\{ e^{-|k| \ln \lambda}, e^{-|k-q_n| \ln \lambda \frac{q_{n+1}}{\bar{r}_1^n}} \right\} e^{-\varepsilon|k|}.$$

THEOREM 8.4. *For any $q_n^{\frac{8}{9}} \leq k \leq \frac{q_n}{2}$, let n_0 be the smallest positive integer such that $q_{n-n_0} \leq k < q_{n-n_0+1}$. Suppose $jq_{n-n_0} \leq k < (j+1)q_{n-n_0+1}$ with $j \geq 1$. Then the following hold:*

$$(132) \quad \|A_k\| \leq \max \left\{ e^{-|k-jq_{n-n_0}| \ln \lambda \frac{q_{n-n_0+1}}{\bar{r}_j^{n-n_0}}}, e^{-|k-(j+1)q_{n-n_0}| \ln \lambda \frac{q_{n-n_0+1}}{\bar{r}_{j+1}^{n-n_0}}} \right\} e^{\varepsilon|k|}$$

and

$$(133) \quad \|A_k\| \geq \max \left\{ e^{-|k-jq_{n-n_0}| \ln \lambda \frac{q_{n-n_0+1}}{\bar{r}_j^{n-n_0}}}, e^{-|k-(j+1)q_{n-n_0}| \ln \lambda \frac{q_{n-n_0+1}}{\bar{r}_{j+1}^{n-n_0}}} \right\} e^{-\varepsilon|k|}.$$

Proof. As in the proof of Theorem 6.3, we split into the same two cases: Cases 1 and 2. Case 1 can be done directly by Theorem 8.3. For Case 2, as in the proof of Case 2 of Theorem 6.3, it suffices to show that

$$(134) \quad e^{(\ln \lambda - \varepsilon)k} \leq \|A_k\| \leq e^{(\ln \lambda + \varepsilon)k},$$

which follows directly from (93), (116) and (28). □

Proof of (13). The arguments are similar to the proof of (12) and consist of collecting the already proved facts, with the same cases.

Case (i): $\frac{q_n}{2} \leq q_{n+1}^{\frac{8}{9}}$. For $\frac{q_n}{2} \leq k \leq 4q_{n+1}^{\frac{8}{9}}$, the result follows from Theorem 8.3. For $4q_{n+1}^{\frac{8}{9}} \leq k \leq \frac{q_{n+1}}{2}$, (13) follows from Theorem 8.4. (Notice that now $k \geq 2q_n$, and thus $n_0 = 1$.)

Case (ii): $q_{n+1}^{\frac{8}{9}} \leq \frac{q_n}{2}$.

Case (ii.1): $\frac{q_n}{2} \leq k \leq \min\{q_n, \frac{q_{n+1}}{2}\}$. If $q_n = q_{n-1} + q_{n-2}$, then $q_{n-1} \geq \frac{q_n}{2}$. This is Case 2 of Theorem 8.4. By (134), for any $q_{n-1} \leq k \leq \min\{q_n, \frac{q_{n+1}}{2}\}$, one has

$$\|A_k\| \geq e^{(\ln \lambda - \varepsilon)k}.$$

This leads to

$$\|A_k\| \geq e^{(\ln \lambda - \varepsilon)k}$$

for $\frac{q_n}{2} \leq k \leq \min\{q_n, \frac{q_{n+1}}{2}\}$. This also implies (13).

If $q_n = jq_{n-1} + q_{n-2}$ with $j \geq 2$, then $\frac{q_n}{2} \geq q_{n-1}$. (13) follows directly from Theorem 8.4. (Notice that now $n+1-n_0 = n-1$.)

Case (ii.2): $q_n \leq k \leq \frac{q_{n+1}}{2}$. In this case (13) follows directly from Theorem 8.4. (Notice that now $n+1-n_0 = n$.) □

9. Proof of the corollaries

Proof of Corollary 2.12. Due to (28), (i) follows from (iii). By Theorem 2.1, to prove (iii) it is enough to show that for any $\varepsilon > 0$, sufficiently large n and $\varepsilon q_n < k < q_n$, we have

$$(135) \quad e^{(\ln \lambda - C\varepsilon)k} \leq g(k) \leq e^{(\ln \lambda + C\varepsilon)k}.$$

Let $m \leq n + 1$ be such that $q_m/2 \leq k < q_{m+1}/2$. If $q_m \leq k$, we are in Case 1 of the definition of g with $m \leq n - 1$. Notice that $\ell q_m \geq \varepsilon q_n \geq \varepsilon q_{m+1}$, which leads to $\frac{\ln \frac{q_{m+1}}{\ell}}{q_m}$ being small. Then (135) follows from (6). If $q_m/2 < k < q_m$, then (135) is automatic by the definition of g .

It remains to establish (ii). First by (98), we must have

$$(136) \quad \liminf_{k \rightarrow \infty} \frac{\ln \|A_k\|}{k} \geq \ln \lambda - \beta.$$

Let $j_k = \lfloor q_{n_k}^\varepsilon \rfloor$, where sequence q_{n_k} is given by (99). Then

$$\begin{aligned} g(j_k q_{n_k}) &= \frac{q_{n_k} + 1}{\bar{r}_{j_k}^{n_k}} \\ &= e^{(\ln \lambda - \frac{\ln q_{n_k} + 1}{q_{n_k}} + \frac{\ln j_k}{q_{n_k}}) j_k q_{n_k}} q_{n_k + 1} \\ &\leq e^{(\ln \lambda - \beta + C\varepsilon) j_k q_{n_k}}. \end{aligned}$$

Combining with Theorem 2.1, we must have

$$(137) \quad \liminf_{k \rightarrow \infty} \frac{\ln \|A_k\|}{k} \leq \ln \lambda - \beta.$$

(ii) holds by (136) and (137). □

Proof of Corollary 2.13. (i) follows from (iii), and (ii) follows from (98) and (100). To establish (iii), we only need to show that for any $\varepsilon q_n \leq k \leq q_n - \frac{\beta}{2 \ln \lambda} q_n - \varepsilon q_n$,

$$e^{-(\ln \lambda + C\varepsilon)k} \leq \|U(k)\| \leq e^{-(\ln \lambda - C\varepsilon)k}.$$

Let $m \leq n + 1$ be such that $q_m/2 \leq k < q_{m+1}/2$.

Case 1: $m \leq n - 1$. Then by Theorem 2.1, the statement is not immediate only in Case 1, but then it follows from (5) and (7) since in that case, $\bar{r}_\ell^m \leq e^{-(\ln \lambda - \varepsilon)\ell q_m}$.

Case 2: $m = n$ or $m = n + 1$. In this case we have $\frac{q_m}{2} \leq k \leq q_m - \frac{\beta}{2 \ln \lambda} q_m - \varepsilon q_m$ and $\frac{q_m}{2} \leq k < \frac{q_{m+1}}{2}$. Then the statement holds by Theorem 2.1 and, in Case 1, (7).

To prove (iv) it suffices to show that for any $q_{n_j} - \frac{\beta}{2 \ln \lambda} q_{n_j} + \varepsilon q_{n_j} \leq k \leq q_{n_j} - \varepsilon q_{n_j}$, we have

$$\|U(k)\| \geq e^{-(\ln \lambda - c\varepsilon)k}$$

where q_{n_j} is a subsequence satisfying (99). Indeed, under this assumption we are in Case 1 of the definition of f and the second addend dominates in (7) leading to the statement. \square

Remarks.

- If we take for q_{n_k} a subsequence with *any* bounded away from zero exponential growth, we still get non-Lyapunov behavior on intervals of the form $[q_{n_k} - cq_{n_k}, q_{n_k} + \varepsilon q_{n_k}]$ for some $c < \frac{\beta}{2 \ln \lambda}$.
- In fact, in all the arguments β can be replaced with $\ln q_{n+1}/q_n$.

Proof of Corollary 2.14. First by (24), one has

$$(138) \quad \|U(k)\| \|\tilde{U}(k)\| \sin \delta_k = \frac{1}{2}.$$

Combining with (114), we have

$$(139) \quad \frac{1}{2\|U(k)\| \|A_k\|} \leq \sin \delta_k \leq \frac{1}{\|U(k)\| \|A_k\|}.$$

We first prove (21). Clearly, it suffices to show

$$(140) \quad \limsup_{k \rightarrow \infty} \frac{\ln \delta_k}{k} \geq 0.$$

Let $k_j = \lfloor \frac{1}{4}q_{n_j+1} \rfloor$, where the sequence q_{n_j} is given by (99). By Theorem 2.1, we must have

$$\|U(k_j)\| \leq e^{-(\ln \lambda - \varepsilon)k_j}$$

and

$$\|A_{k_j}\| \leq e^{(\ln \lambda + \varepsilon)k_j}.$$

Combining with (139), we must have

$$(141) \quad \delta_{k_n} \geq e^{-\varepsilon k_n}.$$

This implies (140) and also (21).

Now we verify (22). By the definition of $f(k), g(k)$, for any large k , we have

$$(142) \quad f(k)g(k) \leq e^{(\beta + \varepsilon)k}.$$

Then by (139) and Theorem 2.1 again, one has

$$(143) \quad \liminf_{k \rightarrow \infty} \frac{\ln \delta_k}{k} \geq -\beta.$$

Let $k_j = q_{n_j}$. One has

$$(144) \quad f(k_j)g(k_j) = q_{n_j+1}.$$

Combining with (139) and Theorem 2.1 again, we get

$$(145) \quad \lim_{j \rightarrow \infty} \frac{\ln \delta_{k_j}}{k_j} = -\beta.$$

Equation (22) follows directly from (143) and (145). \square

Proof of Corollary 2.11. This follows directly from (13) and (114). \square

Appendix A. Gordon arguments for $\lambda \leq e^\beta$

PROPOSITION A.1. *The almost Mathieu operator*

$$(H_{\lambda,\alpha,\theta}u)(n) = u(n + 1) + u(n - 1) + 2\lambda \cos 2\pi(\theta + n\alpha)u(n)$$

has no localized eigenfunctions if $|\lambda| \leq e^\beta$.¹⁴

Proof. Otherwise, there exists a solution $\{u(n)\}_{n \in \mathbb{Z}}$ of $H_{\lambda,\alpha,\theta}u = Eu$ such that

$$(146) \quad |u(n)| \leq Ce^{-5c|n|},$$

where $c > 0$. Without loss of generality, assume that the vector $\begin{pmatrix} u(0) \\ u(-1) \end{pmatrix}$ is unit.

Let $\varphi(n) = \begin{pmatrix} u(n) \\ u(n-1) \end{pmatrix}$. For simplicity, denote $\varphi = \varphi(0)$. By the definition of $\beta(\alpha)$, there exists a subsequence \tilde{q}_k of q_n such that

$$(147) \quad \|\tilde{q}_k\alpha\|_{\mathbb{R}/\mathbb{Z}} \leq e^{-(\beta - \frac{\epsilon}{4})\tilde{q}_k}.$$

Denote $B = A_{\tilde{q}_k}(\theta)$. Then we have

$$(148) \quad B^2 + (\text{Tr } B)B + I = 0.$$

Case 1: If $\text{Tr } B \leq e^{2c\tilde{q}_k}$, then one has either $\|B^2\varphi\| \geq \frac{1}{2}$ or $\|B\varphi\| \geq \frac{1}{2}e^{-2c\tilde{q}_k}$. By (146), we must have $\|B^2\varphi\| \geq \frac{1}{2}$. This is impossible. Indeed, from the following estimate:

$$\begin{aligned} \|\varphi(2\tilde{q}_k) - B^2\varphi\| &= \|A_{\tilde{q}_k}(\theta + \tilde{q}_k\alpha) - A_{\tilde{q}_k}(\theta)\| \|\varphi(\tilde{q}_k)\| \\ &\leq Ce^{\frac{3}{2}c\tilde{q}_k} e^{-5c\tilde{q}_k} \\ &\leq e^{-3c\tilde{q}_k}, \end{aligned}$$

where the first inequality holds by (71). Then

$$\|\varphi(2\tilde{q}_k)\| \geq \frac{1}{4},$$

contradicting $\|\varphi(2\tilde{q}_k)\| \leq Ce^{-10c\tilde{q}_k}$.

Case 2: If $\text{Tr } B \geq e^{2c\tilde{q}_k}$, then from (148), it is easy to see that either $\|B\varphi\| \geq \frac{1}{2}e^{2c\tilde{q}_k}$ or $\|B^{-1}\varphi\| \geq \frac{1}{2}e^{2c\tilde{q}_k}$ holds. By (146) again, we must have $\|B^{-1}\varphi\| \geq \frac{1}{2}e^{2c\tilde{q}_k}$. By (72), the following holds:

$$\begin{aligned} \|\varphi(-\tilde{q}_k) - B^{-1}\varphi\| &= \|A_{\tilde{q}_k}^{-1}(\theta - \tilde{q}_k\alpha) - A_{\tilde{q}_k}^{-1}(\theta)\| \|\varphi\| \\ &\leq e^{\frac{3}{2}c\tilde{q}_k}. \end{aligned}$$

¹⁴Localized here means exponentially decaying. One can exclude any decaying solutions for $|\lambda| < e^\beta$ [AYZ17] but not for $|\lambda| = e^\beta$ [AJZ18].

Thus

$$\|\varphi(-\tilde{q}_k)\| \geq \frac{1}{4}e^{2c\tilde{q}_k}.$$

This is also impossible. □

Appendix B. Uniformity

We start with some basic facts. Let $\frac{p_n}{q_n}$ be the continued fraction approximants to α . Then

$$(149) \quad \forall 1 \leq k < q_{n+1}, \text{dist}(k\alpha, \mathbb{Z}) \geq |q_n\alpha - p_n|$$

and

$$(150) \quad \frac{1}{2q_{n+1}} \leq \Delta_n := |q_n\alpha - p_n| \leq \frac{1}{q_{n+1}}.$$

LEMMA B.1 ([AJ09, Lemma 9.7]). *Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, $x \in \mathbb{R}$ and $0 \leq \ell_0 \leq q_n - 1$ be such that $|\sin \pi(x + \ell_0\alpha)| = \inf_{0 \leq \ell \leq q_n - 1} |\sin \pi(x + \ell\alpha)|$. Then for some absolute constant $C > 0$,*

$$(151) \quad -C \ln q_n \leq \sum_{\ell=0, \ell \neq \ell_0}^{q_n-1} \ln |\sin \pi(x + \ell\alpha)| + (q_n - 1) \ln 2 \leq C \ln q_n.$$

We now prove

LEMMA B.2. *For any $|i|, |j| \leq 50C_*b_{n+1}$, if θ is n -Diophantine with respect to α , then the following estimate holds:*

$$(152) \quad \ln |\sin \pi(2\theta + (j + i)\alpha)| \geq -C \ln q_n.$$

Proof. By the Diophantine condition on θ , (2), one has that there exist $\kappa > 0$ and $\nu > 0$ such that

$$(153) \quad \min_{j, i \in [-q_n, q_n]} |\sin \pi(2\theta + (j + i)\alpha)| \geq \frac{\kappa}{q_n^\nu}.$$

Let $\ell_i, \ell_j \in \mathbb{Z}$ be such that $\text{dist}(i, q_n\mathbb{Z}) = |i - \ell_i q_n|$ and $\text{dist}(j, q_n\mathbb{Z}) = |j - \ell_j q_n|$. Then $|\ell_i|, |\ell_j| \leq 50C_* \frac{b_{n+1}}{q_n} + 1$. Let $i' = i - \ell_i q_n$ and $j' = j - \ell_j q_n$. Then $i', j' \in [-q_n, q_n]$.

If $q_{n+1}^{1-t} > \frac{100C_*}{\kappa} q_n^{\nu+2}$, it is easy to verify that $|\ell_k \Delta_n| < \frac{\kappa}{q_n^{\nu+1}}$. Combining with (153), for any $|i|, |j| \leq 50C_*b_{n+1}$, we have

$$\begin{aligned} |\sin \pi(2\theta + (j + i)\alpha)| &= \left| \sin \pi(2\theta + (j' + i')\alpha) \cos \pi(\ell_i + \ell_j)\Delta_n \right. \\ &\quad \left. \pm \cos \pi(2\theta + (j' + i')\alpha) \sin \pi(\ell_i + \ell_j)\Delta_n \right| \\ &\geq \frac{\kappa}{100q_n^\nu}. \end{aligned}$$

(The choice of \pm depends on the sign of $q_n\alpha - p_n$.)

If $q_{n+1}^{1-t} \leq \frac{100C_*}{\kappa} q_n^{\nu+2}$, we also have, for any $|i|, |j| \leq 50C_*b_{n+1}$,

$$|\sin \pi(2\theta + (j + i)\alpha)| \geq \frac{\kappa^{1+\frac{t\nu}{1-t}}}{(100C_*)^{\frac{\nu}{1-t}} q_n^{\frac{\nu t(\nu+2)}{1-t}}}.$$

Thus in both cases, we have

$$(154) \quad \min_{|i|, |j| \leq 50C_*b_{n+1}} \ln |\sin \pi(2\theta + (j + i)\alpha)| \geq -C \ln q_n. \quad \square$$

LEMMA B.3. *Assume $|i|, |j| \leq 50C_*b_{n+1}$, and $i - j \neq q_n\mathbb{Z}$. Then*

$$(155) \quad \ln |\sin \pi(j - i)\alpha| \geq -C \ln q_n.$$

Proof. By assumption, $|j - i| = \ell q_n + r$ with $0 \leq \ell \leq 100C_* \frac{b_{n+1}}{q_n}$ and $0 < r < q_n$. Then by (149) and (150) again, we also have

$$\begin{aligned} \|(j - i)\alpha\|_{\mathbb{R}/\mathbb{Z}} &\geq \|r\alpha\|_{\mathbb{R}/\mathbb{Z}} - |\ell| \|q_n\alpha\|_{\mathbb{R}/\mathbb{Z}} \\ &\geq \frac{1}{2q_n} - \frac{|\ell|}{q_{n+1}} \\ &\geq \frac{1}{2q_n} - \frac{100C_*}{q_{n+1}} \frac{1}{q_n} \\ &\geq \frac{1}{4q_n}. \end{aligned}$$

This implies (155). □

We are now ready to study the behavior at non-resonant points. For an n -non-resonant y , as before let n_0 be the least positive integer such that $4q_{n-n_0} \leq \text{dist}(y, q_n\mathbb{Z})$. Let s be the largest positive integer such that $4sq_{n-n_0} \leq \text{dist}(y, q_n\mathbb{Z})$. Recall that, automatically, $n_0 \leq C(\alpha)$. Set $I_1, I_2 \subset \mathbb{Z}$ as follows:

$$\begin{aligned} I_1 &= [-sq_{n-n_0}, sq_{n-n_0} - 1], \\ I_2 &= [y - sq_{n-n_0}, y + sq_{n-n_0} - 1]. \end{aligned}$$

We have

THEOREM B.4. *For an n -non-resonant y , assume that*

$$(156) \quad \min_{j, i \in I_1 \cup I_2} \ln |\sin \pi(2\theta + (j + i)\alpha)| \geq -C \ln q_n$$

and

$$(157) \quad \min_{i \neq j; i, j \in I_1 \cup I_2} \ln |\sin \pi(j - i)\alpha| \geq -C \ln q_n.$$

Then for any $\varepsilon > 0$ and n large enough, we have

y is $(\ln \lambda + 8 \ln(sq_{n-n_0}/q_{n-n_0+1})/q_{n-n_0} - \varepsilon, 4sq_{n-n_0} - 1)$ regular with $\delta = \frac{1}{4}$.

Proof. Without loss of generality, assume $y > 0$. By the definition of s and n_0 , we have $4sq_{n-n_0} \leq \text{dist}(y, q_n\mathbb{Z})$ and $4q_{n-n_0+1} > \text{dist}(y, q_n\mathbb{Z})$. This leads to $sq_{n-n_0} \leq q_{n-n_0+1}$. Let $\theta_j = \theta + j\alpha$ for $j \in I_1 \cup I_2$. The set $\{\theta_j\}_{j \in I_1 \cup I_2}$ consists of $4sq_{n-n_0}$ elements.

In (34), let $x = \cos 2\pi a$, $k = 4sq_{n-n_0} - 1$ and take the logarithm. Then

$$\begin{aligned} \ln \prod_{j \in I_1 \cup I_2, j \neq i} \frac{|\cos 2\pi a - \cos 2\pi \theta_j|}{|\cos 2\pi \theta_i - \cos 2\pi \theta_j|} &= \sum_{j \in I_1 \cup I_2, j \neq i} \ln |\cos 2\pi a - \cos 2\pi \theta_j| \\ &\quad - \sum_{j \in I_1 \cup I_2, j \neq i} \ln |\cos 2\pi \theta_i - \cos 2\pi \theta_j|. \end{aligned}$$

First, we estimate $\sum_{j \in I_1 \cup I_2, j \neq i} \ln |\cos 2\pi a - \cos 2\pi \theta_j|$. Obviously,

$$\begin{aligned} &\sum_{j \in I_1 \cup I_2, j \neq i} \ln |\cos 2\pi a - \cos 2\pi \theta_j| \\ &= \sum_{j \in I_1 \cup I_2, j \neq i} \ln |\sin \pi(a + \theta_j)| \\ &\quad + \sum_{j \in I_1 \cup I_2, j \neq i} \ln |\sin \pi(a - \theta_j)| + (4sq_{n-n_0} - 1) \ln 2 \\ &= \Sigma_+ + \Sigma_- + (4sq_{n-n_0} - 1) \ln 2. \end{aligned}$$

Both Σ_+ and Σ_- consist of $4s$ terms of the form of (151), plus $4s$ terms of the form

$$\ln \min_{j=0,1,\dots,q_{n-n_0}} |\sin \pi(x + j\alpha)|,$$

minus $\ln |\sin \pi(a \pm \theta_i)|$. Thus, using (151) $4s$ times for Σ_+ and Σ_- respectively, one has

$$(158) \quad \sum_{j \in I_1 \cup I_2, j \neq i} \ln |\cos 2\pi a - \cos 2\pi \theta_j| \leq -4sq_{n-n_0} \ln 2 + Cs \ln q_{n-n_0}.$$

If $a = \theta_i$, we obtain

$$\begin{aligned} &\sum_{j \in I_1 \cup I_2, j \neq i} \ln |\cos 2\pi \theta_i - \cos 2\pi \theta_j| \\ &= \sum_{j \in I_1 \cup I_2, j \neq i} \ln |\sin \pi(\theta_i + \theta_j)| \\ (159) \quad &\quad + \sum_{j \in I_1 \cup I_2, j \neq i} \ln |\sin \pi(\theta_i - \theta_j)| + (4sq_{n-n_0} - 1) \ln 2 \\ &= \Sigma_+ + \Sigma_- + (4sq_{n-n_0} - 1) \ln 2, \end{aligned}$$

where

$$\Sigma_+ = \sum_{j \in I_1 \cup I_2, j \neq i} \ln |\sin \pi(2\theta + (i + j)\alpha)|$$

and

$$\Sigma_- = \sum_{j \in I_1 \cup I_2, j \neq i} \ln |\sin \pi(i - j)\alpha|.$$

We will estimate Σ_+ . Set $J_1 = [-s, s - 1]$ and $J_2 = [s, 3s - 1]$, which are two adjacent disjoint intervals of length $2s$. Then $J_1 \cup J_2$ can be represented as a disjoint union of segments B_j , $j \in J_1 \cup J_2$, each of length q_{n-n_0} . Applying (151) to each B_j , we obtain

$$(160) \quad \Sigma_+ \geq -4sq_{n-n_0} \ln 2 + \sum_{j \in J_1 \cup J_2} \ln |\sin \pi \hat{\theta}_j| - Cs \ln q_{n-n_0} - \ln |\sin 2\pi(\theta + i\alpha)|,$$

where

$$(161) \quad |\sin \pi \hat{\theta}_j| = \min_{\ell \in B_j} |\sin \pi(2\theta + (\ell + i)\alpha)|.$$

Next we estimate $\sum_{j \in J_1} \ln |\sin \pi \hat{\theta}_j|$. Assume that $\hat{\theta}_{j+1} = \hat{\theta}_j + q_{n-n_0}\alpha$ for every $j, j + 1 \in J_1$. In this case, for any $i, j \in J_1$ and $i \neq j$, we have

$$(162) \quad \|\hat{\theta}_i - \hat{\theta}_j\|_{\mathbb{R}/\mathbb{Z}} \geq \|q_{n-n_0}\alpha\|_{\mathbb{R}/\mathbb{Z}}.$$

Applying the Stirling formula, (156) and (162), one has

$$(163) \quad \begin{aligned} \sum_{j \in J_1} \ln |\sin 2\pi \hat{\theta}_j| &> 2 \sum_{j=1}^s \ln(j\Delta_{n-n_0}) - C \ln q_n \\ &> 2s \ln \frac{s}{q_{n-n_0+1}} - C \ln q_n - Cs. \end{aligned}$$

In the other cases, decompose J_1 in maximal intervals T_κ such that for $j, j + 1 \in T_\kappa$, we have $\hat{\theta}_{j+1} = \hat{\theta}_j + q_{n-n_0}\alpha$. Notice that the boundary points of an interval T_κ are either boundary points of J_1 or satisfy $\|\hat{\theta}_j\|_{\mathbb{R}/\mathbb{Z}} + \Delta_{n-n_0} \geq \frac{\Delta_{n-n_0-1}}{2}$. This follows from the fact that if $0 < |z| < q_{n-n_0}$, then

$$\|\hat{\theta}_j + q_{n-n_0}\alpha\|_{\mathbb{R}/\mathbb{Z}} \leq \|\hat{\theta}_j\|_{\mathbb{R}/\mathbb{Z}} + \Delta_{n-n_0}$$

and

$$\begin{aligned} \|\hat{\theta}_j + (z + q_{n-n_0})\alpha\|_{\mathbb{R}/\mathbb{Z}} &\geq \|z\alpha\|_{\mathbb{R}/\mathbb{Z}} - \|\hat{\theta}_j + q_{n-n_0}\alpha\|_{\mathbb{R}/\mathbb{Z}} \\ &\geq \Delta_{n-n_0-1} - \|\hat{\theta}_j\|_{\mathbb{R}/\mathbb{Z}} - \Delta_{n-n_0}. \end{aligned}$$

Assuming $T_\kappa \neq J_1$, then there exists $j \in T_\kappa$ such that $\|\hat{\theta}_j\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{\Delta_{n-n_0-1}}{2} - \Delta_{n-n_0}$.

If T_κ contains some j with $\|\hat{\theta}_j\|_{\mathbb{R}/\mathbb{Z}} < \frac{\Delta_{n-n_0-1}}{10}$, then

$$(164) \quad \begin{aligned} |T_\kappa| &\geq \frac{\frac{\Delta_{n-n_0-1}}{2} - \Delta_{n-n_0} - \frac{\Delta_{n-n_0-1}}{10}}{\Delta_{n-n_0}} \\ &\geq \frac{1}{4} \frac{\Delta_{n-n_0-1}}{\Delta_{n-n_0}} - 1 \geq \frac{s}{8} - 1, \end{aligned}$$

since $sq_{n-n_0} \leq q_{n-n_0+1}$, where $|T_\kappa| = b - a + 1$ if $T_\kappa = [a, b]$. For such T_κ , a similar estimate to (163) gives

$$(165) \quad \begin{aligned} \sum_{j \in T_\kappa} \ln |\sin \pi \hat{\theta}_j| &\geq |T_\kappa| \ln \frac{|T_\kappa|}{q_{n-n_0+1}} - Cs - C \ln q_n \\ &\geq |T_\kappa| \ln \frac{s}{q_{n-n_0+1}} - Cs - C \ln q_n. \end{aligned}$$

If T_κ does not contain any j with $\|\hat{\theta}_j\|_{\mathbb{R}/\mathbb{Z}} < \frac{\Delta_{n-n_0-1}}{10}$, then by (150),

$$(166) \quad \begin{aligned} \sum_{j \in T_\kappa} \ln |\sin \pi \hat{\theta}_j| &\geq -|T_\kappa| \ln q_{n-n_0} - C|T_\kappa| \\ &\geq |T_\kappa| \ln \frac{s}{q_{n-n_0+1}} - C|T_\kappa|. \end{aligned}$$

By (165) and (166), one has

$$(167) \quad \sum_{j \in J_1} \ln |\sin \pi \hat{\theta}_j| \geq 2s \ln \frac{s}{q_{n-n_0+1}} - Cs - C \ln q_n.$$

Similarly,

$$(168) \quad \sum_{j \in J_2} \ln |\sin \pi \hat{\theta}_j| \geq 2s \ln \frac{s}{q_{n-n_0+1}} - Cs - C \ln q_n.$$

Putting (160), (167) and (168) together, we have

$$(169) \quad \Sigma_+ > -4sq_{n-n_0} \ln 2 + 4s \ln \frac{s}{q_{n-n_0+1}} - Cs \ln q_{n-n_0} - C \ln q_n.$$

Now we start to estimate Σ_- . Replacing (156) with (157), and following the proof of (169), we obtain,

$$(170) \quad \Sigma_- > -4sq_{n-n_0} \ln 2 + 4s \ln \frac{s}{q_{n-n_0+1}} - Cs \ln q_{n-n_0} - C \ln q_n.$$

From (159), (169) and (170), one has

$$(171) \quad \begin{aligned} \sum_{j \in I_1 \cup I_2, j \neq i} \ln |\cos 2\pi \theta_i - \cos 2\pi \theta_j| &\geq -4sq_{n-n_0} \ln 2 \\ &\quad + 8s \ln \frac{s}{q_{n-n_0+1}} - Cs \ln q_{n-n_0} - C \ln q_n. \end{aligned}$$

By (158) and (171), we have

$$\max_{i \in I_1 \cup I_2} \prod_{j \in I_1 \cup I_2, j \neq i} \frac{|x - \cos 2\pi \theta_j|}{|\cos 2\pi \theta_i - \cos 2\pi \theta_j|} < e^{4sq_{n-n_0}(-2 \ln(s/q_{n-n_0+1})/q_{n-n_0} + \varepsilon)}.$$

Combining with Lemma 3.4, there exists some j_0 with $j_0 \in I_1 \cup I_2$ such that

$$\theta_{j_0} \notin A_{4sq_{n-n_0}-1, \ln \lambda + 2 \ln(s/q_{n-n_0+1})/q_{n-n_0} - \varepsilon}.$$

First, we assume $j_0 \in I_2$. Set $I = [j_0 - 2sq_{n-n_0} + 1, j_0 + 2sq_{n-n_0} - 1] = [x_1, x_2]$. By (30), (31) and (32), it is easy to verify that

$$|G_I(y, x_i)| \leq \exp \left\{ (\ln \lambda + \varepsilon)(4sq_{n-n_0} - 1 - |y - x_i|) - 4sq_{n-n_0}(\ln \lambda + 2 \ln(s/q_{n-n_0+1})/q_{n-n_0} - \varepsilon) \right\}.$$

Notice that $|y - x_i| \geq sq_{n-n_0}$, so we obtain

$$(172) \quad |G_I(y, x_i)| \leq \exp\{-(\ln \lambda + 8 \ln(s/q_{n-n_0+1})/q_{n-n_0} - 2\varepsilon)|y - x_i|\}.$$

If $j_0 \in I_1$, we may let $y = 0$ or $y = 1$ in (172). Combining with (33), we get

$$|\phi(0)|, |\phi(-1)| \leq 6sq_{n-n_0} \exp\{-(\ln \lambda + 8 \ln(s/q_{n-n_0+1})/q_{n-n_0} - 2\varepsilon)sq_{n-n_0}\}.$$

This is in contradiction to $|\phi(0)|^2 + |\phi(-1)|^2 = 1$. Thus $j_0 \in I_2$, and the theorem follows from (172). \square

Proof of Theorem 3.5. In Case (i), (156) and (157) are obtained correspondingly from Lemmas B.2 and B.3. Thus Theorem 3.5 follows from Theorem B.4. In Case (ii), it is easy to see that (156) and (157) also hold, so Theorem B.4 applies as well. \square

Assume $b_{n+1} \geq \frac{q_n}{2}$. For any $1 \leq j \leq 48C_* \frac{b_{n+1}}{q_n}$, we construct $I_1, I_2 \subset \mathbb{Z}$ as follows:

$$I_1 = \left[-\left\lfloor \frac{1}{2}q_n \right\rfloor, q_n - \left\lfloor \frac{1}{2}q_n \right\rfloor - 1 \right],$$

$$I_2 = \left[jq_n - \left\lfloor \frac{1}{2}q_n \right\rfloor, (j+1)q_n - \left\lfloor \frac{1}{2}q_n \right\rfloor - 1 \right].$$

Let $\theta_m = \theta + m\alpha$ for $m \in I_1 \cup I_2$. Then

THEOREM B.5. *Suppose θ is n -Diophantine with respect to α . Then for any $\varepsilon > 0$, the set $\{\theta_m\}_{m \in I_1 \cup I_2}$ is $\frac{\ln q_{n+1} - \ln j}{2q_n} + \varepsilon$ -uniform for sufficiently large n .*

Proof. In (34), let $x = \cos 2\pi a$, $k = 2q_n - 1$ and take the logarithm. Thus in order to prove the theorem, it suffices to show that

$$\begin{aligned} \ln \prod_{m \in I_1 \cup I_2, m \neq i} \frac{|\cos 2\pi a - \cos 2\pi \theta_m|}{|\cos 2\pi \theta_i - \cos 2\pi \theta_m|} &= \sum_{m \in I_1 \cup I_2, m \neq i} \ln |\cos 2\pi a - \cos 2\pi \theta_m| \\ &\quad - \sum_{m \in I_1 \cup I_2, m \neq i} \ln |\cos 2\pi \theta_i - \cos 2\pi \theta_m| \\ &\leq (2q_n - 1) \left(\frac{\ln q_{n+1} - \ln j}{2q_n} + \varepsilon \right). \end{aligned}$$

First, we estimate $\sum_{m \in I_1 \cup I_2, m \neq i} \ln |\cos 2\pi a - \cos 2\pi \theta_m|$. Obviously,

$$\begin{aligned} & \sum_{m \in I_1 \cup I_2, m \neq i} \ln |\cos 2\pi a - \cos 2\pi \theta_m| \\ &= \sum_{m \in I_1 \cup I_2, m \neq i} \ln |\sin \pi(a + \theta_m)| \\ &+ \sum_{m \in I_1 \cup I_2, m \neq i} \ln |\sin \pi(a - \theta_m)| + (2q_n - 1) \ln 2 \\ &= \Sigma_+ + \Sigma_- + (2q_n - 1) \ln 2. \end{aligned}$$

Both Σ_+ and Σ_- consist of two terms of the form of (151), plus two terms of the form

$$\min_{k=1, \dots, q_n} \ln |\sin \pi(x + k\alpha)|,$$

minus $\ln |\sin \pi(a \pm \theta_i)|$. Thus one has

$$\sum_{m \in I_1 \cup I_2, m \neq i} \ln |\cos 2\pi a - \cos 2\pi \theta_m| \leq -2q_n \ln 2 + C \ln q_n.$$

Setting $a = \theta_i$ and using the first inequality of (151) two times, we obtain

$$\begin{aligned} & \sum_{m \in I_1 \cup I_2, m \neq i} \ln |\cos 2\pi \theta_i - \cos 2\pi \theta_m| \\ (173) \quad & \geq -2q_n \ln 2 - C \ln q_n + 2 \min_{m, i \in I_1 \cup I_2} \ln |\sin \pi(2\theta + (m + i)\alpha)| \\ & + \min_{m \in I_1 \cup I_2, m \neq i} \ln |\sin \pi(m - i)\alpha|. \end{aligned}$$

By Lemma B.2, we also have

$$(174) \quad \min_{m, i \in I_1 \cup I_2} \ln |\sin \pi(2\theta + (m + i)\alpha)| \geq -C \ln q_n.$$

By (149) and (150), the corresponding minimum term of

$$\min_{m \in I_1 \cup I_2, m \neq i} \ln |\sin \pi((m - i)\alpha)|$$

is achieved at jq_n . It is easy to check that

$$(175) \quad \min \{ \ln |\sin \pi jq_n \alpha| \} > -\ln \frac{q_{n+1}}{j} - C,$$

since $\Delta_n \geq \frac{1}{2q_{n+1}}$.

Putting (173), (174) and (175) together, we obtain

$$\max_{x \in [-1, 1]} \max_{i=1, \dots, k+1} \prod_{m=1, m \neq i}^{k+1} \frac{|x - \cos 2\pi \theta_m|}{|\cos 2\pi \theta_i - \cos 2\pi \theta_m|} \leq e^{(2q_n - 1) \left(\frac{\ln q_{n+1} - \ln j}{2q_n} + \varepsilon \right)}. \quad \square$$

Acknowledgments. We would like to thank Ya. Pesin and Q. Zhou for comments on earlier versions of the manuscript and Ya. Pesin also for the related discussions of non-regular dynamics. S.J. was supported by the Simons Foundation and W.L. was supported by the AMS-Simons Travel Grant 2016-2018. This research was partially supported by NSF DMS-1401204 and NSF DMS-1700314. We are grateful to the Isaac Newton Institute for Mathematical Sciences, Cambridge, for its hospitality, supported by EPSRC Grant Number EP/K032208/1, during the programme Periodic and Ergodic Spectral Problems, where this work was started.

References

- [ATW98] A. G. ABANOV, J. C. TALSTRA, and P. B. WIEGMANN, Hierarchical structure of Azbel-Hofstadter problem: strings and loose ends of Bethe ansatz, *Nuclear Phys. B* **525** no. 3 (1998), 571–596. MR 1639312. Zbl 1047.82511. [https://doi.org/10.1016/S0550-3213\(98\)00346-0](https://doi.org/10.1016/S0550-3213(98)00346-0).
- [AA80] S. AUBRY and G. ANDRÉ, Analyticity breaking and Anderson localization in incommensurate lattices, in *Group Theoretical Methods in Physics* (Proc. Eighth Internat. Colloq., Kiryat Anavim, 1979), *Ann. Israel Phys. Soc.* **3**, Hilger, Bristol, 1980, pp. 133–164. MR 0626837. Zbl 0943.82510.
- [Avi08] A. AVILA, The absolutely continuous spectrum of the almost Mathieu operator, 2008. arXiv 0810.2965.
- [Avi10] A. AVILA, Almost reducibility and absolute continuity I, 2010. arXiv 1006.0704.
- [AFK11] A. AVILA, B. FAYAD, and R. KRIKORIAN, A KAM scheme for $SL(2, \mathbb{R})$ cocycles with Liouvillean frequencies, *Geom. Funct. Anal.* **21** no. 5 (2011), 1001–1019. MR 2846380. Zbl 1277.37089. <https://doi.org/10.1007/s00039-011-0135-6>.
- [AJ09] A. AVILA and S. JITOMIRSKAYA, The Ten Martini Problem, *Ann. of Math.* (2) **170** no. 1 (2009), 303–342. MR 2521117. Zbl 1166.47031. <https://doi.org/10.4007/annals.2009.170.303>.
- [AJ10] A. AVILA and S. JITOMIRSKAYA, Almost localization and almost reducibility, *J. Eur. Math. Soc. (JEMS)* **12** no. 1 (2010), 93–131. MR 2578605. Zbl 1185.47028. <https://doi.org/10.4171/JEMS/191>.
- [AJZ18] A. AVILA, S. JITOMIRSKAYA, and Q. ZHOU, Second phase transition line, *Math. Ann.* **370** no. 1–2 (2018), 271–285. MR 3747487. <https://doi.org/10.1007/s00208-017-1543-1>.
- [AYZ] A. AVILA, J. YOU, and Q. ZHOU, Dry Ten Martini Problem in the non-critical case, preprint.
- [AYZ17] A. AVILA, J. YOU, and Q. ZHOU, Sharp phase transitions for the almost Mathieu operator, *Duke Math. J.* **166** no. 14 (2017), 2697–2718. MR 3707287. Zbl 06803180. <https://doi.org/10.1215/00127094-2017-0013>.
- [AS82] J. AVRON and B. SIMON, Singular continuous spectrum for a class of almost periodic Jacobi matrices, *Bull. Amer. Math. Soc. (N.S.)* **6** no. 1

- (1982), 81–85. MR 0634437. Zbl 0491.47014. <https://doi.org/10.1090/S0273-0979-1982-14971-0>.
- [Azb64] M. Y. AZBEL, Energy spectrum of a conduction electron in a magnetic field, *Sov. Phys. JETP* **19** no. 3 (1964), 634–645. Available at http://www.jetp.ac.ru/cgi-bin/dn/e_019_03_0634.pdf.
- [BP07] L. BARREIRA and Y. PESIN, *Nonuniform Hyperbolicity*, *Encyclopedia Math. Appl.* **115**, Cambridge Univ. Press, Cambridge, 2007, Dynamics of systems with nonzero Lyapunov exponents. MR 2348606. Zbl 1144.37002. <https://doi.org/10.1017/CBO9781107326026>.
- [BS00] L. BARREIRA and J. SCHMELING, Sets of “non-typical” points have full topological entropy and full Hausdorff dimension, *Israel J. Math.* **116** (2000), 29–70. MR 1759398. Zbl 0988.37029. <https://doi.org/10.1007/BF02773211>.
- [BLT83] J. BELLISSARD, R. LIMA, and D. TESTARD, A metal-insulator transition for the almost Mathieu model, *Comm. Math. Phys.* **88** no. 2 (1983), 207–234. MR 0696805. Zbl 0542.35059. <https://doi.org/10.1007/BF01209477>.
- [BC91] M. BENEDICKS and L. CARLESON, The dynamics of the Hénon map, *Ann. of Math. (2)* **133** no. 1 (1991), 73–169. MR 1087346. Zbl 0724.58042. <https://doi.org/10.2307/2944326>.
- [Ber68] J. M. BEREZANS’KIĬ, *Expansions in Eigenfunctions of Selfadjoint Operators*, *Transl. Math. Monogr.* **17**, American Mathematical Society, Providence, R.I., 1968, translated from the Russian by R. Bolstein, J. M. Danskin, J. Rovnyak and L. Shulman. MR 0222718.
- [Bje15] K. BJERKLÖV, The dynamics of a class of quasi-periodic Schrödinger cocycles, *Ann. Henri Poincaré* **16** no. 4 (2015), 961–1031. MR 3317790. Zbl 1312.81061. <https://doi.org/10.1007/s00023-014-0330-8>.
- [BG00] J. BOURGAIN and M. GOLDSTEIN, On nonperturbative localization with quasi-periodic potential, *Ann. of Math. (2)* **152** no. 3 (2000), 835–879. MR 1815703. Zbl 1053.39035. <https://doi.org/10.2307/2661356>.
- [BJ02] J. BOURGAIN and S. JITOMIRSKAYA, Continuity of the Lyapunov exponent for quasiperiodic operators with analytic potential, *J. Statist. Phys.* **108** no. 5-6 (2002), 1203–1218, Dedicated to David Ruelle and Yasha Sinai on the occasion of their 65th birthdays. MR 1933451. Zbl 1039.81019. <https://doi.org/10.1023/A:1019751801035>.
- [Dam17] D. DAMANIK, Schrödinger operators with dynamically defined potentials, *Ergodic Theory Dynam. Systems* **37** no. 6 (2017), 1681–1764. MR 3681983. Zbl 06823048. <https://doi.org/10.1017/etds.2015.120>.
- [Eli97] L. H. ELIASSON, Discrete one-dimensional quasi-periodic Schrödinger operators with pure point spectrum, *Acta Math.* **179** no. 2 (1997), 153–196. MR 1607554. Zbl 0908.34072. <https://doi.org/10.1007/BF02392742>.
- [Fed13] A. A. FEDOTOV, The monodromization method in the theory of almost periodic equations, *Algebra i Analiz* **25** no. 2 (2013), 203–233. MR 3114856. <https://doi.org/10.1090/S1061-0022-2014-01292-7>.

- [FK08] A. FEDOTOV and F. KLOPP, Pointwise existence of the Lyapunov exponent for a quasi-periodic equation, in *Mathematical Results in Quantum Mechanics*, World Sci. Publ., Hackensack, NJ, 2008, pp. 66–78. MR 2466679. Zbl 1156.81370. https://doi.org/10.1142/9789812832382_0006.
- [FSW90] J. FRÖHLICH, T. SPENCER, and P. WITTWER, Localization for a class of one-dimensional quasi-periodic Schrödinger operators, *Comm. Math. Phys.* **132** no. 1 (1990), 5–25. MR 1069198. Zbl 0722.34070. <https://doi.org/10.1007/BF02277997>.
- [Fur97] A. FURMAN, On the multiplicative ergodic theorem for uniquely ergodic systems, *Ann. Inst. H. Poincaré Probab. Statist.* **33** no. 6 (1997), 797–815. MR 1484541. Zbl 0892.60011. [https://doi.org/10.1016/S0246-0203\(97\)80113-6](https://doi.org/10.1016/S0246-0203(97)80113-6).
- [Gor76] A. YA. GORDON, The point spectrum of the one-dimensional Schrödinger operator, *Uspekhi Mat. Nauk* **31** no. 4(190) (1976), 257–258. MR 0458247. Zbl 0342.34012. Available at <http://www.mathnet.ru/php/archive.phtml?wshow=paper&jrnid=rm&paperid=3813&option.lang=eng>.
- [HJY] R. HAN, S. JITOMIRSKAYA, and F. YANG, Universal hierarchical structure of eigenfunctions of the Maryland model, in preparation.
- [HS89] B. HELFFER and J. SJÖSTRAND, Semiclassical analysis for Harper’s equation. III. Cantor structure of the spectrum, *Mém. Soc. Math. France (N.S.)* no. 39 (1989), 1–124. MR 1041490. Zbl 0725.34099.
- [Hof76] D. R. HOFSTADTER, Energy levels and wave functions of Bloch electrons in rational and irrational magnetic fields, *Physical Review B* **14** no. 6 (1976), 2239. <https://doi.org/10.1103/PhysRevB.14.2239>.
- [HY12] X. HOU and J. YOU, Almost reducibility and non-perturbative reducibility of quasi-periodic linear systems, *Invent. Math.* **190** no. 1 (2012), 209–260. MR 2969277. Zbl 1294.37027. <https://doi.org/10.1007/s00222-012-0379-2>.
- [Jit95] S. JITOMIRSKAYA, Almost everything about the almost Mathieu operator. II, in *XIth International Congress of Mathematical Physics* (Paris, 1994), Int. Press, Cambridge, MA, 1995, pp. 373–382. MR 1370694. Zbl 1052.82539.
- [Jit99] S. JITOMIRSKAYA, Metal-insulator transition for the almost Mathieu operator, *Ann. of Math. (2)* **150** no. 3 (1999), 1159–1175. MR 1740982. Zbl 0946.47018. <https://doi.org/10.2307/121066>.
- [Jit07] S. JITOMIRSKAYA, Ergodic Schrödinger operators (on one foot), in *Spectral Theory and Mathematical Physics: A Festschrift in Honor of Barry Simon’s 60th Birthday, Proc. Sympos. Pure Math.* **76**, Amer. Math. Soc., Providence, RI, 2007, pp. 613–647. MR 2307750. Zbl 1129.82018. <https://doi.org/10.1090/pspum/076.2/2307750>.
- [JK16] S. JITOMIRSKAYA and I. KACHKOVSKIY, L^2 -reducibility and localization for quasiperiodic operators, *Math. Res. Lett.* **23** no. 2 (2016), 431–444. MR 3512893. Zbl 1351.81048. <https://doi.org/10.4310/MRL.2016.v23.n2.a7>.

- [JKL] S. JITOMIRSKAYA, H. KRÜGER, and W. LIU, Exact dynamical exponent for the almost Mathieu operator, preprint.
- [JL] S. JITOMIRSKAYA and W. LIU, in preparation.
- [JL18] S. JITOMIRSKAYA and W. LIU, Universal reflective-hierarchical structure of quasiperiodic eigenfunctions and sharp spectral transition in phase, 2018. arXiv 1802.00781.
- [JLT] S. JITOMIRSKAYA, W. LIU, and S. TCHEREMCHANTZEV, Wavepacket spreading and fractal spectral dimension of quasiperiodic operators with singular continuous spectrum, preprint.
- [JS94] S. JITOMIRSKAYA and B. SIMON, Operators with singular continuous spectrum. III. Almost periodic Schrödinger operators, *Comm. Math. Phys.* **165** no. 1 (1994), 201–205. MR 1298948. Zbl 0830.34074. <https://doi.org/10.1007/BF02099743>.
- [Las05] Y. LAST, Spectral theory of Sturm-Liouville operators on infinite intervals: a review of recent developments, in *Sturm-Liouville Theory*, Birkhäuser, Basel, 2005, pp. 99–120. MR 2145079. Zbl 1098.39011. https://doi.org/10.1007/3-7643-7359-8_5.
- [LS99] Y. LAST and B. SIMON, Eigenfunctions, transfer matrices, and absolutely continuous spectrum of one-dimensional Schrödinger operators, *Invent. Math.* **135** no. 2 (1999), 329–367. MR 1666767. Zbl 0931.34066. <https://doi.org/10.1007/s002220050288>.
- [LY15a] W. LIU and X. YUAN, Anderson localization for the almost Mathieu operator in the exponential regime, *J. Spectr. Theory* **5** no. 1 (2015), 89–112. MR 3340177. Zbl 1319.47029. <https://doi.org/10.4171/JST/92>.
- [LY15b] W. LIU and X. YUAN, Anderson localization for the completely resonant phases, *J. Funct. Anal.* **268** no. 3 (2015), 732–747. MR 3292353. Zbl 1333.47025. <https://doi.org/10.1016/j.jfa.2014.10.024>.
- [MJ17] C. A. MARX and S. JITOMIRSKAYA, Dynamics and spectral theory of quasiperiodic Schrödinger-type operators, *Ergodic Theory Dynam. Systems* **37** no. 8 (2017), 2353–2393. MR 3719264. Zbl 06823073. <https://doi.org/10.1017/etds.2016.16>.
- [Pei33] R. PEIERLS, Zur Theorie des Diamagnetismus von Leitungselektronen, *Z. Physik* **80** no. 11-12 (1933), 763–791. Zbl 0006.19204. <https://doi.org/10.1007/BF01342591>.
- [Sim85] B. SIMON, Almost periodic Schrödinger operators. IV. The Maryland model, *Ann. Physics* **159** no. 1 (1985), 157–183. MR 0776654. Zbl 0595.35032. [https://doi.org/10.1016/0003-4916\(85\)90196-4](https://doi.org/10.1016/0003-4916(85)90196-4).
- [Sim00] B. SIMON, Schrödinger operators in the twenty-first century, in *Mathematical Physics 2000*, Imp. Coll. Press, London, 2000, pp. 283–288. MR 1773049. Zbl 1074.81521. https://doi.org/10.1142/9781848160224_0014.
- [Sin87] YA. G. SINAI, Anderson localization for one-dimensional difference Schrödinger operator with quasiperiodic potential, *J. Statist. Phys.* **46** no. 5-6 (1987), 861–909. MR 0893122. Zbl 0682.34023. <https://doi.org/10.1007/BF01011146>.

- [Tho84] D. J. THOULESS, Quantized Hall conductance, particle transport and topological invariants, in *XIIIth International Colloquium on Group Theoretical Methods in Physics* (College Park, Md., 1984), World Sci. Publishing, Singapore, 1984, pp. 319–323. MR 0815710.
- [WZ94] P. B. WIEGMANN and A. V. ZABRODIN, Quantum group and magnetic translations Bethe ansatz for the Asbel-Hofstadter problem, *Nuclear Phys. B* **422** no. 3 (1994), 495–514. MR 1287576. Zbl 0990.82506. [https://doi.org/10.1016/0550-3213\(94\)90443-X](https://doi.org/10.1016/0550-3213(94)90443-X).
- [YZ13] J. YOU and Q. ZHOU, Embedding of analytic quasi-periodic cocycles into analytic quasi-periodic linear systems and its applications, *Comm. Math. Phys.* **323** no. 3 (2013), 975–1005. MR 3106500. Zbl 1286.37004. <https://doi.org/10.1007/s00220-013-1800-4>.
- [You97] L.-S. YOUNG, Lyapunov exponents for some quasi-periodic cocycles, *Ergodic Theory Dynam. Systems* **17** no. 2 (1997), 483–504. MR 1444065. Zbl 0873.28013. <https://doi.org/10.1017/S0143385797079170>.
- [Zhi91] S. Y. ZHITOMIRSKAYA, Singular spectral properties of a one-dimensional Schrödinger operator with almost periodic potential, in *Dynamical Systems and Statistical Mechanics* (Moscow, 1991), *Adv. Soviet Math.* **3**, Amer. Math. Soc., Providence, RI, 1991, Translated from the Russian by V. E. Nazaïkinskii, pp. 215–254. MR 1118164. Zbl 0734.34075.

(Received: September 27, 2016)

(Revised: September 29, 2017)

UNIVERSITY OF CALIFORNIA, IRVINE, CA
E-mail: szhitomi@math.uci.edu

UNIVERSITY OF CALIFORNIA, IRVINE, CA
E-mail: liuwencai1226@gmail.com