

# Uniform Manin-Mumford for a family of genus 2 curves

By LAURA DEMARCO, HOLLY KRIEGER, and HEXI YE

## Abstract

We introduce a general strategy for proving quantitative and uniform bounds on the number of common points of height zero for a pair of inequivalent height functions on  $\mathbb{P}^1(\overline{\mathbb{Q}})$ . We apply this strategy to prove a conjecture of Bogomolov, Fu, and Tschinkel asserting uniform bounds on the number of common torsion points of elliptic curves in the case of two Legendre curves over  $\mathbb{C}$ . As a consequence, we obtain two uniform bounds for a two-dimensional family of genus 2 curves: a uniform Manin-Mumford bound for the family over  $\mathbb{C}$ , and a uniform Bogomolov bound for the family over  $\overline{\mathbb{Q}}$ .

## 1. Introduction

In this article, we use the Arakelov-Zhang intersection of adelically-metrized line bundles on  $\mathbb{P}^1(\overline{\mathbb{Q}})$  to prove a uniform Manin-Mumford bound for a two-dimensional family of genus 2 curves over  $\mathbb{C}$ . The Manin-Mumford Conjecture, proved by Raynaud [Ray83], asserts the following: Let  $X$  be any smooth complex projective curve of genus  $g \geq 2$ ,  $P \in X(\mathbb{C})$  any point,  $j_P : X \hookrightarrow J(X)$  the Abel-Jacobi embedding of  $X$  into its Jacobian  $J(X)$  based at  $P$ , and  $J(X)^{\text{tors}}$  the set of torsion points of the Jacobian. Then

$$(1.1) \quad |j_P(X) \cap J(X)^{\text{tors}}| < \infty.$$

In the case of genus  $g = 2$ , the curve is hyperelliptic, and the fixed points of the hyperelliptic involution provide geometrically natural choices of base point for the Abel-Jacobi map. We show there is a uniform bound on the number of torsion images under such a map, provided the curve is also bielliptic, meaning that it admits a degree-two branched covering to an elliptic curve.

---

Keywords: arithmetic intersection, Manin-Mumford, torsion points, elliptic curves, non-archimedean potential theory, Lattès maps, preperiodic points

AMS Classification: Primary: 14H40; Secondary: 11G50, 37F45, 37P50.

© 2020 Department of Mathematics, Princeton University.

THEOREM 1.1. *There exists a uniform constant  $B$  such that*

$$|j_P(X) \cap J(X)^{\text{tors}}| \leq B$$

for all smooth, bielliptic curves  $X$  over  $\mathbb{C}$  of genus 2 and all Weierstrass points  $P$  on  $X$ .

The curves satisfying the hypothesis of [Theorem 1.1](#) form a complex surface  $\mathcal{L}_2$  in the moduli space  $\mathcal{M}_2$  of genus 2 curves. These  $X$  are also characterized by the property that their Jacobians admit real multiplication by the real quadratic order of discriminant 4. Further details on  $\mathcal{L}_2$  are given in [Section 9](#).

*Remark 1.2.* We do not give an explicit value for the  $B$  of [Theorem 1.1](#), but this bound can be made effective by estimating the continuity constants of [Section 4](#). Poonen showed that there are infinitely many curves  $X \in \mathcal{L}_2$  for which  $|j_P(X) \cap J(X)^{\text{tors}}|$  is at least 22, taking  $P$  to be a Weierstrass point on  $X$  [[Poo00](#), Th. 1]. More recently, Stoll found an example with  $|j_P(X) \cap J(X)^{\text{tors}}| = 34$  for Weierstrass point  $P$  on  $X \in \mathcal{L}_2$  [[Sto](#)]; the curve  $X$  is defined over  $\mathbb{Q}$ . We know of no curve  $X \in \mathcal{M}_2(\mathbb{C})$  and point  $P \in X$  satisfying  $|j_P(X) \cap J(X)^{\text{tors}}| > 34$ .

*Remark 1.3.* Although [Theorem 1.1](#) provides a uniform bound on the number of torsion points on  $X$  in its Jacobian, there cannot be a uniform bound on the *order* of these torsion points. See [Proposition 9.2](#).

The question of uniformity in (1.1) was raised by Mazur in [[Maz86](#)], who asked if a bound could be given that depends only on the genus  $g$  of the curve  $X$ . Quantitative bounds on torsion points on curves have been obtained when the curve is defined over a number field, notably by Coleman [[Col85](#)], Buium [[Bui96](#)], Hrushovski [[Hru01](#)], and more recently by Katz, Rabinoff, and Zureick-Brown [[KRZB16](#)]. By quantifying the  $p$ -adic approach to (1.1), these authors achieve bounds for general families of curves; however, these bounds all involve dependence on field of definition or the choice of a prime for the family of curves, so they are not uniform for families over  $\overline{\mathbb{Q}}$  or  $\mathbb{C}$ .

Our new technique that yields [Theorem 1.1](#) is a quantification of the approach of Szpiro, Ullmo, and Zhang [[SUZ97](#)], [[Ull98](#)], [[Zha98](#)] to proving (1.1), utilizing adelic equidistribution theory. We first reduce to the setting where the curve is defined over  $\overline{\mathbb{Q}}$ . Over  $\overline{\mathbb{Q}}$ , we build on the proof of the quantitative equidistribution theorem for height functions on  $\mathbb{P}^1(\overline{\mathbb{Q}})$  of Favre and Rivera-Letelier [[FRL06](#)].

In fact, we deduce [Theorem 1.1](#) from a case of the following conjecture, discussed by Bogomolov and Tschinkel [[BT07](#)] and stated formally as [[BFT18](#), Conjs. 2 and 12], which asserts uniform bounds on common torsion points for pairs of elliptic curves. By a *standard projection*  $\pi : E \rightarrow \mathbb{P}^1$  of an elliptic curve

$E$  over  $\mathbb{C}$ , we mean any degree-two quotient that identifies a point  $P$  and its inverse  $-P$ . Note that every standard projection  $\pi$  will have a simple critical point at each of the four elements of the 2-torsion subgroup  $E[2]$ .

CONJECTURE 1.4 ([BFT18]). *There exists a uniform constant  $B$  such that*

$$|\pi_1(E_1^{\text{tors}}) \cap \pi_2(E_2^{\text{tors}})| \leq B$$

*for any pair of elliptic curves  $E_i$  over  $\mathbb{C}$  and any pair of standard projections  $\pi_i$  for which  $\pi_1(E_1[2]) \neq \pi_2(E_2[2])$ .*

Note that if  $\pi_1(E_1[2]) = \pi_2(E_2[2])$ , then  $E_1$  is isomorphic to  $E_2$  and  $\pi_1(E_1^{\text{tors}}) = \pi_2(E_2^{\text{tors}})$ . The finiteness of the set  $\pi_1(E_1^{\text{tors}}) \cap \pi_2(E_2^{\text{tors}})$ , under the assumption that  $\pi_1(E_1[2]) \neq \pi_2(E_2[2])$ , follows from the main theorem of Raynaud in [Ray83]; indeed, the diagonal in  $\mathbb{P}^1 \times \mathbb{P}^1$  lifts to a (singular) curve  $C \subset E_1 \times E_2$  via  $\pi_1 \times \pi_2$  with normalization of genus  $g \geq 2$  [BT07].

We prove [Conjecture 1.4](#) in the case of maximal overlap of the 2-torsion points; i.e., when

$$|\pi_1(E_1[2]) \cap \pi_2(E_2[2])| = 3.$$

This setting corresponds to the case where the (normalization of the) curve  $C$  in  $E_1 \times E_2$  has genus 2. By fixing coordinates on  $\mathbb{P}^1$ , it suffices to work with the Legendre family of elliptic curves

$$(1.2) \quad E_t : y^2 = x(x-1)(x-t)$$

with  $t \in \mathbb{C} \setminus \{0, 1\}$  and the standard projection  $\pi(x, y) = x$  on  $E_t$ . (See [Corollary 8.2](#).)

THEOREM 1.5. *There exists a uniform constant  $B$  such that*

$$|\pi(E_{t_1}^{\text{tors}}) \cap \pi(E_{t_2}^{\text{tors}})| \leq B,$$

*for all  $t_1 \neq t_2$  in  $\mathbb{C} \setminus \{0, 1\}$ , for the curves  $E_t$  defined by (1.2) and projection  $\pi(x, y) = x$ .*

To prove [Theorem 1.5](#), we introduce a general strategy for bounding the number of common height-zero points for any pair of distinct height functions  $h_1, h_2 : \mathbb{P}^1(\overline{\mathbb{Q}}) \rightarrow \mathbb{R}$  that arise from continuous, semipositive, adelic metrics on the line bundle  $\mathcal{O}_{\mathbb{P}^1}(1)$ . There is a natural Arakelov-Zhang pairing between any two such heights, given by the intersection number of the associated metrized line bundles. Our heights are normalized so this intersection number, which we denote by  $h_1 \cdot h_2$ , will satisfy

$$h_1 \cdot h_2 \geq 0 \text{ with equality if and only if } h_1 = h_2.$$

Details on these heights and the pairing are given in [Section 2](#). The value of  $h_1 \cdot h_2$  provides a notion of distance between the two heights (as was observed by

Fili in [Fil17]). It follows from equidistribution [CL06], [FRL06], [BR10] that

$$(1.3) \quad \lim_{n \rightarrow \infty} h_2(x_n) = h_1 \cdot h_2$$

for any infinite sequence of distinct points  $x_n \in \mathbb{P}^1(\overline{\mathbb{Q}})$  such that  $h_1(x_n) \rightarrow 0$  as  $n \rightarrow \infty$ , suggesting that large numbers of common zeroes between  $h_1$  and  $h_2$  will imply that  $h_1$  and  $h_2$  are close. However, this measure of closeness between two heights is not generally uniform in families of heights, because the rate of equidistribution is not uniform. Nevertheless, by bounding the height pairing  $h_1 \cdot h_2$  from below, we can obtain an upper bound on the number of common zeroes for certain families.

In the context of [Theorem 1.5](#), we consider the family of height functions  $\hat{h}_t$  on  $\mathbb{P}^1(\overline{\mathbb{Q}})$  induced from the Néron-Tate canonical height on the elliptic curve  $E_t$  for  $t \in \overline{\mathbb{Q}} \setminus \{0, 1\}$ ; its zeroes are precisely the elements of  $\pi(E_t^{\text{tors}})$ . We implement this general strategy by proving three bounds on the intersection pairing  $\hat{h}_{t_1} \cdot \hat{h}_{t_2}$ . We prove a uniform lower bound on the pairing:

**THEOREM 1.6.** *There exists  $\delta > 0$  such that*

$$\hat{h}_{t_1} \cdot \hat{h}_{t_2} \geq \delta$$

*for all  $t_1 \neq t_2 \in \overline{\mathbb{Q}} \setminus \{0, 1\}$ .*

We also prove an asymptotic lower bound for parameters  $t_1$  and  $t_2$  with large height:

**THEOREM 1.7.** *There exist constants  $\alpha, \beta > 0$  such that*

$$\hat{h}_{t_1} \cdot \hat{h}_{t_2} \geq \alpha h(t_1, t_2) - \beta$$

*for all  $t_1 \neq t_2$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$ . Here  $h(t_1, t_2)$  is the naive logarithmic height on  $\mathbb{A}^2(\overline{\mathbb{Q}})$ .*

We find an upper bound that depends on the number of common zeroes of  $\hat{h}_{t_1}$  and  $\hat{h}_{t_2}$  as well as the heights of the parameters  $t_1$  and  $t_2$ :

**THEOREM 1.8.** *For all  $\varepsilon > 0$ , there exists a constant  $C(\varepsilon) > 0$  such that*

$$\hat{h}_{t_1} \cdot \hat{h}_{t_2} \leq \left( \varepsilon + \frac{C(\varepsilon)}{N(t_1, t_2)} \right) (h(t_1, t_2) + 1)$$

*for all  $t_1$  and  $t_2$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$ , where  $N(t_1, t_2) := |\pi(E_{t_1}^{\text{tors}}) \cap \pi(E_{t_2}^{\text{tors}})|$ .*

The three theorems combine to give a uniform bound on the number  $N(t_1, t_2)$  of common zeroes of  $\hat{h}_{t_1}$  and  $\hat{h}_{t_2}$  for all  $t_1 \neq t_2$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$ .

[Theorems 1.6](#) and [1.7](#) follow from estimates on the local height functions and the local equilibrium measures on the  $v$ -adic Berkovich projective line at each place  $v$  of a number field containing  $t_1$  and  $t_2$ , computed using the dynamical Lattès map  $f_t : \mathbb{P}^1 \rightarrow \mathbb{P}^1$  induced by multiplication by 2 on a

Legendre curve  $E_t$ . The non-archimedean contributions to  $\hat{h}_{t_1} \cdot \hat{h}_{t_2}$  turn out to be straightforward to compute for these heights. Significant technical issues arise when  $v$  is archimedean and both parameters  $t_i$  are tending to the singularity set  $\{0, 1, \infty\}$  for this family; we resolve these issues by appealing to the theory of degenerations of complex dynamical systems on  $\mathbb{P}^1(\mathbb{C})$ , in which a family of complex rational maps degenerates to a non-archimedean dynamical system acting on a Berkovich space, as in the work of DeMarco-Faber [DMF14] and Favre [Fav16], using the formalism of hybrid space as discussed by Boucksom-Jonsson in [BJ17].

For [Theorem 1.8](#), we expand upon the quantitative equidistribution results of Favre-Rivera-Letelier [FRL06] and Fili [Fil17] to analyze the rates of convergence of measures supported on finite sets of zeroes of a height  $h$  to the associated equilibrium measures at each place  $v$ . To do so requires control on the modulus of continuity of the local heights, and again we rely on estimates from the hybrid space to treat the cases where a parameter  $t$  is tending to one of the singularities for the family  $E_t$ .

Although [Theorem 1.6](#) alone was not enough to prove [Theorem 1.5](#), it implies a uniform bound of a different sort, when combined with Zhang's inequality on the essential minimum of a height function [Zha95]:

**PROPOSITION 1.9.** *Choose any  $b$  satisfying  $0 < b < \delta/2$  for the  $\delta$  of [Theorem 1.6](#). Then the set*

$$S(b, t_1, t_2) := \{x \in \mathbb{P}^1(\overline{\mathbb{Q}}) : \hat{h}_{t_1}(x) + \hat{h}_{t_2}(x) \leq b\}$$

*is finite for each pair  $t_1 \neq t_2 \in \overline{\mathbb{Q}} \setminus \{0, 1\}$ .*

The complete proof of [Theorem 1.5](#), however, gives a much stronger statement: we obtain a uniform bound on the size of the set  $S(b, t_1, t_2)$  defined in [Proposition 1.9](#) over all pairs  $t_1 \neq t_2 \in \overline{\mathbb{Q}} \setminus \{0, 1\}$ ; see [Theorem 8.1](#). This in turn provides a uniform version of the Bogomolov Conjecture for the associated family of genus 2 curves. The Bogomolov Conjecture was proved for each individual curve  $X$  over  $\overline{\mathbb{Q}}$  in [Ull98], [Zha98]. To state our result precisely, we fix ample and symmetric line bundles on the family of Jacobians  $J(X)$  for the genus 2 curves  $X$  defined over  $\overline{\mathbb{Q}}$  that we consider in [Theorem 1.1](#). Specifically, we take  $L_X = \Phi^* L_D$  for the isogeny  $\Phi : J(X) \rightarrow E_1 \times E_2$  of [Proposition 9.1](#), with  $L_D$  the line bundle associated to the divisor  $D = \{O_1\} \times E_2 + E_1 \times \{O_2\}$ , where  $O_i$  is the identity element of  $E_i$ .

**THEOREM 1.10.** *There exist constants  $B$  and  $b > 0$  such that*

$$|\{x \in j_P(X)(\overline{\mathbb{Q}}) : \hat{h}_{L_X}(x) \leq b\}| \leq B$$

*for all smooth curves  $X$  over  $\overline{\mathbb{Q}}$  of genus 2 admitting a degree-two map to an elliptic curve and all Weierstrass points  $P$  on  $X$ , where  $\hat{h}_{L_X}$  is the Néron-Tate canonical height on the Jacobian  $J(X)$ .*

Finally, we mention that we implement this general strategy towards uniform boundedness in a follow-up article [DKY19] in another setting, providing a uniform bound on the number of common preperiodic points for distinct polynomials of the form  $f_c(z) = z^2 + c$  with  $c \in \mathbb{C}$ .

*Remark 1.11.* We have chosen to work with the Arakelov-Zhang pairing  $\hat{h}_{t_1} \cdot \hat{h}_{t_2}$  to measure proximity of the two height functions, with  $t_1 \neq t_2$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$ , but there are other choices we could have made. For example, Kawaguchi and Silverman in [KS09] study

$$\delta(\hat{h}_{t_1}, \hat{h}_{t_2}) := \sup_{x \in \mathbb{P}^1(\overline{\mathbb{Q}})} |\hat{h}_{t_1}(x) - \hat{h}_{t_2}(x)|.$$

It turns out that the two quantities are comparable for this family of heights. The upper bound  $\hat{h}_{t_1} \cdot \hat{h}_{t_2} \leq \delta(\hat{h}_{t_1}, \hat{h}_{t_2})$  can be seen as a corollary of arithmetic equidistribution and (1.3), and therefore holds for any pair of normalized heights coming from continuous, semipositive adelic metrics on  $\mathcal{O}_{\mathbb{P}^1}(1)$ . A lower bound of the form  $\hat{h}_{t_1} \cdot \hat{h}_{t_2} \geq C_1 \delta(\hat{h}_{t_1}, \hat{h}_{t_2}) - C_2$  for positive constants  $C_1, C_2$ , and for all  $t_1 \neq t_2$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$ , is a consequence of [Theorem 1.7](#), when combined with [KS09, Th. 1]. However, such a lower bound does not hold for all pairs of heights coming from metrics on  $\mathcal{O}_{\mathbb{P}^1}(1)$ . A comparison of these two pairings is addressed further in [DKY19] for the canonical heights associated to morphisms of  $\mathbb{P}^1$ .

*Outline of the paper.* We fix our notation and provide background in [Section 2](#). [Sections 3, 4](#), and [5](#) provide the estimates on local height functions and local measures needed to prove all of our theorems. [Theorem 1.7](#) is proved in [Section 6](#), and from it we deduce [Theorem 1.6](#) and [Proposition 1.9](#). A generalization of [Theorem 1.8](#) is proved in [Section 7](#) that treats points of small height, not only of height 0. We prove [Theorem 1.5](#) in [Section 8](#) and finally [Theorems 1.1](#) and [1.10](#) in [Section 9](#).

*Acknowledgements.* We thank the American Institute of Mathematics, where the initial work for this paper took place as part of an AIM SQuaRE. Special thanks go to Ken Jacobs, Mattias Jonsson, Curt McMullen, and Khoa Nguyen for helpful discussions. We would also like to thank the anonymous referees for their many suggestions and careful reading of the article. During the preparation of this paper, L. DeMarco was supported by the National Science Foundation (DMS-1600718), H. Krieger was supported by Isaac Newton Trust (RG74916), and H. Ye was partially supported by ZJNSF (LR18A010001) and NSFC (11701508).

## 2. Heights, measures, and energies

This section develops the background and notation needed for the proofs that follow. Throughout,  $K$  is a number field and  $M_K$  its set of places.

2.1. *The canonical height.* Fix  $t \in \overline{\mathbb{Q}} \setminus \{0, 1\}$ . Let  $E_t$  be the Legendre elliptic curve and  $\pi : E_t \rightarrow \mathbb{P}^1$  the projection defined by  $\pi(x, y) = x$ . The multiplication-by-two endomorphism on  $E_t$  descends via  $\pi$  to a morphism of degree 4 on  $\mathbb{P}^1$  given by

$$(2.1) \quad f_t(x) = \frac{(x^2 - t)^2}{4x(x-1)(x-t)}.$$

The canonical height on the elliptic curve

$$\hat{h}_{E_t} : E_t(\overline{\mathbb{Q}}) \rightarrow \mathbb{R}$$

can be defined via the projection  $\pi$  and the iteration of  $f_t$  as  $h_{E_t}(P) := \frac{1}{2} \hat{h}_t(\pi(P))$ , where

$$\hat{h}_t : \mathbb{P}^1(\overline{\mathbb{Q}}) \rightarrow \mathbb{R}$$

is the dynamical canonical height defined by

$$(2.2) \quad \hat{h}_t(x) = \lim_{n \rightarrow \infty} \frac{1}{4^n} h(f_t^n(x)).$$

Here,  $h$  is the (logarithmic) Weil height on  $\mathbb{P}^1(\overline{\mathbb{Q}})$ . Note that  $\hat{h}_t(x) \geq 0$  for all  $x \in \mathbb{P}^1(\overline{\mathbb{Q}})$ , and

$$\hat{h}_t(x) = 0 \iff x \in \pi(E_t^{\text{tors}})$$

[Sil09], [CS93].

The height  $\hat{h}_t$  has a local decomposition as follows: for any number field  $K$  containing  $t$ , and for each place  $v \in M_K$ , there exists a local height function  $\lambda_{t,v}$  such that

$$\hat{h}_t(x) = \sum_{v \in M_K} \frac{r_v}{|\text{Gal}(\overline{K}/K) \cdot x|} \sum_{y \in \text{Gal}(\overline{K}/K) \cdot x} \lambda_{t,v}(y)$$

for all  $x \in \overline{K}$ , where

$$r_v := \frac{[K_v : \mathbb{Q}_v]}{[K : \mathbb{Q}]}.$$

The local heights  $\lambda_{t,v}$  can be chosen to extend continuously to  $\mathbb{P}^1(\mathbb{C}_v) \setminus \{\infty\}$ , where  $\mathbb{C}_v$  is the completion (with respect to  $v$ ) of an algebraic closure of the completion  $K_v$ , and to satisfy

$$\lambda_{t,v}(x) = \log |x|_v + \mathcal{O}(1)$$

as  $|x|_v \rightarrow \infty$ .

2.2. *Local heights and escape rates.* To compute the local heights, we will often express the maps  $f_t : \mathbb{P}^1 \rightarrow \mathbb{P}^1$  of (2.1) in homogeneous coordinates, as

$$F_t(z, w) := ((z^2 - tw^2)^2, 4zw(z - w)(z - tw))$$

for  $z$  and  $w$  in  $\mathbb{C}_v$ . As observed in [BR10, Ch. 10], its escape-rate function

$$(2.3) \quad G_{F_t, v}(z, w) := \lim_{n \rightarrow \infty} \frac{1}{4^n} \log \|F_t^n(z, w)\|_v,$$

where  $\|(z, w)\|_v = \max\{|x|_v, |y|_v\}$ , satisfies

$$\hat{h}_t(x) = \sum_{v \in M_K} \frac{r_v}{|\text{Gal}(\overline{K}/K) \cdot \tilde{x}|} \sum_{\tilde{y} \in \text{Gal}(\overline{K}/K) \cdot \tilde{x}} G_{F_t, v}(\tilde{y})$$

for  $x \in \mathbb{P}^1(\overline{K})$  and  $\tilde{x}$  any choice of lift of  $x$  to  $\overline{K}^2 \setminus \{(0, 0)\}$ . In particular, we may take

$$(2.4) \quad \lambda_{t, v}(x) = G_{F_t, v}(x, 1)$$

as a local height for  $\hat{h}_t$ .

The elliptic curves  $E_t$  and  $E_{1-t}$  and  $E_{1/t}$  are isomorphic, with the following transformation formulas for the local heights:

**PROPOSITION 2.1.** *Fix any number field  $K$  and  $v \in M_K$ . Then, for all  $t \in K \setminus \{0, 1\}$ , we have*

$$G_{F_{1-t}, v}(1 - z, 1) = G_{F_t, v}(z, 1) = G_{F_{1/t}, v}(z, t) = G_{F_{1/t}, v}(z/t, 1) + \log |t|_v.$$

*Proof.* Let  $A$  be the automorphism  $A(z, w) = (w - z, w)$ . Then

$$A \circ F_t^n = -F_{1-t}^n \circ A$$

for all iterates, proving the first equality. Similarly, let  $B(z, w) = (z, tw)$ . Then

$$B \circ F_t^n = F_{1/t}^n \circ B$$

for all iterates, proving the second equality. The final equality follows from the logarithmic homogeneity of  $G$ .  $\square$

2.3. *The Berkovich projective line.* Let  $K$  be a number field. For each  $v \in M_K$ , let  $\mathbb{A}_v^{1, an}$  denote the Berkovich affine line over  $\mathbb{C}_v$ . For non-archimedean  $v$ , the points of  $\mathbb{A}_v^{1, an}$  come in four types. The Type I points in  $\mathbb{A}_v^{1, an}$  are, by definition, the elements of the field  $\mathbb{C}_v$ . The Type II points are in one-to-one correspondence with disks  $D(a, r) = \{x \in \mathbb{C}_v : |x - a|_v \leq r\}$  with  $r > 0$  rational, and these are the branch points for the underlying tree structure on  $\mathbb{A}_v^{1, an}$ . The Type III points correspond to disks  $D(a, r)$  with  $r$  irrational. (We will not need the Type IV points in this article.) A Type II or III point corresponding to  $D(a, r)$  will be denoted by  $\zeta_{a, r}$ . The *Gauss point*  $\zeta_{0, 1}$  is the Type II point identified with the unit disk. The Berkovich projective line  $\mathbb{P}_v^{1, an} = \mathbb{A}_v^{1, an} \cup \{\infty\}$

is the one-point compactification of  $\mathbb{A}_v^{1,an}$ , which is a canonically-defined path-connected compact Hausdorff space containing  $\mathbb{P}^1(\mathbb{C}_v)$  as a dense subspace. If  $v$  is archimedean, then  $\mathbb{C}_v \simeq \mathbb{C}$  and  $\mathbb{P}_v^{1,an} = \mathbb{P}^1(\mathbb{C})$ .

For each  $v \in M_K$ , there is a distribution-valued Laplacian operator  $\Delta$  on  $\mathbb{P}_v^{1,an}$ . The function  $\log^+ |z|_v$  on  $\mathbb{P}^1(\mathbb{C}_v)$  extends naturally to a continuous real valued function  $\mathbb{P}_v^{1,an} \rightarrow \mathbb{R} \cup \{\infty\}$ , and the Laplacian is normalized such that

$$\Delta \log^+ |z|_v = \omega_v - \delta_\infty$$

on  $\mathbb{P}_v^{1,an}$ , where  $\omega_v = m_{S^1}$  is the Lebesgue probability measure on the unit circle when  $v$  is archimedean, and  $\omega_v = \delta_G$  is a point mass at the Gauss point of  $\mathbb{P}_v^{1,an}$  when  $v$  is non-archimedean. A probability measure  $\mu_v$  on  $\mathbb{P}_v^{1,an}$  is said to have *continuous potentials* if  $\mu_v - \omega_v = \Delta g$  with  $g : \mathbb{P}_v^{1,an} \rightarrow \mathbb{R}$  continuous. The function  $g$  for  $\mu_v$  is unique up to the addition of a constant. See [BR10, Ch. 5] for more details. Note that the Laplacian used here is the negative of the one appearing in [PST12] and [BR10], but it agrees with the usual Laplacian (up to a factor of  $2\pi$ ) at the archimedean places.

For  $v$  non-archimedean, we set

$$\mathbb{H} := \mathbb{A}_v^{1,an} \setminus \mathbb{C}_v.$$

The hyperbolic distance  $d_{\text{hyp}}$  on  $\mathbb{H}$  gives it the structure of a metrized  $\mathbb{R}$ -tree and satisfies

$$d_{\text{hyp}}(\zeta_{a,r_1}, \zeta_{a,r_2}) = \log(r_1/r_2)$$

for any  $a \in \mathbb{C}_v$  and any  $r_1 \geq r_2 > 0$ . We will say that a probability measure  $\mu_v$  on  $\mathbb{H}$  is an *interval measure* if it is the uniform distribution on an interval  $[\zeta_1, \zeta_2] \subset \mathbb{H}$  with respect to the linear structure induced from the hyperbolic metric  $d_{\text{hyp}}$ .

**2.4. Canonical measures and good reduction.** For each Legendre elliptic curve  $E_t$  with  $t$  in a number field  $K$  and each  $v \in M_K$ , the local height  $\lambda_{t,v}$  of (2.4) extends to define a continuous and subharmonic function on  $\mathbb{A}_v^{1,an}$  with logarithmic singularity at  $\infty$ . We have

$$\Delta \lambda_{t,v} = \mu_{t,v} - \delta_\infty$$

on  $\mathbb{P}_v^{1,an}$ , where  $\mu_{t,v}$  is the canonical probability measure for the dynamical system  $f_t$  at  $v$  [FRL06], [BR10, Th. 10.2].

For archimedean  $v \in M_K$ , the measure  $\mu_{t,v}$  is the unique  $f_t$ -invariant measure on  $\mathbb{P}^1(\mathbb{C})$  achieving the maximal entropy  $\log 4$ . It is the push-forward of the Haar measure on  $E_t(\mathbb{C})$  via the projection  $\pi$  introduced in Section 2.1. See, for example, [Mil06] for a dynamical discussion of the maps  $f_t$  on the Riemann sphere.

For non-archimedean  $v \in M_K$ , if the curve  $E_t$  and the map  $f_t$  have good reduction, the measure  $\mu_{t,v}$  is the point mass  $\delta_G$  supported on the Gauss

point  $\zeta_{0,1}$ . The map  $f_t$  has potential good reduction, meaning that it has good reduction under a suitable change of coordinates on  $\mathbb{P}^1$ , if and only if the measure  $\mu_{t,v}$  is supported at a single Type II point in  $\mathbb{H}$ . In general, the support of  $\mu_{t,v}$  is equal to the Julia set of  $f_t$  in  $\mathbb{P}_v^{1,an}$ .

Recall that the  $j$ -invariant of the elliptic curve  $E_t$  over  $\mathbb{C}$  is given by

$$(2.5) \quad j(t) = \frac{256(1-t+t^2)^3}{(1-t)^2 t^2}.$$

For  $t \in K$  and non-archimedean  $v \in M_K$ , the map  $f_t$  has potential good reduction at  $v$  if and only if the curve  $E_t$  has potential good reduction at  $v$ . This equivalence can be proved via equidistribution of torsion points on  $E_t$  at all places [BP05, Th. 1] (thus implying that the measure  $\mu_{t,v}$  will also be supported at a single point of  $\mathbb{P}_v^{1,an}$ ) or via a direct calculation showing that the Julia set of  $f_t$  is a singleton if and only if  $|j(t)|_v \leq 1$ .

**2.5. The height as an adelic metric.** Suppose  $t \in K \setminus \{0, 1\}$ . The height  $\hat{h}_t$  on  $\mathbb{P}^1(\overline{\mathbb{Q}})$ , introduced in Section 2.1, is induced from an adelic metric on  $\mathcal{O}_{\mathbb{P}^1}(1)$ , in the sense of Zhang [Zha95]. Fixing coordinates on  $\mathbb{P}^1$  and a section  $s$  of  $\mathcal{O}_{\mathbb{P}^1}(1)$  with  $(s) = (\infty)$ , then a metric  $\|\cdot\|_{t,v}$  can be defined at each place  $v$  of  $K$  by setting

$$-\log \|s(z)\|_{t,v} = \lambda_{t,v}(z) = G_{F_t,v}(z, 1)$$

for the function  $G_{F_t,v}$  of (2.3). The height  $\hat{h}_t$  satisfies

$$\hat{h}_t(x) = \sum_{v \in M_K} \frac{r_v}{|\text{Gal}(\overline{K}/K) \cdot x|} \sum_{y \in \text{Gal}(\overline{K}/K) \cdot x} (-\log \|s(y)\|_{t,v})$$

for all  $x \neq \infty$  in  $\mathbb{P}^1(\overline{\mathbb{Q}})$ . Writing  $\lambda_{t,v}(z) = \log |z|_v + c_v + o(1)$  as  $|z|_v \rightarrow \infty$  with a constant  $c_v$  at each place  $v$  of  $K$ , we may compute that

$$(2.6) \quad 0 = \hat{h}_t(\infty) = \sum_{v \in M_K} r_v c_v,$$

because  $\infty$  is the projection of the origin of  $E_t$ .

**2.6. The intersection pairing.** For these heights  $\hat{h}_t$  coming from the Legendre family of elliptic curves, with  $t \in \overline{\mathbb{Q}} \setminus \{0, 1\}$ , we have

$$(2.7) \quad \hat{h}_{t_1} = \hat{h}_{t_2} \iff t_1 = t_2.$$

Indeed, any height coming from an adelic metric on  $\mathcal{O}_{\mathbb{P}^1}(1)$  is uniquely determined, up to an additive constant, by the associated curvature distributions; see, for example, the construction of a height function from the measures in [FRL06]. For heights of the form  $\hat{h}_t$ , at each archimedean place  $v$  of a number field containing  $t$ , the curvature distribution  $\mu_{t,v}$  on  $\mathbb{P}^1(\mathbb{C})$  is the push-forward

of the Haar measure on  $E_t(\mathbb{C})$  by  $\pi$ ; it therefore has a greater density at the four branch points  $\{0, 1, t, \infty\}$  of  $\pi$  and thus determines  $t$ .

There is a well-defined intersection number between any pair of such heights, as in [Zha95] (see also [CL11]); more precisely, it is the arithmetic intersection number of the two associated adelically metrized line bundles. By the non-degeneracy of this height pairing and (2.7),

$$(2.8) \quad \hat{h}_{t_1} \cdot \hat{h}_{t_2} \geq 0 \text{ with equality if and only if } t_1 = t_2,$$

as computed in [PST12].

To define the pairing  $\hat{h}_{t_1} \cdot \hat{h}_{t_2}$ , we fix sections  $s$  and  $u$  of  $\mathcal{O}_{\mathbb{P}^1}(1)$  such that their divisors do not intersect. Given  $t_1$  and  $t_2$  in a number field  $K$ , and a place  $v$  of  $K$ , we set

$$\langle \hat{h}_{t_1}, \hat{h}_{t_2} \rangle_v^{s,u} := \int \log \|s\|_{t_1,v}^{-1} \Delta(\log \|u\|_{t_2,v}^{-1}) = \langle \hat{h}_{t_2}, \hat{h}_{t_1} \rangle_v^{u,s}.$$

The integral is over the Berkovich analytification  $\mathbb{P}_v^{1,an}$  of  $\mathbb{P}^1$ , over the field  $\mathbb{C}_v$ . The metrics satisfy

$$\Delta(\log \|s\|_{t,v}^{-1}) = \mu_{t,v} - \delta_{(s)},$$

and  $\mu_{t,v}$  is the associated curvature distribution.

The height pairing is then defined as

$$(2.9) \quad \hat{h}_{t_1} \cdot \hat{h}_{t_2} := \hat{h}_{t_1}((u)) + \hat{h}_{t_2}((s)) + \sum_{v \in M_K} r_v \langle \hat{h}_{t_1}, \hat{h}_{t_2} \rangle_v^{s,u},$$

which is independent of the choices of  $s$  and  $u$ . This pairing is easily seen to be symmetric, and since  $\hat{h}_t(\infty) = 0$  for all  $t$ , it can be expressed as

(2.10)

$$\begin{aligned} \hat{h}_{t_1} \cdot \hat{h}_{t_2} &= \hat{h}_{t_2}(\infty) + \sum_{v \in M_K} r_v \int (\log \|s\|_{t_1,v}^{-1}) d\mu_{t_2,v} = \sum_{v \in M_K} r_v \int \lambda_{t_1,v} d\mu_{t_2,v} \\ &= \hat{h}_{t_1}(\infty) + \sum_{v \in M_K} r_v \int (\log \|s\|_{t_2,v}^{-1}) d\mu_{t_1,v} = \sum_{v \in M_K} r_v \int \lambda_{t_2,v} d\mu_{t_1,v} \end{aligned}$$

when  $(s) = (\infty)$ .

As  $\hat{h}_t \cdot \hat{h}_t = 0$  for all  $t \in \overline{\mathbb{Q}} \setminus \{0, 1\}$  from (2.8), note that

$$(2.11) \quad \sum_{v \in M_K} r_v \int \lambda_{t,v} d\mu_{t,v} = 0 = \sum_{v \in M_K} r_v c_v,$$

by combining (2.10) and (2.6). The pairing can be rewritten as:

(2.12)

$$\begin{aligned}\hat{h}_{t_1} \cdot \hat{h}_{t_2} &= \frac{1}{2} \left( \hat{h}_{t_2}(\infty) + \hat{h}_{t_1}(\infty) + \sum_{v \in M_K} r_v \left( \int \lambda_{t_1,v} d\mu_{t_2,v} + \int \lambda_{t_2,v} d\mu_{t_1,v} \right) \right) \\ &= \frac{1}{2} \sum_{v \in M_K} r_v \left( \int (\lambda_{t_1,v} - \lambda_{t_2,v}) d\mu_{t_2,v} + \int (\lambda_{t_2,v} - \lambda_{t_1,v}) d\mu_{t_1,v} \right).\end{aligned}$$

The advantage of working with (2.12) is the following local version of the non-degeneracy property (2.8):

PROPOSITION 2.2 ([FRL06, Props. 2.6 and 4.5]). *Let  $K$  be a number field and  $v \in M_K$ . For any  $t_1, t_2 \in K \setminus \{0, 1\}$ , the local energy*

$$E_v(t_1, t_2) := \frac{1}{2} \left( \int (\lambda_{t_1,v} - \lambda_{t_2,v}) d\mu_{t_2,v} + \int (\lambda_{t_2,v} - \lambda_{t_1,v}) d\mu_{t_1,v} \right)$$

*is non-negative; it is equal to 0 if and only if  $\mu_{t_1,v} = \mu_{t_2,v}$ .*

PROPOSITION 2.3. *Let  $v \in M_K$ , and fix  $t_1, t_2 \in K \setminus \{0, 1\}$ . We have*

$$E_v(t_2, t_1) = E_v(t_1, t_2) = E_v(1 - t_1, 1 - t_2) = E_v(1/t_1, 1/t_2).$$

*Proof.* Given measures  $\mu_{t_1,v}$  and  $\mu_{t_2,v}$ , the local energy  $E_v(t_1, t_2)$  can be expressed as

$$-\frac{1}{2} \int g d(\mu_{t_1,v} - \mu_{t_2,v})$$

for any continuous potential  $g$  of the signed measure  $\mu_{t_1,v} - \mu_{t_2,v}$ , because  $g = \lambda_{t_1,v} - \lambda_{t_2,v} + c$  for some constant  $c$ . We have

$$f_{1-t} = \alpha \circ f_t \circ \alpha^{-1}$$

for  $\alpha(z) = 1 - z = \alpha^{-1}(z)$ , such that  $\mu_{1-t,v} = \alpha_* \mu_{t,v}$  and  $g = (\lambda_{t_1,v} - \lambda_{t_2,v}) \circ \alpha^{-1}$  is a potential for the measure  $\mu_{1-t_1,v} - \mu_{1-t_2,v}$ . Therefore,  $E_v(1 - t_1, 1 - t_2) = E_v(t_1, t_2)$ . Similarly, we have  $f_{1/t}(z) = \alpha \circ f_t \circ \alpha^{-1}(z)$  for  $\alpha(z) = z/t$ , so  $E_v(1/t_1, 1/t_2) = E_v(t_1, t_2)$ .  $\square$

2.7. *Measures and mutual energy.* Suppose that  $\nu_1$  and  $\nu_2$  are signed measures on  $\mathbb{P}^1(\mathbb{C})$  with trace measures  $|\nu_i|$  for which the function  $\log |z - w| \in L^1(|\nu_1| \otimes |\nu_2|)$  on  $\mathbb{C}^2 \setminus \text{Diag}$ . The mutual energy of  $\nu_1$  and  $\nu_2$  is defined in [FRL06] by

$$(2.13) \quad (\nu_1, \nu_2) := - \int_{\mathbb{C}^2 \setminus \text{Diag}} \log |z - w| d\nu_1 \otimes d\nu_2.$$

This definition extends to the non-archimedean setting by replacing  $|z - w|$  with the Hsia kernel  $\delta_v(z, w)$  based at the point at  $\infty$ . In this way, for  $v \in M_K^0$ , a

pairing is defined similarly as

$$(2.14) \quad (\nu_1, \nu_2)_v := - \int_{\mathbb{A}_v^{1,an} \times \mathbb{A}_v^{1,an} \setminus \text{Diag}} \log \delta_v(z, w) d\nu_1 \otimes d\nu_2.$$

See [FRL06, §4.4] and [BR10, Ch. 4].

For measures  $\nu_i$  of total mass 0 with continuous potentials on  $\mathbb{P}_v^{1,an}$ , we have

$$(\nu_1, \nu_2)_v = - \int g_1 d\nu_2$$

for any choice of continuous potential  $g_1$  for  $\nu_1$ . Further,  $(\nu_1, \nu_2)_v \geq 0$  with equality if and only if  $\nu_1 = \nu_2$  [FRL06, Props. 2.6 and 4.5]. Note that [Proposition 2.2](#) is a special case of this fact. Indeed, in this notation, the local energy  $E_v(t_1, t_2)$  defined in [Proposition 2.2](#) is given by

$$(2.15) \quad E_v(t_1, t_2) = \frac{1}{2} (\mu_{t_1,v} - \mu_{t_2,v}, \mu_{t_1,v} - \mu_{t_2,v})_v$$

at each place  $v$  of a number field containing  $t_1$  and  $t_2$  for the canonical measures introduced in [Section 2.4](#).

The mutual energy  $(\cdot, \cdot)_v$  of (2.13) and (2.14) can also be defined for discrete measures. If  $F = \{x_1, \dots, x_n\}$  is any finite set in a number field  $K$ , and  $v \in M_K$ , then denote by  $[F]_v$  the probability measure supported equally on the elements of  $F \subset \mathbb{C}_v$ . Then

$$(2.16) \quad \sum_v r_v ([F]_v, [F]_v)_v = \sum_v r_v \frac{1}{|F|^2} \sum_{i \neq j} \log |x_i - x_j|_v = 0$$

by the product formula.

**2.8. A metric on the space of adelic heights.** The height pairing gives rise to a metric on the space of continuous, semipositive, adelic metrics on  $\mathcal{O}_{\mathbb{P}^1}(1)$  [Fil17, Th. 1]. Given a number field  $K$  and any collection of probability measures  $\{\mu_v\}_{v \in M_K}$  on  $\mathbb{P}_v^{1,an}$  with continuous potentials for which  $\mu_v = \omega_v$  at all but finitely many places (where  $\omega_v$  is a point mass supported on the Gauss point), then there is a unique metric on  $\mathcal{O}_{\mathbb{P}^1}(1)$  with curvature distributions given by  $\{\mu_v\}_{v \in M_K}$ , normalized such that its associated height function  $h : \mathbb{P}^1(\mathbb{Q}) \rightarrow \mathbb{R}$  satisfies  $h \cdot h = 0$  [FRL06]. The height pairing between any two such heights is computed as

$$h_1 \cdot h_2 = \frac{1}{2} \sum_{v \in M_K} r_v (\mu_{1,v} - \mu_{2,v}, \mu_{1,v} - \mu_{2,v})_v.$$

Fili observed that a distance between  $h_1$  and  $h_2$  can be defined by

$$\text{dist}(h_1, h_2) := (h_1 \cdot h_2)^{1/2}.$$

Indeed, we have already seen that  $h_1 \cdot h_2 = 0$  if and only if  $h_1 = h_2$  because of the non-degeneracy of the mutual energy  $(\cdot, \cdot)_v$  at each place. Furthermore,

$\text{dist}(\cdot, \cdot)$  satisfies a triangle inequality: at each place, the mutual energy induces a non-degenerate, symmetric, bilinear form on the vector space of measures of mass 0 with continuous potentials on  $\mathbb{P}_v^{1,an}$ , and so the triangle inequality for  $\text{dist}(\cdot, \cdot)$  follows from an  $\ell^2$  triangle inequality.

### 3. Non-archimedean energy

Throughout this section, we fix a number field  $K$  and a non-archimedean place  $v \in M_K$ , and we provide a lower bound on the non-archimedean local energy defined in [Proposition 2.2](#):

**THEOREM 3.1.** *For  $t_1, t_2 \in K \setminus \{0, 1\}$ , we have*

$$E_v(t_1, t_2) - \frac{4}{3} \log |2|_v \geq \begin{cases} \frac{\log^2 |t_1/t_2|_v}{6 \log \max\{|t_2|_v, |t_1|_v\}} & \text{for } \min\{|t_2|_v, |t_1|_v\} > 1, \\ \frac{\log^2 |t_1/t_2|_v}{-6 \log \min\{|t_2|_v, |t_1|_v\}} & \text{for } \max\{|t_2|_v, |t_1|_v\} < 1, \\ \frac{|\log |t_1/t_2|_v|}{6} & \text{otherwise.} \end{cases}$$

*Equality holds for  $v \nmid 2$  with  $\min\{|t_1 - 1|_v, |t_2 - 1|_v\} \geq 1$ .*

#### 3.1. Measure and escape rate for $v \nmid 2$ .

**PROPOSITION 3.2.** *Suppose  $t \in K \setminus \{0, 1\}$  and  $v \nmid 2$ . Then  $f_t$  has good reduction at  $v$  if and only if  $|t(t-1)|_v = 1$ . If  $|t(t-1)|_v \neq 1$ , then  $f_t$  fails to have potential good reduction at  $v$ , and the canonical measure  $\mu_{t,v}$  on  $\mathbb{P}_v^{1,an}$  of  $f_t$  is the interval measure supported on*

$$I = \begin{cases} [\zeta_{0,1}, \zeta_{0,|t|_v}] & \text{for } |t|_v > 1 \text{ or } |t|_v < 1, \\ [\zeta_{0,1}, \zeta_{1,|1-t|_v}] & \text{for } |1-t|_v < 1. \end{cases}$$

*Proof.* By [Proposition 2.1](#), it suffices to treat the cases with  $|t|_v > 1$ . By [\[FRL10, §5.1\]](#),  $f_t^{-1}(I) = I$  and the action of  $f_t$  on  $I$  is by a tent map of degree 2. That is,

$$f_t(\zeta_{0,|t|_v^r}) = \begin{cases} \zeta_{0,|t|_v^{2r-1}} & \text{for } 1/2 \leq r \leq 1, \\ \zeta_{0,|t|_v^{1-2r}} & \text{for } 0 \leq r \leq 1/2. \end{cases}$$

The proposition follows.  $\square$

We may now compute the local height  $\lambda_{t,v}(z) = G_{F_{t,v}}(z, 1)$  on  $\mathbb{A}_v^{1,an}$ , which is locally constant away from the interval  $[0, \infty) \subset \mathbb{A}_v^{1,an}$ .

PROPOSITION 3.3. *Suppose  $v \nmid 2$  is non-archimedean and  $|t(t-1)|_v \geq 1$ . The escape-rate function  $G_{F_t,v}$  satisfies*

$$(3.1) \quad G_{F_t,v}(z, 1) = \begin{cases} \log |z|_v & \text{for } |z|_v \geq |t|_v, \\ \frac{1}{2} \left( \frac{\log^2 |z|_v}{\log |t|_v} + \log |t|_v \right) & \text{for } 1 < |z|_v < |t|_v, \\ \frac{1}{2} \log |t|_v & \text{for } |z|_v \leq 1 \end{cases}$$

for all  $z \in \mathbb{C}_v$ .

*Proof.* Let  $\lambda$  be the continuous extension of the expression on the right-hand side of formula (3.1) to  $\mathbb{A}_v^{1,an}$ . By Proposition 3.2,  $\mu_{t,v}$  is the interval measure corresponding to  $[\zeta_{0,1}, \zeta_{0,|t|_v}]$ , and a direct computation shows that

$$\Delta\lambda = \mu_{t,v} - \delta_\infty.$$

Thus it suffices to show that  $G_{F_t,v}(\cdot, 1)$  and  $\lambda$  agree at a single point. For any  $z_0 \in \mathbb{C}_v$  with  $|z_0|_v > |t|_v$ , define  $(z_n, w_n) := F_t^n(z_0, 1)$ , so that

$$(3.2) \quad (z_{n+1}, w_{n+1}) = F_t(z_n, w_n) = ((z_n^2 - tw_n^2)^2, 4z_n w_n (z_n - tw_n)).$$

Inductively,

$$|z_n|_v = |z_0|_v^{4^n} > |t|_v |w_n|_v > |w_n|_v.$$

Consequently,

$$G_{F_t,v}(z_0, 1) = \lim_{n \rightarrow \infty} \frac{1}{4^n} \log \|F_t^n(z_0, 1)\|_v = \log |z_0|_v = \lambda(z_0). \quad \square$$

A similar application of Proposition 3.2 yields

PROPOSITION 3.4. *Suppose  $v \nmid 2$  is non-archimedean and  $|t|_v < 1$ . The escape-rate function  $G_{F_t,v}$  satisfies*

$$(3.3) \quad G_{F_t,v}(z, 1) = \begin{cases} \log |z|_v & \text{for } |z|_v \geq 1, \\ -\frac{\log^2 |z|_v}{2 \log |t|_v} + \log |z|_v & \text{for } |t|_v < |z|_v < 1, \\ \frac{1}{2} \log |t|_v & \text{for } |z|_v \leq |t|_v \end{cases}$$

for all  $z \in \mathbb{C}_v$ .

3.2. *Proof of Theorem 3.1 for  $v \nmid 2$ .* We compute the local energy  $E_v(t_1, t_2)$  by cases.

*Case (1):*  $|t_1|_v > 1$  and  $|t_2|_v < 1$ . Recall the local energy can be expressed as

$$2E_v(t_1, t_2) = \int_{\mathbb{P}_v^{1,an}} (\lambda_{t_1,v} - \lambda_{t_2,v}) d\mu_{t_2,v} + \int_{\mathbb{P}_v^{1,an}} (\lambda_{t_2,v} - \lambda_{t_1,v}) d\mu_{t_1,v}.$$

Therefore by [Proposition 3.2](#), [3.3](#) and [3.4](#),

$$\begin{aligned} 2E_v(t_1, t_2) &= \int_{\mathbb{P}_v^{1,an}} (\lambda_{t_1,v} - \lambda_{t_2,v}) d\mu_{t_2,v} + \int_{\mathbb{P}_v^{1,an}} (\lambda_{t_2,v} - \lambda_{t_1,v}) d\mu_{t_1,v} \\ &= \int_{\log|t_2|_v}^0 \left( \frac{\log|t_1|_v}{2} - \left( -\frac{x^2}{2\log|t_2|_v} + x \right) \right) \frac{dx}{-\log|t_2|_v} \\ &\quad + \int_0^{\log|t_1|_v} \left( x - \frac{1}{2} \left( \frac{x^2}{\log|t_1|_v} + \log|t_1|_v \right) \right) \frac{dx}{\log|t_1|_v} \\ &= \frac{\log|t_1/t_2|_v}{3}. \end{aligned}$$

*Case (2):*  $|t_1|_v > 1$  and  $|t_2|_v > 1$ . Without loss of generality, we assume that  $|t_1|_v = \max\{|t_1|_v, |t_2|_v\}$ . By [Proposition 3.2](#) and [3.3](#),

$$\begin{aligned} 2E_v(t_1, t_2) &= \int_{\mathbb{P}_v^{1,an}} (\lambda_{t_1,v} - \lambda_{t_2,v}) d\mu_{t_2,v} + \int_{\mathbb{P}_v^{1,an}} (\lambda_{t_2,v} - \lambda_{t_1,v}) d\mu_{t_1,v} \\ &= \int_0^{\log|t_2|_v} \left( \frac{1}{2} \left( \frac{x^2}{\log|t_1|_v} + \log|t_1|_v \right) - \frac{1}{2} \left( \frac{x^2}{\log|t_2|_v} + \log|t_2|_v \right) \right) \frac{dx}{\log|t_2|_v} \\ &\quad + \int_0^{\log|t_2|_v} \left( \frac{1}{2} \left( \frac{x^2}{\log|t_2|_v} + \log|t_2|_v \right) - \frac{1}{2} \left( \frac{x^2}{\log|t_1|_v} + \log|t_1|_v \right) \right) \frac{dx}{\log|t_1|_v} \\ &\quad + \int_{\log|t_2|_v}^{\log|t_1|_v} \left( x - \frac{1}{2} \left( \frac{x^2}{\log|t_1|_v} + \log|t_1|_v \right) \right) \frac{dx}{\log|t_1|_v} \\ &= \frac{\log^2|t_1/t_2|_v}{3\log\max\{|t_1|_v, |t_2|_v\}}. \end{aligned}$$

*Case (3):*  $|t_2(t_2-1)|_v = 1$  and  $|t_1-1|_v \geq 1$ . In this case,  $f_{t_2}$  has good reduction, so  $\mu_{t_2,v}$  is a point mass supported on the Gauss point  $\zeta_{0,1}$ . Hence

$$2E_v(t_1, t_2) = \frac{|\log|t_1|_v|}{3} = \frac{|\log|t_1/t_2|_v|}{3}.$$

*Case (4):* The remaining cases reduce to the above three by the symmetry relations of [Proposition 2.3](#). This completes the proof of [Theorem 3.1](#) under the assumption that  $v \nmid 2$ .

### 3.3. Measure and escape rate for $v \mid 2$ .

**PROPOSITION 3.5.** *Suppose  $v \mid 2$  is non-archimedean. The canonical measure  $\mu_{t,v}$  on  $\mathbb{P}_v^{1,an}$  of  $f_t$  is the interval measure corresponding to the interval  $I$*

with

$$I = \begin{cases} [\zeta_{0,|t/4|_v}, \zeta_{0,|4|_v}] & \text{for } |t|_v < |16|_v, \\ [\zeta_{0,|1/4|_v}, \zeta_{0,|4t|_v}] & \text{for } |t|_v > 1/|16|_v, \\ [\zeta_{1,|1-t|_v/|4|_v}, \zeta_{1,|4|_v}] & \text{for } |1-t|_v < |16|_v. \end{cases}$$

For  $|16|_v \leq |t|_v \leq 1/|16|_v$  with  $|1-t|_v \geq |16|_v$ ,  $f_t(z)$  has potential good reduction, and  $\mu_{t,v}$  is supported on a single point in  $\mathbb{H}$ .

*Proof.* We proceed as in the computations of [FRL10, §5.1], though the authors had assumed for simplicity that the residue characteristic of their field is not 2. If  $|t|_v > |1/16|_v$ , the interval  $[\zeta_{0,|1/4|_v}, \zeta_{0,|4t|_v}]$  is totally invariant by  $f_t$ , and

$$f_t(\zeta_{0,|4t|_v|16t|_v^{-r}}) = \zeta_{0,|4t|_v|16t|_v^{-2r}} \quad \text{and} \quad f_t(\zeta_{0,|4t|_v|16t|_v^{r-1}}) = \zeta_{0,|4t|_v|16t|_v^{-2r}}$$

for  $r \in [0, 1/2]$ . Thus  $\mu_{t,v}$  is the interval measure on  $[\zeta_{0,|1/4|_v}, \zeta_{0,|4t|_v}]$ . The cases  $|t|_v < |16|_v$  or  $|1-t|_v < |16|_v$  can then be deduced from [Proposition 2.1](#).

For all  $|16|_v \leq |t|_v \leq 1/|16|_v$  with  $|1-t|_v \geq |16|_v$ , we have  $|j(t)|_v \leq 1$ , so  $f_t$  has potential good reduction.  $\square$

Following the proofs of [Propositions 3.3](#) and [3.4](#), from [Proposition 3.5](#) we obtain

**PROPOSITION 3.6.** *Suppose  $v \mid 2$  is non-archimedean. We have*

$$(3.4) \quad G_{F_t,v}(z, 1) = \begin{cases} \log |z|_v & \text{for } |z|_v \geq |4t|_v, \\ \frac{1}{2} \left( \frac{\log^2 |4z|_v}{\log |16t|_v} + \log |t|_v \right) & \text{for } 1/|4|_v < |z|_v < |4t|_v, \\ \frac{1}{2} \log |t|_v & \text{for } |z|_v \leq 1/|4|_v \end{cases}$$

for  $t$  with  $|t|_v \geq 1/|16|_v$ , and

$$(3.5) \quad G_{F_t,v}(z, 1) = \begin{cases} \log |z|_v & \text{for } |z|_v \geq |4|_v, \\ \frac{1}{2} \left( \frac{\log^2 |4z/t|_v}{\log |16/t|_v} + \log |t|_v \right) & \text{for } |4|_v < |z|_v < |t/4|_v, \\ \frac{1}{2} \log |t|_v & \text{for } |z|_v \leq |t/4|_v \end{cases}$$

for  $t$  with  $|t|_v \leq |16|_v$ .

3.4. *Proof of Theorem 3.1 for  $v \mid 2$ .* We compute as in the case where  $v \nmid 2$ .

*Case (1):*  $\{t_1, t_2\}$  with  $\min\{|t_1|_v, |t_2|_v\} \geq 1/|16|_v$  and  $\max\{|t_1|_v, |t_2|_v\} > 1/|16|_v$ . [Proposition 3.6](#) yields

$$2E_v(t_1, t_2) = \frac{\log^2 |t_1/t_2|_v}{3 \log \max\{|16t_1|_v, |16t_2|_v\}} \geq \frac{\log^2 |t_1/t_2|_v}{3 \log \max\{|t_1|_v, |t_2|_v\}}.$$

*Case (2):*  $|t_1|_v \geq 1/|16|_v$  and  $|t_2|_v \leq |16|_v$ . Again by [Proposition 3.6](#),

$$2E_v(t_1, t_2) = \frac{\log |16t_1|_v - \log |t_2/16|_v}{3} - \log |16|_v \geq \frac{\log |t_1/t_2|_v}{3}.$$

*Case (3):*  $|t_1|_v > 1/|16|_v$ ,  $|16|_v \leq |t_2|_v \leq 1/|16|_v$  and  $|1 - t_2|_v \geq |16|_v$ . Let  $\zeta_{t_2} \in \mathbb{H}$  be the support of  $\mu_{t_2, v}$ . For any  $z \in \mathbb{C}_v$  with  $|z|_v > 1/|4|_v$ ,

$$|f_{t_2}(z)|_v = \frac{|(z^2 - t_2)^2|_v}{|4z(z-1)(z-t_2)|_v} > |z|_v.$$

Hence  $\zeta_{0,1/|4|_v} \in [\zeta_{t_2}, \infty)$ . Let  $z_0 \in \mathbb{C}_v$  with  $|z_0|_v > 1/|4|_v$ , and let  $(z_n, w_n) := F_{t_2}^n(z_0, 1)$ . From the recursive formula (3.2), inductively we have  $|z_n| = |z_0|_v^{4^n} > |w_n|_v/|4|_v$ . Consequently

$$\lambda_{t_2, v}(z) = G_{F_{t_2, v}}(z, 1) = \lim_{n \rightarrow \infty} \frac{\log \|F_{t_2}^n\|_v}{4^n} = \log |z|_v$$

for  $z$  with  $|z|_v > 1/|4|_v$ , and then  $\lambda_{t_2, v}(\zeta_{0, r}) = \log r$  for  $r \geq 1/|4|_v$ . Moreover, as  $\Delta \lambda_{t_2, v} = \delta_{\zeta_{t_2}} - \delta_\infty$ , the function  $\lambda_{t_2, v}$  is increasing at a constant rate along the ray  $[\zeta_{t_2}, \infty)$ , with respect to the hyperbolic metric. Therefore  $\lambda_{t_2, v}(\zeta_{t_2}) \leq \lambda_{t_2, v}(\zeta_{0,1/|4|_v}) = -\log |4|_v$ . Hence by [Propositions 3.5](#) and [3.6](#),

$$\begin{aligned} 2E_v(t_1, t_2) &= \int_{\mathbb{P}_v^{1, an}} (\lambda_{t_1, v} - \lambda_{t_2, v}) \, d\mu_{t_2, v} + \int_{\mathbb{P}_v^{1, an}} (\lambda_{t_2, v} - \lambda_{t_1, v}) \, d\mu_{t_1, v} \\ &= (\lambda_{t_1, v}(\zeta_{t_2}) - \lambda_{t_2, v}(\zeta_{t_2})) \\ &\quad + \int_{\log |1/4|_v}^{\log |4t_1|_v} \left( x - \frac{1}{2} \left( \frac{(x + \log |4|_v)^2}{\log |16t_1|_v} + \log |t_1|_v \right) \right) \frac{dx}{\log |16t_1|_v} \\ &\geq \frac{\log |16t_1|_v}{3}. \end{aligned}$$

Here we have used  $\lambda_{t_2, v}(\zeta_{t_2}) \leq -\log |4|_v$  and  $\lambda_{t_1, v}(\zeta_{t_2}) = \frac{1}{2} \log |t_1|_v$  for the last inequality.

Of course, for  $|16|_v \leq |t_i|_v \leq 1/|16|_v$  and  $|1 - t_i|_v \geq |16|_v$  for  $i = 1, 2$ , we have

$$2E_v(t_1, t_2) \geq 0 \geq \frac{|\log|t_1/t_2||_v}{3} + \frac{8}{3} \log|2|_v.$$

*Case (4):* The remaining cases reduce to the above three by the symmetry relations of [Proposition 2.3](#). This completes the proof of [Theorem 3.1](#).

#### 4. Archimedean places and the hybrid space

In this section, we provide some of the estimates we need to control the archimedean contributions to the height pairings. Throughout this section, we assume our parameter  $t \in \mathbb{C} \setminus \{0, 1\}$  is complex. We let  $\mu_t$  denote the probability measure on  $\mathbb{P}^1(\mathbb{C})$  that is the push-forward of the Haar measure on the Legendre elliptic curve  $E_t(\mathbb{C})$  via  $\pi(x, y) = x$ . This measure is also the unique measure of maximal entropy for the dynamical system defined by the Lattès map

$$f_t(z) = \frac{(z^2 - t)^2}{4z(z - 1)(z - t)},$$

as noted in [\[Mil06, §7\]](#). We study degenerations of the probability measures  $\mu_t$  and their potentials as  $t \rightarrow 0$ . (The cases of  $t \rightarrow 1$  and  $t \rightarrow \infty$  are similar.) To this end, we consider the action of  $f_t$  sending  $(t, z)$  to  $(t, f_t(z))$  on the complex surface  $X = \mathbb{D}^* \times \mathbb{P}^1(\mathbb{C})$ , where  $\mathbb{D}^*$  is the punctured unit disk. We make use of the hybrid space  $X^{\text{hyb}}$ , in which the Berkovich projective line over the field of formal Laurent series  $\mathbb{C}((t))$  creates a central fiber of  $X$  over  $t = 0$  in the unit disk  $\mathbb{D}$ . We appeal to the topological description of the hybrid space from [\[BJ17\]](#) and the associated dynamical degenerations described in [\[Fav16\]](#).

**4.1. The family of Lattès maps and their escape rates.** In homogeneous coordinates on  $\mathbb{C}^2$ , recall that the maps  $f_t$  may be presented as

$$F_t(z, w) := ((z^2 - tw^2)^2, 4zw(z - w)(z - tw))$$

for  $t \in \mathbb{C} \setminus \{0, 1\}$ . They have escape-rate functions

$$(4.1) \quad G_{F_t}(z, w) := \lim_{t \rightarrow \infty} \frac{1}{4^n} \log \|F_t^n(z, w)\|,$$

as in [\(2.3\)](#).

View the families  $f_t$  and  $F_t$  as maps  $f = f_T$  and  $F = F_T$  defined over the field  $k = \mathbb{C}(T)$ , and consider the non-archimedean absolute value  $|\cdot|_0$  on  $k$  satisfying  $|g(T)|_0 = e^{-\text{ord}_0 g}$ . Let  $k_0 = \mathbb{C}((T))$  denote the completion of  $\mathbb{C}(T)$  with respect to this absolute value. Let  $\mathbb{L}$  denote a (minimal) complete and algebraically closed field containing  $k_0$ . The non-archimedean escape rate  $\hat{G}_F$

on  $\mathbb{L}^2$  is defined as in (2.3). Since  $|T|_0 < 1$ , it is given for  $x \in \mathbb{L}$  by the following formula, exactly as in [Proposition 3.4](#):

$$(4.2) \quad \begin{aligned} \hat{g}_f(x) := \hat{G}_F(x, 1) &= \begin{cases} \log|x|_0 & \text{for } |x|_0 \geq 1, \\ \log|x|_0 - \frac{(\log|x|_0)^2}{2 \log|T|_0} & \text{for } |T|_0 < |x|_0 < 1, \\ \frac{1}{2} \log|T|_0 & \text{for } |x|_0 \leq |T|_0 \end{cases} \\ &= \begin{cases} -a & \text{for } |x|_0 = |T|_0^a \text{ with } a \leq 0 \\ -a + \frac{1}{2} a^2 & \text{for } |x|_0 = |T|_0^a \text{ with } 0 \leq a \leq 1 \\ -\frac{1}{2} & \text{for } |x|_0 = |T|_0^a \text{ with } a \geq 1. \end{cases} \end{aligned}$$

The function  $\hat{g}_f$  extends naturally to the Berkovich space  $\mathbb{P}_{\mathbb{L}}^{1,an}$ ; away from the point at  $\infty$ , it is a continuous potential for the equilibrium measure  $\hat{\mu}_f$  of  $f$ .

The potential  $\hat{g}_f$  and the measure  $\hat{\mu}_f$  are invariant under the action of  $\text{Gal}(\mathbb{L}/k_0)$  on  $\mathbb{P}_{\mathbb{L}}^{1,an}$ . They descend to define a function and probability measure — that we will also denote by  $\hat{g}_f$  and  $\hat{\mu}_f$  — on the quotient Berkovich line  $\mathbb{P}_{k_0}^{1,an}$ . (See [\[Ber90, §4.2\]](#) for details on this quotient map.) As computed in [Proposition 3.2](#), the measure  $\hat{\mu}_f$  is supported on the interval  $[\zeta_{0,|T|_0}, \zeta_{0,1}]$ , and it is uniform with respect the linear structure from the hyperbolic metric.

**4.2. Convergence of measures.** The family  $f_t$  acts on the product space  $\mathbb{D}^* \times \mathbb{P}^1$  sending  $(t, z)$  to  $(t, f_t(z))$ . It extends meromorphically to  $X_0 := \mathbb{D} \times \mathbb{P}^1$ , or indeed to any model complex surface  $X \rightarrow \mathbb{D}$  that is isomorphic to  $\mathbb{D}^* \times \mathbb{P}^1$  over  $\mathbb{D}^*$  and has a simple normal crossings divisor as its central fiber.

Fixing a surface  $X \rightarrow \mathbb{D}$  and letting  $t \rightarrow 0$ , the degeneration of the measures  $\mu_t$  of maximal entropy for  $f_t$  — or indeed for any meromorphic family of rational maps on  $\mathbb{P}^1$  — to the central fiber of  $X$  is now well understood. In [\[DMF14\]](#), [\[DF16\]](#), the limit of the measures  $\mu_t$  is computed for any choice of model  $X$ , and a relation is shown between these limits and the non-archimedean measure  $\hat{\mu}_f$ . In particular, if we define the annulus

$$A_t(a, b, C) := \{z \in \mathbb{C} : C^{-1}|t|^a \leq |z| \leq C|t|^b\}$$

for  $t \in \mathbb{D}^*$ ,  $C > 1$ , and real numbers  $a \geq b$ , then

$$(4.3) \quad \mu_t(A_t(a, b, C)) \rightarrow \hat{\mu}_f([\zeta_{0,|T|_0^a}, \zeta_{0,|T|_0^b}]) = \text{length}_{\mathbb{R}}([0, 1] \cap [b, a])$$

as  $t \rightarrow 0$ . This follows from [\[DMF14, Th. B\]](#) (allowing for changes of coordinates on  $\mathbb{P}^1$  and base changes, passing to covers of the punctured disk  $\mathbb{D}^*$ ) or from the computations described in [\[DF16, Th. D\]](#) (taking  $\Gamma$  to be a vertex set

in the interval  $[\zeta_{0,1}, \zeta_{0,|T|_0}]$ ). Another proof is described below in [Section 4.3](#). In particular, this convergence implies

LEMMA 4.1. *Given any  $\varepsilon > 0$  and integer  $n \geq 1$ , there exists  $\delta > 0$  such that*

$$\frac{1}{n} - \varepsilon < \mu_t(\{|t|^{(i+1)/n} \leq |z| \leq |t|^{i/n}\}) < \frac{1}{n} + \varepsilon$$

for all  $0 < |t| < \delta$  and  $i = 0, \dots, n-1$ .

Taking  $\varepsilon = 1/n^2$  in [Lemma 4.1](#), we observe that for any given  $n$ , there is a  $\delta > 0$  such that we also have

$$(4.4) \quad \mu_t\left(\{|z| \geq 1\} \cup \{|z| \leq |t|\}\right) < \frac{1}{n}$$

for all  $0 < |t| < \delta$ .

4.3. *Convergence in the hybrid space.* In [\[Fav16\]](#), Favre gives an alternate proof of [\(4.3\)](#) by showing that

$$(4.5) \quad \mu_t \rightarrow \hat{\mu}_f$$

weakly in the hybrid space  $X^{\text{hyb}}$  [\[Fav16, Th. B\]](#). The hybrid space consists of replacing the central fiber in the models  $X$  above with the Berkovich line  $\mathbb{P}_{k_0}^{1,an}$ , carrying an appropriate topology. The convergence of measures follows from the convergence of their potentials to the potential of the measure  $\hat{\mu}_f$  in the Berkovich line. We describe this convergence here, as we will use it for proving our main result.

Let  $m_1$  denote the Lebesgue measure on the unit circle in  $\mathbb{C}$ , normalized to have total length 1. Let  $\Phi_t(z)$  denote a continuous potential on  $\mathbb{P}^1(\mathbb{C})$  for the measure  $\mu_t - m_1$ . Explicitly, in local coordinates  $z \in \mathbb{C} \subset \mathbb{P}^1$ , we can take

$$(4.6) \quad \Phi_t(z) = G_{F_t}(z, 1) - \log^+ |z|$$

with  $G_{F_t}$  as in [\(4.1\)](#). In [\[Fav16\]](#), Favre proves that the function

$$(4.7) \quad \varphi(t, z) := \frac{\Phi_t(z)}{\log |t|^{-1}}$$

extends to define a continuous function on  $X^{\text{hyb}}$ , taking the values of a potential of the limiting measure  $\hat{\mu}_f - \omega_0$  on the central fiber. Here  $\omega_0$  is the delta mass on the Gauss point  $\zeta_{0,1}$  of the Berkovich line  $\mathbb{P}_{k_0}^{1,an}$ . More precisely, we consider

the function

$$\begin{aligned}
 \hat{\varphi}_f(x) &:= \begin{cases} 0 & \text{for } |x|_0 \geq 1, \\ \log |x|_0 - \frac{(\log |x|_0)^2}{2 \log |T|_0} & \text{for } |T|_0 < |x|_0 < 1, \\ \frac{1}{2} \log |T|_0 & \text{for } |x|_0 \leq |T|_0 \end{cases} \\
 (4.8) \quad &= \begin{cases} 0 & \text{for } |x|_0 = |T|_0^a \text{ with } a \leq 0, \\ -a + \frac{1}{2} a^2 & \text{for } |x|_0 = |T|_0^a \text{ with } 0 \leq a \leq 1, \\ -\frac{1}{2} & \text{for } |x|_0 = |T|_0^a \text{ with } a \geq 1 \end{cases}
 \end{aligned}$$

for  $x \in \mathbb{L}$ , similar to the formula for  $\hat{g}_f$  in (4.2). This function  $\hat{\varphi}_f$  extends continuously to all of  $\mathbb{P}_{\mathbb{L}}^{1,an}$ ; it is Galois invariant over  $k_0$ ; and it descends to the quotient  $\mathbb{P}_{k_0}^{1,an}$ . Favre's theorem implies that the function  $\varphi$  of (4.7) extends continuously to  $X^{\text{hyb}}$ , coinciding with  $\hat{\varphi}_f$  over  $t = 0$ :

PROPOSITION 4.2. *Given any  $\varepsilon > 0$ , there exists  $\delta > 0$  such that*

$$|\varphi(t, z) - \hat{\varphi}_f(\zeta_{0,|T|_0^a})| < \varepsilon$$

for all  $0 < |t| < \delta$ , for all  $a \in \mathbb{R}$ , and all  $z$  for which

$$\left| \frac{\log |z|}{\log |t|} - a \right| < \delta.$$

*Proof.* Recall that the absolute value  $|\cdot|_0$  on  $\mathbb{L}$  induces a continuous function on the Berkovich space that we will also denote by  $|\cdot|_0 : \mathbb{P}_{k_0}^{1,an} \rightarrow \mathbb{R}_{\geq 0} \cup \{\infty\}$ . We use the standard absolute value  $|\cdot|$  on  $\mathbb{C}$ , extended to a continuous function  $\mathbb{P}^1(\mathbb{C}) \rightarrow \mathbb{R}_{\geq 0} \cup \{\infty\}$ .

The topology on  $\mathbb{P}_{k_0}^{1,an}$  is such that annuli of the form

$$A(r_1, r_2) := \{x \in \mathbb{P}_{k_0}^{1,an} : r_1 < |x|_0 < r_2\}$$

are open for any choice of  $0 \leq r_1 < r_2 \leq \infty$ , as are the Berkovich disks of the form

$$D_0(r) := \{x \in \mathbb{P}_{k_0}^{1,an} : |x|_0 < r\} \text{ and } D_{\infty}(r) := \{x \in \mathbb{P}_{k_0}^{1,an} : |x|_0 > r\}$$

for any  $0 < r < \infty$ . The topology on  $X^{\text{hyb}}$  is such that an annular set of the form

$$\{(t, z) \in \mathbb{D}^* \times \mathbb{P}^1(\mathbb{C}) : |t|^{a+\delta} < |z| < |t|^{a-\delta} \text{ and } 0 < |t| < \delta\} \cup A(|T|_0^{a+\delta}, |T|_0^{a-\delta})$$

is an open neighborhood of  $\zeta_{0,|T|_0^a}$  on the central fiber for any  $a$  and any  $\delta > 0$ . Similarly, the disk-like sets

$$\{(t, z) \in \mathbb{D}^* \times \mathbb{P}^1(\mathbb{C}) : |z| < |t|^a \text{ and } 0 < |t| < \delta\} \cup D_0(|T|_0^a)$$

and

$$\{(t, z) \in \mathbb{D}^* \times \mathbb{P}^1(\mathbb{C}) : |z| > |t|^a \text{ and } 0 < |t| < \delta\} \cup D_\infty(|T|_0^a)$$

are open for any  $a \in \mathbb{R}$ , and allowing  $a$  and  $\delta$  to vary provides open neighborhoods at 0 and  $\infty$  respectively in the central fiber. See [BJ17, §2.2 and Def. 4.9] for details on the hybrid topology. Note in particular that the hybrid topology restricted to the central fiber induces the usual (weak) Berkovich topology.

By the continuity statement of [Fav16, Th. 2.10] and exhibiting  $\varphi$  as a uniform limit of model functions ([Fav16, §4.3] provides the details in the dynamical case) the function  $\varphi$  extends to define a continuous function on  $X^{\text{hyb}}$ , taking the values of  $\hat{\varphi}_f$  on the central fiber. Let  $L$  denote the closed segment in  $\mathbb{P}_{k_0}^{1,an}$  between 0 and  $\infty$ . We may by compactness cover  $L$  by finitely many neighborhoods on which  $|\varphi(x) - \varphi(y)| \leq \epsilon$ . As the values of  $\hat{\varphi}_f$  depend only on the values of  $\varphi$  on  $L$ , each open neighborhood of a point in the interior of  $L$  contains an open interval in  $L$ , and  $\hat{\varphi}$  is constant near 0 and  $\infty$ , we may assume these neighborhoods are annular or disk-like as defined above. Thus we obtain a uniform  $\delta$  as claimed.  $\square$

As  $\Phi_t(z) = G_{F_t}(z, 1) - \log^+ |z|$ , we also have a uniform continuity statement for  $G$  when  $|z|$  is bounded from above:

**PROPOSITION 4.3.** *Given any  $\epsilon > 0$  and  $M > 1$ , there exists  $\delta > 0$  such that*

$$\left| \frac{G_{F_t}(z, 1)}{\log |t|^{-1}} - \hat{g}_f(\zeta_{0,|T|_0^a}) \right| < \epsilon$$

for all  $0 < |t| < \delta$ , for all  $a \in \mathbb{R}$ , and all  $|z| \leq M$  for which

$$\left| \frac{\log |z|}{\log |t|} - a \right| < \delta.$$

**4.4. Discrete measures and regularizations.** Let  $F$  be any finite set in  $\mathbb{C}$ . Denote by  $[F]$  the probability measure supported equally on the elements of  $F$ , and for  $r > 0$ , denote by  $[F]_r$  the probability measure supported equally and uniformly on circles of radius  $r$  about each element of  $F$ .

**PROPOSITION 4.4.** *For every  $\epsilon > 0$ , there exists  $c = c(\epsilon) > 0$  such that*

$$|(\mu_t, [F]) - (\mu_t, [F]_r)| < \epsilon \max\{\log |t|^{-1}, \log |t - 1|^{-1}, \log |t|, 1\}$$

for all  $t \in \mathbb{C} \setminus \{0, 1\}$  and any finite set  $F$  in  $\mathbb{C}$  and any

$$r \leq c \min\{|t|^2, |t - 1|^2, |t|^{-2}\}.$$

*Proof.* For any  $x \in \mathbb{C}$  and any  $r > 0$ , let  $m_{x,r}$  be the probability measure supported on the circle of radius  $r$  around  $x$ . Recall that

$$(\rho, \sigma) := - \iint_{\mathbb{C} \times \mathbb{C} \setminus \Delta} \log |z - w| d\rho(z) d\sigma(w).$$

For each fixed  $t$ , the function  $G_{F_t}(\cdot, 1)$  is a potential for  $\mu_t$  in  $\mathbb{C}$ , and therefore, there exists a constant  $C_t$  such that

$$\int_{\mathbb{C}} \log |z - w| d\mu_t(z) = G_{F_t}(w, 1) + C_t.$$

Now let  $F$  be any finite set in  $\mathbb{C}$ . Then, assuming  $r < 1$ , we have

$$\begin{aligned} (\mu_t, [F]_r) - (\mu_t, [F]) &= \frac{1}{|F|} \sum_{x \in F} \left( G_{F_t}(x, 1) - \int G_{F_t}(\zeta, 1) dm_{x,r}(\zeta) \right) \\ &= \frac{1}{|F|} \sum_{\{x \in F : |x| > 2\}} \left( \Phi_t(x) - \int \Phi_t(\zeta) dm_{x,r}(\zeta) \right) \\ &\quad + \frac{1}{|F|} \sum_{\{x \in F : |x| \leq 2\}} \left( G_{F_t}(x, 1) - \int G_{F_t}(\zeta, 1) dm_{x,r}(\zeta) \right) \end{aligned}$$

because the function  $\log^+ |z|$  is harmonic away from the unit circle on  $\mathbb{C}$ .

By [Proposition 4.3](#), there exists  $\delta > 0$  such that

$$(4.9) \quad \left| \frac{G_{F_t}(z, 1)}{\log |t|^{-1}} - \hat{g}_f(\zeta_{0,|T|^a_0}) \right| < \varepsilon/2$$

for all  $|z| \leq 2$  satisfying

$$\left| \frac{\log |z|}{\log |t|} - a \right| < \delta$$

and all  $|t| < \delta$  and any  $a \in \mathbb{R}$ . Shrinking  $\delta$  if needed, we have

$$(4.10) \quad |\varphi(t, z)| < \varepsilon/2$$

for  $|z| \geq 1$  and all  $|t| < \delta$ , by [Proposition 4.2](#).

Let  $C_\delta$  be the compact subset of  $\mathbb{C} \setminus \{0, 1\}$  consisting of all  $t$  with  $|t| \geq \delta$  and  $|t - 1| \geq \delta$  and  $|1/t| \geq \delta$ . Over  $C_\delta \times \mathbb{P}^1$ , the family of potentials  $\{\Phi_t\}$  is uniformly continuous. So there exists  $c_1 = c_1(\delta)$  such that

$$|\Phi_t(z) - \Phi_t(z')| < \varepsilon$$

whenever  $\text{dist}(z, z') < c_1$  and for all  $t \in C_\delta$ . Here,  $\text{dist}$  represents the chordal distance on  $\mathbb{P}^1$ . Furthermore, we may take  $c_1$  such that we also have

$$|G_{F_t}(z, 1) - G_{F_t}(z', 1)| < \varepsilon$$

for all  $|z - z'| < c_1$  with  $|z| \leq 2$ , and all  $t \in C_\delta$ . Thus

$$|(\mu_t, [F]_r) - (\mu_t, [F])| < \varepsilon$$

for any choice of finite set  $F$ ,  $t \in C_\delta$ , and  $r < c_1$ .

Now assume that  $|t| < \delta$ . We will consider three cases. First, suppose  $|t|^{1+\delta} \leq |z| \leq 2$ . Choose any  $c_2 = c_2(\delta)$  such that

$$(4.11) \quad \left| \log(1 \pm c_2 \delta^{1-\delta}) \right| < \delta(\log \delta^{-1}).$$

Then

$$\begin{aligned}
|\log |z'/z|| &= \left| \log \left| \frac{z' - z}{z} + 1 \right| \right| \\
&\leq \max \left\{ \log \left( 1 + \left| \frac{z' - z}{z} \right| \right), \left| \log \left( 1 - \left| \frac{z' - z}{z} \right| \right) \right| \right\} \\
&\leq \max \left| \log \left( 1 \pm \frac{c_2 |t|^2}{|z|} \right) \right| \\
&\leq \max \left| \log \left( 1 \pm c_2 \delta^{1-\delta} \right) \right| \\
&< \delta (\log \delta^{-1}) \leq \delta \log |t|^{-1}
\end{aligned}$$

for all  $|t| < \delta$ . This is equivalent to

$$(4.12) \quad \left| \frac{\log |z|}{\log |t|} - \frac{\log |z'|}{\log |t|} \right| < \delta$$

for all  $|z - z'| < c_2 |t|^2$  with  $|t|^{1+\delta} \leq |z| \leq 2$ . Combined with (4.9) and setting  $a = (\log |z|)/(\log |t|)$ , this implies that

$$|G_{F_t}(z, 1) - G_{F_t}(z', 1)| < \varepsilon \log |t|^{-1}$$

for such pairs  $z$  and  $z'$ .

Second, suppose that  $|z| \leq |t|^{1+\delta}$ . By shrinking  $c_2$  further if necessary, we have  $c_2 < (1 - \delta^\delta)/\delta$ , and therefore if  $|z| \leq |t|^{1+\delta}$  and  $|z - z'| < c_2 |t|^2$ , with  $|t| < \delta$ , we also have  $|z'| \leq |t|$ . Applying the convergence (4.9) where  $\hat{g}_f = -1/2$ , for all  $|z| \leq |t|^{1+\delta}$  we have

$$|G_{F_t}(z, 1) - G_{F_t}(z', 1)| < \varepsilon \log |t|^{-1}$$

for  $z'$  satisfying  $|z - z'| < c_2 |t|^2$  and for all  $|t| < \delta$ .

Third, for  $|z| \geq 2$ , by the convergence (4.10),

$$|\Phi_t(z) - \Phi_t(z')| < \varepsilon \log |t|^{-1}$$

for all  $|z| \geq 2$  and  $|z - z'| < c_2 |t|^2$  and  $|t| < \delta$ .

Together these three cases yield

$$|(\mu_t, [F]_r) - (\mu_t, [F])| < \varepsilon \log |t|^{-1}$$

for any choice of finite set  $F$  and all  $|t| < \delta$ , with  $r < c_2 |t|^2$ .

If  $|t - 1| < \delta$ , the arguments above go through by replacing  $z$  with  $1 - z$ , as

$$G_{F_{1-t}}(1 - z, 1) = G_{F_t}(z, 1)$$

by Proposition 2.1. It follows that

$$|(\mu_t, [F]_r) - (\mu_t, [F])| < \varepsilon \log |t - 1|^{-1}$$

for any choice of finite set  $F$ ,  $|t - 1| < \delta$ , and  $r < c_2 |t - 1|^2$ , with  $\delta$  and  $c_2$  as above.

For  $t$  near  $\infty$ , more care is needed, as

$$G_{F_t}(z, 1) = G_{F_{1/t}}(z, t) = G_{F_{1/t}}(z/t, 1) + \log|t|$$

by [Proposition 2.1](#). Setting  $s = 1/t$ ,

$$\frac{G_{F_t}(z, 1)}{\log|t|} = \frac{G_{F_s}(s z, 1) - \log|s|}{\log|s|^{-1}} = \frac{G_{F_s}(s z, 1)}{\log|s|^{-1}} + 1.$$

From [\(4.9\)](#), we have

$$\left| \frac{G_{F_s}(s z, 1)}{\log|s|^{-1}} - \hat{g}_f(\zeta_{0,|T|_0^{1-a}}) \right| < \varepsilon/2$$

for  $|s z| \leq 2$ ,  $|s| < \delta$ , and

$$\left| \frac{\log|s z|}{\log|s|} - (1-a) \right| < \delta$$

for any choice of  $a \in \mathbb{R}$ . Thus,

$$(4.13) \quad \left| \frac{G_{F_t}(z, 1)}{\log|t|} - \left( \hat{g}_f(\zeta_{0,|T|_0^{1-a}}) + 1 \right) \right| < \varepsilon/2$$

for all  $|z| \leq 2|t|$  satisfying

$$\left| \frac{\log|z|}{\log|t|} - a \right| < \delta$$

with  $|t| > \delta^{-1}$  and any  $a \in \mathbb{R}$ . As in [\(4.10\)](#), we also have

$$|\varphi(t, z)| < \varepsilon/2$$

for  $|z| \geq |t|$  and  $|t| > \delta^{-1}$ , because  $\hat{g}_f(\zeta_{0,|T|_0^{1-a}}) + 1 = a$  for all  $a \geq 1$ , from the formula given in [\(4.2\)](#). The choice of  $c_2$  in [\(4.11\)](#) is similar. It follows that

$$|(\mu_t, [F]_r) - (\mu_t, [F])| < \varepsilon \log|t|$$

for any choice of finite set  $F$  and all  $|t| > 1/\delta$ , with  $r < c_2 |t|^{-2}$ .

Let  $c := \min\{c_1, c_2\}$  to complete the proof.  $\square$

## 5. Archimedean energy

As in [Section 4](#), assume  $t \in \mathbb{C} \setminus \{0, 1\}$  is a complex parameter, with  $\mu_t$  on  $\mathbb{P}^1(\mathbb{C})$  the push-forward of the Haar measure on  $E_t(\mathbb{C})$ , and  $\lambda_t(z) = G_{F_t}(z, 1)$  a potential for  $\mu_t - \delta_\infty$  on  $\mathbb{P}^1(\mathbb{C})$ . In this section we provide estimates on the archimedean local energy (introduced in [Proposition 2.2](#))

$$E_\infty(s, t) := \frac{1}{2} \left( \int (\lambda_s - \lambda_t) d\mu_t + \int (\lambda_t - \lambda_s) d\mu_s \right)$$

for  $s, t \in \mathbb{C} \setminus \{0, 1\}$  as one or both of the parameters tends to 0, 1, or  $\infty$ . We treat three cases separately: where only one parameter escapes into a cusp, where both parameters escape into a cusp, and where the two parameters

head to two different cusps. By the symmetry established in [Proposition 2.3](#), we focus on the case where  $s$  tends to 0.

Throughout, we work in hybrid space and make use of the convergence of potentials to  $\hat{g}_f$  and measures to  $\hat{\mu}_f$  as  $t \rightarrow 0$ , as proved in [\[Fav16, Th. B\]](#) and explained in [Section 4.3](#).

### 5.1. A single escaping parameter.

**THEOREM 5.1.** *Given  $\varepsilon > 0$  and any compact set  $C \subset \mathbb{C} \setminus \{0, 1\}$ , there exists  $\delta > 0$  such that*

$$\left(\frac{1}{6} - \varepsilon\right) \log |s|^{-1} \leq E_\infty(s, t) \leq \left(\frac{1}{6} + \varepsilon\right) \log |s|^{-1}$$

for all  $s$  satisfying  $0 < |s| < \delta$  and all  $t \in C$ .

*Proof.* Recall that for any  $s \in \mathbb{C} \setminus \{0, 1\}$ , we have defined  $\Phi_s(z) = G_{F_s}(z, 1) - \log^+ |z|$  in [\(4.6\)](#) and  $\varphi(s, z) = \Phi_s(z) / (\log |s|^{-1})$ . For any pair  $s, t \in \mathbb{C} \setminus \{0, 1\}$ , the local energy  $E_\infty(s, t)$  satisfies

$$\begin{aligned} \frac{E_\infty(s, t)}{\log |s|^{-1}} &= \frac{1}{2 \log |s|^{-1}} \left( \int (\Phi_s - \Phi_t) d\mu_t + \int (\Phi_t - \Phi_s) d\mu_s \right) \\ &= \frac{1}{2} \left( \int \left( \varphi(s, z) - \frac{\Phi_t}{\log |s|^{-1}} \right) d\mu_t + \int \frac{\Phi_t}{\log |s|^{-1}} d\mu_s - \int \varphi(s, z) d\mu_s \right). \end{aligned}$$

Fix  $\varepsilon > 0$ , and suppose that  $C \subset \mathbb{C} \setminus \{0, 1\}$  is compact. The  $\Phi_t$  functions are uniformly bounded for all  $t \in C$  and all  $z \in \mathbb{P}^1(\mathbb{C})$ , so there is a  $\delta$  such that

$$\left| \int \frac{\Phi_t}{\log |s|^{-1}} d\mu_s \right| < \varepsilon$$

for all  $|s| < \delta$  and all  $t \in C$ . We can also find a small  $r = r(C)$  such that

$$\mu_t(\{|z| \leq r\}) < \varepsilon$$

for all  $t \in C$ . By [Proposition 4.2](#) (shrinking  $\delta$  if needed),

$$|\varphi(s, z)| < \varepsilon$$

for all  $|z| > r$  and  $|s| < \delta$ , and

$$|\varphi(s, z)| < 1$$

for all  $z$  and all  $|s| < \delta$ . Consequently,

$$\begin{aligned} \left| \int \left( \varphi(s, z) - \frac{\Phi_t}{\log |s|^{-1}} \right) d\mu_t \right| &\leq \int_{\{|z| \leq r\}} |\varphi(s, z)| d\mu_t + \int_{\{|z| > r\}} |\varphi(s, z)| d\mu_t \\ &\quad + \int \left| \frac{\Phi_t}{\log |s|^{-1}} \right| d\mu_t \\ &< 3\varepsilon. \end{aligned}$$

Finally, by the weak convergence of  $\mu_s \rightarrow \hat{\mu}_f$  and convergence of  $\varphi(s, z)$  to  $\hat{\varphi}_f$ , we can shrink  $\delta$  again such that

$$\left| \int \varphi(s, z) d\mu_s - \int \hat{\varphi}_f d\hat{\mu}_f \right| < \varepsilon$$

for all  $|s| < \delta$ . Recalling the formula for  $\hat{\varphi}_f$  from (4.8), we have

$$\int \hat{\varphi}_f d\hat{\mu}_f = \int_0^1 \left( -a + \frac{a^2}{2} \right) da = -\frac{1}{3},$$

since the measure  $\hat{\mu}_f$  is the uniform distribution on the interval  $[0, 1]$  in the  $a$  coordinates, as described in Section 4.1. Therefore,

$$\left( \frac{1}{6} - 4\varepsilon \right) \log |s|^{-1} \leq E_\infty(s, t) \leq \left( \frac{1}{6} + 4\varepsilon \right) \log |s|^{-1}$$

for all  $|s| < \delta$  and all  $t \in C$ . □

### 5.2. Both parameters escaping to the same cusp.

**THEOREM 5.2.** *Given  $\varepsilon > 0$ , there exists  $\delta > 0$  such that*

$$\left( \frac{1}{6} \left( 1 - \frac{1}{b} \right)^2 - \varepsilon \right) \log |s|^{-1} \leq E_\infty(s, t) \leq \left( \frac{1}{6} \left( 1 - \frac{1}{b} \right)^2 + \varepsilon \right) \log |s|^{-1}$$

for all  $s, t$  satisfying  $0 < |s| \leq |t| < \delta$ , where  $b = (\log |s|)/(\log |t|) \geq 1$ .

For each real number  $b \geq 1$ , consider the function

$$\begin{aligned} \hat{\varphi}_b(x) &:= \begin{cases} 0 & \text{for } |x|_0 \geq 1, \\ \log |x|_0 - \frac{(\log |x|_0)^2}{2b \log |T|_0} & \text{for } |T|_0^b \leq |x|_0 \leq 1, \\ \frac{b}{2} \log |T|_0 & \text{for } |x|_0 \leq |T|_0^b \end{cases} \\ &= \begin{cases} 0 & \text{for } |x|_0 = |T|_0^a \text{ with } a \leq 0, \\ -a + a^2/(2b) & \text{for } |x|_0 = |T|_0^a \text{ with } 0 \leq a \leq b, \\ -b/2 & \text{for } |x|_0 = |T|_0^a \text{ with } a \geq b \end{cases} \end{aligned}$$

for all  $x \in \mathbb{L}$ . Note that  $\hat{\varphi}_1 = \hat{\varphi}_f$  from (4.8). As with  $\hat{\varphi}_1$ , each  $\hat{\varphi}_b$  extends naturally to a function on the Berkovich projective line  $\mathbb{P}_{k_0}^{1,an}$  and is a potential of the measure  $\hat{\mu}_b - \delta_G$ , where  $\hat{\mu}_b$  is interval measure on  $[\zeta_{0,|T|_0^b}, \zeta_{0,1}]$  and  $\delta_G$  is the delta-mass at the Gauss point  $\zeta_{0,1}$ .

For each  $b \geq 1$ , the non-archimedean local energy  $E_0(\hat{\mu}_1, \hat{\mu}_b)$  is given by

$$\begin{aligned} E_0(\hat{\mu}_1, \hat{\mu}_b) &:= \frac{1}{2} \left( \int (\hat{\varphi}_1 - \hat{\varphi}_b) d\hat{\mu}_b + \int (\hat{\varphi}_b - \hat{\varphi}_1) d\hat{\mu}_1 \right) \\ &= \frac{(\log(|T|_0/|T|_0^b))^2}{-6 \log \min\{|T|_0, |T|_0^b\}} \\ &= \frac{(b-1)^2}{6b}, \end{aligned}$$

as computed in [Theorem 3.1](#) (in the case  $v \nmid 2$ ).

For  $s$  and  $t$  in  $\mathbb{C} \setminus \{0, 1\}$ , if both  $s$  and  $t$  are close to one of the three cusps, we can estimate the archimedean local energy  $E_\infty(s, t)$  in terms of the non-archimedean pairing using the degeneration description in hybrid space. We first prove a special case of [Theorem 5.2](#):

**PROPOSITION 5.3.** *Given  $\varepsilon > 0$  and  $B > 2$ , there exists  $\delta > 0$  such that*

$$(E_0(\hat{\mu}_1, \hat{\mu}_b) - \varepsilon) \log |t|^{-1} \leq E_\infty(s, t) \leq (E_0(\hat{\mu}_1, \hat{\mu}_b) + \varepsilon) \log |t|^{-1}$$

for all  $s, t$  satisfying  $0 < |t|^B \leq |s| \leq |t| < \delta$ , where  $b = (\log |s|)/(\log |t|) \leq B$ .

This proposition is an immediate consequence of the weak convergence of measures  $\mu_t \rightarrow \hat{\mu}_1$  in the hybrid space, and the convergence of potentials as described in [Section 4.3](#). We give the details to clarify how the bound  $b \leq B$  is used.

*Proof.* Fix  $\varepsilon > 0$  and  $B > 2$ .

For  $s$  and  $t$  in the punctured unit disk  $\mathbb{D}^*$ , and for any  $1 \leq b \leq B$ , consider

$$(5.1) \quad \varphi(t, z) := \frac{\Phi_{F_t}(z, 1)}{\log |t|^{-1}} \quad \text{and} \quad b\varphi(s, z) = \frac{b\Phi_{F_s}(z, 1)}{\log |s|^{-1}} = \frac{\Phi_{F_s}(z, 1)}{\log |t|^{-1}},$$

viewed as functions on the fiber  $\{t\} \times \mathbb{C}$  in the hybrid space. By [Proposition 4.2](#), there exists  $\delta_1 > 0$  such that

$$(5.2) \quad \left| \varphi(t, z) - \hat{\varphi}_1(\zeta_{0, |T|_0^a}) \right| < \varepsilon/(4B)$$

for all  $|t| < \delta_1$ ,  $a \geq 0$ , and all  $|t|^{a+\delta_1} < |z| < |t|^{a-\delta_1}$ . In particular,

$$|\varphi(t, z)| < \varepsilon/(4B)$$

for all  $|z| \geq 1$  and all  $|t| < \delta_1$ . It follows that

$$b\varphi(s, z) \rightarrow \hat{\varphi}_b(\zeta_{|T|_0^a})$$

in the hybrid space as  $s$  and  $t$  tend to 0 with  $|s| = |t|^b$  and  $(\log |z|)/(\log |t|) \rightarrow a$ , uniformly in  $b$  for  $1 \leq b \leq B$ . This is because the annulus

$$A_t(a, \delta) := \{z \in \mathbb{C} : |t|^{a+\delta} < |z| < |t|^{a-\delta}\}$$

for each fixed  $t \in \mathbb{D}^*$ ,  $\delta > 0$  and  $a \in \mathbb{R}$  can be written in terms of  $s$  as

$$A_t(a, \delta) = \{z \in \mathbb{C} : |s|^{a/b+\delta/b} < |z| < |s|^{a/b-\delta/b}\}$$

whenever  $|s| = |t|^b$ . Therefore,

$$(5.3) \quad \left| b\varphi(s, z) - \hat{\varphi}_b(\zeta_{0, |T|_0^a}) \right| = \left| \frac{b\Phi_{F_s}(z, 1)}{\log |s|^{-1}} - b\hat{\varphi}_1(\zeta_{0, |T|_0^{a/b}}) \right| < b\varepsilon/(4B) < \varepsilon/4$$

for all  $z \in A_t(a, \delta_1)$ , as a consequence of (5.2). In particular,

$$|b\varphi(s, z)| < \varepsilon/4$$

for all  $|z| \geq 1$  and all  $|t| < \delta_1$ .

Recall that the measures  $\mu_t$  on the fiber over  $t$  converge weakly in  $X^{\text{hyb}}$  to the measure  $\hat{\mu}_1$  on the central fiber. For each  $s$  with  $|s| = |t|^b$ , let  $\mu_t^s$  denote the measure associated to  $f_s$  but viewed in the fiber  $\{t\} \times \mathbb{P}^1$ . The measures  $\mu_t^s$  converge to the measure  $\hat{\mu}_b$  as  $t \rightarrow 0$  with  $|s| = |t|^b$ , and this convergence can also be made uniform in  $b$  with  $b \leq B$ . That is, by Lemma 4.1, for any  $n$  there exists  $\delta_2 > 0$  such that

$$(5.4) \quad \frac{1}{n} - \frac{1}{n^2} < \mu_t(\{|t|^{(i+1)/n} \leq |z| \leq |t|^{i/n}\}) < \frac{1}{n} + \frac{1}{n^2}$$

for all  $|t| < \delta_2$  and each  $i = 0, \dots, n$ . Note that this implies that

$$\mu_t(\{|z| \leq |t|\} \cup \{|z| \geq 1\}) < \frac{1}{n}.$$

Therefore, we also have

$$(5.5) \quad \frac{1}{n} - \frac{1}{n^2} < \mu_t^s(\{|t|^{b(i+1)/n} \leq |z| \leq |t|^{bi/n}\}) < \frac{1}{n} + \frac{1}{n^2}$$

and

$$\mu_t^s(\{|z| \leq |t|^b\} \cup \{|z| \geq 1\}) < \frac{1}{n}$$

for all  $|t| < \delta_2$ . Thus, the measure  $\hat{\mu}_b$  on small sub-annuli of the annulus  $\{|t|^b \leq |z| \leq 1\}$  is controlled uniformly for all  $1 \leq b \leq B$ .

Putting all the pieces together,

$$\frac{E_\infty(s, t)}{\log |t|^{-1}} = \frac{1}{2} \left( \int (b\varphi(s, z) - \varphi(t, z)) \, d\mu_t + \int (\varphi(t, z) - b\varphi(s, z)) \, d\mu_t^s \right)$$

is within  $\varepsilon$  of

$$E_0(\hat{\mu}_1, \hat{\mu}_b) = \frac{1}{2} \left( \int (\hat{\varphi}_b - \hat{\varphi}_1) \, d\hat{\mu}_1 + \int (\hat{\varphi}_1 - \hat{\varphi}_b) \, d\hat{\mu}_b \right) = \frac{(b-1)^2}{6b}$$

for all  $t$  sufficiently small and all  $s$  with  $|s| = |t|^b$  for any  $1 \leq b \leq B$ .  $\square$

Here is an equivalent restatement of Theorem 5.2, expressed in terms of the growth of  $|t|$ :

THEOREM 5.4. *Given  $\varepsilon > 0$ , there exists  $\delta > 0$  such that*

$$\left( \frac{(b-1)^2}{6b} - b\varepsilon \right) \log |t|^{-1} \leq E_\infty(s, t) \leq \left( \frac{(b-1)^2}{6b} + b\varepsilon \right) \log |t|^{-1}$$

for all  $s, t$  satisfying  $0 < |s| \leq |t| < \delta$ , where  $b = (\log |s|)/(\log |t|) \geq 1$ .

Comparing [Theorem 5.4](#) to the statement of [Proposition 5.3](#), we see that we lose the ability to bound the energy within a uniform  $\varepsilon$  when  $b$  becomes large.

*Proof of Theorems 5.4 and 5.2.* Fix  $\varepsilon > 0$ .

As in the proof of [Proposition 5.3](#), we make use of the weak convergence of measures  $\mu_t \rightarrow \hat{\mu}_1$  and convergence of the potentials  $\varphi(t, z) \rightarrow \hat{\varphi}_1$  in the hybrid space as  $t \rightarrow 0$ . Recalling the formula for  $\hat{\varphi}_f$  from (4.8), we have

$$\int \hat{\varphi}_f d\hat{\mu}_f = \int_0^1 \left( -a + \frac{a^2}{2} \right) da = -\frac{1}{3},$$

since the measure  $\hat{\mu}_f$  is the uniform distribution on the interval  $[0, 1]$  in the  $a$  coordinates, as described in [Section 4.1](#).

Choose  $r$  satisfying  $0 < r < \varepsilon/100$ . There is a  $\delta_2$  such that

$$\begin{aligned} \mu_t(\{|z| \leq |t|\}) &< \varepsilon/50, \\ |\varphi(t, z)| &< \varepsilon/50 \text{ for } |z| \geq |t|^r, \end{aligned}$$

and

$$|\varphi(t, z)| < \frac{1}{2} + \varepsilon/50 \text{ for all } z$$

for all  $|t| < \delta_2$ . Thus, for  $s \in \mathbb{C}^*$  with  $|s| = |t|^b$  and  $b > 1/r$ , we have

$$|b\varphi(s, z)| < b\varepsilon/50 \text{ for } |z| \geq |t|$$

and

$$|b\varphi(s, z)| < \frac{b}{2} + b\varepsilon/50 \text{ for all } z$$

for all  $|t| < \delta_2$ . By shrinking  $\delta_2$  further if necessary, we appeal to the weak convergence of measures  $\mu_t \rightarrow \hat{\mu}_1$  in the hybrid space to deduce that

$$(5.6) \quad \left| \int \varphi(t, z) d\mu_t + \frac{1}{3} \right| = \left| \int \varphi(t, z) d\mu_t - \int \hat{\varphi}_1 d\hat{\mu}_1 \right| < \varepsilon/10$$

for all  $|t| < \delta_2$ .

Now fix  $B > 1/r$ , and recall that  $r < \varepsilon/100$ , so that

$$(5.7) \quad \left| \frac{(b-1)^2}{6b} - \frac{b}{6} \right| < b\varepsilon/50$$

for all  $b \geq B$ . For this  $B$ , we can find a  $\delta_B > 0$  such that [Proposition 5.3](#) is satisfied for all  $0 < |s| = |t|^b \leq |t| < \delta_B$  with  $1 \leq b \leq B$ . Choose any  $\delta \leq \min\{\delta_B, \delta_2\}$ , and we obtain the theorem for  $b \leq B$ .

Now suppose  $b \geq B$ . We will estimate

$$\frac{E_\infty(s, t)}{\log |t|^{-1}} = \frac{1}{2} \left( \int (b\varphi(s, z) - \varphi(t, z)) d\mu_t + \int (\varphi(t, z) - b\varphi(s, z)) d\mu_t^s \right)$$

for all  $|t| < \delta$  and any  $s$  with  $|s| = |t|^b$  by estimating the two integrals separately.

As shown above,

$$|b\varphi(s, z) - \varphi(t, z)| \leq 1/2 + \varepsilon/50 + b\varepsilon/50 < b\varepsilon/10$$

for all  $|z| \geq |t|$  and  $0 < |s| = |t|^b \leq |t| < \delta$  with  $b \geq B$ , and

$$|b\varphi(s, z) - \varphi(t, z)| \leq 1/2 + \varepsilon/50 + b/2 + b\varepsilon/50$$

for all  $z$  and  $0 < |s| = |t|^b \leq |t| < \delta$ . Writing the first integral as

$$\begin{aligned} \int (b\varphi(s, z) - \varphi(t, z)) d\mu_t &= \int_{|z| \geq |t|} (b\varphi(s, z) - \varphi(t, z)) d\mu_t \\ &\quad + \int_{|z| \leq |t|} (b\varphi(s, z) - \varphi(t, z)) d\mu_t, \end{aligned}$$

it follows that

$$\left| \int (b\varphi(s, z) - \varphi(t, z)) d\mu_t \right| \leq b\varepsilon/10 + (1/2 + \varepsilon/50 + b/2 + b\varepsilon/50)(\varepsilon/50) < b\varepsilon/5$$

for all  $b \geq B$  and  $0 < |s| = |t|^b \leq |t| < \delta$ .

Write the second integral as

$$\int (\varphi(t, z) - b\varphi(s, z)) d\mu_t^s = \int \varphi(t, z) d\mu_t^s - \int b\varphi(s, z) d\mu_t^s.$$

As  $|\varphi(t, z)|$  is bounded by  $1/2 + \varepsilon/50$ , we have

$$\left| \int \varphi(t, z) d\mu_t^s \right| \leq \frac{1}{2} + \varepsilon/50 < b\varepsilon/25$$

for all  $b \geq B$ . On the other hand, we have

$$\int b\varphi(s, z) d\mu_t^s = b \int \varphi(s, z) d\mu_s$$

so that

$$\left| \int b\varphi(s, z) d\mu_t^s + \frac{b}{3} \right| < b\varepsilon/10$$

for all  $0 < |s| = |t|^b \leq |t| < \delta$  from (5.6).

We conclude that

$$\left| \frac{E_\infty(s, t)}{\log |t|^{-1}} - \frac{b}{6} \right| < b\varepsilon/2$$

for all  $b$  sufficiently large and all  $|t| < \delta$ . On the other hand, we also have

$$\left| \frac{b}{6} - \frac{(b-1)^2}{6b} \right| < b\varepsilon/50$$

for all  $b \geq B$  by our choice of  $B$ , so the theorem is proved.  $\square$

### 5.3. Parameters escaping to different cusps.

THEOREM 5.5. *Given  $\varepsilon > 0$ , there exists  $\delta > 0$  such that*

$$\left(\frac{1}{6}\left(1 + \frac{1}{b}\right) - \varepsilon\right) \log |s|^{-1} \leq E_\infty(s, t) \leq \left(\frac{1}{6}\left(1 + \frac{1}{b}\right) + \varepsilon\right) \log |s|^{-1}$$

for all  $s, t \in \mathbb{C}$  satisfying  $|t| > 1/\delta$  and  $0 < |s| \leq 1/|t|$ , where

$$b = -(\log |s|)/(\log |t|).$$

*Proof.* The proof is nearly identical to that of [Theorem 5.2](#), working in the hybrid space over a unit disk that we will parametrize by  $u \in \mathbb{D}$ . For fixed  $b \geq 1$ ,  $t = 1/u$  and any  $s$  satisfying  $|s| = |u|^b$ , consider the functions

$$g_u(z) = \frac{G_{F_{1/u}}(z, 1)}{\log |u|^{-1}} = \frac{G_{F_t}(z, 1)}{\log |t|}$$

and

$$bg_s(z) = b \frac{G_{F_s}(z, 1)}{\log |s|^{-1}}$$

in the fiber  $\{u\} \times \mathbb{P}^1$ .

As computed in [Proposition 3.3](#), the limit of  $g_u(z)$  as  $u \rightarrow 0$  with  $|z| = |u|^a$  is

$$\hat{g}_\infty(a) := \begin{cases} -a & \text{for } a \leq -1, \\ (a^2 + 1)/2 & \text{for } -1 \leq a \leq 0, \\ 1/2 & \text{for } a \geq 0. \end{cases}$$

As  $u \rightarrow 0$ , the measures  $\mu_{1/u}$  on  $\{u\} \times \mathbb{P}^1$  will converge the canonical measure  $\hat{\mu}_\infty$  for the map  $f = f_{1/U}$  on the Berkovich projective line, working over the field  $\mathbb{C}((U))$ ; the measure  $\hat{\mu}_\infty$  is uniformly distributed on the interval  $[\zeta_{0,1}, \zeta_{0,|U|_0^{-1}}]$ .

As  $s \rightarrow 0$  with  $|s| = 1/|t|^b = |u|^b$ ,  $b \geq 1$ , we have

$$bg_s(z) \rightarrow \hat{g}_b(\zeta_{0,|U|_0^a})$$

for  $|z| = |u|^a$ , exactly as in [\(5.3\)](#). The non-archimedean local energy is computed in [Theorem 3.1](#) as

$$E(\hat{\mu}_\infty, \hat{\mu}_b) = \frac{b+1}{6}.$$

We conclude as in the proof of [Theorem 5.2](#) that, for all given  $\varepsilon > 0$ , there exists  $\delta > 0$  such that

$$\left(\frac{b+1}{6} - b\varepsilon\right) \log |t| \leq E_\infty(s, t) \leq \left(\frac{b+1}{6} + b\varepsilon\right) \log |t|$$

for all  $|t| > 1/\delta$  and  $|s| = 1/|t|^b$ . This completes the proof of [Theorem 5.5](#).  $\square$

## 6. Proof of Theorems 1.6 and 1.7

In this section, we first prove [Theorem 1.7](#), which states there exist constants  $\alpha, \beta > 0$  such that

$$\hat{h}_{t_1} \cdot \hat{h}_{t_2} \geq \alpha h(t_1, t_2) - \beta$$

for all  $t_1 \neq t_2$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$ . We then use this lower bound to prove [Theorem 1.6](#) and [Proposition 1.9](#).

6.1. *Balancing local contributions.* Fix any  $r$  such that

$$0 < r \leq 1/16.$$

Fix  $t_1, t_2 \in \overline{\mathbb{Q}} \setminus \{0, 1\}$ , and let  $K$  be any number field containing  $t_1$  and  $t_2$ . We split the places  $M_K$  into “good” and “bad” subsets, depending on the pair  $t_1, t_2$  and the choice of  $r$ . Let  $M_{\text{good}}(t_1, t_2)$  be the set of places  $v \in M_K$  with

$$|\log |t_2/t_1|_v| \geq r \cdot \max\{|\log |t_2|_v|, |\log |t_1|_v|\},$$

and set  $M_{\text{bad}}(t_1, t_2) = M_K \setminus M_{\text{good}}(t_1, t_2)$ . We further decompose  $M_{\text{good}}(t_1, t_2)$  into its archimedean ( $M_{\text{good}}^\infty$ ) and non-archimedean ( $M_{\text{good}}^0$ ) places.

LEMMA 6.1. *There exists a constant  $C_0 > 0$  such that*

$$6 E_v(t_1, t_2) \geq \frac{3r}{4} |\log |t_1/t_2|_v| - C_0$$

for any choice of  $t_1$  and  $t_2$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$  and for all  $v \in M_{\text{good}}^\infty(t_1, t_2)$ .

*Proof.* Let

$$\epsilon = \frac{r^2}{24},$$

and let  $\delta_1$  be the minimum of the  $\delta$ ’s from [Theorems 5.2](#) and [5.5](#) for this choice of  $\epsilon$ . Let  $\delta_2$  be the  $\delta$  of [Theorem 5.1](#) for the compact set

$$\{t \in \mathbb{C} : \delta_1 \leq |t| \leq 1/\delta_1 \text{ and } |t - 1| \geq \delta_1\}$$

in  $\mathbb{C} \setminus \{0, 1\}$ . Let  $\delta_0$  be the minimum of  $\delta_1$  and  $\delta_2$ , and let  $C_0$  be any real number larger than  $\log(1/\delta_0)$ .

Now fix  $t_1, t_2$  and any number field  $K$  containing  $t_1$  and  $t_2$ , and fix a place  $v \in M_{\text{good}}^\infty(t_1, t_2) \subset M_K$ . If  $\delta_0 \leq |t_i|_v \leq 1/\delta_0$  for  $i = 1, 2$ , we have

$$6 E_v(t_1, t_2) \geq 0 \geq \frac{3r}{4} |\log |t_1/t_2|_v| - C_0.$$

As  $v \in M_{\text{good}}^\infty$ , if  $|t_2|_v \leq |t_1|_v < 1$ , then

$$|t_2|_v = |t_1|_v^b < |t_1|_v \text{ for } \frac{b-1}{b} \geq r$$

and therefore, by [Theorem 5.2](#), if additionally  $|t_1|_v < \delta_1$ , then we have

$$\begin{aligned} 6E_v(t_1, t_2) &\geq \left( \frac{(b-1)^2}{b^2} - 6\epsilon \right) |\log |t_1|_v^b| \\ &\geq \left( \frac{(b-1)r}{b} - \frac{r^2}{4} \right) |\log |t_1|_v^b| \\ &\geq \frac{3r}{4} \frac{b-1}{b} |\log |t_1|_v^b| = \frac{3r}{4} |\log |t_1/t_2|_v|. \end{aligned}$$

If  $|t_1|_v < \delta_1$  and  $|t_2|_v = 1/|t_1|_v^b$  for some  $b \geq 1$ , then by [Theorem 5.5](#),

$$\begin{aligned} 6E_v(t_1, t_2) &\geq \left( \frac{(b+1)}{b} - 6\epsilon \right) |\log |t_1|_v^b| \\ &\geq \frac{3(b+1)r}{4} |\log |t_1|_v| = \frac{3r}{4} |\log |t_1/t_2|_v|; \end{aligned}$$

if  $\delta_1 \leq |t_1|_v \leq 1/\delta_1$ ,  $|t_1 - 1|_v \geq \delta_1$  and  $|t_2|_v < \delta_0$ , we have by [Theorem 5.1](#) that

$$6E_v(t_1, t_2) \geq (1 - 6\epsilon) |\log |t_2|_v| \geq \frac{3r}{4} |\log |t_1/t_2|_v| - C_0.$$

Combining the above inequalities with the symmetry relations of [Proposition 2.3](#), we obtain

$$6E_v(t_1, t_2) \geq \frac{3r}{4} |\log |t_1/t_2|_v| - C_0. \quad \square$$

LEMMA 6.2. *There is a constant  $C > 0$  such that*

$$\sum_{v \in M_{\text{good}}(t_1, t_2)} 3r_v E_v(t_1, t_2) \geq \frac{3r}{4} h(t_2/t_1) - \frac{3r^2}{2} h(t_1, t_2) - C$$

for any  $t_1 \neq t_2 \in \overline{\mathbb{Q}} \setminus \{0, 1\}$ .

*Proof.* Fix  $t_1$  and  $t_2$  and any number field  $K$  containing them. For the non-archimedean places  $v \in M_{\text{good}}^0(t_1, t_2)$ , by [Theorem 3.1](#), we have

$$6E_v(t_1, t_2) \geq r \cdot |\log |t_2/t_1|_v| - 8 \log^+ |1/2|_v,$$

and thus

$$(6.1) \quad \sum_{v \in M_{\text{good}}^0(t_1, t_2)} 6r_v E_v(t_1, t_2) \geq \sum_{v \in M_{\text{good}}^0(t_1, t_2)} r_v (r \cdot |\log |t_2/t_1|_v| - 8 \log^+ |1/2|_v).$$

Now choose any integer  $N_0$  so that  $\log N_0$  is larger than the  $C_0$  of [Lemma 6.1](#) for each archimedean  $v \in M_{\text{good}}^\infty(t_1, t_2)$ . We have

$$6E_v(t_1, t_2) \geq \frac{3r}{4} |\log |t_2/t_1|_v| - \log N_0$$

for all  $v \in M_{\text{good}}^\infty(t_1, t_2)$ . With  $h$  the naive logarithmic height on  $\overline{\mathbb{Q}}$ , we set

$$C = 4h(2) + \frac{1}{2}h(N_0) = \frac{1}{2}\log(2^8 N_0).$$

Then, we have

$$\begin{aligned} \sum_{v \in M_{\text{good}}(t_1, t_2)} 6r_v E_v(t_1, t_2) &= \sum_{v \in M_{\text{good}}^\infty(t_1, t_2)} 6r_v E_v(t_1, t_2) \\ &\quad + \sum_{v \in M_{\text{good}}^0(t_1, t_2)} 6r_v E_v(t_1, t_2) \\ &\geq \sum_{v \in M_{\text{good}}(t_1, t_2)} r_v \frac{3r}{4} |\log|t_2/t_1|_v| - \sum_{v \in M_K} r_v (8\log^+|2|_v + \log^+|N_0|_v) \\ &= \sum_{v \in M_{\text{good}}(t_1, t_2)} r_v \frac{3r}{4} |\log|t_2/t_1|_v| - 2C \\ &= \sum_{v \in M_K} r_v \frac{3r}{4} |\log|t_2/t_1|_v| - \sum_{v \in M_{\text{bad}}(t_1, t_2)} r_v \frac{3r}{4} |\log|t_2/t_1|_v| - 2C \\ &\geq \sum_{v \in M_K} \frac{3r_v r}{4} |\log|t_2/t_1|_v| \\ &\quad - \sum_{v \in M_{\text{bad}}(t_1, t_2)} \frac{3r^2 r_v}{4} \max\{|\log|t_2|_v|, |\log|t_1|_v|\} - 2C \\ &\geq \frac{3r}{4} 2h(t_2/t_1) - \frac{3r^2}{4} \cdot 4h(t_2, t_1) - 2C. \end{aligned}$$

For the last inequality, we use the facts that  $2h(x) = \sum_{v \in M_K} r_v |\log|x|_v|$  for nonzero  $x \in K$  and

$$\sum_{v \in M_K} r_v \max\{|\log|t_2|_v|, |\log|t_1|_v|\} \leq 2(h(t_2) + h(t_1)) \leq 4h(t_2, t_1). \quad \square$$

6.2. *Proof of Theorem 1.7.* We begin with a standard lemma.

LEMMA 6.3. *There is a constant  $C > 0$ , such that*

$$h\left(\frac{t_2}{t_1}, \frac{1-t_2}{1-t_1}\right) \geq \frac{1}{2}h(t_1, t_2) - C$$

for  $t_1 \neq t_2 \in \overline{\mathbb{Q}} \setminus \{0, 1\}$ . Here the  $h$  is the naive logarithmic height on  $\mathbb{A}^2(\overline{\mathbb{Q}})$ .

*Proof.* Consider the birational transformation  $g : \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$  defined in affine coordinates by  $g(x_1, x_2) = (x_2/x_1, (1-x_2)/(1-x_1))$ , with inverse

$$g^{-1}(y_1, y_2) = \left(\frac{1-y_2}{y_1-y_2}, \frac{y_1(1-y_2)}{y_1-y_2}\right)$$

of degree  $d = 2$ . There exists a constant  $C$  such that

$$h(g^{-1}(x : y : z)) \leq (\deg g^{-1}) h(x : y : z) + C = 2 h(x : y : z) + C$$

outside of the indeterminacy set for  $g^{-1}$  in  $\mathbb{P}^2$  [HS00, Th. B.2.5]. The indeterminacy set for  $g^{-1}$  is  $\{(0 : 1 : 0), (1 : 0 : 0), (1 : 1 : 1)\}$ . Therefore, letting  $(t_1 : t_2 : 1) = g^{-1}(x : y : 1)$  for some point  $(x : y : 1)$ , we obtain

$$h(t_1, t_2) \leq 2 h(g(t_1, t_2)) + C$$

for all  $t_1 \neq t_2$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$ . In other words,

$$h\left(\frac{t_2}{t_1}, \frac{t_2 - 1}{t_1 - 1}\right) \geq \frac{1}{2} h(t_1, t_2) - \frac{1}{2} C. \quad \square$$

Now fix  $t_1 \neq t_2$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$ . From [Lemma 6.2](#), we know there is a constant  $C$  (independent of  $t_1$  and  $t_2$ ) such that

$$\begin{aligned} 3 \hat{h}_{t_1} \cdot \hat{h}_{t_2} &= 3 \sum_{v \in M_K} r_v E_v(t_1, t_2) \\ (6.2) \quad &\geq 3 \sum_{v \in M_{\text{good}}(t_1, t_2)} r_v E_v(t_1, t_2) \\ &\geq \frac{3r}{4} h(t_2/t_1) - \frac{3r^2}{2} h(t_2, t_1) - C \end{aligned}$$

for any  $t_1 \neq t_2 \in \overline{\mathbb{Q}} \setminus \{0, 1\}$ . Replacing  $t_i$  in [inequality \(6.2\)](#) with  $1 - t_i$ , for  $i = 1, 2$ , we also have

$$3 \hat{h}_{1-t_1} \cdot \hat{h}_{1-t_2} \geq \frac{3r}{4} h\left(\frac{1-t_2}{1-t_1}\right) - \frac{3r^2}{2} h(1-t_2, 1-t_1) - C.$$

Combining this with [Proposition 2.3](#), we find that

$$(6.3) \quad 3 \hat{h}_{t_1} \cdot \hat{h}_{t_2} \geq \frac{3r}{4} h\left(\frac{1-t_2}{1-t_1}\right) - \frac{3r^2}{2} h(1-t_2, 1-t_1) - C.$$

Consequently, by adding the [inequalities \(6.2\)](#) and [\(6.3\)](#), we have

$$\begin{aligned} 6 \hat{h}_{t_1} \cdot \hat{h}_{t_2} &\geq \frac{3r}{4} \left( h(t_2/t_1) + h\left(\frac{1-t_2}{1-t_1}\right) \right) \\ &\quad - \frac{3r^2}{2} (h(t_2, t_1) + h(1-t_2, 1-t_1)) - 2C. \end{aligned}$$

Observe that there is a constant  $C' > 0$  from [Lemma 6.3](#) such that

$$h(t_2/t_1) + h\left(\frac{1-t_2}{1-t_1}\right) \geq h\left(\frac{t_2}{t_1}, \frac{1-t_2}{1-t_1}\right) \geq \frac{1}{2} \cdot h(t_1, t_2) - C'.$$

Since  $|h(1-t_2, 1-t_1) - h(t_1, t_2)|$  is uniformly bounded over all pairs  $t_1, t_2 \in \overline{\mathbb{Q}}$ , we may combine the above inequality with the previous to conclude that

$$6 \hat{h}_{t_1} \cdot \hat{h}_{t_2} \geq \frac{3r}{8} h(t_1, t_2) - 3r^2 \cdot h(t_1, t_2) - 6C''.$$

In other words,

$$\hat{h}_{t_1} \cdot \hat{h}_{t_2} \geq (r/16 - r^2/2)h(t_2, t_1) - C''',$$

and the proof of [Theorem 1.7](#) is complete by taking  $\alpha = r/16 - r^2/2$  and  $\beta = C'''$ .  $\square$

*Remark 6.4.* If we set  $r = 1/16$ , the constant  $\alpha > 0$  in [Theorem 1.7](#) can be taken to be  $\alpha = 1/512$ .

**6.3. Proof of Theorem 1.6.** We will use [Theorem 1.7](#) to deduce a uniform lower bound on the height pairing  $\hat{h}_s \cdot \hat{h}_t$  for all  $s \neq t$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$ .

Suppose there exist parameters  $s_n \neq t_n \in \overline{\mathbb{Q}}$  such that

$$\hat{h}_{s_n} \cdot \hat{h}_{t_n} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Fix  $\epsilon > 0$ . For each  $n$ , choose a number field  $K_n$  containing  $s_n$  and  $t_n$ . By assumption and non-negativity of the local energies  $E_v$  ([Proposition 2.2](#)), there is  $N \in \mathbb{N}$  such that for all  $n > N$ , the archimedean contribution to the pairing is less than  $\epsilon$ ; that is, for  $n > N$ ,

$$\sum_{v \in M_{K_n}^\infty} r_v E_v(s_n, t_n) < \epsilon,$$

recalling that  $r_v = \frac{[K_{n,v} : \mathbb{Q}_v]}{[K_n : \mathbb{Q}]}$  now depends on  $n$ .

Let  $M_n$  be the set of archimedean places  $v$  in  $M_{K_n}^\infty$  such that  $E_v(s_n, t_n) < 2\epsilon$ , noting that for  $n > N$ , we have

$$\sum_{v \in M_n} r_v \geq \frac{1}{2}.$$

Recall that the local energy  $E_v(s, t)$  is continuous in  $s$  and  $t$ , and it vanishes if and only if  $s = t$ . So there exists a  $\delta$ , depending only on  $\epsilon$ , so that, for each  $n > N$  and for each place  $v \in M_n$ , one of the following must hold:

- (1)  $|t_n - s_n|_v < \delta$ ;
- (2)  $\min\{|s_n|_v, |t_n|_v\} < \delta$ ;
- (3)  $\min\{|s_n - 1|_v, |t_n - 1|_v\} < \delta$ ;
- (4)  $\max\{|s_n|_v, |t_n|_v\} > 1/\delta$ .

Note that we can take  $\delta \rightarrow 0$  as  $\epsilon \rightarrow 0$ . We may then, for each  $n > N$ , choose a subset  $M'_n$  of  $M_n$  for which  $s_n$  and  $t_n$  satisfy the same one of the four conditions at all places  $v \in M'_n$ , and such that

$$\sum_{v \in M'_n} r_v \geq \frac{1}{8}.$$

We conclude by the product formula that

$$\max\{h(s_n - t_n), h(s_n, t_n), h(s_n - 1, t_n - 1)\} > \frac{1}{8} \log \frac{1}{\delta}.$$

It then follows from the triangle inequality, combined with shrinking our choice of  $\epsilon$ , that we have  $h(s_n, t_n) \rightarrow \infty$ . The inequality of [Theorem 1.7](#) implies that  $\hat{h}_{s_n} \cdot \hat{h}_{t_n} \rightarrow \infty$  as well, a contradiction. This completes the proof of [Theorem 1.6](#).

**6.4. Proof of [Proposition 1.9](#).** Fix a number field  $K$ , and fix  $t_1 \neq t_2$  in  $K \setminus \{0, 1\}$ . Let  $\|\cdot\|_i$  denote the adelic metric on the line bundle  $\mathcal{O}_{\mathbb{P}^1}(1)$  associated to the height  $\hat{h}_{t_i}$ . Let  $\bar{L}$  denote the line bundle  $\mathcal{O}_{\mathbb{P}^1}(1)$  equipped with the metric  $(\|\cdot\|_1 \|\cdot\|_2)^{1/2}$ ; its associated height function is

$$h_{\bar{L}}(x) = \frac{1}{2} (\hat{h}_{t_1}(x) + \hat{h}_{t_2}(x)).$$

Zhang's inequality on the essential minimum of a height function implies that

$$\liminf_{n \rightarrow \infty} h_{\bar{L}}(x_n) \geq (h_{\bar{L}} \cdot h_{\bar{L}})/(2 \deg L) = \frac{1}{2} (h_{\bar{L}} \cdot h_{\bar{L}})$$

along any infinite sequence of distinct points  $x_n \in \mathbb{P}^1(\overline{\mathbb{Q}})$  [[Zha95](#), Th. 1.10]. In particular, the set

$$\{x \in \mathbb{P}^1(\overline{\mathbb{Q}}) : \hat{h}_{t_1}(x) + \hat{h}_{t_2}(x) \leq b\}$$

is finite for any choice of  $b < h_{\bar{L}} \cdot h_{\bar{L}}$ .

By the linearity of the intersection pairing, we see that

$$h_{\bar{L}} \cdot h_{\bar{L}} = \frac{1}{4} \hat{h}_{t_1} \cdot \hat{h}_{t_1} + \frac{1}{2} \hat{h}_{t_1} \cdot \hat{h}_{t_2} + \frac{1}{4} \hat{h}_{t_2} \cdot \hat{h}_{t_2} = \frac{1}{2} \hat{h}_{t_1} \cdot \hat{h}_{t_2}.$$

Therefore, we may choose any  $b < \delta/2$  for the  $\delta$  of [Theorem 1.6](#), and the proposition is proved.

## 7. Proof of [Theorem 1.8](#)

Fix any  $b \geq 0$  such that  $b < \delta/2$  for the  $\delta$  of [Theorem 1.6](#). Recall from [Proposition 1.9](#) that the set

$$(7.1) \quad S(b, t_1, t_2) := \{x \in \mathbb{P}^1(\overline{\mathbb{Q}}) : \hat{h}_{t_1}(x) + \hat{h}_{t_2}(x) \leq b\}$$

is finite for every pair  $t_1 \neq t_2 \in \overline{\mathbb{Q}} \setminus \{0, 1\}$ . Note that  $\{0, 1, \infty\} \subset S(0, t_1, t_2)$  so that  $|S(b, t_1, t_2)| \geq 3$  for all  $t_1 \neq t_2$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$  and all  $b \geq 0$ . In this section, we prove the following generalization of [Theorem 1.8](#).

**THEOREM 7.1.** *Let  $b \geq 0$  be chosen so that  $b < \delta/2$  for the  $\delta$  of [Theorem 1.6](#). For all  $\varepsilon > 0$ , there exists a constant  $C(\varepsilon)$  so that*

$$\hat{h}_{t_1} \cdot \hat{h}_{t_2} \leq 4b + \left( \varepsilon + \frac{C(\varepsilon)}{|S(b, t_1, t_2)|} \right) (h(t_1, t_2) + 1),$$

for all  $t_1 \neq t_2$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$ , for the set  $S(b, t_1, t_2)$  defined by (7.1).

Note that [Theorem 1.8](#) follows from [Theorem 7.1](#) by setting  $b = 0$ .

7.1. *Adelic measures and heights associated to a finite set.* Fix a number field  $K$ , and suppose that  $F$  is a finite set in  $\overline{K}$  that is  $\text{Gal}(\overline{K}/K)$ -invariant. Let  $\eta$  be a collection of positive real numbers

$$\eta := \{\eta_v\}_{v \in M_K}$$

with  $\eta_v = 1$  for all but finitely many  $v \in M_K$ . For archimedean  $v \in M_K$  and  $x \in F$ , we let  $m_{x,v}$  denote the Lebesgue probability measure on the circle of radius  $\eta_v$  centered at the point  $x \in F$ . We then set

$$m_{F,\eta,v} = \frac{1}{|F|} \sum_{x \in F} m_{x,v}.$$

Similarly, for each non-archimedean  $v \in M_K$ , we let  $m_{F,\eta,v}$  denote the probability measure distributed uniformly on the points  $\zeta_{x,\eta_v}$  in  $\mathbb{A}_v^{1,an}$  over all  $x \in F$ . Then  $m_{F,\eta} = \{m_{F,\eta,v}\}_{v \in M_K}$  is an adelic measure in the sense of [FRL06]. It gives rise to a unique height  $h_{F,\eta}$  on  $\mathbb{P}^1(\overline{\mathbb{Q}})$  associated to a continuous and semi-positive adelic metric on  $\mathcal{O}_{\mathbb{P}^1}(1)$  with curvature distributions given by  $m_{F,\eta,v}$  and satisfying

$$(7.2) \quad h_{F,\eta} \cdot h_{F,\eta} = 0.$$

Its local heights are given by

$$\lambda_{F,\eta,v}(z) = \alpha_v + \frac{1}{|F|} \sum_{x \in F} \log \max\{|z - x|_v, \eta_v\}$$

for  $z \in \mathbb{C}_v$  and suitable constants  $\alpha_v$ ; taking

$$\alpha_v = -\frac{1}{2|F|} \sum_{x \in F} \int \log \max\{|z - x|_v, \eta_v\} dm_{F,\eta,v}$$

gives (7.2).

*Remark 7.2.* The height  $h_{F,\eta}$  will generally not admit sequences of “small” points, meaning sequences  $x_n \in \mathbb{P}^1(\overline{\mathbb{Q}})$  with  $h_{F,\eta}(x_n) \rightarrow 0$ . In fact, for any choices of  $F$  and  $\eta$  such that  $\sum_v r_v \alpha_v \neq 0$ , the essential minimum of  $h_{F,\eta}$  is positive.

7.2. *An upper bound on the height pairing.* Now suppose that  $t_1$  and  $t_2$  lie in  $K \setminus \{0, 1\}$ . Recall that  $\mu_t$  and  $\hat{h}_t$  respectively denote the measure and height associated to the curve  $E_t$ . By the triangle inequality for the distance function of Section 2.8, we have

$$(7.3) \quad (\hat{h}_{t_1} \cdot \hat{h}_{t_2})^{1/2} \leq (\hat{h}_{t_1} \cdot h_{F,\eta})^{1/2} + (\hat{h}_{t_2} \cdot h_{F,\eta})^{1/2}$$

for any choice of  $F$  and  $\eta$ . By symmetry and bilinearity of the mutual energy,

$$\begin{aligned}\hat{h}_{t_i} \cdot h_{F,\eta} &= \frac{1}{2} \sum_{v \in M_K} r_v (\mu_{t_i,v} - m_{F,\eta,v}, \mu_{t_i,v} - m_{F,\eta,v})_v \\ &= \frac{1}{2} \sum_{v \in M_K} r_v ((\mu_{t_i,v}, \mu_{t_i,v})_v - 2(m_{F,\eta,v}, \mu_{t_i,v})_v + (m_{F,\eta,v}, m_{F,\eta,v})_v)\end{aligned}$$

for  $i = 1, 2$ . For fixed  $i$ , writing the local height for  $\hat{h}_{t_i}$  as  $\lambda_{t_i,v} = \log |z|_v + c_v + o(1)$  for  $|z|_v \rightarrow \infty$  yields

$$\sum_v r_v (\mu_{t_i,v}, \mu_{t_i,v})_v = - \sum_v r_v \int (\lambda_{t_i,v} - c_v) d\mu_{t_i,v} = 0$$

from (2.11). Therefore,

$$\hat{h}_{t_i} \cdot h_{F,\eta} = \frac{1}{2} \sum_{v \in M_K} r_v (-2(\mu_{t_i,v}, m_{F,\eta,v})_v + (m_{F,\eta,v}, m_{F,\eta,v})_v).$$

Recall from Section 2.7 that  $[F]_v$  is the probability measure on  $\mathbb{P}_v^{1,an}$  distributed equally on the elements of  $F$  for each  $v \in M_K$ . By [FRL06, Lemma 4.11] and [Fil17, Lemma 12], we have

$$(m_{F,\eta,v}, m_{F,\eta,v})_v \leq ([F]_v, [F]_v)_v + \frac{-\log \eta_v}{|F|}.$$

It follows that

$$\begin{aligned}(7.4) \quad \hat{h}_{t_i} \cdot h_{F,\eta} &\leq \frac{1}{2} \sum_{v \in M_K} r_v \cdot \left( -2(\mu_{t_i,v}, m_{F,\eta,v})_v + ([F]_v, [F]_v)_v + \frac{-\log \eta_v}{|F|} \right) \\ &= \frac{1}{2} \sum_{v \in M_K} r_v \cdot \left( -2(\mu_{t_i,v}, m_{F,\eta,v})_v + \frac{-\log \eta_v}{|F|} \right)\end{aligned}$$

with the final equality following from (2.16).

**PROPOSITION 7.3.** *Suppose  $t \neq 0, 1$  lies in a number field  $K$ . Assume that  $F$  is a finite,  $\text{Gal}(\overline{K}/K)$ -invariant set of points. Then*

$$\hat{h}_t \cdot h_{F,\eta} \leq \hat{h}_t(F) + \sum_{v \in M_K} r_v \left( -(\mu_{t,v}, m_{F,\eta,v})_v + (\mu_{t,v}, [F]_v)_v + \frac{-\log \eta_v}{2|F|} \right)$$

for any choice of  $\eta = \{\eta_v\}_v$  with  $\eta_v = 1$  for all but finitely many  $v \in M_K$ .

*Proof.* The height of  $F$  is computed as

$$\hat{h}_t(F) = \frac{1}{|F|} \sum_{x \in F} \hat{h}_t(x) = \hat{h}_t(\infty) - \sum_{v \in M_K} r_v (\mu_{t,v}, [F]_v)_v = - \sum_{v \in M_K} r_v (\mu_{t,v}, [F]_v)_v,$$

and therefore we may add  $\hat{h}_t(F) + \sum_v r_v (\mu_{t,v}, [F]_v)_v$  to the right-hand side of (7.4).  $\square$

7.3. *Proof of Theorem 7.1.* Fix  $0 \leq b < \delta/2$  so that [Proposition 1.9](#) is satisfied for all  $t_1 \neq t_2$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$ . Now fix  $t_1 \neq t_2$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$  and a number field  $K$  containing  $t_1$  and  $t_2$ . Set

$$F = \{x \in \mathbb{P}^1(\overline{\mathbb{Q}}) : \hat{h}_{t_1}(x) + \hat{h}_{t_2}(x) \leq b\} \setminus \{\infty\} = S(b, t_1, t_2) \setminus \{\infty\},$$

so  $F$  is a finite,  $\text{Gal}(\overline{K}/K)$ -invariant set with

$$\hat{h}_{t_i}(F) \leq b$$

for  $i = 1, 2$ . At each non-archimedean place  $v$  of  $K$ , we set

$$\eta_v := \min\{1, |t_1(t_1 - 1)|_v, |t_2(t_2 - 1)|_v\}.$$

Now fix  $\varepsilon' > 0$ . For each archimedean  $v$ , we set

$$\eta_v := c(\varepsilon') \min_{i=1,2} \min\{|t_i|_v^2, |t_i - 1|_v^2, |t_i|_v^{-2}\},$$

where the constant  $c(\varepsilon')$  is from [Proposition 4.4](#). Let  $\eta = \{\eta_v\}_v$ ; observe that  $\eta_v = 1$  for all but finitely many  $v$ , and

$$(7.5) \quad \sum_{v \in M_K} -r_v \log \eta_v \leq 2(h(t_1) + h(1 - t_1) + h(t_2) + h(1 - t_2)) - \frac{1}{2} \log c(\varepsilon').$$

For non-archimedean  $v$ , the explicit form of the measure  $\mu_{t_i, v}$  (described in [Section 3](#)) implies that

$$(\mu_{t_i, v}, m_{F, \eta, v})_v = (\mu_{t_i, v}, [F]_v)_v$$

for this choice of  $\eta$ , because the potentials for  $\mu_{t_i, v}$  are constant on disks of radius  $\eta_v$ .

We thus obtain from [Proposition 7.3](#) that

$$\begin{aligned} \hat{h}_{t_i} \cdot h_{F, \eta} &\leq b + \sum_{v \in M_K^0} r_v \frac{-\log \eta_v}{2|F|} \\ &\quad + \sum_{v \in M_K^\infty} r_v \left( -(\mu_{t_i, v}, m_{F, \eta, v})_v + (\mu_{t_i, v}, [F]_v)_v + \frac{-\log \eta_v}{2|F|} \right) \end{aligned}$$

for  $i = 1, 2$ , where  $M_K^0$  denotes the non-archimedean places and  $M_K^\infty$  the archimedean places.

We have for  $v \in M_K^\infty$  that

$$-(\mu_{t_i, v}, m_{F, \eta, v})_v + (\mu_{t_i, v}, [F]_v)_v \leq \varepsilon' \log \max\{|t_i|_v, |t_i|_v^{-1}, |t_i - 1|_v^{-1}\}$$

for  $i = 1, 2$  by [Proposition 4.4](#).

Since the logarithmic Weil height satisfies  $2h(x) = \sum_v |\log |x|_v|$ , we thus obtain

$$\begin{aligned} \hat{h}_{t_i} \cdot h_{F, \eta} &\leq b + 2\varepsilon' (h(t_i) + h(t_i - 1)) \\ &\quad + \frac{2(h(t_1) + h(1 - t_1) + h(t_2) + h(1 - t_2)) - \frac{1}{2} \log c(\varepsilon')}{|F|} \end{aligned}$$

for  $i = 1, 2$ . Since  $h(1 - t_i) \leq h(t_i) + \log 2 \leq h(t_1, t_2) + \log 2$  for  $i = 1, 2$ , this inequality becomes

$$\hat{h}_{t_i} \cdot h_{F,\eta} \leq b + 2\varepsilon' (2h(t_1, t_2) + \log 2) + \frac{1}{|F|} \left( 8h(t_1, t_2) + 4\log 2 - \frac{1}{2} \log c(\varepsilon') \right)$$

for  $i = 1, 2$ .

By the triangle inequality (7.3), we have

$$\begin{aligned} (\hat{h}_{t_1} \cdot \hat{h}_{t_2})^{1/2} &\leq (\hat{h}_{t_1} \cdot h_{F,\eta})^{1/2} + (\hat{h}_{t_2} \cdot h_{F,\eta})^{1/2} \\ &\leq 2 \left( b + 2\varepsilon' (2h(t_1, t_2) + \log 2) + \frac{1}{|F|} (8h(t_1, t_2) + 4\log 2 - \frac{1}{2} \log c(\varepsilon')) \right)^{1/2}, \end{aligned}$$

so

(7.6)

$$\begin{aligned} \hat{h}_{t_1} \cdot \hat{h}_{t_2} &\leq 4 \left( b + 2\varepsilon' (2h(t_1, t_2) + \log 2) + \frac{1}{|F|} (8h(t_1, t_2) + 4\log 2 - \frac{1}{2} \log c(\varepsilon')) \right) \\ &= 4b + \left( \frac{32}{|F|} + 16\varepsilon' \right) h(t_1, t_2) + \frac{16\log 2 - 2\log c(\varepsilon')}{|F|} + 8\varepsilon' \log 2. \end{aligned}$$

Fix any  $\varepsilon > 0$ , and choose  $\varepsilon' < \varepsilon/16$ . Since  $|F| = |S(b, t_1, t_2)| - 1 \geq 2$ , we can find a large constant  $C(\varepsilon)$  satisfying

$$\frac{32}{|F|} + 16\varepsilon' \leq \varepsilon + \frac{C(\varepsilon)}{|S(b, t_1, t_2)|}$$

and

$$\frac{16\log 2 - 2\log c(\varepsilon')}{|F|} + 8\varepsilon' \log 2 \leq \varepsilon + \frac{C(\varepsilon)}{|S(b, t_1, t_2)|}.$$

The inequality (7.6) then yields

$$\hat{h}_{t_1} \cdot \hat{h}_{t_2} \leq 4b + \left( \varepsilon + \frac{C(\varepsilon)}{|S(b, t_1, t_2)|} \right) (h(t_1, t_2) + 1),$$

concluding the proof of Theorem 7.1.

## 8. Proof of Theorem 1.5

In this section, we deduce Theorem 1.5 from Theorems 1.6, 1.7 and 1.8 for algebraic values of  $t_1$  and  $t_2$ ; we then extend the result to hold for parameters  $t_i$  in  $\mathbb{C}$ , via a specialization argument. In fact, we prove the following stronger result over  $\overline{\mathbb{Q}}$ :

**THEOREM 8.1.** *There exist constants  $B$  and  $b > 0$  so that*

$$|\{x \in \mathbb{P}^1(\overline{\mathbb{Q}}) : \hat{h}_{t_1}(x) + \hat{h}_{t_2}(x) \leq b\}| \leq B$$

for all  $t_1 \neq t_2$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$ .

Finally, in [Section 8.4](#), we observe that there is no uniform bound on the order of the torsion points on  $E_{t_1}$  and  $E_{t_2}$  that can share an  $x$ -coordinate, even if [Theorem 1.5](#) provides a uniform bound on the total number of such points.

8.1. *Proof of Theorem 8.1.* Let  $\delta > 0$  be as in [Theorem 1.6](#) so that

$$\hat{h}_{t_1} \cdot \hat{h}_{t_2} \geq \delta$$

for all  $t_1 \neq t_2$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$ . Fix

$$0 < b < \delta/8$$

so that, from [Proposition 1.9](#), the set

$$S(b, t_1, t_2) = \{x \in \mathbb{P}^1(\overline{\mathbb{Q}}) : \hat{h}_{t_1}(x) + \hat{h}_{t_2}(x) \leq b\}$$

is finite for all  $t_1 \neq t_2$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$ . Let  $h(t_1, t_2)$  be the naive logarithmic height on  $\mathbb{A}^2(\overline{\mathbb{Q}})$ . Fix  $H > \frac{2\beta}{\alpha}$  for the  $\alpha, \beta$  of [Theorem 1.7](#) and such that

$$(8.1) \quad \frac{H - 8b/\alpha}{H + 1} > 3/4.$$

Suppose that  $t_1 \neq t_2 \in \overline{\mathbb{Q}}$  satisfy  $h(t_1, t_2) \geq H$ . Then for  $\varepsilon = \frac{\alpha}{4}$ , there exists by [Theorem 7.1](#) a constant  $C$  such that

$$\hat{h}_{t_1} \cdot \hat{h}_{t_2} \leq 4b + \left( \frac{\alpha}{4} + \frac{C}{|S(b, t_1, t_2)|} \right) (h(t_1, t_2) + 1).$$

On the other hand, by [Theorem 1.7](#) and the choice of  $H$ , we have

$$\frac{\alpha}{2} h(t_1, t_2) \leq \alpha h(t_1, t_2) - \beta \leq \hat{h}_{t_1} \cdot \hat{h}_{t_2}.$$

Therefore

$$\frac{\alpha}{2} h(t_1, t_2) \leq 4b + \left( \frac{\alpha}{4} + \frac{C}{|S(b, t_1, t_2)|} \right) (h(t_1, t_2) + 1),$$

and so

$$|S(b, t_1, t_2)| \leq \frac{C}{\left( \frac{\alpha h/2 - 4b}{h+1} \right) - \frac{\alpha}{4}} = \frac{C}{\frac{\alpha}{2} \left( \frac{h - 8b/\alpha}{h+1} \right) - \frac{\alpha}{4}} \leq \frac{8C}{\alpha}$$

for  $h := h(t_1, t_2) \geq H$ , from (8.1).

Suppose now that  $t_1 \neq t_2 \in \overline{\mathbb{Q}}$  satisfy  $h(t_1, t_2) < H$ . Set  $\varepsilon' = \frac{\delta}{4(H+1)}$ , and find a constant  $C'$  as in [Theorem 7.1](#) so that

$$\hat{h}_{t_1} \cdot \hat{h}_{t_2} \leq 4b + \left( \varepsilon' + \frac{C'}{|S(b, t_1, t_2)|} \right) (h(t_1, t_2) + 1),$$

and thus, since  $b < \delta/8$ , we have

$$\delta/2 < \delta - 4b \leq \left( \varepsilon' + \frac{C'}{|S(b, t_1, t_2)|} \right) (h(t_1, t_2) + 1).$$

We conclude that

$$|S(b, t_1, t_2)| \leq \frac{4(H+1)C'}{\delta},$$

providing a uniform bound also for  $t_1$  and  $t_2$  satisfying  $h(t_1, t_2) < H$ . This completes the proof of [Theorem 8.1](#).

**8.2. Specialization: proof of Theorem 1.5.** We implement a standard specialization argument to deduce [Theorem 1.5](#) from [Theorem 8.1](#). Note that the division polynomials for the Legendre curve  $E_t$  have coefficients in  $\mathbb{Q}[t]$ ; see, for example, [[Sil09](#), Exer. 3.7]. Let  $B$  be the uniform bound obtained in [Theorem 8.1](#), so that

$$(8.2) \quad |\pi(E_{t_1}^{\text{tors}}) \cap \pi(E_{t_2}^{\text{tors}})| = |\{x \in \mathbb{P}^1(\overline{\mathbb{Q}}) : \hat{h}_{t_1}(x) = \hat{h}_{t_2}(x) = 0\}| \leq B$$

for all  $t_1 \neq t_2 \in \overline{\mathbb{Q}} \setminus \{0, 1\}$ . Assume that there exist  $t_1 \neq t_2 \in \mathbb{C} \setminus \{0, 1\}$  with

$$N(t_1, t_2) := |\pi(E_{t_1}^{\text{tors}}) \cap \pi(E_{t_2}^{\text{tors}})| > B$$

and  $t_1$  transcendental. If  $x \in \pi(E_{t_1}^{\text{tors}}) \cap \mathbb{P}^1(\overline{\mathbb{Q}})$ , then  $x \in \pi(E_t^{\text{tors}})$  for all  $t \in \mathbb{C} \setminus \{0, 1\}$  as it is a root of a division polynomial. It follows that there is at least one non-algebraic point  $x \in \pi(E_{t_1}^{\text{tors}}) \cap \pi(E_{t_2}^{\text{tors}})$ , as only  $x = 0, 1, \infty$  are torsion images for all  $t \in \mathbb{C} \setminus \{0, 1\}$  [[DWY16](#), Prop. 1.4].

Now let

$$S := \{x_1, x_2, \dots, x_N\} = \pi(E_{t_1}^{\text{tors}}) \cap \pi(E_{t_2}^{\text{tors}}),$$

where  $N = N(t_1, t_2)$ , and assume that  $x_1$  is transcendental. Because it is a torsion image for both parameters,  $\mathbb{Q}(x_1, t_1, t_2)$  and therefore also the field

$$L := \mathbb{Q}(t_1, t_2, x_1, \dots, x_N)$$

are of transcendence degree one. Consequently  $L$  is isomorphic to a function field  $k = K(X)$  for a number field  $K$  and an algebraic curve  $X$  defined over  $\overline{\mathbb{Q}}$ . Via the identification of  $L$  with  $k$ , there exists an algebraic point  $\gamma \in X(\overline{K})$  with distinct specializations  $x_i(\gamma) \in \mathbb{P}^1(\overline{\mathbb{Q}})$  for  $i = 1, \dots, N$  and

$$t_1(\gamma) \neq t_2(\gamma) \in \overline{\mathbb{Q}} \setminus \{0, 1\}.$$

The division relations in  $L$  imply that the specializations  $E_{t_1(\gamma)}$  and  $E_{t_2(\gamma)}$  have at least  $N$  common torsion images, contradicting (8.2). Therefore, we must have

$$|\pi(E_{t_1}^{\text{tors}}) \cap \pi(E_{t_2}^{\text{tors}})| \leq B$$

for all  $t_1 \neq t_2 \in \mathbb{C} \setminus \{0, 1\}$ , and the proof of [Theorem 1.5](#) is complete.  $\square$

**8.3. Common torsion images.** We obtain the following immediate corollary of [Theorem 1.5](#), which is a special case of [Conjecture 1.4](#). Recall that a standard projection from elliptic curve  $E$  to  $\mathbb{P}^1$  is any degree-two branched cover that identifies each point  $P \in E$  with its inverse  $-P$ .

**COROLLARY 8.2.** *There exists a uniform bound  $B$  such that*

$$|\pi_1(E_1^{\text{tors}}) \cap \pi_2(E_2^{\text{tors}})| \leq B$$

for any pair of elliptic curves  $E_i$  over  $\mathbb{C}$  and any pair of standard projections  $\pi_i$  for which

$$|\pi_1(E_1[2]) \cap \pi_2(E_2[2])| = 3.$$

*Proof.* By fixing coordinates on  $\mathbb{P}^1$ , we may assume that  $\pi_1(E_1[2]) \cap \pi_2(E_2[2]) = \{0, 1, \infty\}$ . For each  $e \in E_i[2]$ , the composition  $\pi_i^e(P) = \pi_i(P+e)$  is again a standard projection and satisfies  $\pi_i^e(E_i^{\text{tors}}) = \pi_i(E_i^{\text{tors}})$ . Therefore, we may assume that  $\pi_i(O_i) = \infty$  for the origin  $O_i$  of  $E_i$ ,  $i = 1, 2$ . Putting each  $E_i$  into Legendre form now shows that the corollary follows from [Theorem 1.5](#).  $\square$

8.4. *No uniform bound on the torsion order.* We conclude this section by observing that there cannot be a uniform bound on the order of the torsion points that lie in the intersections in [Theorem 1.5](#).

**PROPOSITION 8.3.** *Let  $E_t$  be the Legendre elliptic curve with equation [\(1.2\)](#) and  $\pi(x, y) = x$ . For every  $N > 0$  and for every  $x_0 \in \overline{\mathbb{Q}} \setminus \{0, 1\}$ , there exist  $t_1 \neq t_2$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$  so that the points*

$$P_i = \left( x_0, \pm \sqrt{x_0(x_0 - 1)(x_0 - t_i)} \right)$$

*are torsion points on  $E_{t_i}$  of order  $\geq N$  for  $i = 1, 2$ .*

*Proof.* Fix  $x_0 \in \overline{\mathbb{Q}} \setminus \{0, 1\}$  and  $N > 0$ . Let

$$P_t = \left( x_0, \pm \sqrt{x_0(x_0 - 1)(x_0 - t)} \right)$$

be the points on  $E_t$  with  $x$ -coordinate equal to  $x_0$ . The set

$$\text{Tor}(x_0) := \{t \in \overline{\mathbb{Q}} \setminus \{0, 1\} : P_t \text{ is torsion on } E_t\}$$

is an infinite but proper subset of  $\overline{\mathbb{Q}} \setminus \{0, 1\}$ . See, for example, [\[DWY16, Th. 1.2\]](#) for a proof, taking  $a = b = x_0$  in the statement of the theorem. The fact that  $P_t$  is not torsion for every  $t$  is a consequence of [Proposition 1.4](#) of [\[DWY16\]](#) and the remark that follows it; it follows that  $\text{Tor}(x_0)$  is a set of bounded height, and so not equal to all of  $\overline{\mathbb{Q}} \setminus \{0, 1\}$ . Consequently, there are only finitely many elements of  $\text{Tor}(x_0)$  corresponding to points  $P_t$  of order  $\leq N$ , so we can find  $t_1 \neq t_2$  where the point  $P_{t_i}$  has order at least  $N$ . This proves the proposition.  $\square$

## 9. Proof of Theorems 1.1 and 1.10

Throughout this section, we let  $\mathcal{L}_2$  denote the hypersurface in the moduli space  $\mathcal{M}_2$  consisting of all genus 2 curves  $X$  over  $\mathbb{C}$  that admit a degree-two map to an elliptic curve; see, e.g., [\[SV04\]](#) for details on  $\mathcal{L}_2$ . The surface  $\mathcal{L}_2$  consists of all  $X$  whose Jacobians admit real multiplication by the real quadratic order of discriminant 4, as explained in the proof of [\[McM07, Th. 4.10\]](#).

For any smooth, compact, genus 2 curve over  $\mathbb{C}$ , and for any Weierstrass point  $P$  on  $X$ ,

$$|j_P(X) \cap J(X)^{\text{tors}}| \geq 6$$

as the difference of two Weierstrass points is torsion. On the other hand, any curve  $X$  of genus  $g \geq 2$  has  $|j_P(X) \cap J(X)^{\text{tors}}| \leq 2$  for all but finitely many  $P$ , by Baker and Poonen [BP01], so an Abel-Jacobi map based at a Weierstrass point has in this sense a large number of torsion images.

In this section we deduce [Theorem 1.1](#) from [Corollary 8.2](#), providing a uniform upper bound on  $|j_P(X) \cap J(X)^{\text{tors}}|$  for all  $X$  in  $\mathcal{L}_2$ . We also deduce [Theorem 1.10](#) from [Theorem 8.1](#).

**9.1. Genus 2 curve from a pair of elliptic curves.** Suppose  $\pi_1 : E_1 \rightarrow \mathbb{P}^1$  and  $\pi_2 : E_2 \rightarrow \mathbb{P}^1$  are standard projections on elliptic curves  $E_i$  such that

$$|\pi_1(E_1[2]) \cap \pi_2(E_2[2])| = 3,$$

as in [Corollary 8.2](#). Recall that standard projections are degree-two branched covers  $\pi : E \rightarrow \mathbb{P}^1$  such that  $\pi(P) = \pi(-P)$  for all points  $P \in E$ , and so they have simple critical points at the four points of  $E[2]$ . Consider the diagonal  $D \subset \mathbb{P}^1 \times \mathbb{P}^1$ , and lift  $D$  to a curve  $C \subset E_1 \times E_2$  via  $\Pi = \pi_1 \times \pi_2$ . Let  $\nu : X \rightarrow C$  normalize  $C$ , noting that the degree four map  $\Pi \circ \nu : X \rightarrow D$  has branch locus  $\pi_1(E_1[2]) \cup \pi_2(E_2[2])$ , with each branch point the image of two points in  $X$ , each of multiplicity two. By Riemann-Hurwitz, the genus of  $X$  is 2, and by construction, the curve  $X$  is in  $\mathcal{L}_2$  in  $\mathcal{M}_2$ . Note that  $X$  maps to both of the elliptic curves  $E_1$  and  $E_2$  with degree 2.

**9.2. A pair of elliptic curves from a genus 2 curve.** Here we observe that every  $X \in \mathcal{L}_2$  arises from the construction described in [Section 9.1](#). In particular, admitting a degree-two branched cover  $X \rightarrow E_1$  to an elliptic curve  $E_1$  implies that  $X$  also admits a second degree-two branched cover  $X \rightarrow E_2$ . The proof of the following proposition shows how the curve  $E_2$  arises:

**PROPOSITION 9.1.** *Every  $X \in \mathcal{L}_2$  is the lift of the diagonal under a product of standard projections  $\pi_i$  on elliptic curves  $E_i$  for which*

$$|\pi_1(E_1[2]) \cap \pi_2(E_2[2])| = 3.$$

*Moreover, there are a Weierstrass point  $Q \in X(\mathbb{C})$  and a degree-four isogeny  $\Phi : J(X) \rightarrow E_1 \times E_2$  such that*

$$\Phi \circ j_Q(X) = (\pi_1 \times \pi_2)^{-1}D \text{ in } E_1 \times E_2,$$

*where  $D$  is the diagonal in  $\mathbb{P}^1 \times \mathbb{P}^1$ ,  $J(X)$  is the Jacobian of  $X$ , and  $j_Q$  is the Abel-Jacobi embedding associated to  $Q$ .*

*Proof.* As noted by [SV04] and attributed to Jacobi [Jac32], each curve  $X \in \mathcal{L}_2$  has an affine model

$$C : y^2 = x^6 - s_1x^4 + s_2x^2 - 1,$$

where the polynomial on the right has non-zero discriminant. Here  $C$  admits degree two maps  $(x, y) \mapsto (x^2, y)$  and  $(x, y) \mapsto (1/x^2, iy/x^3)$  to elliptic curves with affine presentation

$$E_1 : y^2 = x^3 - s_1x^2 + s_2x - 1$$

and

$$E_2 : y^2 = x^3 - s_2x^2 + s_1x - 1,$$

respectively, defining a map  $\nu : X \rightarrow E_1 \times E_2$ . For each of these curves, the  $x$ -coordinate projection  $\pi_x$  is standard, so  $\pi_1 := \pi_x$  and  $\pi_2 := 1/\pi_x$  are standard projections for  $E_1$  and  $E_2$  respectively. The projection  $\pi_1$  ramifies over  $\{\infty, r_1, r_2, r_3\}$  and  $\pi_2$  ramifies over  $\{0, r_1, r_2, r_3\}$ , where  $\{r_1, r_2, r_3\}$  are the distinct, nonzero roots of  $x^3 - s_1x^2 + s_2x - 1$ . Thus

$$|\pi_1(E_1[2]) \cap \pi_2(E_2[2])| = 3.$$

Define  $\Pi := \pi_1 \times \pi_2$ , noting that for  $(x, y) \in C$ , we have

$$\Pi \circ \nu(x, y) = \Pi(x^2, 1/x^2) = (x^2, x^2).$$

Thus  $\Pi \circ \nu(X) = D$ , where  $D \subset \mathbb{P}^1 \times \mathbb{P}^1$  is the diagonal.

Fix  $r \in \pi_1(E_1[2]) \cap \pi_2(E_2[2])$ , and equip each  $E_i$  with a group structure such that the identity lies above  $r$ . Observe that the  $[-1]$ -involution on  $E_1 \times E_2$  induces the hyperelliptic involution on  $X$ . In particular, the Weierstrass points on  $X$  are the six preimages of  $\pi_1(E_1[2]) \cap \pi_2(E_2[2])$  under  $\Pi \circ \nu$ . Choose  $Q \in X$  such that  $\Pi(\nu(Q)) = (r, r)$ , so that  $Q$  is Weierstrass and  $\nu$  factors as  $\Phi \circ j_Q$  for some isogeny  $\Phi : J(X) \rightarrow E_1 \times E_2$ . The nontrivial elements of the kernel of  $\Phi$  are precisely the three 2-torsion points in  $J(X)$  that are differences of Weierstrass points mapping to the same point in the diagonal  $D \subset \mathbb{P}^1 \times \mathbb{P}^1$ . Thus  $\Phi$  is degree four as claimed, completing the proof.  $\square$

**9.3. Proof of Theorem 1.1.** Fix  $X \in \mathcal{L}_2$ . From [Proposition 9.1](#), we have elliptic curves  $E_1$  and  $E_2$  and a Weierstrass point  $Q \in X$  such that

$$|j_Q(X) \cap J(X)^{\text{tors}}| \leq 16 |\pi_1(E_1^{\text{tors}}) \cap \pi_2(E_2^{\text{tors}})|$$

for a pair of standard projections  $\pi_i : E_i \rightarrow \mathbb{P}^1$  satisfying  $|\pi_1(E_1[2]) \cap \pi_2(E_2[2])| = 3$ . Given any other Weierstrass point  $P \in X$ , we have  $[P - Q] \in J(X)^{\text{tors}}$ , so we conclude that

$$|j_P(X) \cap J(X)^{\text{tors}}| = |j_Q(X) \cap J(X)^{\text{tors}}| \leq 16B,$$

where  $B$  is the constant of [Corollary 8.2](#).

9.4. *Proof of Theorem 1.10.* Fix  $X \in \mathcal{L}_2 \subset \mathcal{M}_2$ , defined over  $\overline{\mathbb{Q}}$ . From [Proposition 9.1](#) there are a pair  $t_1 \neq t_2$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$  and an isogeny  $\Phi : J(X) \rightarrow E_{t_1} \times E_{t_2}$  of degree 4 so that  $\Pi \circ \Phi \circ j_Q(X)$  is the diagonal in  $\mathbb{P}^1 \times \mathbb{P}^1$ , where  $\Pi = \pi \times \pi$  and  $Q$  is a Weierstrass point on  $X$ . Recall from [Section 2.1](#) that the Néron-Tate canonical height on  $\hat{h}_{E_t}$  on  $E_t$  satisfies

$$\hat{h}_{E_t}(P) = \frac{1}{2} \hat{h}_t(\pi(P))$$

for all  $P \in E_t(\overline{\mathbb{Q}})$  and each  $t \in \overline{\mathbb{Q}} \setminus \{0, 1\}$ .

Let

$$D = \{O_1\} \times E_{t_2} + E_{t_1} \times \{O_2\}$$

be a divisor on  $E_{t_1} \times E_{t_2}$ , where  $O_i$  denotes the identity element of  $E_{t_i}$ , and let  $L_D$  be the associated line bundle. Let  $L_X = \Phi^* L_D$  on  $J(X)$ , and let  $\hat{h}_{L_X}$  be the associated Néron-Tate canonical height on  $J(X)(\overline{\mathbb{Q}})$ . By the functoriality of canonical heights [[HS00](#), Th. B.5.6], we have

$$\begin{aligned} \hat{h}_{L_X}(x) &= \hat{h}_{L_D}(\Phi(x)) \\ &= \hat{h}_{E_{t_1}}(\Phi(x)_1) + \hat{h}_{E_{t_2}}(\Phi(x)_2) \\ &= \frac{1}{2} \left( \hat{h}_{t_1}(\pi(\Phi(x)_1)) + \hat{h}_{t_2}(\pi(\Phi(x)_2)) \right), \end{aligned}$$

where  $\Phi(x) = (\Phi(x)_1, \Phi(x)_2)$  in  $E_{t_1} \times E_{t_2}$ . Restricting to the points  $x \in j_P(X)(\overline{\mathbb{Q}})$ , so that  $\pi(\Phi(x)_1) = \pi(\Phi(x)_2)$  in  $\mathbb{P}^1$ , the theorem now follows from [Theorem 8.1](#).  $\square$

9.5. *No uniform bound on the torsion order.* We conclude the article with the observation that there cannot be a uniform bound on the *order* of the torsion points that lie in the intersections in [Theorem 1.1](#). It is an immediate consequence of [Propositions 8.3](#) and [9.1](#).

**PROPOSITION 9.2.** *Given any  $N > 0$ , there exist a curve  $X \in \mathcal{L}_2$  defined over  $\overline{\mathbb{Q}}$  and a Weierstrass point  $P \in X$  so that the intersection*

$$j_P(X) \cap J(X)^{\text{tors}}$$

*contains a point of order  $\geq N$  in the Jacobian  $J(X)$ .*

*Proof.* Fix  $N > 0$ . From [Proposition 8.3](#), we can find  $t_1 \neq t_2$  in  $\overline{\mathbb{Q}} \setminus \{0, 1\}$  and torsion points  $P_{t_i} \in E_{t_i}$ ,  $i = 1, 2$ , having the same  $x$ -coordinate and torsion orders  $\geq N$ . Let  $x_0$  denote the  $x$ -coordinate of  $P_{t_i}$ . Let  $X$  be the normalization of the curve  $C \subset E_{t_1} \times E_{t_2}$  that is the lift of the diagonal in  $\mathbb{P}^1 \times \mathbb{P}^1$  under the projection  $\pi \times \pi$ . From [Proposition 9.1](#), we know that  $X$  is of genus 2 and lies in  $\mathcal{L}_2$ , and there is a Weierstrass point  $Q \in X$  so that  $j_Q(X)$  maps to the curve  $C$  under a degree-four isogeny  $\Phi : J(X) \rightarrow E_{t_1} \times E_{t_2}$ . Let  $w$  be a preimage in  $J(X)$  of the point  $(x_0, x_0)$  in  $\mathbb{P}^1 \times \mathbb{P}^1$  under the map  $(\pi \times \pi) \circ \Phi$ . Then  $w$  has order at least  $N$ .  $\square$

## References

- [BP05] M. BAKER and C. PETSCHE, Global discrepancy and small points on elliptic curves, *Int. Math. Res. Not.* **61** (2005), 3791–3834. [MR 2205235](#). [Zbl 1120.11025](#). <https://doi.org/10.1155/IMRN.2005.3791>.
- [BP01] M. BAKER and B. POONEN, Torsion packets on curves, *Compositio Math.* **127** no. 1 (2001), 109–116. [MR 1832989](#). [Zbl 0987.14020](#). <https://doi.org/10.1023/A:1017557230785>.
- [BR10] M. BAKER and R. RUMELY, *Potential Theory and Dynamics on the Berkovich Projective Line*, *Math. Surveys Monogr.* **159**, Amer. Math. Soc., Providence, RI, 2010. [MR 2599526](#). [Zbl 1196.14002](#). <https://doi.org/10.1090/surv/159>.
- [Ber90] V. G. BERKOVICH, *Spectral Theory and Analytic Geometry over Non-Archimedean Fields*, *Math. Surveys Monogr.* **33**, Amer. Math. Soc., Providence, RI, 1990. [MR 1070709](#). [Zbl 0715.14013](#). <https://doi.org/10.1090/surv/033>.
- [BFT18] F. BOGOMOLOV, H. FU, and Y. TSCHINKEL, Torsion of elliptic curves and unlikely intersections, in *Geometry and Physics. Vol. I*, Oxford Univ. Press, Oxford, 2018, pp. 19–37. [MR 3932255](#). [Zbl 1423.14214](#). <https://doi.org/10.1093/oso/9780198802013.003.0002>.
- [BT07] F. BOGOMOLOV and Y. TSCHINKEL, Algebraic varieties over small fields, in *Diophantine Geometry, CRM Series* **4**, Ed. Norm., Pisa, 2007, pp. 73–91. [MR 2349648](#). [Zbl 1142.14016](#).
- [BJ17] S. BOUCKSOM and M. JONSSON, Tropical and non-Archimedean limits of degenerating families of volume forms, *J. Éc. Polytech. Math.* **4** (2017), 87–139. [MR 3611100](#). [Zbl 1401.32019](#). <https://doi.org/10.5802/jep.39>.
- [Bui96] A. BUIUM, Geometry of  $p$ -jets, *Duke Math. J.* **82** no. 2 (1996), 349–367. [MR 1387233](#). [Zbl 0882.14007](#). <https://doi.org/10.1215/S0012-7094-96-08216-2>.
- [CS93] G. S. CALL and J. H. SILVERMAN, Canonical heights on varieties with morphisms, *Compositio Math.* **89** no. 2 (1993), 163–205. [MR 1255693](#). [Zbl 0826.14015](#). Available at [http://www.numdam.org/item?id=CM\\_1993\\_89\\_2\\_163\\_0](http://www.numdam.org/item?id=CM_1993_89_2_163_0).
- [CL06] A. CHAMBERT-LOIR, Mesures et équidistribution sur les espaces de Berkovich, *J. Reine Angew. Math.* **595** (2006), 215–235. [MR 2244803](#). [Zbl 1112.14022](#). <https://doi.org/10.1515/CRELLE.2006.049>.
- [CL11] A. CHAMBERT-LOIR, Heights and measures on analytic spaces. A survey of recent results, and some remarks, in *Motivic Integration and its Interactions with Model Theory and Non-Archimedean Geometry. Vol. II*, *London Math. Soc. Lecture Note Ser.* **384**, Cambridge Univ. Press, Cambridge, 2011, pp. 1–50. [MR 2885340](#). [Zbl 1279.14027](#). <https://doi.org/10.1017/CBO9780511984433.002>.
- [Col85] R. F. COLEMAN, Torsion points on curves and  $p$ -adic abelian integrals, *Ann. of Math.* (2) **121** no. 1 (1985), 111–168. [MR 0782557](#). [Zbl 0578.14038](#). <https://doi.org/10.2307/1971194>.

- [DMF14] L. DE MARCO and X. FABER, Degenerations of complex dynamical systems, *Forum Math. Sigma* **2** (2014), e6, 36. [MR 3264250](#). [Zbl 1308.37023](#). <https://doi.org/10.1017/fms.2014.8>.
- [DF16] L. DEMARCO and X. FABER, Degenerations of complex dynamical systems II: analytic and algebraic stability, *Math. Ann.* **365** no. 3-4 (2016), 1669–1699, With an appendix by Jan Kiwi. [MR 3521102](#). [Zbl 1343.37089](#). <https://doi.org/10.1007/s00208-015-1331-8>.
- [DKY19] L. DEMARCO, H. KRIEGER, and H. YE, Common preperiodic points of quadratic polynomial, 2019. [arXiv 1911.02458v1](#).
- [DWY16] L. DEMARCO, X. WANG, and H. YE, Torsion points and the Lattès family, *Amer. J. Math.* **138** no. 3 (2016), 697–732. [MR 3506383](#). [Zbl 1364.37101](#). <https://doi.org/10.1353/ajm.2016.0026>.
- [Fav16] C. FAVRE, Degeneration of endomorphisms of the complex projective space in the hybrid space, 2016. [arXiv 1611.08490v1](#).
- [FRL06] C. FAVRE and J. RIVERA-LETELIER, Équidistribution quantitative des points de petite hauteur sur la droite projective, *Math. Ann.* **335** no. 2 (2006), 311–361. [MR 2221116](#). [Zbl 1175.11029](#). <https://doi.org/10.1007/s00208-006-0751-x>.
- [FRL10] C. FAVRE and J. RIVERA-LETELIER, Théorie ergodique des fractions rationnelles sur un corps ultramétrique, *Proc. Lond. Math. Soc.* (3) **100** no. 1 (2010), 116–154. [MR 2578470](#). [Zbl 1254.37064](#). <https://doi.org/10.1112/plms/pdp022>.
- [Fil17] P. FILI, A metric of mutual energy and unlikely intersections for dynamical systems, 2017. [arXiv 1708.08403v1](#).
- [HS00] M. HINDRY and J. H. SILVERMAN, *Diophantine Geometry, Grad. Texts in Mathematics* **201**, Springer-Verlag, New York, 2000, An introduction. [MR 1745599](#). [Zbl 0948.11023](#). <https://doi.org/10.1007/978-1-4612-1210-2>.
- [Hru01] E. HRUSHOVSKI, The Manin–Mumford conjecture and the model theory of difference fields, *Ann. Pure Appl. Logic* **112** no. 1 (2001), 43–115. [MR 1854232](#). [Zbl 0987.03036](#). [https://doi.org/10.1016/S0168-0072\(01\)00096-3](https://doi.org/10.1016/S0168-0072(01)00096-3).
- [Jac32] C. G. J. JACOBI, Nachrichten von Büchern, *J. Reine Angew. Math.* **8** (1832), 413–417. <https://doi.org/10.1515/crll.1832.8.413>.
- [KRZB16] E. KATZ, J. RABINOFF, and D. ZUREICK-BROWN, Uniform bounds for the number of rational points on curves of small Mordell-Weil rank, *Duke Math. J.* **165** no. 16 (2016), 3189–3240. [MR 3566201](#). [Zbl 06666955](#). <https://doi.org/10.1215/00127094-3673558>.
- [KS09] S. KAWAGUCHI and J. H. SILVERMAN, Canonical heights and the arithmetic complexity of morphisms on projective space, *Pure Appl. Math. Q.* **5** no. 4, Special Issue: In honor of John Tate. Part 1 (2009), 1201–1217. [MR 2560315](#). [Zbl 1221.11152](#). <https://doi.org/10.4310/PAMQ.2009.v5.n4.a2>.

- [Maz86] B. MAZUR, Arithmetic on curves, *Bull. Amer. Math. Soc. (N.S.)* **14** no. 2 (1986), 207–259. [MR 0828821](#). [Zbl 0593.14021](#). <https://doi.org/10.1090/S0273-0979-1986-15430-3>.
- [McM07] C. T. McMULLEN, Dynamics of  $\mathrm{SL}_2(\mathbb{R})$  over moduli space in genus two, *Ann. of Math. (2)* **165** no. 2 (2007), 397–456. [MR 2299738](#). [Zbl 1131.14027](#). <https://doi.org/10.4007/annals.2007.165.397>.
- [Mil06] J. MILNOR, On Lattès maps, in *Dynamics on the Riemann Sphere*, Eur. Math. Soc., Zürich, 2006, pp. 9–43. [MR 2348953](#). [Zbl 1235.37015](#). <https://doi.org/10.4171/011-1/1>.
- [PST12] C. PETSCHE, L. SZPIRO, and T. J. TUCKER, A dynamical pairing between two rational maps, *Trans. Amer. Math. Soc.* **364** no. 4 (2012), 1687–1710. [MR 2869188](#). [Zbl 1300.37052](#). <https://doi.org/10.1090/S0002-9947-2011-05350-X>.
- [Poo00] B. POONEN, Genus-two curves with 22 torsion points, *C. R. Acad. Sci. Paris Sér. I Math.* **330** no. 7 (2000), 573–576. [MR 1760441](#). [Zbl 0966.11026](#). [https://doi.org/10.1016/S0764-4442\(00\)00222-6](https://doi.org/10.1016/S0764-4442(00)00222-6).
- [Ray83] M. RAYNAUD, Courbes sur une variété abélienne et points de torsion, *Invent. Math.* **71** no. 1 (1983), 207–233. [MR 0688265](#). [Zbl 0564.14020](#). <https://doi.org/10.1007/BF01393342>.
- [SV04] T. SHASKA and H. VÖLKLEIN, Elliptic subfields and automorphisms of genus 2 function fields, in *Algebra, Arithmetic and Geometry with Applications* (West Lafayette, IN, 2000), Springer, Berlin, 2004, pp. 703–723. [MR 2037120](#). [Zbl 1093.14036](#). [https://doi.org/10.1007/978-3-642-18487-1\\_42](https://doi.org/10.1007/978-3-642-18487-1_42).
- [Sil09] J. H. SILVERMAN, *The Arithmetic of Elliptic Curves*, second ed., *Grad. Texts in Math.* **106**, Springer, Dordrecht, 2009. [MR 2514094](#). [Zbl 1194.11005](#). <https://doi.org/10.1007/978-0-387-09494-6>.
- [Sto] M. STOLL, Another New Record, available on the author’s website.
- [SUZ97] L. SZPIRO, E. ULLMO, and S. ZHANG, Équirépartition des petits points, *Invent. Math.* **127** no. 2 (1997), 337–347. [MR 1427622](#). [Zbl 0991.11035](#). <https://doi.org/10.1007/s002220050123>.
- [Ull98] E. ULLMO, Positivité et discrétion des points algébriques des courbes, *Ann. of Math. (2)* **147** no. 1 (1998), 167–179. [MR 1609514](#). [Zbl 0934.14013](#). <https://doi.org/10.2307/120987>.
- [Zha98] S.-W. ZHANG, Equidistribution of small points on abelian varieties, *Ann. of Math. (2)* **147** no. 1 (1998), 159–165. [MR 1609518](#). [Zbl 0991.11034](#). <https://doi.org/10.2307/120986>.
- [Zha95] S. ZHANG, Small points and adelic metrics, *J. Algebraic Geom.* **4** no. 2 (1995), 281–300. [MR 1311351](#). [Zbl 0861.14019](#).

(Received: January 27, 2019)  
(Revised: November 26, 2019)

NORTHWESTERN UNIVERSITY, EVANSTON, IL, USA  
*E-mail:* [demarco@northwestern.edu](mailto:demarco@northwestern.edu)

UNIVERSITY OF CAMBRIDGE, CAMBRIDGE, UK  
*E-mail:* [hkrieger@dpmms.cam.ac.uk](mailto:hkrieger@dpmms.cam.ac.uk)

ZHEJIANG UNIVERSITY, HANGZHOU, CHINA  
*E-mail:* [yehexi@gmail.com](mailto:yehexi@gmail.com)