

Minimal p -divisible groups

By FRANS OORT

Introduction

A p -divisible group X can be seen as a tower of building blocks, each of which is isomorphic to the same finite group scheme $X[p]$. Clearly, if X_1 and X_2 are isomorphic then $X_1[p] \cong X_2[p]$; however, conversely $X_1[p] \cong X_2[p]$ does in general not imply that X_1 and X_2 are isomorphic. Can we give, over an algebraically closed field in characteristic p , a condition on the p -kernels which ensures this converse? Here are two known examples of such a condition: consider the case that X is *ordinary*, or the case that X is *superspecial* (X is the p -divisible group of a product of supersingular elliptic curves); in these cases the p -kernel uniquely determines X .

These are special cases of a surprisingly complete and simple answer:

If G is “minimal”, then $X_1[p] \cong G \cong X_2[p]$ implies $X_1 \cong X_2$;

see (1.2); for a definition of “minimal” see (1.1). This is “necessary and sufficient” in the sense that for any G that is *not minimal* there exist infinitely many mutually nonisomorphic p -divisible groups with p -kernel isomorphic to G ; see (4.1).

Remark (motivation). You might wonder why this is interesting.

EO. In [7] we defined a natural *stratification* of the moduli space of polarized abelian varieties in positive characteristic: moduli points are in the same stratum if and only if the corresponding p -kernels are geometrically isomorphic. Such strata are called EO-strata.

Fol. In [8] we define in the same moduli spaces a *foliation*: Moduli points are in the same leaf if and only if the corresponding p -divisible groups are geometrically isomorphic; in this way we obtain a foliation of every open Newton polygon stratum.

Fol \subset **EO.** The observation $X \cong Y \Rightarrow X[p] \cong Y[p]$ shows that any leaf in the second sense is contained in precisely one stratum in the first sense; the main result of this paper, “ X is minimal if and only if $X[p]$ is minimal”,

shows that *a stratum* (in the first sense) *and a leaf* (in the second sense) *are equal* in the minimal, principally polarized situation.

In this paper we consider p -divisible groups and finite group schemes over an *algebraically closed* field k of characteristic p .

An apology. In (2.5) and in (3.5) we fix notation, used for the proof of (2.2), respectively (3.1); according to the need, the notation in these two different cases is different. We hope this difference in notation in Section 2 versus Section 3 will not cause confusion.

Group schemes considered are supposed to be commutative. We use *covariant* Dieudonné module theory and write $W = W_\infty(k)$ for the ring of infinite Witt vectors with coordinates in k . Finite products in the category of W -modules are denoted “ \times ” or by “ \prod ”, while finite products in the category of Dieudonné modules are denoted by “ \oplus ”; for finite products of p -divisible groups we use “ \times ” or “ \prod ”. We write F and V , as usual, for “Frobenius” and “Verschiebung” on commutative group schemes and let $\mathcal{F} = \mathbb{D}(V)$ and $\mathcal{V} = \mathbb{D}(F)$; see [7, 15.3], for the corresponding operations on Dieudonné modules.

Acknowledgments. Part of the work for this paper was done while visiting Université de Rennes, and the Massachusetts Institute of Technology; I thank the Mathematics Departments of these universities for hospitality and stimulating working environment. I thank Bas Edixhoven and Johan de Jong for discussions on ideas necessary for this paper. I thank the referee for helpful, critical remarks.

1. Notation and the main result

(1.1) *Definitions and notation.*

$H_{m,n}$. We define the p -divisible group $H_{m,n}$ over the prime field \mathbb{F}_p in case m and n are coprime nonnegative integers; see [2, 5.2]. This p -divisible group $H_{m,n}$ is of dimension m , its Serre-dual X^t is of dimension n , it is isosimple, and *its endomorphism ring* $\text{End}(H_{m,n} \otimes \overline{\mathbb{F}_p})$ *is the maximal order in the endomorphism algebra* $\text{End}^0(H_{m,n} \otimes \overline{\mathbb{F}_p})$ (and these properties characterize this p -divisible group over $\overline{\mathbb{F}_p}$). We will use the notation $H_{m,n}$ over any base S in characteristic p ; i.e., we write $H_{m,n}$ instead of $H_{m,n} \times_{\text{Spec}(\mathbb{F}_p)} S$, if no confusion can occur.

The ring $\text{End}(H_{m,n} \otimes \mathbb{F}_p) = R'$ is commutative; write L for the field of fractions of R' . Consider integers x, y such that for the coprime positive integers m and n we have $x \cdot m + y \cdot n = 1$. In L we define the element $\pi = \mathcal{F}^y \cdot \mathcal{V}^x \in L$. Write $h = m + n$. Note that $\pi^h = p$ in L . Here $R' \subset L$ is the maximal order; hence R' is integrally closed in L , and we conclude that $\pi \in R'$.

This element π will be called the uniformizer in this endomorphism ring. In fact, $W_\infty(\mathbb{F}_p) = \mathbb{Z}_p$, and $R' \cong \mathbb{Z}_p[\pi]$. In L we have:

$$m + n =: h, \quad \pi^h = p, \quad \mathcal{F} = \pi^n, \quad \mathcal{V} = \pi^m.$$

For a further description of π , of $R = \text{End}(H_{m,n} \otimes k)$ and of $D = \text{End}^0(H_{m,n} \otimes k)$, see [2, 5.4]; note that $\text{End}^0(H_{m,n} \otimes k)$ is noncommutative if $m > 0$ and $n > 0$. Note that R is a “discrete valuation ring” (terminology sometimes also used for noncommutative rings).

Newton polygons. Let β be a Newton polygon. By definition, in the notation used here, this is a lower convex polygon in \mathbb{R}^2 starting at $(0, 0)$, ending at (h, c) and having break points with integral coordinates; it is given by h slopes in nondecreasing order; every slope λ is a rational number, $0 \leq \lambda \leq 1$.

To each ordered pair of nonnegative integers (m, n) we assign a set of $m + n = h$ slopes equal to $n/(m + n)$; this Newton polygon ends at $(h, c = n)$.

In this way a Newton polygon corresponds with a set of ordered pairs; such a set we denote symbolically by $\sum_i (m_i, n_i)$; conversely such a set determines a Newton polygon. Usually we consider only coprime pairs (m_i, n_i) ; we write $H(\beta) := \times_i H_{m_i, n_i}$ in case $\beta = \sum_i (m_i, n_i)$. A p -divisible group X over a field of positive characteristic defines a Newton polygon where h is the height of X and c is the dimension of its Serre-dual X^t . By the Dieudonné-Manin classification, see [5, Th. 2.1, p. 32], we know: *Two p -divisible groups over an algebraically closed field of positive characteristic are isogenous if and only if their Newton polygons are equal.*

Definition. A p -divisible group X is called *minimal* if there exists a Newton polygon β and an isomorphism $X_k \cong H(\beta)_k$, where k is an algebraically closed field.

Note that in every isogeny class of p -divisible groups over an algebraically closed field there is precisely one minimal p -divisible group.

Truncated p -divisible groups. A finite group scheme G (finite and flat over some base, but in this paper we will soon work over a field) is called a BT_1 , see [1, p. 152], if $G[\mathbb{F}] := \text{Ker}F_G = \text{Im}V_G =: V(G)$ and $G[V] = F(G)$ (in particular this implies that G is annihilated by p). Such group schemes over a perfect field appear as the p -kernel of a p -divisible group, see [1, Prop. 1.7, p. 155]. The abbreviation “ BT_1 ” stand for “1-truncated Barsotti-Tate group”; the terms “ p -divisible group” and “Barsotti-Tate group” indicate the same concept.

The Dieudonné module of a BT_1 over a perfect field K is called a DM_1 ; for $G = X[p]$ we have $\mathbb{D}(G) = \mathbb{D}(X)/p\mathbb{D}(X)$. In other terms: such a Dieudonné module $M_1 = \mathbb{D}(X[p])$ is a finite dimensional vector space over K , on which

\mathcal{F} and \mathcal{V} operate (with the usual relations), with the property that $M_1[\mathcal{V}] = \mathbb{F}(M_1)$ and $M_1[\mathcal{F}] = \mathbb{V}(M_1)$.

Definition. Let G be a BT_1 group scheme; we say that G is *minimal* if there exists a Newton polygon β such that $G_k \cong H(\beta)[p]_k$. A DM_1 is called *minimal* if it is the Dieudonné module of a minimal BT_1 .

(1.2) **THEOREM.** *Let X be a p -divisible group over an algebraically closed field k of characteristic p . Let β be a Newton polygon. Then*

$$X[p] \cong H(\beta)[p] \implies X \cong H(\beta).$$

In particular: if X_1 and X_2 are p -divisible groups over k , with $X_1[p] \cong G \cong X_2[p]$, where G is minimal, then $X_1 \cong X_2$.

Remark. We have no *a priori* condition on the Newton polygon of X , nor do we *a priori* assume that X_1 and X_2 have the same Newton polygon.

Remark. In general an isomorphism $\varphi_1 : X[p] \rightarrow H(\beta)[p]$ does not lift to an isomorphism $\varphi : X \rightarrow H(\beta)$.

(1.3) Here is another way of explaining the result of this paper. Consider the map

$$[p] : \{X \mid \text{a } p\text{-divisible group}\} / \cong_k \longrightarrow \{G \mid \text{a } \mathrm{BT}_1\} / \cong_k, \quad X \mapsto X[p].$$

This map is surjective; e.g., see [1, 1.7]; also see [7, 9.10].

- By results of this paper we know: For every Newton polygon β there is an isomorphism class $X := H(\beta)$ such that the fiber of the map $[p]$ containing X consists of one element.
- For every X not isomorphic to some $H(\beta)$ the fiber of $[p]$ containing X is infinite; see (4.1)

Convention. The slope $\lambda = 0$, given by the pair $(1, 0)$, defines the p -divisible group $G_{1,0} = \mathbb{G}_m[p^\infty]$, and its p -kernel is μ_p . The slope $\lambda = 1$, given by the pair $(0, 1)$, defines the p -divisible group $G_{0,1} = \mathbb{Q}_p/\mathbb{Z}_p$ and its p -kernel is $\mathbb{Z}/p\mathbb{Z}$. These p -divisible groups and their p -kernels split off naturally over a perfect field; see [6, 2.14]. The theorem is obvious for these minimal BT_1 group schemes over an algebraically closed field. Hence it suffices to prove the theorem in case all group schemes considered are of local-local type, i.e. all slopes considered are strictly between 0 and 1; from now on we make this assumption.

(1.4) We give now one explanation about notation and method of proof. Let $m, n \in \mathbb{Z}_{>0}$ be coprime. Start with $H_{m,n}$ over \mathbb{F}_p . Let $Q' = \mathbb{D}(H_{m,n} \otimes \mathbb{F}_p)$. In the terminology of [2, 5.6 and §6], a semi-module of $H_{m,n}$ equals $[0, \infty) = \mathbb{Z}_{\geq 0}$. Choose a nonzero element in $Q'/\pi Q'$; this is a one-dimensional vector space over \mathbb{F}_p , and lift this element to $A_0 \in Q'$. Write $A_i = \pi^i A_0$ for every $i \in \mathbb{Z}_{>0}$. Note that

$$\pi A_i = A_{i+1}, \quad \mathcal{F}A_i = A_{i+n}, \quad \mathcal{V}A_i = A_{i+m}.$$

Fix an algebraically closed field k ; we write $Q = \mathbb{D}(H_{m,n} \otimes k)$. Clearly $A_i \in Q' \subset Q$, and the same relations as given above hold. Note that $\{A_i \mid i \in \mathbb{Z}_{\geq 0}\}$ generate Q as a W -module. *The fact that a semi-module of the minimal p -divisible group $H_{m,n}$ does not contain “gaps” is the essential (but sometimes hidden) argument in the proofs below.*

The set $\{A_0, \dots, A_{m+n-1}\}$ is a W -basis for Q . If $m \geq n$ we see that $\{A_0, \dots, A_{n-1}\}$ is a set of generators for Q as a Dieudonné module; the structure of this Dieudonné module can be described as follows: For this set of generators we consider another numbering $\{C_1, \dots, C_n\} = \{A_0, \dots, A_{n-1}\}$ and define positive integers γ_i by $C_1 = A_0$ and $\mathcal{F}^{\gamma_1} C_1 = \mathcal{V}C_2, \dots, \mathcal{F}^{\gamma_n} C_n = \mathcal{V}C_1$ (note that we assume $m \geq n$), which gives a “cyclic” set of generators for Q/pQ in the sense of [3]. This notation will be repeated and explained more in detail in (2.5) and (3.5).

2. A slope filtration

(2.1) We consider a Newton polygon β given by $r_1(m_1, n_1), \dots, r_t(m_t, n_t)$; here $r_1, \dots, r_t \in \mathbb{Z}_{>0}$, and every (m_j, n_j) is an ordered pair of coprime positive integers; we write $h_j = m_j + n_j$ and suppose the ordering is chosen in such a way that $\lambda_1 := n_1/h_1 < \dots < \lambda_t := n_t/h_t$. Now,

$$H := H(\beta) = \prod_{1 \leq j \leq t} (H_{m_j, n_j})^{r_j}; \quad G := H(\beta)[p].$$

The following proposition uses this notation; suppose that $t > 0$.

(2.2) PROPOSITION. *Suppose X is a p -divisible group over an algebraically closed field k , that $X[p] \cong H(\beta)[p]$, and that $\lambda_1 = n_1/h_1 \leq 1/2$. Then there exists a p -divisible subgroup $X_1 \subset X$ and isomorphisms*

$$X_1 \cong (H_{m_1, n_1})^{r_1} \quad \text{and} \quad (X/X_1)[p] \cong \prod_{j>1} (H_{m_j, n_j}[p])^{r_j}.$$

(2.3) Remark. The condition that $X[p]$ is *minimal* is essential; e.g. it is easy to give an example of a p -divisible group X which is isosimple, such that $X[p]$ is decomposable; see [9].

(2.4) COROLLARY. For X with $X[p] \cong H(\beta)[p]$, with β as in (2.1), there exists a filtration by p -divisible subgroups

$$X_0 := 0 \subset X_1 \subset \cdots \subset X_t = X$$

such that

$$X_j/X_{j-1} \cong (H_{m_j, n_j})^{r_j},$$

for $1 \leq j \leq t$.

Proof of the corollary. Assume by induction that the result has been proved for all p -divisible groups where $Y[p] = H(\beta')[p]$ is minimal such that β' has at most $t - 1$ different slopes; induction starting at $t - 1 = 0$, i.e. $Y = 0$. If on the one hand the smallest slope of X is at most $1/2$, the proposition gives $0 \subset X_1 \subset X$, and using the induction hypothesis on $Y = X/X_1$ we derive the desired filtration. If on the other hand all slopes of X are bigger than $1/2$, we apply the proposition to the Serre-dual of X , using the fact that the Serre-dual of $H_{m,n}$ is $H_{n,m}$; dualizing back we obtain $0 \subset X_{t-1} \subset X$, and using the induction hypothesis on $Y = X_{t-1}$ we derive the desired filtration. Hence we see that the proposition gives the induction step; this proves the corollary.

$$\square(2.2) \Rightarrow (2.4)$$

(2.5) We use notation as in (2.1) and (2.2), and fix further notation which will be used in the proof of (2.2). Let $M = \mathbb{D}(X)$. We write $Q_j = \mathbb{D}(H_{m_j, n_j})$. Hence

$$M/pM \cong \bigoplus_{1 \leq j \leq t} (Q_j/pQ_j)^{r_j}.$$

Using this isomorphism we construct a map

$$v : M \longrightarrow \mathbb{Q}_{\geq 0} \cup \{\infty\}.$$

We use notation as in (1.1) and in (1.4). Let π_j be the uniformizer of $\text{End}(Q_j)$. We choose $A_{i,s}^{(j)} \in Q_j$ with $i \in \mathbb{Z}_{\geq 0}$ and $1 \leq s \leq r_j$ (which generate Q_j) such that $\pi_j \cdot A_{i,s}^{(j)} = A_{i+1,s}^{(j)}$, $\mathcal{F} \cdot A_{i,s}^{(j)} = A_{i+n_j,s}^{(j)}$ and $\mathcal{V} \cdot A_{i,s}^{(j)} = A_{i+m_j,s}^{(j)}$. Now, $Q_j/pQ_j = \times_{0 \leq i < h_j} k \cdot (A_{i,s}^{(j)} \text{ mod } pQ_j)$ and

$$A_i^{(j)} = (A_{i,s}^{(j)} \mid 1 \leq s \leq r_j) \in (Q_j)^{r_j}$$

for the vector with coordinate $A_{i,s}^{(j)}$ in the summand on the s^{th} place.

For $B \in M$ we uniquely write

$$B \text{ mod } pM = a = \sum_{j, 0 \leq i < h_j, 1 \leq s \leq r_j} b_{i,s}^{(j)} \cdot (A_{i,s}^{(j)} \text{ mod } pQ_j), \quad b_{i,s}^{(j)} \in k;$$

if moreover $B \notin pM$ we define

$$v(B) = \min_{j, i, s, b_{i,s}^{(j)} \neq 0} \frac{i}{h_j}.$$

If $B' \in p^\beta M$ and $B' \notin p^{\beta+1} M$ we define $v(B') = \beta + v(p^{-\beta} \cdot B')$ and then write $v(0) = \infty$. This ends the construction of $v : M \rightarrow \mathbb{Q}_{\geq 0} \cup \{\infty\}$.

For any $\rho \in \mathbb{Q}$ we define

$$M_\rho = \{B \mid v(B) \geq \rho\};$$

note that $pM_\rho \subset M_{\rho+1}$. Let T be the least common multiple of h_1, \dots, h_t . Note that, in fact, $v : M - \{0\} \rightarrow \frac{1}{T}\mathbb{Z}_{\geq 0}$ and that, by construction, $v(B) \geq d \in \mathbb{Z}$ if and only if p^d divides B in M . Hence $\bigcap_{\rho \rightarrow \infty} M_\rho = \{0\}$.

The basic assumption $X[p] \cong H(\beta)[p]$ of (1.2) is:

$$M/pM = \bigoplus_{1 \leq j \leq t, 1 \leq s \leq r_j} \prod_{0 \leq i < h_j} k \cdot ((A_{i,s}^{(j)} \bmod pQ_j))$$

(we write this isomorphism of Dieudonné modules as an equality). For $0 \leq i < h_j$ and $1 \leq s \leq r_j$ we choose $B_{i,s}^{(j)} \in M$ such that:

$$B_{i,s}^{(j)} \bmod pM = A_{i,s}^{(j)} \bmod pQ_j^{r_j}.$$

Define $B_{i+\beta \cdot h_j, s}^{(j)} = p^\beta \cdot B_{i,s}^{(j)}$. By construction we have: $v(B_{i,s}^{(j)}) = i/h_j$ for all $i \geq 0$, all j and all s . Note that M_ρ is generated over $W = W_\infty(k)$ by all elements $B_{i,s}^{(j)}$ with $v(B_{i,s}^{(j)}) \geq \rho$. Using shorthand we write

$$B_i^{(j)} \text{ for the vector } (B_{i,s}^{(j)} \mid 1 \leq s \leq r_j) \in M^{r_j}.$$

Next, $P \subset M$ for the sub- W -module generated by all $B_{i,s}^{(j)}$ with $j \geq 2$ and $i < h_j$; also, $N \subset M$ for the sub- W -module generated by all $B_{i,s}^{(1)}$ with $i < h_1$. Note that $M = N \times P$, a direct sum of W -modules and that $M_\rho = (N \cap M_\rho) \times (P \cap M_\rho)$.

In the proof the W -submodule $P \subset M$ will be fixed; its W -complement $N \subset M$ will change eventually if it is not already a Dieudonné submodule.

We write $m_1 = m$, $n_1 = n$, $h = h_1 = m + n$, and $r = r_1$. Note that we assumed $0 < \lambda_1 \leq 1/2$; hence $m \geq n > 0$. For $i \geq 0$ we define integers δ_i by:

$$i \cdot h \leq \delta_i \cdot n < (i+1) \cdot n = ih + n.$$

Also, there are nonnegative integers γ_i such that

$$\delta_0 = 0, \delta_1 = \gamma_1 + 1, \dots, \delta_i = \gamma_1 + 1 + \gamma_2 + 1 + \dots + \gamma_i + 1, \dots;$$

note that $\delta_n = h = m + n$; hence $\gamma_1 + \dots + \gamma_n = m$. For $1 \leq i \leq n$ we write

$$f(i) = \delta_{i-1} \cdot n - (i-1) \cdot h;$$

this means that $0 \leq f(i) < n$ is the remainder after dividing $\delta_{i-1}n$ by h ; note that $f(1) = 0$. As $\gcd(n, h) = 1$ we see that

$$f : \{1, \dots, n\} \rightarrow \{0, \dots, n-1\}$$

is a bijective map. The inverse map f' is given by:

$$f' : \{0, \dots, n-1\} \rightarrow \{1, \dots, n\}, \quad f'(x) \equiv 1 - \frac{x}{h} \pmod{n}, \quad 1 \leq f'(x) \leq n.$$

In $(Q_1)^r$ we have the vectors $A_i^{(1)}$. We choose $C'_1 := A_0^{(1)}$ and we choose $\{C'_1, \dots, C'_n\} = \{A_0^{(1)}, \dots, A_{n-1}^{(1)}\}$ by

$$C'_i := A_{f(i)}^{(1)}, \quad C'_{f'(x)} = A_x^{(1)};$$

this means that:

$$\mathcal{F}^{\gamma_i} C'_i = \mathcal{V} C'_{i+1}, \quad 1 \leq i < n, \quad \mathcal{F}^{\gamma_n} C'_n = \mathcal{V} C'_1;$$

hence

$$\mathcal{F}^{\delta_i} C'_1 = p^i \cdot C'_{i+1}, \quad 1 \leq i < n.$$

Note that $\mathcal{F}^h C'_1 = p^n \cdot C'_1$. With these choices we see that

$$\{\mathcal{F}^j C'_i \mid 1 \leq i \leq n, 0 \leq j \leq \gamma_i\} = \{A_\ell^{(1)} \mid 0 \leq \ell < h\}.$$

For later reference we state:

(2.6) *Suppose Q is a nonzero Dieudonné module with an element $C \in Q$, such that there exist coprime integers n and $n + m = h$ as above such that $\mathcal{F}^h \cdot C = p^n \cdot C$ and such that Q , as a W -module, is generated by $\{p^{-[jn/h]} \mathcal{F}^j C \mid 0 \leq j < h\}$, then $Q \cong \mathbb{D}(H_{m,n})$.*

This is proved by explicitly writing out the required isomorphism. Note that \mathcal{F}^n is injective on Q ; hence $\mathcal{F}^h \cdot C = p^n \cdot C$ implies $\mathcal{F}^m \cdot C = \mathcal{V}^n \cdot C$.

(2.7) Accordingly we choose $C_{i,s} := B_{f(i),s}^{(1)} \in M$ with $1 \leq i \leq n$. Note that

$$\{\mathcal{F}^j C_{i,s} \mid 1 \leq i \leq n, 0 \leq j \leq \gamma_i, 1 \leq s \leq r\}$$
 is a W -basis for N ,

$$\mathcal{F}^{\gamma_i} C_{i,s} - \mathcal{V} C_{i+1,s} \in pM, \quad 1 \leq i < n, \quad \mathcal{F}^{\gamma_n} C_{n,s} - \mathcal{V} C_{1,s} \in pM.$$

We write $C_i = (C_{i,s} \mid 1 \leq s \leq r)$. As a reminder, we sum up some of the notation constructed:

$$\begin{array}{ccc} N \subset M & & \bigoplus_j (Q_j)^{r_j} \\ \downarrow & & \downarrow \\ M/pM & = & \bigoplus_j (Q_j/pQ_j)^{r_j}, \\ \\ B_{i,s}^{(j)} \in M, & & A_{i,s}^{(j)} \in Q_j \subset (Q_j)^{r_j}, \\ C_{i,s} \in N, & & C'_{i,s} \in Q_1 \subset (Q_1)^{r_1}. \end{array}$$

(2.8) LEMMA. *Use the notation fixed up to now.*

- (1) For every $\rho \in \mathbb{Q}_{\geq 0}$ the map $p : M_\rho \rightarrow M_{\rho+1}$, multiplication by p , is surjective.
- (2) For every $\rho \in \mathbb{Q}_{\geq 0}$ there exists $\mathcal{F}M_\rho \subset M_{\rho+(n/h)}$.
- (3) For every i and s , $\mathcal{F}B_{i,s}^{(1)} \in M_{(i+n)/h}$; for every i and s and every $j > 1$, $\mathcal{F}B_{i,s}^{(j)} \in M_{(i/h_j)+(n/h)+(1/T)}$.
- (4) For every $1 \leq i \leq n$ there is $\mathcal{F}^{\delta_i}C_1 - p^i B_{f(i+1)}^{(1)} \in (M_{i+(1/T)})^r$; moreover $\mathcal{F}^{\delta_n}C_1 - p^n C_1 \in (M_{n+(1/T)})^r$.
- (5) If u is an integer with $u > Tn$, and $\xi_N \in (N \cap M_{u/T})^r$, there exists

$$\eta_N \in N \cap (M_{(u/T)-n})^r$$

such that

$$(\mathcal{F}^h - p^n)\eta_N \equiv \xi_N \pmod{(M_{(u+1)/T})^r}.$$

Proof. We know that $M_{\rho+1}$ is generated by the elements $B_{i,s}^{(j)}$ with $i/h_j \geq \rho + 1$; because $\rho \geq 0$ such elements satisfy $i \geq h_j$. Note that $p \cdot B_{i-h_j,s}^{(j)} = B_{i,s}^{(j)}$. This proves the first property. □(1)

At first we show $\mathcal{F}M \subset M_{n/h}$. Note that for all $1 \leq j \leq t$ and all $\beta \in \mathbb{Z}_{\geq 0}$

$$(*) \quad \beta h_j \leq i < \beta h_j + m_j \quad \Rightarrow \quad \mathcal{F}B_i^{(j)} = B_{i+n_j}^{(j)},$$

and

$$(**) \quad \beta h_j + m_j \leq i < (\beta + 1)h_j \quad \Rightarrow \quad B_i^{(j)} = \mathcal{V}B_{i-m_j}^{(j)} + p^{(\beta+1)}\xi, \quad \xi \in M^{r_j}.$$

From these properties, using $n/h \leq n_j/h_j$, we know: $\mathcal{F}M \subset M_{n/h}$.

Further we see by (*) that

$$v(\mathcal{F}B_{i,s}^{(j)}) = v(B_{i+n_j,s}^{(j)}) = (i + n_j)/h_j,$$

and

$$\frac{i + n_j}{h_j} = \frac{i + n}{h} \quad \text{if } j = 1; \quad \frac{i + n_j}{h_j} > \frac{i}{h_j} + \frac{n}{h} \quad \text{if } j > 1.$$

By (**) it suffices to consider only $m_j \leq i < h_j$, and hence $\mathcal{F}B_{i,s}^{(j)} = pB_{i-m_j,s}^{(j)} + p\mathcal{F}\xi_s$; thus

$$v(\mathcal{F}B_{i,s}^{(j)}) \geq \min \left(v(pB_{i-m_j,s}^{(j)}), v(p\mathcal{F}\xi_s) \right).$$

For $j = 1$ we have $v(pB_{i-m_1,s}^{(1)}) = (i+n)/h \geq 1$ and $v(p\mathcal{F}\xi) \geq 1 + (n/h) > (i/h) + (n/h)$; for $j > 1$ we have $v(pB_{i-m_j,s}^{(j)}) > (i/h_j) + (n/h)$ and $(i/h_j) + (n/h) < 1 + (n/h) \leq v(p\mathcal{F}\xi_s)$. Hence $v(\mathcal{F}B_{i,s}^{(j)}) > (i/h_j) + (n/h)$ if $j > 1$. This ends the proof of (3). Using (3) we see that (2) follows. $\square(2)+(3)$

From $\mathcal{F}^{\gamma_i}C_i = \mathcal{V}C_{i+1} + p\xi_i$ for $i < n$ and $\mathcal{F}^{\gamma_n}C_n = \mathcal{V}C_1 + p\xi_n$, here $\xi_i \in M^r$ for $i \leq n$, we have:

$$\mathcal{F}^{\delta_i}C_1 = p^i C_{i+1} + \sum_{1 \leq \ell \leq i} p^\ell \mathcal{F}^{\delta_i - \delta_\ell} \mathcal{F}\xi_\ell, \quad \text{for } i < n,$$

and the analogous formula for $i = n$ (write $C_{n+1} = C_1$). Note that

$$ih \leq \delta_i n \quad \text{and} \quad \delta_\ell n < \ell m + (\ell + 1)n = \ell h + n;$$

this shows that

$$\ell h + (\delta_i - \delta_\ell)n + n > ih;$$

using (2) we have proved (4). $\square(4)$

Note that $h = h_1$ divides T . If ℓ is an integer such that $(\ell - 1)/h < u/T < \ell/h$ then $u < u + 1 \leq \ell \frac{T}{h}$ and we see that $N \cap M_{u/T} = N \cap M_{(u+1)/T}$ and we choose $\eta_N = 0$.

Suppose that ℓ is an integer with $u/T = \ell/h$. Then $N \cap M_{u/T} = N_{\ell/h} \supset N_{(\ell+1)/h} = N \cap M_{(u+1)/T}$. We consider the image of $N \cap M_{(\ell/h)-n}$ under $\mathcal{F}^h - p^n$ and see, using previous results, that this image is in $N_{\ell/h} + M_{(u+1)/T}$ (here “+” stands for the span as W -modules). We obtain a factorization and an isomorphism

$$\mathcal{F}^h - p^n : N \cap M_{(\ell/h)-n} \longrightarrow (N_{\ell/h} + M_{(u+1)/T}) / M_{(u+1)/T} \cong N_{\ell/h} / N_{(\ell+1)/h}.$$

We claim that this map is surjective. The factor space $N_{\ell/h} / N_{(\ell+1)/h}$ is a vector space over k spanned by the residue classes of the elements $B_{\ell,s}^{(1)}$. For the residue class of $y_s B_{\ell,s}^{(1)}$ we solve the equation $x_s^{p^n} - x_s = y_s$ in k ; lifting these x_s to W (denoting the lifts by the same symbol), we see that $\eta_N := \sum_s x_s B_{\ell-nh,s}^{(1)}$ has the required properties. This proves the claim, and gives a proof of part (5) of the lemma. $\square(5),(2.8)$

(2.9) LEMMA (the induction step). *Let $u \in \mathbb{Z}$ with $u \geq nT + 1$. Suppose $D_1 \in M^r$ such that $D_1 \equiv C_1 \pmod{(M_{1/T})^r}$, and such that $\xi := \mathcal{F}^h D_1 - p^n D_1 \in (M_{u/T})^r$. Then there exists $\eta \in (M_{(u/T)-n})^r$ such that for $E_1 := D_1 - \eta$ there exist $\mathcal{F}^h E_1 - p^n E_1 \in (M_{(u+1)/T})^r$ and $E_1 \equiv C_1 \pmod{(M_{1/T})^r}$.*

Proof. We write $\xi = \xi_N + \xi_P$ according to $M = N \times P$ and conclude that $\xi_N \in (N \cap M_{u/T})^r$ and $\xi_P \in (P \cap M_{u/T})^r$. Using (2.8), (5), we construct $\eta_N \in (N \cap M_{1/T})^r$ such that $(\mathcal{F}^h - p^n)\eta_N \equiv \xi_N \pmod{(M_{(u+1)/T})^r}$. As $M_{u/T} \subset M_n$

we can choose $\eta_P := -p^{-n}\xi_P$; we have $\eta_P \in M_{(u/T)-n}^r \subset (M_{1/T})^r$. With $\eta := \eta_N + \eta_P$ we see that

$$(\mathcal{F}^h - p^n)\eta \equiv \xi \pmod{(M_{(u+1)/T})^r} \quad \text{and} \quad \eta \in (M_{1/T})^r.$$

Hence $(\mathcal{F}^h - p^n)(D_1 - \eta) \in (M_{(u+1)/T})^r$ and we see that $E_1 := D_1 - \eta$ has the required properties. This proves the lemma. $\square(2.9)$

(2.10) *Preparation for the Proof of (2.2).*

(1) *There exists $E_1 \in M^r$ such that $(\mathcal{F}^h - p^n)E_1 = 0$ and $E_1 \equiv C_1 \pmod{(M_{1/T})^r}$.*

Proof. For $u \in \mathbb{Z}_{\geq nT+1}$ we write $D_1(u) \in M^r$ for a vector such that

$$D_1(u) \equiv C_1 \pmod{(M_{1/T})^r} \quad \text{and} \quad \mathcal{F}^h D_1(u) - p^n D_1(u) \in (M_{u/T})^r.$$

By (2.8), (4), the vector $C_1 =: D_1(nT+1)$ satisfies this condition for $u = nT+1$. Here we start induction. By repeated application of (2.9) we conclude there exists a sequence

$$\{D_1(u) \mid u \in \mathbb{Z}_{\geq nT+1}\} \quad \text{such that} \quad D_1(u) - D_1(u+1) \in (M_{(u/T)-n})^r$$

satisfying the conditions above. As $\cap_{\rho \rightarrow \infty} M_\rho = \{0\}$ this sequence converges. Writing $E_1 := D_1(\infty)$ we achieve the conclusion. $\square(1)$

(2) *Choose E_1 as in (1). For every $j \geq 0$,*

$$p^{-[\frac{jn}{h}]} \mathcal{F}^j E_1 \in M \quad \text{and} \quad N' := \prod_{1 \leq j < h} W \cdot p^{-[\frac{jn}{h}]} \mathcal{F}^j E_1 \subset M.$$

This is a Dieudonné submodule and a W -module direct summand of M . Moreover there is an isomorphism

$$\mathbb{D}((H_{m,n})^r) \cong N',$$

the map $N' \prod P \rightarrow N' + P$ is an isomorphism of W -modules, and $N' + P = M$. Thus $X_1 \subset X$, with

$$\mathbb{D}(X_1 \subset X) = (N' \subset M) \quad \text{such that} \quad (X/X_1)[p] \cong \prod_{j>1} (M_{m_j, n_j})^{r_j}.$$

Proof of (2) and of Proposition 2.2. By (2.8), (2), we see that $\mathcal{F}^j E_1 \in M_{[jn/h]}$; hence the first statement follows. As $\mathcal{F}^h E_1 = p^n E_1$ it follows that $N' \subset M$ is a Dieudonné submodule; by (2.6) this shows $\mathbb{D}((H_{m,n})^r) \cong N'$.

CLAIM. *The images $N' \twoheadrightarrow N' \otimes k = N'/pN' \subset M/pM$ and $P \twoheadrightarrow P/pP \subset M/pM$ inside M/pM have zero intersection and $N' \otimes k + P \otimes k = M/pM$. Here $- \otimes k = - \otimes_W (W/pW)$.*

For $y \in \mathbb{Z}_{\geq 0}$ we write $g(y) := yn - h \cdot \lfloor \frac{yn}{h} \rfloor$; recall that in the notation in (2.5),

$$p^{-\lfloor \frac{in}{h} \rfloor} \mathcal{F}^j C'_1 = A_{g(j)}^{(1)}.$$

Suppose

$$\tau := \sum_{0 \leq j < h} \beta_{j,s} p^{-\lfloor \frac{jn}{h} \rfloor} \mathcal{F}^j \cdot (E_{1,s} \bmod pM) \in (N' \otimes k \cap P \otimes k) \subset M/pM, \quad \beta_j \in k,$$

such that $\tau \neq 0$. Let x, s be a pair of indices such that $\beta := \beta_{x,s} \neq 0$ and for every y with $g(y) < g(x)$ we have $\beta_{y,s} = 0$. Project inside M/pM on the factor N_s . Then

$$\tau_s \equiv \beta \cdot B_{g(x),s}^{(1)} \pmod{M_{\frac{g(x)}{h} + \frac{1}{T}} + P},$$

which is a contradiction to the fact that $N \cap P = 0$ and to the fact that the residue class of

$$B_{g(x),s}^{(1)} \text{ generates } \left((M_{\frac{g(x)}{h}} + P) / (M_{\frac{g(x)}{h} + \frac{1}{T}} + P) \right)_s = N_{\frac{g(x)}{h},s} / N_{\frac{g(x)}{h} + \frac{1}{h},s}.$$

We see that $\tau \neq 0$ leads to a contradiction. This shows that $N' \otimes k \cap P \otimes k = 0$ and $N' \otimes k + P \otimes k = M/pM$. Hence the claim is proved.

As $(N' \cap P) \otimes k \subset N' \otimes k \cap P \otimes k = 0$ this shows $(N' \cap P) \otimes k = 0$. By Nakayama's lemma this implies $N' \cap P = 0$. The proof of the remaining statements follows; in particular we see that N' is a W -module direct summand of M . This finishes the proof of (2), and ends the proof of the proposition. □(2.2)

3. Split extensions and proof of the theorem

In this section we prove a proposition on split extensions. We will see that Theorem (1.2) follows.

(3.1) PROPOSITION. *Let (m, n) and (d, e) be ordered pairs of pairwise coprime positive integers. Suppose that $n/(m + n) < e/(d + e)$. Let*

$$0 \rightarrow Z := H_{m,n} \rightarrow T \rightarrow Y := H_{d,e} \rightarrow 0$$

be an exact sequence of p -divisible groups such that the induced sequence of the p -kernels splits:

$$0 \rightarrow Z[p] \xrightarrow{\leftarrow} T[p] \xrightarrow{\leftarrow} Y[p] \rightarrow 0.$$

Then the sequence of p -divisible groups splits: $T \cong Z \oplus Y$.

(3.2) Remark. It is easy to give examples of a nonsplit extension $T/Z \cong Y$ of p -divisible groups, with Z nonminimal or Y nonminimal, such that the extension $T[p]/Z[p] \cong Y[p]$ does split.

(3.3) *Proof of Theorem (1.2).* The theorem follows from (2.4) and (3.1). □(1.2)

(3.4) In order to prove (3.1) it suffices to prove it under the extra condition that $\frac{1}{2} \leq e/(d + e)$. In fact, if $n/(m + n) < e/(d + e) < \frac{1}{2}$, we consider the exact sequence

$$0 \rightarrow H_{d,e}^t = H_{e,d} \rightarrow T^t \rightarrow H_{m,n}^t = H_{n,m} \rightarrow 0$$

with $\frac{1}{2} < d/(e + d) < m/(n + m)$. □(3.4)

From now on we assume that $\frac{1}{2} \leq e/(d + e)$.

(3.5) We fix notation to be used in the proof of (3.1). Writing the Dieudonné modules as $\mathbb{D}(Z) = N$, $\mathbb{D}(T) = M$ and $\mathbb{D}(Y) = Q$, we obtain an exact sequence of Dieudonné modules $M/N = Q$, which is a split exact sequence of W -modules, where $W = W_\infty(k)$. Now, $m + n = h$ and $d + e = g$. We know that Q is generated by elements A_i , with $i \in \mathbb{Z}_{\geq 0}$ such that $\pi(A_i) = A_{i+1}$, where $\pi \in \text{End}(Q)$ is the uniformizer, and $\mathcal{V} \cdot A_i = A_{i+d}$, $\mathcal{F} \cdot A_i = A_{i+e}$. Also, $\{A_i \mid 0 \leq i < g = d + e\}$ is a W -basis for Q . Because $\frac{1}{2} \leq e/(d + e)$, and $e \geq d$ we can choose generators for the Dieudonné module Q in the following way. We choose integers δ_i by:

$$i \cdot g \leq \delta_i \cdot d < (i + 1) \cdot d + i \cdot e = ig + d$$

and integers γ_i such that:

$$\delta_1 = \gamma_1 + 1, \dots, \delta_i = \gamma_1 + 1 + \gamma_2 + 1 + \dots + \gamma_i + 1;$$

note that $\delta_d = g = d + e$. We choose $C = A_0 = C_1$ and $\{C_1, \dots, C_d\} = \{A_0, \dots, A_{d-1}\}$ such that:

$$\mathcal{V}^{\gamma_i} C_i = \mathcal{F} C_{i+1}, \quad 1 \leq i < d, \quad \mathcal{V}^{\gamma_d} C_d = \mathcal{F} C_1;$$

hence

$$\mathcal{V}^{\delta_i} C = p^i \cdot C_{i+1}, \quad 1 \leq i < d.$$

Note that $\mathcal{V}^g C = p^d \cdot C$. With these choices we see that

$$\{p^{-[\frac{jd}{g}]} \mathcal{V}^j C \mid 0 \leq j < g\} = \{\mathcal{V}^j C_i \mid 1 \leq i \leq d, \quad 0 \leq j \leq \gamma_i\} = \{A_\ell \mid 0 \leq \ell < g\}.$$

Choose an element $B = B_1 \in M$ such that

$$M \rightarrow Q \quad \text{gives} \quad B_1 = B \mapsto (B \bmod N) = C = C_1.$$

Let π' be the uniformizer of $\text{End}(N)$. Consider the filtration $N = N^{(0)} \supset \dots \supset N^{(i)} \supset N^{(i+1)} \supset \dots$ defined by $(\pi')^i(N^{(0)}) = N^{(i)}$. Note that $\mathcal{F}N^{(i)} = N^{(i+n)}$, and $\mathcal{V}N^{(i)} = N^{(i+m)}$, and $p^i N = N^{(i \cdot h)}$ for $i \geq 0$.

(3.6) *Proof of Proposition (3.1).*

(1) Construction of $\{B_1, \dots, B_d\}$. For every choice of $B = B_1 \in M$ with $(B \bmod N) = C$, and every $1 \leq i < d$ we claim that $\mathcal{V}^{\delta_i} B$ is divisible by p^i . Defining $B_{i+1} := p^{-i} \mathcal{V}^{\delta_i} B$, we see that $B_i \bmod N = C_i$ for $1 \leq i \leq d$. Moreover, we claim:

$$\mathcal{V}^g B - p^d \cdot B \in N^{(dh+1)}.$$

Choose $B''_i \in M$ with $B''_i \bmod N = C_i$. Then $\mathcal{V}^{\gamma_i} B''_i - \mathcal{F} B''_{i+1} =: p \cdot \xi_i \in pN$; hence $\mathcal{V}^{\gamma_i+1} B''_i - p \cdot B''_{i+1} = p \mathcal{V} \xi_i \in p \mathcal{V} N$. For $1 < i \leq d$ we obtain

$$\mathcal{V}^{\delta_i} B - p^i \cdot B = \sum_{1 \leq j < i} \mathcal{V}^{\delta_i - \delta_j} p^j \mathcal{V} \xi_j, \quad \xi_j \in N.$$

From $n/(m+n) < e/(d+e)$ we conclude $g/d > h/m$; since $\delta_i \cdot d \geq ig$ and $\delta_j d < (j+1)d + je$,

$$i > j \quad \text{implies} \quad \delta_i - \delta_j + 1 > (i-j)(g/d) > (i-j)(h/m);$$

hence

$$(\delta_i - \delta_j)m + j(m+n) + m > ih.$$

This shows

$$\mathcal{V}^{\delta_i - \delta_j} p^j \mathcal{V} \xi_j \in p^i N^{(1)}.$$

As $\delta_d = g$ we see that $\mathcal{V}^g B - p^d \cdot B \in p^d N^{(1)} = N^{(dh+1)}$. □(1)

(2) The induction step. Suppose that for a choice $B \in M$ with $(B \bmod N) = C$, there exists an integer $s \geq dh + 1$ such that $\mathcal{V}^g B - p^d \cdot B \in N^{(s)}$; then there exists a choice $B' \in M$ such that $B' - B \in N^{(s-dh)}$ and

$$\mathcal{V}^g B' - p^d \cdot B' \in N^{(s+1)}.$$

In fact, $p^d \cdot B - \mathcal{V}^g B = p^d \cdot \xi$. Then $\xi \in N^{(s-dh)}$. Choose $B' := B - \xi$. Then:

$$\mathcal{V}^g B' - p^d \cdot B' = \mathcal{V}^g B - p^d \cdot B - \mathcal{V}^g \xi + p^d \xi = -\mathcal{V}^g \xi \in N^{(gm-dh+s)}$$

and $gm - dh > 0$. □(2)

(3) For any integer $r \geq d + 1$, and $w \geq rh$ there exists $B = B_1$ as in (3.5) such that $\mathcal{V}^g B - p^d B \in N^{(w)} = p^r \cdot N^{(w-rh)}$. This gives a homomorphism φ_{r-d}

$$M/p^{r-d}M \longleftarrow Q/p^{r-d}Q \quad \text{extending} \quad M/pM \longleftarrow Q/pQ.$$

The induction step (2) proves the first statement, induction starting at $w = (d+1)h > dh + 1$. Having chosen B_1 , using (1) we construct $B_{i+1} := p^{-i} \mathcal{V}^{\delta_i} B_1$ for $1 \leq i < d$. In that case on the one hand $\mathcal{V}^{\gamma_i} B_d - \mathcal{F} B_1 = p \cdot \xi_d$; on the other hand $\mathcal{V}^g B - p^d B \in N^{(w)} \subset p^r N$. Hence $p^d \mathcal{V} \xi_d \in p^r N$; hence $p \xi_d \in p^{r-d} N$. This shows that the residue classes of B_1, \dots, B_d in $M/p^{r-d}M$

generate a Dieudonné module isomorphic to $Q/p^{r-d}Q$ which moreover by (3.5) extends the given isomorphism induced by the splitting. $\square(3)$

By [8, 1.6], for some large r the existence of $M/p^{r-d}M \leftarrow Q/p^{r-d}Q$ as in (3) shows that its restriction $M/pM \leftarrow Q/pQ$ lifts to a homomorphism φ of Dieudonné modules $M \leftarrow Q$; in that case φ_1 is injective. Hence φ splits the extension $M/N \cong Q$. Taking into account (3.4) we have proved the proposition. $\square(3.1)$

Remark. Instead of the last step of the proof above, we could construct an infinite sequence $\{B(u) \mid u \in \mathbb{Z}_{(d+1)h}\}$ such that $\mathcal{V}^g B(u) - p^d B \in N^{(u)}$ and $B(u+1) - B(u) \in N^{(u-dh)}$ for all $u \geq (d+1)h$. This sequence converges and its limit $B(\infty)$ can be used to define the required section.

4. Some comments

(4.1) *Remark.* For any G , a BT_1 over k , which is *not minimal* there exist infinitely many mutually nonisomorphic p -divisible groups X over k such that $X[p] \cong G$. Details will appear in a later publication; see [9].

(4.2) *Remark.* Suppose that G is a *minimal* BT_1 ; we can recover the Newton polygon β with the property $H(\beta)[p] \cong G$ from G . This follows from the theorem, but there are also other ways to prove this fact.

(4.3) For BT_1 group schemes we can define a Newton polygon; let G be a BT_1 group scheme over k , and let $G = \times_i G_i$ be a decomposition into indecomposable ones; see [3]. Let G_i be of rank p^{h_i} , and let n_i be the dimension of the tangent space of G_i^D ; here G_i^D stands for the Cartier dual of G_i . Define $\mathcal{N}'(G_i)$ as the isoclinic polygon consisting of h_i slopes equal to n_i/h_i ; arranging the slopes in nondecreasing order, we have defined $\mathcal{N}'(G)$. For a p -divisible group X we compare $\mathcal{N}(X)$ and $\mathcal{N}'(X[p])$. These polygons have the same endpoints. If X is minimal, equivalently $X[p]$ is minimal, then $\mathcal{N}(X) = \mathcal{N}'(X[p])$. Besides this, I do not see rules describing the relation between $\mathcal{N}(X)$ and $\mathcal{N}'(X[p])$. For Newton polygons β and γ with the same endpoints we write $\beta \prec \gamma$ if every point of β is on or below γ . Note:

- There exists a p -divisible group X such that $\mathcal{N}(X) \not\cong \mathcal{N}'(X[p])$; indeed, when X is isosimple, then $\mathcal{N}(X)$ is isoclinic, such that $X[p]$ is decomposable.
- There exists a p -divisible group X such that $\mathcal{N}(X) \not\cong \mathcal{N}'(X[p])$; indeed, choosing X such that $\mathcal{N}(X)$ is not isoclinic, we have X is not isosimple, all slopes are strictly between 0 and 1 and $a(X) = 1$; then $X[p]$ is indecomposable; hence $\mathcal{N}'(X[p])$ is isoclinic.

Here we use $a(X) := \dim_k \text{Hom}(\alpha_p, X)$. It would be useful to have better insight in the relation between various properties of X and $X[p]$.

UNIVERSITY OF UTRECHT, UTRECHT, THE NETHERLANDS
E-mail address: oort@math.uu.nl

REFERENCES

- [1] L. ILLUSIE, Déformations de Groupes de Barsotti-Tate, Exp. VI in *Séminaire sur les Pinceaux Arithmétiques: la Conjecture de Mordell* (L. Szpiro), *Astérisque* **127**, Soc. Math. France, Paris, 1985.
- [2] A. J. DE JONG and F. OORT, Purity of the stratification by Newton polygons, *J. Amer. Math. Soc.* **13** (2000), 209–241.
- [3] H. KRAFT, Kommutative algebraische p -Gruppen (mit Anwendungen auf p -divisible Gruppen und abelsche Varietäten), Sonderforsch. Bereich Bonn, September 1975, preprint.
- [4] H. KRAFT and F. OORT, Group schemes annihilated by p , in preparation.
- [5] YU. I. MANIN, The theory of commutative formal groups over fields of finite characteristic, *Usp. Math.* **18** (1963), 3–90; *Russian Math. Surveys* **18** (1963), 1–80.
- [6] F. OORT, *Commutative Group Schemes*, *Lecture Notes in Math.* **15**, Springer-Verlag, New York, 1966.
- [7] ———, A stratification of a moduli space of polarized abelian varieties, in *Moduli of Abelian Varieties* (C. Faber, G. van der Geer, F. Oort, eds.), *Progr. Math.* **195**, 345–416, Birkhäuser Verlag, Basel, 2001.
- [8] ———, Foliations in moduli spaces of abelian varieties, *J. Amer. Math. Soc.* **17** (2004), 267–296.
- [9] ———, Simple p -kernels of p -divisible groups, *Advances in Math.*, to appear.

(Received September 3, 2002)

(Revised May 24, 2004)