


# ANNALS OF MATHEMATICS

## The Ten Martini Problem

By ARTUR AVILA and SVETLANA JITOMIRSKAYA



SECOND SERIES, VOL. 170, NO. 1

July, 2009

ANMAAH



# The Ten Martini Problem

By ARTUR AVILA and SVETLANA JITOMIRSKAYA

## Abstract

We prove the conjecture (known as the “Ten Martini Problem” after Kac and Simon) that the spectrum of the almost Mathieu operator is a Cantor set for all nonzero values of the coupling and all irrational frequencies.

## 1. Introduction

The almost Mathieu operator is the Schrödinger operator  $H_{\lambda,\alpha,\theta}$  on  $\ell^2(\mathbb{Z})$  defined by

$$(1-1) \quad (H_{\lambda,\alpha,\theta}u)_n = u_{n+1} + u_{n-1} + 2\lambda \cos 2\pi(\theta + n\alpha)u_n,$$

where  $\lambda, \alpha, \theta \in \mathbb{R}$  are parameters (called the *coupling*, *frequency*, and *phase*, respectively), and one assumes that  $\lambda \neq 0$ . The interest in this particular model is motivated both by its connections to physics and by a remarkable richness of the related spectral theory. This has made the latter a subject of intense research in the last three decades (see [Las05] for a recent historical account and for the physics background). Here we are concerned with the topological structure of the spectrum.

If  $\alpha = p/q$  is rational, it is well known that the spectrum consists of the union of  $q$  intervals called *bands*, possibly touching at the endpoints. In the case of irrational  $\alpha$ , the spectrum  $\Sigma_{\lambda,\alpha}$  (which in this case does not depend on  $\theta$ ) has been conjectured for a long time to be a Cantor set; see a 1964 paper of Azbel [Azb64]. Proving this has been dubbed the Ten Martini Problem by Barry Simon, after an offer of Mark Kac in 1981; see [Sim00, Prob. 4]. For a history of this problem see [Las05]. Earlier partial results include [BS82], [Sin87], [HS89], [CEY90], [Las94], and recent advances include [Pui04] and [AK06]. In this paper, we solve the Ten Martini Problem as stated in [Sim00].

---

Avila is a Clay Research Fellow. Jitomirskaya was supported in part by NSF grant DMS-0300974 and grant 2002068 from the United States-Israel Binational Science Foundation (BSF), Jerusalem.

**MAIN THEOREM.** *The spectrum of the almost Mathieu operator is a Cantor set for all irrational  $\alpha$  and for all  $\lambda \neq 0$ .*

It is important to emphasize that the previous results mentioned above covered a large set of parameters  $(\lambda, \alpha)$ , which is both topologically generic [BS82] and of full Lebesgue measure [Pui04]. As often happens in the analysis of quasiperiodic systems, the “topologically generic” behavior is quite distinct from the “full Lebesgue measure” behavior, and the narrow set of parameters left behind does indeed lie in the interface of two distinct regimes. Furthermore, our analysis seems to indicate an interesting characteristic of the Ten Martini Problem, that the two regimes do not cover nicely the parameter space, and hence there is a nonempty “critical region” of parameters in between; see Remarks 1.1, 5.1, 5.2 and the comments after Theorem 8.2.

This is to some degree reflected in the structure of the proof. While the reasoning outside of the critical region can be made quite effective, in the sense that one essentially identifies specific gaps in the spectrum,<sup>1</sup> in order to be able to cover the critical region we make use of very indirect arguments. As an example, we show that absence of Cantor spectrum enables us to “analytically continue” effective solutions of a small divisor problem, and it is the noneffective solutions thus obtained that can be related to gaps in the spectrum.

This paper builds on a large theory. Especially important for us are [CEY90], [Jit99], [Pui04], whose methods we improve, but several other ingredients are needed (such as Kotani theory [Sim83] and the recent estimates on Lyapunov exponents of [BJ02]). An important new ingredient is the use of analytic continuation techniques in the study of  $m$ -functions and in extending the reach of the analysis of Anderson localization.

1.1. *Strategy.* In this problem, arithmetics of  $\alpha$  rules the game. When  $\alpha$  is not very Liouville, it is reasonable to try to deal with the small divisors. When  $\alpha$  is not very Diophantine, this does not work, and we deal instead with rational approximation arguments. Let  $p_n/q_n$  be the approximants of  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ . Let

$$(1-2) \quad \beta = \beta(\alpha) = \limsup(\ln q_{n+1}/q_n).$$

The relation between  $e^\beta$  and  $\lambda$  will play an important role in our argument, and will decide whether we approach the problem from the Diophantine side or from the Liouvillian side. As discussed before, our analysis indicates that there are parameters that cannot be effectively described from either side, and it is only through the use of indirect arguments that we can enlarge artificially the Diophantine and Liouville

---

<sup>1</sup>These gaps are related either to gaps of periodic approximations or to eigenvalues of a dual almost Mathieu operator.

regimes to cover all parameters. It should be noted that even with such tricks, both sides will just about meet in the middle.

Since  $\Sigma_{\lambda,\alpha} = \Sigma_{-\lambda,\alpha}$ , it is enough to assume  $\lambda > 0$ . It is known that the behavior of the almost Mathieu operator changes drastically at  $\lambda = 1$  (“metal-insulator” transition [Jit99]). Aubry duality shows that  $\Sigma_{\lambda,\alpha} = \lambda \Sigma_{\lambda^{-1},\alpha}$ . So each  $\lambda \neq 1$  admits two lines of attack, and this will be determinant in what follows. The case  $\lambda = 1$  was settled in [AK06] (after several partial results [AvMS90], [HS89], [Las94]), but it is also recovered in our approach.

We will work on  $\lambda < 1$  when approaching from the Liouville side. The approach from the Diophantine side is more delicate. There are actually two classical small divisor problems that apply to the study of the almost Mathieu operator, corresponding to Floquet reducibility (for  $\lambda < 1$ ) and Anderson localization (for  $\lambda > 1$ ). An important point is to attack both problems simultaneously, mixing the best of each problem (“soft” analysis in one case, “hard” analysis in the other).

A key idea in this paper is that absence of Cantor spectrum implies improved regularity of  $m$ -functions in the regime  $0 < \lambda \leq 1$ . This is proved by analytic continuation techniques. The improved regularity of  $m$ -functions (which is fictitious, since we will prove Cantor spectrum) will be used both in the Liouville side and in the Diophantine side. In the Liouville side, it will give improved estimates for the continuity of the spectrum with respect to the frequency. In the Diophantine side, it will allow us to use (again) analytic continuation techniques to solve some small divisor problems in some situations that are beyond what is expected to be possible.

*Remark 1.1.* Since our approach, designed to overcome the difficulties in the interface of the Diophantine and Liouville regimes, works equally well for other ranges of parameters, it will not be necessary in the proof to precisely delimit a critical region. For the reasons discussed in Remarks 5.1 and 5.2 and in the comments after Theorem 8.2, the critical region is believed to contain the parameters such that  $\beta > 0$  and  $\beta \leq |\ln \lambda| \leq 2\beta$ , the parameters such that  $\beta = |\ln \lambda|$  (respectively,  $2\beta = |\ln \lambda|$ ) being seemingly inaccessible (even after artificial extension) by the Diophantine method (respectively, Liouville method). It is reasonable to expect that something should be different in the indicated critical region. For instance, it is the natural place to look for possible counterexamples to the “Dry Ten Martini” conjecture (for a precise formulation see Section 8).

## 2. Background

2.1. *Cocycles, Lyapunov exponents, and fibered rotation number.* A (one-dimensional quasiperiodic  $\mathrm{SL}(2, \mathbb{R})$ ) cocycle is a pair

$$(\alpha, A) \in \mathbb{R} \times C^0(\mathbb{R}/\mathbb{Z}, \mathrm{SL}(2, \mathbb{R})),$$

understood as a *linear skew-product*:

$$(\alpha, A) : \mathbb{R}/\mathbb{Z} \times \mathbb{R}^2 \rightarrow \mathbb{R}/\mathbb{Z} \times \mathbb{R}^2,$$

$$(x, w) \mapsto (x + \alpha, A(x) \cdot w).$$

For  $n \geq 1$ , we let  $A_n(x) = A(x + (n-1)\alpha) \cdots A(x)$  ( $\alpha$  is implicit in this notation).

Given two cocycles  $(\alpha, A)$  and  $(\alpha, A')$ , a *conjugacy* between them is a continuous  $B : \mathbb{R}/\mathbb{Z} \rightarrow \mathrm{SL}(2, \mathbb{R})$  such that

$$B(x + \alpha)A(x)B(x)^{-1} = A'(x).$$

The *Lyapunov exponent* is defined by

$$(2-1) \quad \lim \frac{1}{n} \int \ln \|A_n(x)\| dx,$$

so  $L(\alpha, A) \geq 0$ . It is invariant under conjugacy.

Assume now that  $A : \mathbb{R}/\mathbb{Z} \rightarrow \mathrm{SL}(2, \mathbb{R})$  is homotopic to the identity. Then there exists  $\psi : \mathbb{R}/\mathbb{Z} \times \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}$  and  $u : \mathbb{R}/\mathbb{Z} \times \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}^+$  such that

$$(2-2) \quad A(x) \cdot \begin{pmatrix} \cos 2\pi y \\ \sin 2\pi y \end{pmatrix} = u(x, y) \begin{pmatrix} \cos 2\pi(y + \psi(x, y)) \\ \sin 2\pi(y + \psi(x, y)) \end{pmatrix}.$$

The function  $\psi$  is called a *lift* of  $A$ . Let  $\mu$  be any probability on  $\mathbb{R}/\mathbb{Z} \times \mathbb{R}/\mathbb{Z}$  which is invariant by the continuous map  $T : (x, y) \mapsto (x + \alpha, y + \psi(x, y))$ , projecting over Lebesgue measure on the first coordinate (for instance, take  $\mu$  as any accumulation point of  $\frac{1}{n} \sum_{k=0}^{n-1} T_*^k \nu$ , where  $\nu$  is Lebesgue measure on  $\mathbb{R}/\mathbb{Z} \times \mathbb{R}/\mathbb{Z}$ ). Then the number

$$\rho(\alpha, A) = \int \psi d\mu \bmod \mathbb{Z}$$

does not depend on the choices of  $\psi$  and  $\mu$ , and is called the *fibered rotation number* of  $(\alpha, A)$ ; see [JM82] and [Her83]. It is invariant under conjugacies homotopic to the identity. It immediately follows from the definitions that the fibered rotation number is a continuous function of  $(\alpha, A)$ .

If  $A, A' : \mathbb{R}/\mathbb{Z} \rightarrow \mathrm{SL}(2, \mathbb{R})$  and  $B : \mathbb{R}/\mathbb{Z} \rightarrow \mathrm{SL}(2, \mathbb{R})$  are continuous such that  $A$  is homotopic to the identity and  $B(x + \alpha)A(x)B(x)^{-1} = A'(x)$ , then  $\rho(\alpha, A) = \rho(\alpha, A') - k\alpha$ , where  $k$  is such that  $x \mapsto B(x)$  is homotopic to  $x \mapsto R_{kx}$ , where

$$R_\theta = \begin{pmatrix} \cos 2\pi\theta & -\sin 2\pi\theta \\ \sin 2\pi\theta & \cos 2\pi\theta \end{pmatrix}.$$

2.2. *Almost Mathieu cocycles, integrated density of states, and spectrum.* Let

$$(2-3) \quad S_{\lambda, E} = \begin{pmatrix} E - 2\lambda \cos 2\pi x & -1 \\ 1 & 0 \end{pmatrix}.$$

We call  $(\alpha, S_{\lambda,E})$ , for  $\lambda, \alpha, E \in \mathbb{R}$  and  $\lambda \neq 0$ , *almost Mathieu cocycles*. A sequence  $(u_n)_{n \in \mathbb{Z}}$  is a formal solution of the eigenvalue equation  $H_{\lambda,\alpha,\theta}u = Eu$  if and only if

$$S_{\lambda,E}(\theta + n\alpha) \cdot \begin{pmatrix} u_n \\ u_{n-1} \end{pmatrix} = \begin{pmatrix} u_{n+1} \\ u_n \end{pmatrix}.$$

Let  $L_{\lambda,\alpha}(E) = L(\alpha, S_{\lambda,E})$ . It is easy to see that  $\rho(\alpha, S_{\lambda,E})$  admits a determination  $\rho_{\lambda,\alpha}(E) \in [0, 1/2]$ . We let

$$N_{\lambda,\alpha}(E) = 1 - 2\rho_{\lambda,\alpha}(E) \in [0, 1].$$

It follows that  $E \mapsto N_{\lambda,\alpha}(E)$  is a continuous nondecreasing function. The function  $N$  is the usually defined *integrated density of states* of  $H_{\lambda,\alpha,\theta}$  if  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$  (for  $\alpha \in \mathbb{Q}$ ,  $N$  is the integral of the density of states over different  $\theta$ ); see [AS83] and [JM82]. Thus defining

$$\Sigma_{\lambda,\alpha} = \{E \in \mathbb{R} : N_{\lambda,\alpha} \text{ is not constant in a neighborhood of } E\},$$

we see that (consistently with the introduction)  $\Sigma_{\lambda,\alpha}$  is the spectrum of  $H_{\lambda,\alpha,\theta}$  for  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$  (in this case the spectrum does not depend on  $\theta$ ), while for  $\alpha \in \mathbb{Q}$ ,  $\Sigma_{\lambda,\alpha}$  is the union of the spectra of  $H_{\lambda,\alpha,\theta}$  for  $\theta \in \mathbb{R}$ . One also has

$$\Sigma_{\lambda,\alpha} \subset [-2 - 2|\lambda|, 2 + 2|\lambda|].$$

Continuity of the fibered rotation number implies that  $N_{\lambda,\alpha}$  depends continuously on  $(\lambda, \alpha)$ .

It turns out that there is a relation between  $N$  and  $L$ , the *Thouless formula* (see [AS83])

$$L(E) = \int \ln|E - E'| dN(E').$$

By the Schwarz reflection principle, if  $J \subset \mathbb{R}$  is an open interval where the Lyapunov exponent vanishes, then  $E \mapsto N_{\lambda,\alpha}(E)$  is an increasing analytic function of  $E \in J^2$  (and obviously  $J \subset \Sigma_{\lambda,\alpha}$ ).

We will use several times the following result.

**THEOREM 2.1** [BJ02, Cor. 2]. *Suppose  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$  and  $\lambda \neq 0$ . If  $E \in \Sigma_{\lambda,\alpha}$ , then*

$$L_{\lambda,\alpha}(E) = \max\{0, \ln|\lambda|\}.$$

This result will be mostly important for us for what it says about the range  $0 < \lambda \leq 1$  (zero Lyapunov exponent on the spectrum). It will be also used in very minor way in our proof of localization when  $\lambda > 1$ .

---

<sup>2</sup> This is because  $N + iL/\pi$  is holomorphic in upper half-plane and real on  $J$ . It can also be obtained from the Thouless formula.

2.3. *Kotani theory.* Recall the usual action of  $\mathrm{SL}(2, \mathbb{C})$  on the Riemann sphere  $\bar{\mathbb{C}}$ :  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot z = (az + b)/(cz + d)$ . We can of course define  $\mathrm{SL}(2, \mathbb{C})$  cocycles as pairs  $(\alpha, A) \in \mathbb{R} \times C^0(\mathbb{R}/\mathbb{Z}, \mathrm{SL}(2, \mathbb{C}))$ , but it is convenient to view an  $\mathrm{SL}(2, \mathbb{C})$  cocycle as acting by Möbius transformations:

$$\begin{aligned} (\alpha, A) : \mathbb{R}/\mathbb{Z} \times \bar{\mathbb{C}} &\rightarrow \mathbb{R}/\mathbb{Z} \times \bar{\mathbb{C}}, \\ (x, z) &\mapsto (x + \alpha, A(x) \cdot z). \end{aligned}$$

If one lets  $E$  become a complex number in the definition of the almost Mathieu cocycle, we get an  $\mathrm{SL}(2, \mathbb{C})$  cocycle.

Let  $\mathbb{H}$  be the upper half-plane. Fix  $(\lambda, \alpha)$ . It is well known that there exists a continuous function  $m = m_{\lambda, \alpha} : \mathbb{H} \times \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{H}$  such that  $S_{\lambda, E}(x) \cdot m(E, x) = m(E, x + \alpha)$ , thus defining an invariant section for the cocycle  $(\alpha, S_{\lambda, E})$ :

$$(2-4) \quad (\alpha, S_{\lambda, E})(x, m(E, x)) = (x + \alpha, m(E, x + \alpha)).$$

Moreover,  $E \mapsto m(E, x)$  is holomorphic on  $\mathbb{H}$ .

*Remark 2.1.* In the literature (for instance, in [Sim83]), it is more common to find the definition of a pair  $m_{\pm}(x, E)$  of  $m$ -functions, which is given in terms of nonzero solutions  $(u_{\pm}(n))_{n \in \mathbb{Z}}$  of  $H_{\lambda, \alpha, x} u = Eu$  that are  $\ell^2$  at  $\pm\infty$ , that is,  $m_{\pm}(x, E) = -u_{\pm}(\pm 1)/u_{\pm}(0)$ . In this notation we have  $m(x, E) = -1/m_-(x, E)$ . The relation  $S_{\lambda, E}(x) \cdot m(E, x) = m(E, x + \alpha)$  is an immediate consequence of the definition of  $m_-(x, E)$ .

The following result of Kotani theory [Sim83], [DCJ87] will be important in two key parts of this paper.

**THEOREM 2.2.** *Let  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ , and assume that  $L_{\lambda, \alpha}(E) = 0$  in an open interval  $J \subset \mathbb{R}$ . Then for every  $x \in \mathbb{R}/\mathbb{Z}$ , the function  $E \mapsto m(E, x)$  admits a holomorphic extension to  $\mathbb{C} \setminus (\mathbb{R} \setminus J)$ , with values in  $\mathbb{H}$ . The function  $m : \mathbb{C} \setminus (\mathbb{R} \setminus J) \times \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{H}$  is continuous in both variables.*

2.4. *Polar sets.* Recall one of the possible definitions of a polar set in  $\mathbb{C}$ : it is a set of zero logarithmic capacity. We will need only some properties of polar sets in  $\mathbb{C}$  (see for instance [Hör94]):

- (i) A countable union of polar sets is polar.
- (ii) The image of a polar set by a nonconstant holomorphic function (defined in some domain of  $\mathbb{C}$ ) is a polar set.
- (iii) Polar sets have Hausdorff dimension zero; thus their intersections with  $\mathbb{R}$  have zero Lebesgue measure.
- (iv) Let  $U \subset \mathbb{C}$  be a domain, and let  $f_n : U \rightarrow \mathbb{R}$  be a sequence of subharmonic functions that is uniformly bounded in compacts of  $U$ . Then  $f : U \rightarrow \mathbb{R}$



given by  $f = \limsup f_n$  coincides with its (subharmonic) upper regularization  $f^* : U \rightarrow \mathbb{R}$  (given by  $f^*(z) = \limsup_{w \rightarrow z} f(w)$ ) outside a polar set.

We will say that a subset of  $\mathbb{R}$  is polar if it is polar as a subset of  $\mathbb{C}$ .

The following result on analytic continuation is well known. We will quickly go through the proof, since a similar idea will play a role later in a small divisor problem.

**LEMMA 2.3.** *Let  $W \subset \mathbb{C}$  be a domain, and let  $f : W \times \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{C}$  be a continuous function. If  $z \mapsto f(z, w)$  is holomorphic for all  $w \in \mathbb{R}/\mathbb{Z}$  and  $w \mapsto f(z, w)$  is analytic for some nonpolar set of  $z \in W$ , then  $f$  is analytic.*

*Proof.* We may assume that  $|f(z, w)| < 1$  for  $(z, w) \in W \times \mathbb{R}/\mathbb{Z}$ . Let

$$(2-5) \quad f(z, w) = \sum \hat{f}_z(k) e^{2\pi i k w}.$$

Then  $z \mapsto \hat{f}_z(k)$  is holomorphic, and  $|\hat{f}_z(k)| < 1$ . Using property (i) of polar sets, we obtain that there exists a nonpolar set  $\Delta \subset W$ ,  $\epsilon > 0$ , and  $k > 0$  such that  $|\hat{f}_z(n)| \leq e^{-\epsilon|n|}$  for  $z \in \Delta$  and  $|n| > k$ . Let

$$h(z) = \sup_{|n| > k} \frac{1}{|n|} \ln |\hat{f}_z(n)|.$$

Then, by property (iv) of polar sets,  $h^*$  is a nonpositive subharmonic function satisfying  $h^*(z) \leq -\epsilon$  for  $z \in \Delta \setminus X$ , where  $X$  is polar. Since  $\Delta$  is nonpolar, we conclude that  $h^*$  is not identically 0 in  $W$ . It follows from the maximum principle that  $h^*(z) < 0$  for  $z \in W$ . Thus for any domain  $U \subset W$  compactly contained in  $W$ , there exists a  $\delta = \delta(U) > 0$  such that  $h(z) \leq -\delta$  for  $z \in U$ . Therefore  $(\ln |\hat{f}_z(n)|)/|n| \leq -\delta$  for  $|n| > k$  and  $z \in U$ , which implies that (2-5) converges uniformly on compacts of  $W \times \{w \in \mathbb{C}/\mathbb{Z}, 2\pi |\operatorname{Im} w| < \delta\}$ .  $\square$

### 3. Regularity of the $m$ -functions

**THEOREM 3.1.** *Let  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$  and  $\lambda > 0$ . Let  $m = m_{\lambda, \alpha} : \mathbb{H} \times \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{H}$  be as in Section 2.3. Then  $m$  is analytic.*

*Proof.* Let us show that  $m$  has a holomorphic extension to

$$\Omega_\lambda = \{(E, x) : \operatorname{Im} E > 0, 2\lambda \sinh|2\pi \operatorname{Im} x| < \operatorname{Im} E\}.$$

We have  $S_{\lambda, E}(x) \cdot z = E - 2\lambda \cos(2\pi x) - 1/z$ . For  $(E, t)$  satisfying

$$(3-1) \quad \operatorname{Im} E > 0 \quad \text{and} \quad 2\lambda \sinh|2\pi t| < \operatorname{Im} E,$$

define the half-plane

$$K_{\lambda, E, t}^1 = \{z : \operatorname{Im} z > \operatorname{Im} E - 2\lambda \sinh|2\pi t|\} \subset \mathbb{H},$$

and the disk

$$K_{\lambda,E,t}^2 = \{|z| < |E| + 2\lambda \cosh|2\pi t| + 1/(\operatorname{Im} E - 2\lambda \sinh|2\pi t|)\},$$

and let  $K_{\lambda,E,t} = K_{\lambda,E,t}^1 \cap K_{\lambda,E,t}^2$ , which is a domain compactly contained in  $\mathbb{H}$  depending continuously on  $(E, t)$  satisfying (3-1). If  $(E, x) \in \Omega_\lambda$ , then  $(E, \operatorname{Im} x)$  satisfies (3-1) and one checks directly that

$$S_{\lambda,E}(x) \cdot \mathbb{H} \subset K_{\lambda,E,\operatorname{Im} x}^1 \quad \text{and} \quad S_{\lambda,E}(x) \cdot K_{\lambda,E,\operatorname{Im} x}^1 \subset K_{\lambda,E,\operatorname{Im} x}^2.$$

Since  $\operatorname{Im} x = \operatorname{Im}(x + \alpha)$ , we have

$$S_{\lambda,E}(x + \alpha) \cdot S_{\lambda,E}(x) \cdot \mathbb{H} \subset K_{\lambda,E,\operatorname{Im} x}.$$

Thus, by the Schwarz lemma applied to  $\mathbb{H}$ , for every  $(E, x) \in \Omega_\lambda$ ,

$$S_{\lambda,E}(x - \alpha) \cdots S_{\lambda,E}(x - n\alpha) \cdot \bar{\mathbb{H}}$$

is a sequence of nested compact sets shrinking to a single point  $\hat{m}(E, x)$ . This implies that  $\hat{m}(E, x)$  is the unique solution to (2-4) in  $\mathbb{H}$ . As  $m : \mathbb{H} \times \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{H}$  is a continuous function satisfying (2-4), we must have  $\hat{m}(E, x) = m(E, x)$  for  $(E, x) \in \mathbb{H} \times \mathbb{R}/\mathbb{Z}$ .

Since holomorphic functions  $m^n : \Omega_\lambda \rightarrow \mathbb{H}$  given by

$$m^n(E, x) = S_{\lambda,E}(x - \alpha) \cdots S_{\lambda,E}(x - n\alpha) \cdot i$$

take values in  $\mathbb{H}$ , the sequence  $m^n$  is normal. Since it converges pointwise to  $\hat{m}$ , we conclude that  $\hat{m}$  is holomorphic.  $\square$

**THEOREM 3.2.** *Let  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ , and let  $0 < \lambda \leq 1$ . Let  $m = m_{\lambda,\alpha} : \mathbb{H} \times \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{H}$  be as in Section 2.3. If  $J \subset \Sigma_{\lambda,\alpha}$  is an open interval, then  $m$  admits an analytic extension  $m : \mathbb{C} \setminus (\mathbb{R} \setminus J) \times \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{H}$ .*

*Proof.* By Theorems 2.1 and 2.2, there exists a continuous extension

$$m : \mathbb{C} \setminus (\mathbb{R} \setminus J) \times \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{H}$$

that is analytic in  $E$ . By Theorem 3.1,  $m$  is also analytic in  $x$  for  $E \in \mathbb{H}$ . Analyticity in  $(E, x)$  then follows by Lemma 2.3.  $\square$

#### 4. Analytic continuation

**LEMMA 4.1.** *Let  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ , let  $\phi : \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}$  be analytic, and let  $\theta = \int_{\mathbb{R}/\mathbb{Z}} \phi(x) dx$ . The following are equivalent:*

- (i) *There exists an analytic function  $O : \mathbb{R}/\mathbb{Z} \rightarrow \operatorname{SL}(2, \mathbb{R})$ , homotopic to the identity, such that  $O(x + \alpha) R_{\phi(x)} O(x)^{-1} = R_\theta$ .*
- (ii) *There exists an analytic function  $\psi : \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}$  such that*

$$(4-1) \quad \phi(x) - \theta = \psi(x + \alpha) - \psi(x).$$

*Proof.* Obviously (ii) implies (i): it is enough to take  $O(x) = R_{-\psi(x)}$ .

Let us show that (i) implies (ii). If  $O(x) \cdot i = i$  for all  $x$ , then we have  $O(x) \in \text{SO}(2, \mathbb{R})$  for all  $x$ , and since  $O$  is homotopic to the identity, we have  $O(x) = R_{-\psi(x)}$  for some analytic function  $\psi : \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}$  that must satisfy  $\phi(x) - \theta = \psi(x + \alpha) - \psi(x)$ .

Thus we may assume that  $O(x_0) \cdot i \neq i$  for some  $x_0$ . Notice that

$$O(x_0 + n\alpha) \cdot i = R_{n\theta} O(x_0) \cdot i.$$

It follows that if  $n_k \alpha \rightarrow 0$  in  $\mathbb{R}/\mathbb{Z}$ , then  $2n_k \theta \rightarrow 0$  in  $\mathbb{R}/\mathbb{Z}$ . This implies that  $\theta = (l/2)\alpha$  for some  $l \in \mathbb{Z}$ . We have

$$O(x + \alpha) R_{\phi(x)} O(x)^{-1} = R_{(l/2)(x+\alpha)} R_{-(l/2)x},$$

which implies  $R_{-(l/2)(x+\alpha)} O(x + \alpha) R_{\phi(x)} = R_{-(l/2)x} O(x)$ , and we get

$$R_{-(l/2)(x+\alpha)} O(x + \alpha) \cdot i = R_{-(l/2)x} O(x) \cdot i.$$

It follows that  $R_{-(l/2)x} O(x) \cdot i = z$  does not depend on  $x$ . Let  $Q \in \text{SL}(2, \mathbb{R})$  be such that  $Q \cdot z = i$ , and set

$$S(x) = R_{(l/2)x} Q R_{-(l/2)x} O(x).$$

Since  $O, Q : \mathbb{R}/\mathbb{Z} \rightarrow \text{SL}(2, \mathbb{R})$ , where  $Q(x) = Q$ , are homotopic to the identity,  $S : \mathbb{R}/\mathbb{Z} \rightarrow \text{SL}(2, \mathbb{R})$  is homotopic to the identity and, using that  $\theta = (l/2)\alpha$ , we have  $S(x + \alpha) R_{\phi(x)} S(x)^{-1} = R_\theta$ . Moreover,  $S(x) \cdot i = i$ , so  $S(x) \in \text{SO}(2, \mathbb{R})$ , and thus  $S(x) = R_{-\psi(x)}$  where  $\psi : \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}$ . It follows that  $\psi$  satisfies (4-1).  $\square$

For  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$  and  $0 < \lambda \leq 1$ , let  $\Lambda_{\lambda, \alpha}$  be the set of  $E$  such that there exists an analytic function  $B_E : \mathbb{R}/\mathbb{Z} \rightarrow \text{SL}(2, \mathbb{R})$ , homotopic to the identity, and  $\theta(E) \in \mathbb{R}$ , such that

$$B_E(x + \alpha) S_{\lambda, E}(x) B_E(x)^{-1} = R_{\theta(E)}.$$

**THEOREM 4.2.** *Let  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$  and  $0 < \lambda \leq 1$ . Let  $J \subset \Sigma_{\lambda, \alpha}$  be an open interval. Then*

- (i) if  $\beta = 0$ , then  $\Lambda_{\lambda, \alpha} \cap J = J$ ;
- (ii) if  $\beta < \infty$ , then either  $\Lambda_{\lambda, \alpha} \cap J$  is polar or  $\text{int } \Lambda_{\lambda, \alpha} \cap J \neq \emptyset$ .

*Proof.* Assume that  $J \subset \Sigma_{\lambda, \alpha}$  is an open interval. Let  $m = m_{\lambda, \alpha}$  be given by Theorem 3.2, so that  $m : \mathbb{C} \setminus (\mathbb{R} \setminus J) \times \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{H}$  is continuous,  $E \mapsto m(E, x)$  is holomorphic, and  $S_{\lambda, E} \cdot m(E, x) = m(E, x + \alpha)$ . Let

$$(4-2) \quad C_E(x) = \begin{pmatrix} \frac{\text{Re } m(E, x)}{|m(E, x)|(\text{Im } m(E, x))^{1/2}} & -\frac{|m(E, x)|}{(\text{Im } m(E, x))^{1/2}} \\ \frac{(\text{Im } m(E, x))^{1/2}}{|m(E, x)|} & 0 \end{pmatrix}.$$

Then  $C_E(x + \alpha)S_{\lambda,E}(x)C_E(x)^{-1} \in \text{SO}(2, \mathbb{R})$  for  $E \in J$  and  $x \in \mathbb{R}/\mathbb{Z}$ . Since  $x \mapsto C_E(x)$  is easily verified to be homotopic to the identity for  $E \in J$ , we have  $C_E(x + \alpha)S_{\lambda,E}(x)C_E(x)^{-1} = R_{\phi(E,x)}$  for some real-analytic function  $\phi : J \times \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}$ . It follows that  $\phi$  has a holomorphic extension  $\phi : Z \rightarrow \mathbb{C}$ , where  $Z \subset \mathbb{C} \times \mathbb{C}/\mathbb{Z}$  is some domain containing  $J \times \mathbb{R}/\mathbb{Z}$ . So there exists a domain  $\Delta \subset \mathbb{C}$  such that  $J \subset \Delta$  and  $\Delta \times \mathbb{R}/\mathbb{Z} \subset Z$ . For  $E \in \Delta$ , let

$$\phi(E, x) = \sum \hat{\phi}_E(k)e^{2\pi ikx}.$$

Let  $E \in J$  be such that there exists an analytic function  $\psi_E : \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}$  such that

$$\int_{\mathbb{R}/\mathbb{Z}} \psi_E(x)dx = 0 \quad \text{and} \quad \phi(E, x) - \int_{\mathbb{R}/\mathbb{Z}} \phi(E, x)dx = \psi_E(x + \alpha) - \psi_E(x).$$

Then

$$(4-3) \quad \psi_E(x) = \sum \hat{\psi}_E(k)e^{2\pi ikx},$$

$$\text{where } \hat{\psi}_E(k) = \begin{cases} \hat{\phi}_E(k)/(e^{2\pi ik\alpha} - 1) & \text{if } k \neq 0, \\ 0 & \text{if } k = 0. \end{cases}$$

We can then define an analytic function  $B_E : \mathbb{R}/\mathbb{Z} \rightarrow \text{SL}(2, \mathbb{R})$  by

$$B_E(x) = R_{-\psi_E(x)}C_E(x),$$

which satisfies

$$B_E(x + \alpha)S_{\lambda,E}(x)B_E(x)^{-1} = R_{\theta(E)} \quad \text{and} \quad \theta(E) = \int_{\mathbb{R}/\mathbb{Z}} \phi(E, x)dx.$$

Reciprocally, if there exists an analytic function  $B_E : \mathbb{R}/\mathbb{Z} \rightarrow \text{SL}(2, \mathbb{R})$  homotopic to the identity such that  $B_E(x + \alpha)S_{\lambda,E}(x)B_E(x)^{-1} = R_{\theta(E)}$  for some  $\theta(E) \in \mathbb{R}$ , then we can write

$$O_E(x + \alpha)R_{\phi(E,x)}O_E(x)^{-1} = R_{\theta(E)}, \quad \text{where } O_E(x) = B_E(x)C_E(x)^{-1}.$$

By the previous lemma, there exists an analytic function (having average 0)  $\psi : \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}$  satisfying  $\phi(x) - \int_{\mathbb{R}/\mathbb{Z}} \phi(x)dx = \psi(x + \alpha) - \psi(x)$ .

Notice that

$$(4-4) \quad \limsup_{|k| \rightarrow \infty} \frac{1}{|k|} \ln \frac{1}{|e^{2\pi ik\alpha} - 1|} = \beta,$$

so that if  $\beta = 0$ , then (4-3) really defines an analytic function for any  $E \in J$ ; thus (i) follows.

Let  $a : \Delta \rightarrow [-\infty, \beta]$  be given by

$$a(E) = \limsup_{|k| \rightarrow \infty} \frac{1}{|k|} \ln \left| \frac{\hat{\phi}_E(k)}{e^{2\pi ik\alpha} - 1} \right|.$$

By the previous discussion,  $\Lambda_{\lambda,\alpha} = \{E \in J : a(E) < 0\}$ . If  $\beta < \infty$ , then  $a$  is lim sup of a sequence of subharmonic functions which are uniformly bounded on compacts of  $\Delta$ . It follows that  $a$  coincides with its upper regularization

$$a^*(E) = \limsup_{E' \rightarrow E} a(E')$$

for  $E$  outside some polar exceptional set. Thus the set  $\{E \in J : a(E) < 0\}$  is either polar (contained in the exceptional set) or it has nonempty interior.  $\square$

LEMMA 4.3. *Let  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$  and  $\lambda > 0$ . Then  $\Lambda_{\lambda,\alpha}$  has empty interior.*

*Proof.* We may assume that  $0 < \lambda \leq 1$  (otherwise the Lyapunov exponent is positive on  $\Sigma_{\lambda,\alpha}$ , which easily implies that  $\Lambda_{\lambda,\alpha} = \emptyset$ ). Assume that  $J \subset \Lambda_{\lambda,\alpha}$  is an open interval. Then  $J \subset \Sigma_{\lambda,\alpha}$  (since  $L_{\lambda,\alpha}(E) = 0$  for  $E \in J$ ). Let  $B_E$  be as in the definition of  $\Lambda_{\lambda,\alpha}$ . Then the definition of fibered rotation number (see Section 2.1) implies  $\rho_{\lambda,\alpha}(E) = \theta(E) \pmod{\mathbb{Z}}$ . By the analyticity of  $\rho$  on  $J$  (see note 2), there exist  $E \in J$  and  $l \in \mathbb{Z}$  such that  $\theta(E) = l\alpha \pmod{\mathbb{Z}}$ . Let  $T_E : \mathbb{R}/\mathbb{Z} \rightarrow \text{SL}(2, \mathbb{R})$  be given by  $T_E(x) = R_{-lx} B_E(x)$ . Then

$$T_E(x + \alpha) S_{\lambda,E}(x) T_E(x)^{-1} = \text{id}.$$

The conclusion is as in [Pui04]. For  $v \in \mathbb{R}^2$ ,

$$S_{\lambda,E}(x) T_E(x)^{-1} v = T_E(x + \alpha)^{-1} v.$$

So by (2-3) there exists an analytic  $U_v : \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}$  such that

$$T_E(x + \alpha)^{-1} \cdot v = \begin{pmatrix} U_v(x) \\ U_v(x - \alpha) \end{pmatrix}.$$

Let  $U_v(x) = \sum u_n^v e^{2\pi i n x}$ . It is a standard Aubry duality argument (and can be checked by direct calculation) that  $u_n^v \in \ell^2(\mathbb{Z})$  is an eigenvector of  $H_{\lambda^{-1},\alpha,0}$  with eigenvalue  $\lambda^{-1}E$ . The fact that we get such an eigenvector for every  $v \in \mathbb{R}^2$  contradicts the simplicity of the point spectrum.  $\square$

*Remark 4.1.* Notice that Lemma 4.3 and Theorem 4.2(i) already imply the Ten Martini problem in the case  $\beta = 0$ , and we did not need any localization result (the only recent result we used was Theorem 2.1).

### 5. Localization and Cantor spectrum

We say that the operator  $H_{\lambda,\alpha,\theta}$  displays *Anderson localization* if it has pure point spectrum with exponentially decaying eigenvectors. This requires  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ , and implies that eigenvalues are dense in  $\Sigma_{\lambda,\alpha}$ .

THEOREM 5.1. *Let  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ , and let  $\lambda \geq 1$ . Assume that  $\beta < \infty$ . If  $H_{\lambda,\alpha,\theta}$  displays Anderson localization for a nonpolar set of  $\theta \in \mathbb{R}$ , then  $\Sigma_{\lambda,\alpha}$  is a Cantor set.*

*Proof.* Let  $\Theta$  be the set of  $\theta$  such that  $H_{\lambda,\alpha,\theta}$  displays Anderson localization. If  $\theta \in \Theta$  and  $E$  is an eigenvalue for  $H_{\lambda,\alpha,\theta}$ , let  $(u_n)_{n \in \mathbb{Z}}$  be a nonzero eigenvector. Then

$$S_{\lambda^{-1},\lambda^{-1}E} \cdot W(x) = e^{2\pi i\theta} W(x + \alpha),$$

where

$$W(x) = \begin{pmatrix} U(x)e^{2\pi i\theta} \\ U(x - \alpha) \end{pmatrix} \quad \text{and} \quad U(x) = \sum u_n e^{2\pi i n x}.$$

Let  $M(x)$  be the matrix with columns  $W(x)$  and  $\overline{W(x)}$ . Then

$$S_{\lambda^{-1},\lambda^{-1}E}(x) \cdot M(x) = M(x + \alpha) \begin{pmatrix} e^{2\pi i\theta} & 0 \\ 0 & e^{-2\pi i\theta} \end{pmatrix}.$$

This implies that  $\det M(x)$  is independent of  $x$ , so  $\det M(x) = ci$  for some  $c \in \mathbb{R}$ . Notice that if  $c = 0$ , then

$$V(x + \alpha) = e^{-4\pi i\theta} V(x),$$

with  $V(x) = U(x)/\overline{U(x)}$  (observe that  $U(x) \neq 0$  except at finitely many  $x$  since  $U(x)$  is a nonconstant analytic function), and in particular, if  $n_k \alpha \rightarrow 0$ , then  $2n_k \theta \rightarrow 0$ . So  $2\theta = k\alpha + l$  for some  $k, l \in \mathbb{Z}$ . If  $c > 0$ , we have

$$S_{\lambda^{-1},\lambda^{-1}E}(x) = Q(x + \alpha) R_\theta Q(x)^{-1},$$

where  $Q : \mathbb{R}/\mathbb{Z} \rightarrow \text{SL}(2, \mathbb{R})$  is given by

$$(5-1) \quad Q(x) = \frac{1}{(2c)^{1/2}} M(x) \begin{pmatrix} 1 & i \\ 1 & -i \end{pmatrix}$$

and if  $c < 0$ , we have

$$S_{\lambda^{-1},\lambda^{-1}E}(x) = Q(x + \alpha) R_{-\theta} Q(x)^{-1},$$

where  $Q : \mathbb{R}/\mathbb{Z} \rightarrow \text{SL}(2, \mathbb{R})$  is given by

$$(5-2) \quad Q(x) = \frac{1}{(-2c)^{1/2}} M(x) \begin{pmatrix} 1 & i \\ 1 & -i \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

It follows that in either case  $\lambda^{-1}E \in \Lambda_{\lambda^{-1},\alpha}$ , and moreover,

$$\rho_{\lambda^{-1},\alpha}(\lambda^{-1}E) = \pm \theta + k\alpha \pmod{\mathbb{Z}} \quad \text{for some } k \in \mathbb{Z}$$

(note that  $Q$  defined by (5-1) or (5-2) is not necessarily homotopic to the identity).

Let  $\Theta' \subset \Theta$  be the set of all  $\theta$  such that  $2\theta \neq k\alpha + l$  for all  $k, l \in \mathbb{Z}$ . Let  $J \subset \Sigma_{\lambda,\alpha}$  be an open interval. Then for any  $\theta \in \Theta'$ , there exists some  $E \in J$  such that  $E$  is an eigenvalue for  $H_{\lambda,\alpha,\theta}$ , and by the previous discussion any such  $E$

satisfies

$$N_{\lambda^{-1},\alpha}(\lambda^{-1}E) = 1 - 2\rho_{\lambda^{-1},\alpha}(\lambda^{-1}E) = 1 - 2(\varepsilon\theta + k\alpha + l)$$

for some  $k, l \in \mathbb{Z}$ ,  $\varepsilon \in \{1, -1\}$ , and  $\lambda^{-1}E \in \Lambda_{\lambda^{-1},\alpha}$ .

It follows that

$$\Theta' \subset \left\{ \frac{\varepsilon}{2}(1 - N_{\lambda^{-1},\alpha}(\Lambda_{\lambda^{-1},\alpha} \cap \lambda^{-1}J)) - k\alpha - l : k, l \in \mathbb{Z}, \varepsilon \in \{1, -1\} \right\}.$$

By Theorem 4.2(ii) and Lemma 4.3,  $\Lambda_{\lambda^{-1},\alpha} \cap \lambda^{-1}J$  is polar. Since  $L = 0$  on  $\lambda^{-1}J$ , we have that  $N_{\lambda^{-1},\alpha}$  is a nonconstant analytic function on  $\lambda^{-1}J$ ; thus it follows that  $\Theta'$  is also polar. Therefore  $\Theta \subset \Theta' \cup \{\frac{1}{2}(k\alpha + l) : k, l \in \mathbb{Z}\}$  is polar, which is a contradiction.  $\square$

*Remark 5.1.* In [Pui04], it is shown that if  $\alpha \in DC$ , then Anderson localization of  $H_{\lambda,\alpha,0}$  implies Cantor spectrum. We can not however use the argument of Puig (based on analytic reducibility) to conclude Cantor spectrum in the generality we need. Indeed, we are not able to conclude analytic reducibility from localization of  $H_{\lambda,\alpha,0}$  in our setting (in a sense, we spend all our regularity to take care of small divisors in the localization result, which is half of analytic reducibility, and there is nothing left for the other half). Though this can be bypassed (using Kotani theory to conclude continuous reducibility under the assumption of non-Cantor spectrum), there is a much more serious difficulty in this approach; see Remark 5.2.

The next result gives us a large range of  $\lambda$  and  $\alpha$  where Theorem 5.1 can be applied.

**THEOREM 5.2.** *Let  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$  be such that  $\beta = \beta(\alpha) < \infty$ , and let  $\lambda > e^{16\beta/9}$ . Then  $H_{\lambda,\alpha,\theta}$  displays Anderson localization for almost every  $\theta$ .*

This result improves on [Jit99], where Anderson localization was proved under the assumption that  $\alpha$  is Diophantine. Recall that  $\alpha$  is said to satisfy a Diophantine condition (briefly,  $\alpha \in DC$ ) if  $\ln q_{n+1} = O(\ln q_n)$ , where  $p_n/q_n$  are the rational approximations of  $\alpha$ . In particular  $\alpha \in DC$  implies (but is strictly stronger than)  $\beta(\alpha) = 0$ . The proof in [Jit99] with some modifications can be extended to the case  $\beta(\alpha) = 0$  but not to the case  $\beta(\alpha) > 0$ .

The proof of Theorem 5.2 is the most technical part of this paper, and the considerations involved are independent from our other arguments. We will thus postpone it to Section 9.

*Remark 5.2.* We expect that the operator  $H_{\lambda,\alpha,0}$  does not display Anderson localization for  $1 < \lambda \leq e^{2\beta}$ . The key reason is that in this regime 0 is a very resonant phase, and since  $\alpha$  is Diophantine only in a very weak sense, the compound effect on the small divisors can not be compensated by the Lyapunov exponent. See also Remark 9.1.

## 6. Fictitious results on continuity of the spectrum

The spectrum  $\Sigma_{\lambda,\alpha}$  is a continuous function of  $\alpha$  in the Hausdorff topology. There are several results in the literature about quantitative continuity. The best general result, due to [AvMS90], concerns  $1/2$ -Hölder continuity. Better estimates can be obtained for  $\alpha$  not very Liouville in the region of positive Lyapunov exponent [JK02]. None of those results are enough for our purposes.

The results described above have something in common: they deal with something that actually happens, and it is not clear if it is possible to improve them sufficiently (to the level we need). Thus we will argue by contradiction: assuming the spectrum is not Cantor, we will get very good continuity estimates. This will allow us to proceed the argument, but obviously, since we will eventually conclude that the spectrum is a Cantor set, estimates in this section are not valid for any existing almost Mathieu operator. Those estimates might be useful also when analyzing more general Schrödinger operators.

**THEOREM 6.1.** *Let  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$  and  $0 < \lambda \leq 1$ . Let  $J \subset \mathbb{R}$  be an open interval such that  $\bar{J} \subset \text{int } \Sigma_{\lambda,\alpha}$ . There exists  $K > 0$  such that*

$$|N_{\lambda,\alpha}(E) - N_{\lambda,\alpha'}(E)| \leq K|\alpha - \alpha'| \quad \text{for } E \in J.$$

*Proof.* Let  $m = m_{\lambda,\alpha}$  be as in Theorem 3.2. Define  $x \mapsto C_E(x)$  by (4-2). Then, as discussed in the proof of Theorem 4.2,  $C_E : \mathbb{R}/\mathbb{Z} \rightarrow \text{SL}(2, \mathbb{R})$  is homotopic to the identity, with  $C_E(x + \alpha)S_{\lambda,E}(x)C_E(x)^{-1} \in \text{SO}(2, \mathbb{R})$ . So

$$C_E(x + \alpha)S_{\lambda,E}(x)C_E(x)^{-1} = R_{\phi_E(x)},$$

where  $\phi_E : \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}$  is analytic. Recall the definition of the fibered rotation number in Section 2.1. Then  $\rho(\alpha, S_{\lambda,E}(x)) = \rho(\alpha, R_{\phi(x)})$ . In this case we can take as lift of  $R_{\phi_E(x)}$  the function  $\psi(x, y) = \phi(x)$ .

Write

$$\begin{aligned} \rho(\alpha', S_{\lambda,E}) &= \rho(\alpha', C_E(x + \alpha')S_{\lambda,E}(x)C_E(x)^{-1}) \\ &= \rho(\alpha', C_E(x + \alpha')C_E(x + \alpha)^{-1}R_{\phi_E(x)}). \end{aligned}$$

Since  $m$  is analytic in  $x$ , we can take as lift of  $C_E(x + \alpha')C_E(x + \alpha)^{-1}R_{\phi_E(x)}$  a function  $\tilde{\psi}(x, y)$  satisfying  $|\tilde{\psi}(x, y) - \phi(x)| \leq K|\alpha - \alpha'|$ . Thus

$$\|\rho(\alpha, S_{\lambda,E}) - \rho(\alpha', S_{\lambda,E})\|_{\mathbb{R}/\mathbb{Z}} \leq \int \sup_y |\phi(x) - \tilde{\psi}(x, y)| dx \leq K|\alpha - \alpha'|.$$

The result now follows, since  $N = 1 - 2\rho$  (see Section 2.2) for the determination of  $\rho$  in  $[0, 1/2]$ .  $\square$

*Remark 6.1.* Clearly we also get the fictitious estimate

$$|L_{\lambda,\alpha'}(E) - L_{\lambda,\alpha}(E)| \leq K|\alpha - \alpha'| \quad \text{for } E \in J.$$



### 7. Gaps for rational approximants

It is well known for any  $\lambda \neq 0$  that if  $p/q$  is an irreducible fraction then  $\Sigma_{\lambda, p/q}$  consists of  $q$  bands with disjoint interior. All those bands are actually disjoint, except if  $q$  is even, when there are two bands touching at 0; see [vM89], [CEY90]. The variation of  $N_{\lambda, p/q}$  in each band is precisely  $1/q$ . The connected components of  $\mathbb{R} \setminus \Sigma_{\lambda, p/q}$  are called gaps. Let  $M(\lambda, p/q)$  be the maximum size of the bands of  $\Sigma_{\lambda, p/q}$ .

The following result is well known.

LEMMA 7.1. *Let  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ ,  $\lambda \neq 0$ . If  $p_n/q_n \rightarrow \alpha$ , then  $M(\lambda, p_n/q_n) \rightarrow 0$ . In particular (since  $N_{\lambda, p_n/q_n} \rightarrow N_{\lambda, \alpha}$  uniformly), if one selects a point  $a_{n,i}$  in each band of  $\Sigma_{\lambda, p_n/q_n}$ , then*

$$(7-1) \quad \frac{1}{q_n} \sum_i \delta a_{n,i} \rightarrow dN_{\lambda, \alpha} \quad \text{in the weak* topology.}$$

In [CEY90], a lower bound for the size of gaps of  $\Sigma_{\lambda, p/q}$  is derived of the form  $C(\lambda)^{-q}$ , where, for instance,  $C(1) = 8$ . We will need the following sharpening of this estimate, in the case where  $p/q$  are close to a given irrational number.

THEOREM 7.2. *Let  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$  and let  $0 < \lambda \leq 1$ . Let  $p_n/q_n \rightarrow \alpha$ . For every  $\epsilon > 0$  and for every  $n$  sufficiently large, all gaps of  $\Sigma_{\lambda, p_n/q_n}$  have size at least  $e^{-\epsilon q_n} \lambda^{q_n/2}$ .*

*Proof.* It is known (see the proof of [CEY90, Th. 3.3] for the case  $\lambda = 1$ , the general case being obtained as described in the proof of [CEY90, Cor. 3.4]) that for any bounded gap  $G$  of  $\Sigma_{\lambda, p/q}$ , one can find a sequence  $a_j$  for  $1 \leq j \leq q$ , with one  $a_j$  in each band of  $\Sigma_{\lambda, p/q}$ , such that  $G = (a_i, a_{i+1})$  for some  $1 \leq i \leq q - 1$ , and  $\prod_{j \neq i} |a_j - a_i| \geq \lambda^m$ , where  $q = 2m + 1$  or  $q = 2m + 2$ .

Let  $G_n$  be a bounded gap of  $\Sigma_{\lambda, p_n/q_n}$  of minimal size. Then

$$(7-2) \quad |G_n| \geq \lambda^{q_n/2} \prod_{j \neq i_n, i_n+1} |a_{n,j} - a_{n,i_n}|^{-1},$$

where the  $a_{n,i}$  satisfy the hypothesis of the previous lemma. Passing to a subsequence, we may assume that  $a_{n,i_n} \rightarrow E \in \Sigma_{\lambda, \alpha}$  and  $|G_n| \rightarrow 0$  (otherwise the result is obvious). By (7-2), we get that for  $0 < \delta < 1$  and for  $n$  large we have

$$\begin{aligned} \frac{1}{q_n} \ln(|G_n| \lambda^{-q_n/2}) &\geq -\frac{1}{q_n} \sum_{j \neq i_n, i_n+1} \ln |a_{n,j} - a_{n,i_n}| \\ &\geq -\frac{1}{q_n} \sum_{|a_{n,j} - a_{n,i_n}| > \delta} \ln |a_{n,j} - a_{n,i_n}|, \end{aligned}$$

which implies by (7-1) and the definition of the weak\* topology that

$$\liminf \frac{1}{q_n} \ln(|G_n| \lambda^{-q_n/2}) \geq - \int_{|E'-E|>\delta} \ln|E - E'| dN_{\lambda,\alpha}(E').$$

Thus

$$\liminf \frac{1}{q_n} \ln(|G_n| \lambda^{-q_n/2}) \geq - \int \ln|E - E'| dN_{\lambda,\alpha}(E').$$

By the Thouless formula and Theorem 2.1, this gives

$$\liminf \frac{1}{q_n} \ln(|G_n| \lambda^{-q_n/2}) \geq -L_{\lambda,\alpha}(E) = 0. \quad \square$$

*Remark 7.1.* It is possible to get an estimate on the convergence rate in Lemma 7.1 using [AvMS90]. This implies an estimate on the rate of convergence in Theorem 7.2.

### 8. Proof of the Main Theorem

We now put together the results of the previous sections. Recall that it is enough to consider  $\lambda > 0$ , and that the case  $\lambda = 1$  follows from [AK06, Th. 1.5]. Moreover, Cantor spectrum for  $\lambda$  implies Cantor spectrum for  $1/\lambda$ . Let  $\beta = \beta(\alpha)$ . The Main Theorem follows then from the following.

**THEOREM 8.1.** *Let  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ .*

- (i) *If  $\beta < \infty$  and  $\lambda > e^{16\beta/9}$ , then  $\Sigma_{\lambda,\alpha}$  is a Cantor set.*
- (ii) *If  $\beta = \infty$  or if  $0 < \beta < \infty$  and  $e^{-2\beta} < \lambda \leq 1$ , then  $\Sigma_{\lambda,\alpha}$  is a Cantor set.*

*Proof.*

Item (i) follows from Theorems 5.2 and 5.1.

To get item (ii), we argue by contradiction. Let  $J \subset \text{int } \Sigma$  be a compact interval. Then the density of states satisfies  $dN/dE \geq c > 0$  for  $E \in J$  [AS83]. Let  $p/q$  be close to  $\alpha$  such that  $(1/q) \ln|\alpha - p/q|$  is close to  $-\beta$ . By Lemma 7.1 and Theorem 7.2,  $J \setminus \Sigma_{p/q}$  contains an interval  $G = (a, b)$  of size  $e^{-\epsilon q} \lambda^{q/2}$ . Notice that  $N_{\lambda,p/q}(a) = N_{\lambda,p/q}(b)$ . Theorem 6.1 implies

$$|N_{\lambda,\alpha}(a) - N_{\lambda,\alpha}(b)| \leq K |\alpha - p/q| \leq e^{\epsilon q} e^{-\beta q}.$$

Thus

$$c \leq \frac{N_{\lambda,\alpha}(a) - N_{\lambda,\alpha}(b)}{a - b} \leq e^{2\epsilon q} e^{-\beta q} \lambda^{-q/2}.$$

By taking  $\epsilon \rightarrow 0$ , we conclude that  $\lambda \leq e^{-2\beta}$ . □

Let us point out that  $1/2$ -Hölder continuity of the spectrum [AvMS90] (which holds for every  $\alpha$  and  $\lambda$ ) together with Theorem 7.2 implies the following improvement of [CEY90]. Let us say that all gaps of  $\Sigma_{\lambda,\alpha}$  are open if whenever  $E \in \Sigma_{\lambda,\alpha}$  is

such that  $N_{\lambda,\alpha}(E) = k\alpha + l$  for some  $k \in \mathbb{Z} \setminus \{0\}$  and  $l \in \mathbb{Z}$ , then  $E$  is the endpoint of some bounded gap (this obviously implies Cantor spectrum). The conjecture that  $\Sigma_{\lambda,\alpha}$  has all gaps open for all  $\lambda \neq 0$  and  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$  is sometimes called the “dry” version of the Ten Martini Problem.

**THEOREM 8.2.** *Let  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$  and let  $\beta = \beta(\alpha)$ . If  $\beta = \infty$  or if  $0 < \beta < \infty$  and  $e^{-\beta} < \lambda < e^\beta$ , then  $\Sigma_{\lambda,\alpha}$  has all gaps open.*

The conclusion from Theorem 8.2 appears to be the natural boundary of what can be taken honestly from the Liouvillian method: our computations indicate that although one can get improved estimates on continuity of the spectrum for  $\lambda > e^\beta$  (following [JK02]), things seem to break up at the precise parameter  $\lambda = e^\beta$ . Notice that  $\lambda = e^\beta$  is the expected threshold for localization (for almost every phase)<sup>3</sup> and falls short of the expected threshold for localization with phase  $\theta = 0$  and  $\lambda = e^{2\beta}$ . Thus the use of fictitious estimates does not seem to be an artifact of our estimates, but a rather essential aspect of an approach that tries to cover all parameters with Diophantine and Liouvillian techniques.

*Remark 8.1.* We do not actually need the measure-theoretical result of [AK06] to obtain Cantor spectrum for  $|\lambda| = 1$ . Indeed, Lemma 4.3 and Theorem 4.2(i) imply Cantor spectrum for  $\beta = 0$  (any  $\lambda \neq 0$ ; see Remark 4.1), and Theorem 8.1(ii) implies Cantor spectrum for  $\beta > 0$  (if  $|\lambda| = 1$ ).

### 9. Proof of Theorem 5.2

We will actually prove a slightly more precise version of Theorem 5.2. Let

$$(9-1) \quad \mathcal{R} = \{\theta : |\sin 2\pi(\theta + (k/2)\alpha)| < k^{-2} \text{ holds for infinitely many } ks\} \\ \cup \{s\pi\alpha/2 : s \in \mathbb{Z}\}.$$

$\mathcal{R}$  is easily seen to have zero Lebesgue measure by the Borel-Cantelli lemma.

**THEOREM 9.1.** *Let  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$  be such that  $\beta = \beta(\alpha) < \infty$ , and let  $\lambda > e^{16\beta/9}$ . Then  $H_{\lambda,\alpha,\theta}$  displays Anderson localization for  $\theta \notin \mathcal{R}$ .*

*Remark 9.1.* For  $\beta = 0$  the theorem holds as well for  $\theta = s\pi\alpha/2$ ; however the proof as presented here will not work. See [JKS05] for the details of the argument needed for this case. In general, we believe that for  $\theta$  of the form  $s\pi\alpha/2$  with  $s \in \mathbb{Z}$ , the localization would only hold for  $\lambda > e^{2\beta}$ .

*Remark 9.2.* We believe that for  $\theta \notin \mathcal{R}$ , localization should hold for  $\lambda > e^\beta$ . The proof of this fact would require some additional arguments. Moreover, for  $\lambda \leq e^\beta$ , we do not expect any exponentially decaying eigenvectors.

---

<sup>3</sup>In particular, by Gordon’s argument enhanced with Theorem 2.1,  $H_{\lambda,\alpha,\theta}$  has no eigenvalues for  $\lambda < e^\beta$ , and no localized eigenfunctions for  $\lambda = e^\beta$ .

*Remark 9.3.* The bound  $k^{-2}$  in (9-1) can be replaced by any other sub-exponential function without significant changes in the proof.

We will use the general setup of [Jit99]; however our key technical procedure will have to be quite different.

A formal solution  $\Psi_E(x)$  of the equation  $H_{\lambda,\alpha,\theta}\Psi_E = E\Psi_E$  will be called a *generalized eigenfunction* if

$$(9-2) \quad |\Psi_E(x)| \leq C(1 + |x|) \quad \text{for some } C = C(\Psi_E) < \infty.$$

The corresponding  $E$  is called a *generalized eigenvalue*. It is well known that to prove Theorem 9.1 it suffices to prove that generalized eigenfunctions decay exponentially [Ber68].

We will use the notation  $G_{[x_1,x_2]}(x, y)$  for matrix elements of the Green's function  $(H - E)^{-1}$  of the operator  $H_{\lambda,\alpha,\theta}$  restricted to the interval  $[x_1, x_2]$  with zero boundary conditions at  $x_1 - 1$  and  $x_2 + 1$ . We now fix  $\lambda, \alpha$  as in Theorem 9.1.

Fix a generalized eigenvalue  $E$ , and let  $\Psi$  be the corresponding generalized eigenfunction. Then

$$(9-3) \quad L(E) = \ln \lambda > 0.$$

$\lambda$  will enter into our analysis through  $L$  only and it will be convenient to use  $L$  instead. To simplify notations, in some cases the  $E, \lambda, \alpha$ -dependence of various quantities will be omitted.

Fix  $m > 0$ . A point  $y \in \mathbb{Z}$  will be called  $(m, k)$ -regular if there exists an interval  $[x_1, x_2]$ ,  $x_2 = x_1 + k - 1$ , containing  $y$ , such that

$$|G_{[x_1,x_2]}(y, x_i)| < e^{-m|y-x_i|} \quad \text{and} \quad \text{dist}(y, x_i) \geq k/40 \quad \text{for } i = 1, 2.$$

Otherwise,  $y$  will be called  $(m, k)$ -singular.

It is well known and can be checked easily that values of any formal solution  $\Psi$  of the equation  $H\Psi = E\Psi$  at a point  $x \in I = [x_1, x_2] \subset \mathbb{Z}$  can be reconstructed from the boundary values via

$$(9-4) \quad \Psi(x) = -G_I(x, x_1)\Psi(x_1 - 1) - G_I(x, x_2)\Psi(x_2 + 1).$$

This implies that if  $\Psi_E$  is a generalized eigenfunction, then every point  $y \in \mathbb{Z}$  with  $\Psi_E(y) \neq 0$  is  $(m, k)$ -singular for  $k$  sufficiently large, that is, for  $k$  greater than some  $k_1(E, m, \theta, y)$ . We assume without loss of generality that  $\Psi(0) \neq 0$  and normalize  $\Psi$  so that  $\Psi(0) = 1$ . Our strategy will be to show first that every sufficiently large  $y$  is  $(m, \ell(y))$ -regular for appropriate  $(m, \ell)$ . While  $\ell$  will vary with  $y$ ,  $m$  will have a uniform lower bound. This will be shown in Sections 9.4 and 9.3. Exponential decay will be derived out of this property via a ‘‘patching argument’’ in Section 9.1.

Let us denote

$$P_k(\theta) = \det((H_{\lambda,\alpha,\theta} - E)|_{[0,k-1]}).$$

Then the  $k$ -step transfer-matrix  $A_n(\theta)$  (which is the  $k$ -th iterate of the almost Mathieu cocycle  $A_k(\theta) = S_{\lambda,E}(\theta + (k - 1)\alpha) \cdots S_{\lambda,E}(\theta)$ ) can be written as

$$(9-5) \quad A_k(\theta) = \begin{pmatrix} P_k(\theta) & -P_{k-1}(\theta + \alpha) \\ P_{k-1}(\theta) & -P_{k-2}(\theta + \alpha) \end{pmatrix}.$$

Herman’s subharmonicity trick [Her83] yields  $\int_0^1 \ln|P_k(\theta)|d\theta \geq k \ln \lambda$ ; together with (9-3), this implies that there is a  $\theta \in [0, 1]$  with  $|P_k(\theta)| \geq e^{kL(E)}$ . Note that this is the only place in the proof of localization where we have used (9-3). While this is not really necessary (the rest of the proof can proceed, with only minor technical changes, under the assumption of the lower bound on only one of the four matrix elements, which follows immediately from the positivity of  $L(E)$ ), it simplifies certain arguments in what follows.

By applying Cramer’s rule we have for any  $x_1$  and  $x_2 = x_1 + k - 1$ , with  $x_1 \leq y \leq x_2$ , that

$$(9-6) \quad \begin{aligned} |G_{[x_1,x_2]}(x_1, y)| &= \left| \frac{P_{x_2-y}(\theta + (y + 1)\alpha)}{P_k(\theta + x_1\alpha)} \right|, \\ |G_{[x_1,x_2]}(y, x_2)| &= \left| \frac{P_{y-x_1}(\theta + x_1\alpha)}{P_k(\theta + x_1\alpha)} \right|. \end{aligned}$$

The numerators in (9-6) can be bounded uniformly in  $\theta$  [Jit99], [Fur97]. Namely, for every  $E \in \mathbb{R}$ ,  $\epsilon > 0$ , there exists a  $k_2(\epsilon, E, \alpha)$  such that

$$(9-7) \quad |P_n(\theta)| < e^{(L(E)+\epsilon)n} \quad \text{for all } n > k_2(\epsilon, E, \alpha) \text{ and all } \theta.$$

$P_k(\theta)$  is an even function of  $\theta + \frac{1}{2}(k - 1)\alpha$  and can be written as a polynomial of degree  $k$  in  $\cos 2\pi(\theta + \frac{1}{2}(k - 1)\alpha)$ :

$$P_k(\theta) = \sum_{j=0}^k c_j \cos^j 2\pi\left(\theta + \frac{1}{2}(k - 1)\alpha\right) \stackrel{\text{def}}{=} Q_k\left(\cos 2\pi\left(\theta + \frac{1}{2}(k - 1)\alpha\right)\right).$$

Let  $A_{k,r} = \{\theta \in \mathbb{R} : |Q_k(\cos 2\pi\theta)| \leq e^{(k+1)r}\}$ . The next lemma shows that every singular point “produces” a long piece of the trajectory of the rotation consisting of points belonging to an appropriate  $A_{k,r}$ .

LEMMA 9.2. *Suppose  $y \in \mathbb{Z}$  is  $(L - \epsilon, k)$ -singular and  $0 < \epsilon < L$ . Then for any  $\epsilon_1 > 0$ ,  $1/40 \leq \delta < 1/2$ , for sufficiently large  $k > k(\epsilon, E, \alpha, \epsilon_1, \delta)$ , and for any  $x \in \mathbb{Z}$  such that  $y - (1 - \delta)k \leq x \leq y - \delta k$ , we have that  $\theta + (x + (k - 1)\alpha/2)$  belongs to  $A_{k,L-\epsilon\delta+\epsilon_1}$ .*

*Proof.* This follows immediately from the definition of regularity, (9-6), and (9-7). □

The idea now is to show that  $A_{k,r}$  cannot contain  $k + 1$  uniformly distributed points. In order to quantify this concept of uniformity we introduce the following notion.

*Definition 9.1.* We will say that the set  $\{\theta_1, \dots, \theta_{k+1}\}$  is  $\epsilon$ -uniform if

$$(9-8) \quad \max_{z \in [-1,1]} \max_{j=1, \dots, k+1} \prod_{\substack{\ell=1 \\ \ell \neq j}}^{k+1} \frac{|z - \cos 2\pi \theta_\ell|}{|\cos 2\pi \theta_j - \cos 2\pi \theta_\ell|} < e^{k\epsilon}.$$

Note that we will use this terminology with “large” values of  $\epsilon$  as well.  $\epsilon$ -uniformity (the smaller  $\epsilon$  the better) involves uniformity along with certain cumulative repulsion of the  $\pm\theta_i \pmod 1$ .

LEMMA 9.3. *Let  $\epsilon_1 < \epsilon$ . If  $\theta_1, \dots, \theta_{k+1} \in A_{k,L-\epsilon}$  and  $k > k(\epsilon, \epsilon_1)$  is sufficiently large, then  $\{\theta_1, \dots, \theta_{k+1}\}$  is not  $\epsilon_1$ -uniform.*

*Proof.* Write polynomial  $Q_k(z)$  in the Lagrange interpolation form using  $\cos 2\pi \theta_1, \dots, \cos 2\pi \theta_{k+1}$ :

$$(9-9) \quad |Q_k(z)| = \left| \sum_{j=1}^{k+1} Q_k(\cos 2\pi \theta_j) \frac{\prod_{\ell \neq j} (z - \cos 2\pi \theta_\ell)}{\prod_{\ell \neq j} (\cos 2\pi \theta_j - \cos 2\pi \theta_\ell)} \right|.$$

Let  $\theta_0$  be such that  $|P_k(\theta_0)| \geq \exp(kL)$ . The lemma now follows immediately from (9-9) with  $z = \cos 2\pi(\theta_0 + \frac{1}{2}(k-1)\alpha)$ . □

Suppose we can find two intervals,  $I_1$  around 0 and  $I_2$  around  $y$ , of combined length  $|I_1| + |I_2| = k + 1$ ,<sup>4</sup> such that we can establish the uniformity of  $\{\theta_i\}$  where  $\theta_i = \theta + (x + \frac{1}{2}(k-1)\alpha)$  for  $i = 1, \dots, k + 1$  and  $x$  ranging through  $I_1 \cup I_2$ . Then we can apply Lemma 9.2 and 9.3 to show regularity of  $y$ . This is roughly going to be the framework for our strategy to establish regularity. The implementation will depend highly on the position of  $k$  with respect to the sequence of denominators  $q_n$ .

Assume without loss of generality that  $k > 0$ . Define

$$b_n = \max\{q_n^{8/9}, \frac{1}{20}q_{n-1}\}.$$

Find  $n$  such that  $b_n < k \leq b_{n+1}$ . We will distinguish two cases:

- (i) *Resonant* means  $|k - \ell q_n| \leq b_n$  for some  $\ell \geq 1$ .
- (ii) *Nonresonant* means  $|k - \ell q_n| > b_n$  for all  $\ell \geq 0$ .

---

<sup>4</sup>Here and in what follows, the “length”  $|I|$  of an interval  $I = [a, b] \subset \mathbb{Z}$  denotes cardinality:  $|I| = b - a + 1$ .

We will prove the following estimates.

LEMMA 9.4. *Assume  $\theta \notin \mathcal{R}$ . Suppose  $k$  is nonresonant. Let  $s \in \mathbb{N} \cup \{0\}$  be the largest number such that  $sq_{n-1} \leq \text{dist}(k, \{\ell q_n\}_{\ell \geq 0}) \equiv k_0$ . Then for any  $\epsilon > 0$  and for sufficiently large  $n$ , the following hold:*

- (i) *If  $s \geq 1$  and  $L > \beta$ , then  $k$  is  $(L - \ln q_n/q_{n-1} - \epsilon, 2sq_{n-1} - 1)$ -regular.*
- (ii) *If  $s = 0$ , then  $k$  is either  $(L - \epsilon, 2[q_{n-1}/2] - 1)$  or  $(L - \epsilon, 2[q_n/2] - 1)$  or  $(L - \epsilon, 2q_{n-1} - 1)$ -regular.*

LEMMA 9.5. *Let in addition  $L > \frac{16}{9}\beta$ . Then for sufficiently large  $n$ , every resonant  $k$  is  $(L/50, 2q_n - 1)$ -regular.*

We will prove Lemma 9.4 in Section 9.3 and Lemma 9.5 in Section 9.4. These two sections are not independent: the proof of Lemma 9.5 uses a corollary of the proof of Lemma 9.4 as an important ingredient. As our proofs rely on establishing  $\epsilon$ -uniformity of certain quasiperiodic sequences, we will repeatedly use estimates on trigonometric products proved in Section 9.2.

Theorem 9.1 can be immediately derived from Lemmas 9.4 and 9.5 via a “patching argument”, which we describe now. (A patching argument will also be used in one step of the proof of Lemma 9.5.)

9.1. *Patching. Proof of Theorem 9.1 assuming Lemmas 9.4 and 9.5.* It is an important technical ansatz of the multiscale analysis that the exponential decay of a Green’s function at a scale  $k$  under certain conditions generates exponential decay with the same rate at a larger scale. The proof is usually done using block-resolvent expansion, with the combinatorial factor being killed by the growth of scales. The proof of Theorem 9.1 will consist, roughly, of adapting this type of argument to our situation.

Fix a generalized eigenvalue  $E$  of  $H_{\lambda, \alpha, \theta}$ , and let  $\Psi$  be the corresponding generalized eigenfunction.

Assume without loss of generality that  $k$  is positive. Find  $n$  so that  $k > q_n$ . We assume that  $n$  is sufficiently large. Let  $L_1 = L/50 \leq L - \beta$ . By Lemmas 9.4 and 9.5 and the definition of regularity, for any  $y > b_n$  there exists an interval  $y \in I(y) = [x_1, x_2] \subset \mathbb{Z}$  such that

$$(9-10) \quad \text{dist}(y, \partial I(y)) > |I(y)|/40,$$

$$(9-11) \quad |I(y)| \geq q_n^{8/9} - 2,$$

$$(9-12) \quad G_{I(y)}(k, x_i) < e^{-L_1|k-x_i|} \quad \text{for } i = 1, 2.$$

In addition, if  $b_j < y \leq b_{j+1}$  we have

$$(9-13) \quad |I(y)| \leq 2q_j.$$

We denote the boundary of the interval  $I(y)$ , the set  $\{x_1, x_2\}$ , by  $\partial I(y)$ . For  $z \in \partial I(y)$  we let  $z'$  be the neighbor of  $z$  (i.e.,  $|z - z'| = 1$ ) not belonging to  $I(y)$ .

We now expand  $\Psi(x_2 + 1)$  in (9-4), iterating (9-4) with  $I = I(x_2 + 1)$ . In case  $q_n^{8/9} < x_1 - 1$  we also expand  $\Psi(x_1 - 1)$  using (9-4) with  $I = I(x_1 - 1)$ . We continue to expand each term of the form  $\Psi(z)$  in the same fashion until we arrive to  $z$  such that either  $z \leq b_n$  and  $z > k^2$  or the number of  $G_I$  terms in the product becomes  $\lceil 40k/q_n^{8/9} \rceil$ , whichever comes first. We then obtain an expression of the form

$$(9-14) \quad \Psi(k) = \sum_{s; z_{i+1} \in \partial I(z'_i)} G_{I(k)}(k, z_1) G_{I(z'_1)}(z'_1, z_2) \cdots G_{I(z'_s)}(z'_s, z_{s+1}) \Psi(z'_{s+1}).$$

where in each term of the summation we have  $z_i > b_n$  for  $i = 1, \dots, s$ , and either  $0 < z'_{s+1} \leq b_n$  and  $s \leq 40k/q_n^{8/9}$  or  $z'_{s+1} > k^2$  and  $s \leq 40k/q_n^{8/9}$  or  $s + 1 = \lceil 40k/q_n^{8/9} \rceil$ . By construction, for each  $z'_i$  with  $i \leq s$ , we have that  $I(z'_i)$  is well defined and satisfies (9-12) and (9-13). We now consider the three cases,  $0 < z'_{s+1} \leq b_n$ ,  $z'_{s+1} > k^2$ , and  $s + 1 = \lceil 40k/q_n^{8/9} \rceil$  separately. If  $0 < z'_{s+1} \leq b_n$  we have, by (9-12) and (9-2),

$$\begin{aligned} & |G_{I(k)}(k, z_1) G_{I(z'_1)}(z'_1, z_2) \cdots G_{I(z'_s)}(z'_s, z_{s+1}) \Psi(z'_{s+1})| \\ & \leq C \exp(-L_1(|k - z_1| + \sum_{i=1}^s |z'_i - z_{i+1}|))(1 + b_n) \\ & \leq C \exp(-L_1(|k - z_{s+1}| - (s + 1)))(1 + b_n) \\ & \leq C \exp(-L_1(k - b_n - 40k/q_n^{8/9}))(1 + b_n). \end{aligned}$$

Similarly, if  $z'_{s+1} > k^2$ , we use (9-12) and (9-13) to get

$$\begin{aligned} & |G_{I(k)}(k, z_1) G_{I(z'_1)}(z'_1, z_2) \cdots G_{I(z'_s)}(z'_s, z_{s+1}) \Psi(z'_{s+1})| \\ & \leq C \exp(-L_1(k^2 - k - 40k/q_n^{8/9}))(1 + 3k^{9/4}). \end{aligned}$$

Finally, if  $s + 1 = \lceil 40k/q_n^{8/9} \rceil$ , using again (9-2), (9-12), and also (9-10) we can estimate

$$\begin{aligned} & |G_{I(k)}(k, z_1) G_{I(z'_1)}(z'_1, z_2) \cdots G_{I(z'_s)}(z'_s, z_{s+1}) \Psi(z'_{s+1})| \\ & \leq C \exp(-L_1 \frac{1}{40} q_n^{8/9} \lceil 40k/q_n^{8/9} \rceil)(1 + k^2). \end{aligned}$$

In either case,

$$(9-15) \quad |G_{I(k)}(k, z_1) G_{I(z'_1)}(z'_1, z_2) \cdots G_{I(z'_s)}(z'_s, z_{s+1}) \Psi(z'_{s+1})| \leq e^{-9L_1 k/10}$$

for  $k$  sufficiently large. Finally, we observe that the total number of terms in (9-14) is bounded above by the  $\lceil 40k/q_n^{8/9} \rceil$ -th power of 2. Combining it with (9-14) and (9-15) we obtain

$$|\Psi(k)| \leq 2^{\lceil 40k/q_n^{8/9} \rceil} e^{-9L_1 k/10} < e^{-4L_1 k/5} \quad \text{for large } k. \quad \square$$



9.2. *Estimates on trigonometric products.* We will write  $\|z\|_{\mathbb{R}/\mathbb{Z}}$  for the distance to the nearest integer.

LEMMA 9.6. *Let  $p$  and  $q$  be relatively prime.*

(i) *Let  $1 \leq k_0 \leq q$  be such that*

$$|\sin 2\pi(x + k_0 p/(2q))| = \min_{1 \leq k \leq q} |\sin 2\pi(x + kp/(2q))|.$$

*Then*

$$(9-16) \quad \ln q + \ln 2/\pi < \sum_{\substack{k=1 \\ k \neq k_0}}^q \ln |\sin 2\pi(x + kp/(2q))| + (q - 1) \ln 2 \leq \ln q.$$

$$(ii) \quad \sum_{k=1}^{q-1} \ln |\sin \pi kp/q| = -(q - 1) \ln 2 + \ln q.$$

*Proof.* We use that

$$(9-17) \quad \ln |\sin x/2| = -\ln 2 - \frac{1}{2} \sum_{k \neq 0} \frac{e^{ikx}}{|k|} \quad \text{pointwise for } x \notin 2\pi\mathbb{Z}.$$

Thus, for  $x \neq k/(2q)$ ,

$$\begin{aligned} \sum_{j=1}^q \ln |\sin 2\pi(x + jp/(2q))| &= -q \ln 2 - \frac{1}{2} \sum_{k \neq 0} \frac{1}{|k|} \sum_{j=1}^q e^{2\pi ik(2x + jp/q)} \\ &= -q \ln 2 - \frac{1}{2} \sum_{k \neq 0} \frac{1}{|k|} e^{4\pi ikqx} = -q \ln 2 + \ln 2 + \ln |\sin 2\pi qx|. \end{aligned}$$

Thus

$$(9-18) \quad \sum_{\substack{k=1 \\ k \neq k_0}}^q \ln \left| \sin 2\pi \left( x + \frac{kp}{2q} \right) \right| + (q - 1) \ln 2 = \ln \frac{|\sin 2\pi q(x + k_0 p/(2q))|}{|\sin 2\pi(x + k_0 p/(2q))|}.$$

It is easily checked that if  $0 < qx \leq \pi/2$ , then  $2q/\pi < \sin qx/\sin x < q$ . Since  $\|2x + k_0 p/q\|_{\mathbb{R}/\mathbb{Z}} \leq 1/(2q)$ , (9-18) implies (9-16). Then (ii) follows by taking the limit in (9-18). □

For  $\alpha \notin \mathbb{Q}$ , let  $p_n/q_n$  be its continued fraction approximants. Setting  $\Delta_n = |q_n \alpha - p_n|$ , we recall the basic estimates

$$(9-19) \quad 1/q_n > \Delta_{n-1} > 1/(q_n + q_{n-1}),$$

$$(9-20) \quad \|k\alpha\|_{\mathbb{R}/\mathbb{Z}} > \Delta_{n-1} \quad \text{for } q_{n-1} + 1 \leq k \leq q_n - 1.$$

Notice that if  $z, w \in \mathbb{R}$  are such that  $\cos(z - w) \geq 0$ , then

$$(9-21) \quad \left| \frac{\sin z}{\sin w} - 1 \right| \leq \left| \cos(z - w) - 1 + \frac{\cos w}{\sin w} \sin(z - w) \right| \leq \left| 2 \frac{\sin(z - w)}{\sin w} \right|.$$

LEMMA 9.7. *Let  $1 \leq k_0 \leq q_n$  be such that*

$$|\sin 2\pi(x + k_0\alpha/2)| = \min_{1 \leq k \leq q_n} |\sin 2\pi(x + k\alpha/2)|.$$

Then

$$(9-22) \quad \left| \sum_{\substack{k=1 \\ k \neq k_0}}^{q_n} \ln |\sin 2\pi(x + k\alpha/2)| + (q_n - 1) \ln 2 \right| < C \ln q_n.$$

*Proof.* Let  $1 \leq k_1 \leq q_n$  be such that

$$|\sin 2\pi(x + k_1 p_n / (2q_n))| = \min_{1 \leq k \leq q_n} |\sin 2\pi(x + k p_n / (2q_n))|.$$

We first remark that, by (9-20)

$$(9-23) \quad \|(2x + k\alpha) - (2x + k'\alpha)\|_{\mathbb{R}/\mathbb{Z}} = \|(k - k')\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \Delta_{n-1} \\ \text{for } 1 \leq k, k' \leq q_n \text{ and } k \neq k'.$$

Applying this to the case  $k' = k_0$ , we get, by (9-19),

$$|\ln |\sin 2\pi(x + k\alpha/2)|| < C \ln q_n \quad \text{for } k \neq k_0.$$

An even simpler argument,

$$(9-24) \quad \|(2x + k p_n / q_n) - (2x + k' p_n / q_n)\|_{\mathbb{R}/\mathbb{Z}} = \|(k - k') p_n / q_n\|_{\mathbb{R}/\mathbb{Z}} \geq 1/q_n, \\ \text{for } 1 \leq k, k' \leq q_n \text{ and } k \neq k',$$

also gives that if  $k \neq k_1$  then  $|\ln |\sin 2\pi(x + k p_n / (2q_n))|| < C \ln q_n$ . This and (9-16) show that it is enough to get the estimate

$$(9-25) \quad \sum_{\substack{k=1 \\ k \neq k_0, k_1}}^{q_n} \ln \left| \frac{\sin 2\pi(x + k\alpha/2)}{\sin 2\pi(x + k p_n / (2q_n))} \right| < C \ln q_n.$$

By (9-21),

$$(9-26) \quad \left| \frac{\sin 2\pi(x + k\alpha/2)}{\sin 2\pi(x + k p_n / (2q_n))} - 1 \right| < \frac{C_0 \Delta_n}{|\sin 2\pi(x + k p_n / (2q_n))|},$$

so, if  $C_0 \Delta_n < (1/4) |\sin 2\pi(x + k p_n / (2q_n))|$ , we have

$$\ln \left| \frac{\sin 2\pi(x + k\alpha/2)}{\sin 2\pi(x + k p_n / (2q_n))} \right| < \frac{C \Delta_n}{|\sin 2\pi(x + k p_n / (2q_n))|},$$

Suppose  $s_1, \dots, s_r$  is an enumeration of  $\{1 \leq k \leq q_n, k \neq k_0, k_1\}$  in nondecreasing order of  $|\sin 2\pi(x + kp_n/(2q_n))|$  (so  $r = q_n - 1$  or  $r = q_n - 2$ ). By (9-24), we have  $|\sin 2\pi(x + s_j p_n/(2q_n))| > C_1 j/q_n$ . Then we get (9-25):

$$\begin{aligned} \sum_{\substack{k=1 \\ k \neq k_0, k_1}}^{q_n} \ln \left| \frac{\sin 2\pi(x + k\alpha/2)}{\sin 2\pi(x + kp_n/(2q_n))} \right| &= \sum_{1 \leq j \leq 4C_0/C_1} \ln \left| \frac{\sin 2\pi(x + s_j \alpha/2)}{\sin 2\pi(x + s_j p_n/(2q_n))} \right| \\ &\quad + \sum_{4C_0/C_1 < j \leq r} \ln \left| \frac{\sin 2\pi(x + s_j \alpha/2)}{\sin 2\pi(x + s_j p_n/(2q_n))} \right| \\ &\leq C \ln q_n + \sum_{4C_0/C_1 < j \leq r} C q_n \Delta_n / j \leq C \ln q_n. \quad \square \end{aligned}$$

LEMMA 9.8. *Let  $\ell \in \mathbb{N}$  be such that  $\ell < q_{r+1}/(10q_n)$ , where  $r \geq n$ . Given a sequence  $|\ell_k| \leq \ell - 1$  for  $k = 1, \dots, q_n$ , let  $1 \leq k_0 \leq q_n$  be such that*

$$|\sin 2\pi(x + (k_0 + \ell_{k_0} q_r)\alpha/2)| = \min_{1 \leq k \leq q_n} |\sin 2\pi(x + (k + \ell_k q_r)\alpha/2)|.$$

Then

$$(9-27) \quad \left| \sum_{\substack{k=1 \\ k \neq k_0}}^{q_n} \ln |\sin 2\pi(x + (k + \ell_k q_r)\alpha/2)| + (q_n - 1) \ln 2 \right| < \ln q_n + C(\Delta_n + (\ell - 1)\Delta_r)q_n \ln q_n.$$

*Proof.* Notice that  $\|(k - k')\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \Delta_{n-1} \geq 1/(2q_n)$ , while

$$\|(\ell_k - \ell_{k'})q_r \alpha\|_{\mathbb{R}/\mathbb{Z}} \leq \frac{q_{r+1}}{5q_n} \Delta_r < \frac{1}{5q_n}.$$

This implies

$$(9-28) \quad \|(2x + (k + \ell_k q_r)\alpha) - (2x + (k' + \ell_{k'} q_r)\alpha)\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{1}{5q_n}$$

for  $1 \leq k, k' \leq q_n$  with  $k \neq k'$ .

By (9-21), we have

$$(9-29) \quad \left| \frac{\sin 2\pi(x + (k + \ell_k q_r)\alpha/2)}{\sin 2\pi(x + kp_n/(2q_n))} - 1 \right| \leq \frac{C_0(\Delta_n + (\ell - 1)\Delta_r)}{|\sin 2\pi(x + kp_n/(2q_n))|}.$$

We now argue as in the previous lemma, using (9-28) and (9-29) instead of (9-23) and (9-26). □

9.3. *Nonresonant case. Proof of Lemma 9.4.* In the arguments that follow, we will actually consider a slightly larger range of  $k$ , by assuming a weaker upper bound  $k \leq \max\{\frac{1}{20}q_n, 50q_{n+1}^{8/9}\}$ . The fact that the estimates hold for this larger range will be useful later (when dealing with the resonant case).

We start with the proof of the first part. Let

$$k = mq_n \pm (sq_{n-1} + r) = mq_n \pm k_0, ,$$

with  $s \geq 1$ ,  $0 \leq r < q_{n-1}$  and  $k_0 \leq q_n/2$ , be nonresonant. Notice that  $2sq_{n-1} < q_n$ . Assume without loss of generality that  $k = mq_n + k_0$ , the other case being treated similarly.

Notice that if  $m \geq 1$  then  $k > \frac{1}{20}q_n$ , which implies that  $k \leq 50q_{n+1}^{8/9}$ , and we have

$$(9-30) \quad m \leq 50q_{n+1}^{8/9}/q_n$$

(which is also obviously satisfied if  $m = 0$ ). Set

$$I_1 = [-[sq_{n-1}/2], sq_{n-1} - [sq_{n-1}/2] - 1],$$

$$I_2 = [mq_n + k_0 - [sq_{n-1}/2], mq_n + k_0 + sq_{n-1} - [sq_{n-1}/2] - 1].$$

Set  $\theta_j = \theta + j\alpha$  for  $j \in I_1 \cup I_2$ . The set  $\{\theta_j\}_{j \in I_1 \cup I_2}$  consists of  $2sq_{n-1}$  elements.

LEMMA 9.9. *For any  $\epsilon > 0$  and sufficiently large  $n$ , the set  $\{\theta_j\}_{j \in I_1 \cup I_2}$  is  $(-2 \ln(s/q_n)/q_{n-1} + \epsilon)$ -uniform.*

*Proof.* We will first estimate the numerator in (9-8). We have

$$(9-31) \quad \sum_{\substack{j \in I_1 \cup I_2 \\ j \neq i}} \ln |\cos 2\pi a - \cos 2\pi \theta_j| \\ = \sum_{\substack{j \in I_1 \cup I_2 \\ j \neq i}} \ln \left| \sin 2\pi \frac{a + \theta_j}{2} \right| + \sum_{\substack{j \in I_1 \cup I_2 \\ j \neq i}} \ln \left| \sin 2\pi \frac{a - \theta_j}{2} \right| + (2sq_{n-1} - 1) \ln 2 \\ = \Sigma_+ + \Sigma_- + (2sq_{n-1} - 1) \ln 2.$$

Both  $\Sigma_+$  and  $\Sigma_-$  consist of  $2s$  terms of the form of (9-22) plus  $2s$  terms of the form

$$(9-32) \quad \ln \min_{j=1, \dots, q_{n-1}} |\sin(2\pi(x + j\alpha/2))|,$$

minus  $\ln |\sin(a \pm \theta_i)/2|$ . Therefore, by (9-22)

$$(9-33) \quad \sum_{\substack{j \in I_1 \cup I_2 \\ j \neq i}} \ln |\cos 2\pi a - \cos 2\pi \theta_j| \leq -2sq_{n-1} \ln 2 + Cs \ln q_{n-1}.$$

To estimate the denominator of (9-8), we represent it again in the form (9-31) with  $a = \theta_i$ . Assume that  $i = j_0q_{n-1} + i_0 \in I_1$  with  $0 \leq i_0 < q_{n-1}$ , the other case

being treated similarly. Then

$$(9-34) \quad \Sigma_- = \sum_{\substack{j \in I_1 \cup I_2 \\ j \neq i}} \ln |\sin \pi(i - j)\alpha|.$$

On each interval  $I \subset I_1$  of length  $q_{n-1}$ , the minimum over  $t \in I$  of  $|\sin \pi(t - i)\alpha|$  is achieved at  $t - i$  of the form  $jq_{n-1}$  for some  $j$ . This follows from the fact that if  $0 < |z| < q_{n-1}$  and  $2|j|q_{n-1} < q_n$ , then

$$\|(jq_{n-1} + z)\alpha\|_{\mathbb{R}/\mathbb{Z}} > \|jq_{n-1}\alpha\|_{\mathbb{R}/\mathbb{Z}},$$

since  $\|z\alpha\| \geq \Delta_{n-2}$  and  $\|jq_{n-1}\alpha\| < \Delta_{n-2}/2$ . The possible values of  $j$  form an interval  $[j_-^0, j_+^0]$  of size  $s$  containing  $j_0$ .

Let now  $T$  be an arbitrary interval of length  $q_{n-1}$  contained in  $I_2$ . Notice that  $T$  is contained in  $[i + mq_n + 1, i + (m + 1)q_n - 1]$ . The minimum over  $t \in T$  of  $|\sin \pi(t - i)\alpha|$  is achieved at  $t - i$  of either the form  $mq_n + jq_{n-1}$  or the form  $(m + 1)q_n - jq_{n-1}$  for some  $j \in \mathbb{N}$ .<sup>5</sup> For  $u \in \{0, 1\}$ , let  $t_u \in T$  be (the unique number) of the form  $t_u = i + (m + u)q_n + (-1)^u j_u q_{n-1}$  for some  $j_u \in \mathbb{N}$ . Since  $|t_u - t_{1-u}| < q_{n-1}$  it follows that

$$(9-35) \quad 0 \leq j_{1-u} + j_u - [q_n/q_{n-1}] \leq 1.$$

For all  $j \in [1, [q_n/q_{n-1}]]$ , we have the lower bound

$$\|((-1)^u j q_{n-1} + (m + u)q_n)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \Delta_{n-1}/2.$$

Indeed, by (9-30), if  $m \geq 1$  then  $(m + u)\Delta_n \leq 100(q_n^{8/9}/q_n)\Delta_n \leq \Delta_{n-1}/2$ , while  $\|jq_{n-1}\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \Delta_{n-1}$ . If  $m = 0$  then  $(m + u)q_n + (-1)^u j q_{n-1} \in [1, q_n - 1]$ , and we get the lower bound  $\Delta_{n-1}$ . Those considerations also give the upper bound  $(m + u)\Delta_n \leq \max\{\Delta_{n-1}/2, \Delta_n\}$ . This gives the estimate

$$(9-36) \quad \|((-1)^u j q_{n-1} + (m + u)q_n)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq j\Delta_{n-1}/C$$

for all  $j \in [1, [q_n/q_{n-1}]]$ .

Let  $T$  now run through the set of disjoint segments  $T^p$ , each of length  $q_{n-1}$ , such that  $I_2 = \bigcup_{p=1}^s T^p$ . It is not difficult to see that there exists a  $u$  (possibly both  $u = 0$  and  $u = 1$ ) such that for all  $p$  the corresponding  $j_u$  satisfy  $j_u \leq \frac{3}{4}[q_n/q_{n-1}]$ .<sup>6</sup>

<sup>5</sup>Suppose that  $t \in T$  minimizes  $\|(t - i)\alpha\|_{\mathbb{R}/\mathbb{Z}}$ , and suppose that  $j_u, u \in \{0, 1\}$ , is such that  $t_u = (m + u)q_n + (-1)^u j_u q_{n-1} + i \in T$ . If  $t \neq t_0$  and  $t \neq t_1$ , then  $\|(t - t_u)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \Delta_{n-2}$  as above. Since the  $(t_u - i)\alpha$  minus nearest integer are on opposite sides of 0, this implies that  $\|(t_0 - t_1)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq 2\Delta_{n-2}$ . But one easily checks that  $\|(t_0 - t_1)\alpha\|_{\mathbb{R}/\mathbb{Z}}$  is either equal to  $\Delta_{n-2}$  (if  $t_1 > t_0$ ) or to  $\Delta_{n-1} + \Delta_{n-2}$  (if  $t_1 \leq t_0$ ).

<sup>6</sup>For  $u = 0, 1$  the  $j_u$  form an interval  $[j_-^u, j_+^u]$  of length  $s$  contained in  $[1, [q_n/q_{n-1}]]$ . If  $j_+^u > \frac{3}{4}[q_n/q_{n-1}]$ , then, since  $s \leq [q_n/2q_{n-1}]$ , we have that  $j_-^u > \frac{1}{4}[q_n/q_{n-1}] + 1$ . Then, by (9-35),  $j_+^{1-u} < \frac{3}{4}[q_n/q_{n-1}]$ .

We now fix  $u \in \{0, 1\}$  with this property. Then

$$\begin{aligned} \|((m+u)q_n + (-1)^u j_u q_{n-1})\alpha\|_{\mathbb{R}/\mathbb{Z}} &\leq \frac{3}{4}[q_n/q_{n-1}]\Delta_{n-1} + |m+u|\Delta_n \\ &\leq (\frac{3}{4}[q_n/q_{n-1}] + 1)\Delta_{n-1} \end{aligned}$$

Then, by (9-35),  $j_{1-u} \geq \frac{1}{4}[q_n/q_{n-1}]$  and by (9-36),

$$\|((-1)^{1-u} j_{1-u} q_{n-1} + (m+1-u)q_n)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{1}{4C}[q_n/q_{n-1}]\Delta_{n-1}.$$

Thus  $\|(t_{1-u} - i)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \|(t_u - i)\alpha\|_{\mathbb{R}/\mathbb{Z}}/(7C)$ .

Notice that the  $j_u$  form an interval  $[j_-, j_+]$  of length  $s$ , which is contained in  $[1, [3q_n/(4q_{n-1})]]$ .

Splitting again  $\Sigma_-$  into  $2s$  sums of length  $q_{n-1}$  and applying (9-22) on each, we obtain

$$\begin{aligned} (9-37) \quad \Sigma_- &> -2sq_{n-1} \ln 2 + \sum_{\substack{j_-^0 \leq j \leq j_+^0 \\ j \neq j_0}} \ln |\sin \pi(j - j_0)q_{n-1}\alpha| \\ &+ \sum_{j=j_-}^{j_+} \ln |\sin \pi((-1)^u j q_{n-1} + (m+u)q_n)\alpha| - Cs - Cs \ln q_{n-1}. \end{aligned}$$

Denote the sums in (9-37) by  $\Sigma_1$  and  $\Sigma_2$ . Since  $j_-^0 \leq j_0 \leq j_+^0$  and  $|[j_-^0, j_+^0]| = s$ , we have that

$$(9-38) \quad \Sigma_1 > 2 \sum_{j=1}^{[s/2]} \ln \sin |\pi j q_{n-1}\alpha| > 2 \sum_{j=1}^{[s/2]} \ln 2j\Delta_{n-1} > s(\ln s/q_n - C),$$

where the second inequality follows from the Stirling formula. For  $j \in [j_-, j_+]$ , we use (9-36) to obtain

$$(9-39) \quad \Sigma_2 > \sum_{j=1}^s \ln j\Delta_{n-1} - Cs > s \ln s/q_n - Cs.$$

Therefore,

$$(9-40) \quad \Sigma_- > -2sq_{n-1} \ln 2 + 2s(\ln s/q_n - C \ln q_{n-1}).$$

$\Sigma_+$  is estimated in a similar way. Set  $J_1 = [-(s+1)/2, s - [(s+1)/2] - 1]$  and  $J_2 = [[s/2], s + [s/2] - 1]$ , which are two adjacent disjoint intervals of length  $s$ . Then  $I_1 \cup I_2$  can be represented as a disjoint union of segments  $B_j$  for  $j \in J_1 \cup J_2$ , each of length  $q_{n-1}$ . Applying (9-22) on each  $B_j$  we obtain

$$(9-41) \quad \Sigma_+ > -2sq_{n-1} \ln 2 + \sum_{j \in J_1 \cup J_2} \ln |\sin 2\pi \hat{\theta}_j| - Cs \ln q_{n-1} - \ln |\sin 2\pi(\theta + i\alpha)|,$$

where  $|\sin 2\pi \hat{\theta}_j| = \min_{\ell \in B_j} |\sin 2\pi(\theta + (i + \ell)\alpha/2)|$ .

Let  $\tilde{\theta}_j = \hat{\theta}_j$  for  $j \in J_1$  and  $\tilde{\theta}_j = \hat{\theta}_j - m q_n \alpha/2$  for  $j \in J_2$ . Since  $\theta \notin \mathcal{R}$ , for sufficiently large  $n$ , we have that

$$\min_{j \in J_1 \cup J_2} |\sin 2\pi \tilde{\theta}_j| > 1/(9s^2 q_{n-1}^2).$$

To estimate  $|\sin 2\pi \hat{\theta}_j|$ ,  $j \in J_2$ , we distinguish two cases:

- If  $q_{n+1} > (20s^2 q_{n-1}^2)^9$ , we write

$$(9-42) \quad \begin{aligned} |\sin 2\pi \hat{\theta}_j| &\geq |\sin 2\pi \tilde{\theta}_j \cos \pi m \Delta_n| - |\cos 2\pi \tilde{\theta}_j \sin \pi m \Delta_n| \\ &> (10s^2 q_{n-1}^2)^{-1} - (q_{n+1}^{1/9})^{-1} > (20s^2 q_{n-1}^2)^{-1}. \end{aligned}$$

- If  $q_{n+1} \leq (20s^2 q_{n-1}^2)^9$  we use that since  $\theta \notin \mathcal{R}$ , for large  $n$ ,

$$(9-43) \quad \min_{j \in J_2} |\sin 2\pi \hat{\theta}_j| > ((2m + 2)q_n)^{-2} > (4q_{n+1})^{-2} > (20sq_{n-1})^{-36}$$

In either case,  $\ln \min_{j \in J_2} |\sin 2\pi \hat{\theta}_j| > -C \ln sq_{n-1}$ .

Let  $J = J_1$  or  $J = J_2$  and assume that  $\hat{\theta}_{j+1} = \hat{\theta}_j + (q_{n-1}/2)\alpha$  for every  $j, j + 1 \in J$ . We obtain therefore

$$(9-44) \quad \begin{aligned} \sum_{j \in J} \ln |\sin 2\pi \hat{\theta}_j| &> -C \ln sq_{n-1} + \sum_{j=1}^s \ln j \Delta_{n-1} / C \\ &> s \ln s / q_n - C(\ln sq_{n-1} + s), \end{aligned}$$

where again the last inequality follows from the Stirling formula. In the other case, decompose  $J$  in maximal intervals  $T_\kappa$  such that for  $j, j + 1 \in T_\kappa$  we have  $\hat{\theta}_{j+1} = \hat{\theta}_j + (q_{n-1}/2)\alpha$ . Notice that the boundary points of an interval  $T_\kappa$  are either boundary points of  $J$  or satisfy  $\|2\hat{\theta}_j\|_{\mathbb{R}/\mathbb{Z}} + \Delta_{n-1} \geq \Delta_{n-2}/2$ . Assuming  $T_\kappa \neq J$ , there exists a  $j \in T_\kappa$  such that  $\|2\hat{\theta}_j\|_{\mathbb{R}/\mathbb{Z}} + \Delta_{n-1} \geq \Delta_{n-2}/2$ . An estimate similar to (9-44) gives

$$(9-45) \quad \sum_{j \in T_\kappa} \ln |\sin 2\pi \hat{\theta}_j| > -|T_\kappa| \ln q_{n-1} - C(\ln sq_{n-1} + |T_\kappa|).$$

If  $T_\kappa$  does not contain a boundary point of  $J$  (in particular  $|T_\kappa| \leq |J| - 2 = s - 2$  and  $[q_n/q_{n-1}] \geq 2s \geq 6$ ), then  $T_\kappa$  does not contain any  $j$  with

$$\|2\hat{\theta}_j\|_{\mathbb{R}/\mathbb{Z}} < \Delta_{n-2}/10 < \Delta_{n-2}/2 - \Delta_{n-1}$$

(otherwise  $|T_\kappa| - 1 \geq \Delta_{n-2}/\Delta_{n-1} - 2 \geq q_n/(2q_{n-1}) - 2 \geq s - 2$ , which is impossible) and hence

$$(9-46) \quad \sum_{j \in T_\kappa} \ln |\sin 2\pi \hat{\theta}_j| > -|T_\kappa|(\ln q_{n-1} + C).$$

Putting together all  $T_\kappa$ , using (9-45) for the ones that intersect the boundary of  $J$  and (9-46) for the others, we get in all cases that

$$\sum_{j \in J} \ln |\sin 2\pi \hat{\theta}_j| > s \ln s/q_n - C(\ln sq_{n-1} + s).$$

Putting together  $J = J_1$  and  $J = J_2$  we have

$$\sum_{j \in J_1 \cup J_2} \ln |\sin 2\pi \hat{\theta}_j| > 2s \ln s/q_n - C(\ln sq_{n-1} + s).$$

Combining it with (9-41) we obtain

$$(9-47) \quad \Sigma_+ > -2sq_{n-1} \ln 2 + 2s(\ln s/q_n - C \ln q_{n-1}).$$

Putting together (9-47),(9-40), and (9-31) gives

$$\sum_{\substack{j \in I_1 \cup I_2 \\ j \neq i}} \ln |\cos 2\pi \theta_i - \cos 2\pi \theta_j| > 4s(\ln(sq_{n-1}/q_n) - C \ln q_{n-1}) - 2sq_{n-1} \ln 2.$$

This together with (9-33) yields

$$\max_{j \in I_1 \cup I_2} \prod_{\substack{\ell \in I_1 \cup I_2 \\ \ell \neq j}} \frac{|z - \cos 2\pi \theta_\ell|}{|\cos 2\pi \theta_j - \cos 2\pi \theta_\ell|} < e^{-4s \ln(sq_{n-1}/q_n) + Cs \ln q_{n-1}}. \quad \square$$

By Lemmas 9.3 and 9.9 at least one of  $\theta_j$  for  $j \in I_1 \cup I_2$  is not in

$$(9-48) \quad A(2sq_{n-1} - 1, L + 2 \ln(sq_{n-1}/q_n)/q_{n-1} - \epsilon),$$

where  $\epsilon$  can be made arbitrarily small for large  $n$ . By Lemma 9.2 and singularity of  $0$ ,<sup>7</sup> we have that for all  $j \in I_1$ ,  $\theta_j$  belongs to the set (9-48) (using that  $(s+1)q_{n-1} > q_n^{8/9}$  and the bound  $\ln q_n/q_{n-1} < L$ ). Let  $j_0 \in I_2$  be such that  $\theta_{j_0}$  does not belong to the set (9-48). Set

$$I = [j_0 - sq_{n-1} + 1, j_0 + sq_{n-1} - 1] = [x_1, x_2].$$

Then by (9-6) and (9-7),

$$\begin{aligned} |G_I(k, x_i)| &< e^{(L+\epsilon_1)(2sq_{n-1}-2-|k-x_i|)-2sq_{n-1}(L+2 \ln(sq_{n-1}/q_n)/q_{n-1}-\epsilon)} \\ &< e^{-(L+\epsilon_1)|k-x_i|-4sq_{n-1} \ln(sq_{n-1}/q_n)/q_{n-1}+(\epsilon_1+\epsilon)sq_{n-1}}. \end{aligned}$$

Since

$$(9-49) \quad |k - x_i| \geq [sq_{n-1}/2] - 1,$$

<sup>7</sup>To get what we need here one can take in Lemma 9.2, besides  $y = 0$ , also  $\epsilon = 99/100L$  and  $\delta = 99/400$ .



we obtain that

$$(9-50) \quad |G_I(k, x_i)| < \exp(-(L + 9 \ln(sq_{n-1}/q_n)/q_{n-1} - \epsilon)|k - x_i|)$$

which in view of  $(s + 1)q_{n-1} > q_n^{8/9}$  gives the statement of the first part of Lemma 9.4.

We now assume  $s = 0$ . In this case  $\alpha$  is “Diophantine” on the scale  $q_{n-1}$ ; however some caution is needed as it may not be so on the scale  $q_n$ . Let  $k = mq_n \pm k_0$  with  $\max\{q_{n-1}/20, q_n^{8/9}\} < k_0 < q_{n-1}$ . We will assume that  $m = q_n + k_0$ , the other case being analogous.

We distinguish three cases.

- If  $\frac{1}{20}q_{n-1} < k_0 \leq \frac{4}{5}q_{n-1}$ , set

$$I_1 = [-[\frac{19}{40}q_{n-1}] + 1, [\frac{19}{40}q_{n-1}]],$$

$$I_2 = [mq_n + [\frac{19}{40}q_{n-1}] + 1, mq_n + 2[\frac{1}{2}q_{n-1}] - [\frac{19}{40}q_{n-1}]].$$

- If  $\frac{4}{5}q_{n-1} < k_0 < q_{n-1}$  and  $q_n \leq 2q_{n-1}$ , set

$$I_1 = [-[\frac{1}{4}q_n] + 1, [\frac{1}{4}q_n]],$$

$$I_2 = [mq_n + [\frac{1}{4}q_n] + 1, mq_n + 2[\frac{1}{2}q_n] - [\frac{1}{4}q_n]].$$

- If  $\frac{4}{5}q_{n-1} < k_0 < q_{n-1}$  and  $q_n > 2q_{n-1}$ , set

$$I_1 = [-[\frac{1}{2}q_{n-1}] + 1, q_{n-1} - [\frac{1}{2}q_{n-1}]],$$

$$I_2 = [mq_n + q_{n-1} - [\frac{1}{2}q_{n-1}] + 1, mq_n + 2q_{n-1} - [\frac{1}{2}q_{n-1}]].$$

Let  $\theta_j = \theta + j\alpha$  for  $j \in I_1 \cup I_2$ . The set  $\{\theta_j\}_{j \in I_1 \cup I_2}$  consists of  $2[q_{n-1}/2]$  elements in the first case, of  $2[q_n/2]$  elements in the second case, and of  $2q_{n-1}$  elements in the third case.

LEMMA 9.10. *For any  $\epsilon > 0$  and sufficiently large  $n$ , the set  $\{\theta_j\}_{j \in I_1 \cup I_2}$  is  $\epsilon$ -uniform.*

*Proof.* Consider first the case  $k_0 \leq \frac{4}{5}q_{n-1}$ . We will assume  $q_{n-1}$  is even, the other case needing obvious adjustments. As in the proof of Lemma 9.9 we will first estimate the numerator in (9-8). We have

$$(9-51) \quad \sum_{\substack{j \in I_1 \cup I_2 \\ j \neq i}} \ln |\cos 2\pi a - \cos 2\pi \theta_j|$$

$$= \sum_{\substack{j \in I_1 \cup I_2 \\ j \neq i}} \ln \left| \sin 2\pi \frac{a + \theta_j}{2} \right| + \sum_{\substack{j \in I_1 \cup I_2 \\ j \neq i}} \ln \left| \sin 2\pi \frac{a - \theta_j}{2} \right| + (q_{n-1} - 1) \ln 2$$

$$= \Sigma_+ + \Sigma_- + (q_{n-1} - 1) \ln 2.$$

Both  $\Sigma_+$  and  $\Sigma_-$  are of the form (9-27) with  $\ell_k \in \{0, m\}$ <sup>8</sup> and  $r = n$  plus a minimum term minus  $\ln|\sin 2\pi(a \pm \theta_i)/2|$ , so that the last two cancel each other for the purpose of the upper bound. Therefore, by (9-27)

$$(9-52) \quad \sum_{\substack{j \in I_1 \cup I_2 \\ j \neq i}} \ln|\cos 2\pi a - \cos 2\pi \theta_j| \leq (1 - q_{n-1}) \ln 2 + 2 \ln q_{n-1} + C(\Delta_{n-1} + m \Delta_n) q_{n-1} \ln q_{n-1} \leq -q_{n-1} \ln 2 + C q_{n-1}^{8/9} \ln q_{n-1}.$$

To estimate the denominator of (9-8), we write it in the form (9-51) with  $a = \theta_i$ . Then

$$\Sigma_- = \sum_{\substack{j \in I_1 \cup I_2 \\ j \neq i}} \ln|\sin \pi(i - j)\alpha|$$

is exactly of the form (9-27). Therefore, by (9-27),

$$(9-53) \quad \Sigma_- > (1 - q_{n-1}) \ln 2 - \ln q_{n-1} - C((q_n)^{-1} + (q_{n+1}^{1/9})^{-1}) q_{n-1} \ln q_{n-1} > -q_{n-1} \ln 2 - C q_{n-1}^{8/9} \ln q_{n-1}.$$

Similarly, for  $\Sigma_+$  we have

$$(9-54) \quad \Sigma_+ > (1 - q_{n-1}) \ln 2 + \ln \min_{\ell \in I_1 \cup I_2} |\sin 2\pi(\theta + (i + \ell)\alpha/2)| - C q_{n-1}^{8/9} \ln q_{n-1} > -q_{n-1} \ln 2 - C q_{n-1}^{8/9} \ln q_{n-1}.$$

Here, we use the estimate

$$(9-55) \quad \ln \min_{\ell \in I_1 \cup I_2} |\sin 2\pi(\theta + (i + \ell)\alpha/2)| > -C \ln q_{n-1},$$

which is obtained by considering separately the two cases  $q_{n+1} > q_{n-1}^C$  and  $q_{n+1} < q_{n-1}^C$ , and arguing in the same way as in (9-42) and (9-43). Combining (9-52), (9-51), (9-53) and (9-54), we arrive at

$$(9-56) \quad \max_{j \in I_1 \cup I_2} \prod_{\substack{\ell \in I_1 \cup I_2 \\ \ell \neq j}} \frac{|z - \cos 2\pi \theta_\ell|}{|\cos 2\pi \theta_j - \cos 2\pi \theta_\ell|} < \exp(C q_{n-1}^{8/9} \ln q_{n-1}) < e^{\epsilon q_{n-1}}$$

for any  $\epsilon > 0$  and sufficiently large  $n$ , as stated.

For the other cases,  $k > \frac{4}{5}q_{n-1}$ , the proof is very similar. If  $q_n \leq 2q_{n-1}$ , the argument is the same (replacing  $q_{n-1}$  by  $q_n$ ). We will concentrate on the case

---

<sup>8</sup>Recall that  $m$  is chosen so that  $k = mq_n + k_0$ , where  $k \leq \max\{50q_{n+1}^{8/9}, q_n/20\}$ . We have the bound  $m \leq 50q_{n+1}^{8/9}/q_n$ , so (9-27) really applies.

$q_n > 2q_{n-1}$  where the changes are slightly more substantial. Arguing as above we obtain by (9-27)

$$(9-57) \quad \sum_{\substack{j \in I_1 \cup I_2 \\ j \neq i}} \ln |\cos 2\pi a - \cos 2\pi \theta_j| \leq -2q_{n-1} \ln 2 + 4 \ln q_{n-1} + C(\Delta_{n-1} + m\Delta_n)q_{n-1} \ln q_{n-1} < -2q_{n-1} \ln 2 + Cq_{n-1}^{8/9} \ln q_{n-1}.$$

The denominator in (9-8) can be again split as  $\Sigma_+ + \Sigma_- + (2q_{n-1} - 1) \ln 2$ . Both  $\Sigma_+$  and  $\Sigma_-$  are, up to a constant, the sums of two terms of the form (9-27) plus minimum terms (two for  $\Sigma_+$  and one for  $\Sigma_-$ ). For the minimum terms of  $\Sigma_+$  the estimate (9-55) holds, so we obtain

$$(9-58) \quad \Sigma_+ > -2q_{n-1} \ln 2 - Cq_{n-1}^{8/9} \ln q_{n-1}.$$

For the minimum term of  $\Sigma_-$ , that is,  $\ln \min |\sin \pi(i - j)\alpha|$  (where the minimum is taken over all  $j$  that belong to the interval  $I_1$  or  $I_2$  that does not contain  $i$ ) we observe that it is achieved at  $j_0$  such that  $\|(i - j_0)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \Delta_{n-1} - m\Delta_n$  (since the possible values of  $|i - j|$  are contained in  $[mq_n + 1, mq_n + 2q_{n-1} - 1]$  and  $q_n > 2q_{n-1}$  by hypothesis). Thus, recalling that in the present situation we have  $q_n^{8/9} < q_{n-1}$ ,

$$\ln \min |\sin \pi(i - j)\alpha| > \ln(\Delta_{n-1} - m\Delta_n) > \ln(1/(2q_n) - 50/q_{n+1}^{1/9}q_n) > -C \ln q_n > -C \ln q_{n-1}.$$

Therefore, by (9-27),

$$(9-59) \quad \Sigma_- > -2q_{n-1} \ln 2 - Cq_{n-1}^{8/9} \ln q_{n-1}.$$

Combining (9-57),(9-51),(9-59) and (9-58) gives (9-56), as desired. □

By Lemmas 9.3 and 9.10 at least one of the  $\theta_j$ , for  $j \in I_1 \cup I_2$ , is not in  $A_{2[q_{n-1}/2]-1, L-\epsilon}$  if  $k_0 \leq \frac{4}{5}q_{n-1}$ , not in  $A_{2[q_n/2]-1, L-\epsilon}$  if  $\frac{4}{5}q_{n-1} < k_0 < q_{n-1}$  and  $q_n \leq 2q_{n-1}$ , and not in  $A_{2q_{n-1}-1, L-\epsilon}$  if  $k_0 > \frac{4}{5}q_{n-1}$  and  $q_n > 2q_{n-1}$ , where  $\epsilon$  can be made arbitrarily small for large  $n$ . By Lemma 9.2 and singularity of 0, we have that, in all three cases,  $\theta_j$  belongs to the corresponding  $A_{\cdot, L-\epsilon}$  for all  $j \in I_1$ . Let  $j_0 \in I_2$  be such that  $\theta_{j_0} \notin A_{L-\epsilon}$ .

For  $k_0 \leq \frac{4}{5}q_{n-1}$ , set  $I = [j_0 - [q_{n-1}/2] + 1, j_0 + [q_{n-1}/2]] = [x_1, x_2]$ . We then have

$$(9-60) \quad |k - x_i| > \frac{1}{40}q_{n-1}.$$

Then by (9-6) and (9-7),

$$(9-61) \quad |G_I(k, x_i)| < \exp((L + \epsilon_1)(q_{n-1} - 2 - |k - x_i|) - q_{n-1}(L - \epsilon)) < \exp(-(L + \epsilon_1 - 40(\epsilon_1 + \epsilon))|k - x_i|),$$

as desired.

For  $k_0 > \frac{4}{5}q_{n-1}$  and  $q_n \leq 2q_{n-1}$ , set

$$I = [j_0 - [q_n/2] + 1, j_0 + [q_n/2]] = [x_1, x_2].$$

Then

$$(9-62) \quad |k - x_i| > \frac{1}{10}q_n,$$

since

$$\begin{aligned} k - x_1 &> \frac{4}{5}q_{n-1} - \frac{1}{4}q_n \geq \frac{3}{10}q_{n-1}, \\ x_2 - k &> \frac{3}{4}q_n - q_{n-1} = (3q_{n-2} - q_{n-1})/4 > \frac{1}{5}q_{n-1} \end{aligned}$$

(using that  $\frac{4}{5}q_{n-1} < k_0 \leq \frac{1}{2}q_n = (q_{n-1} + q_{n-2})/2$ ). Thus for any  $\epsilon > 0$  and sufficiently large  $n$ , by (9-6) and (9-7) and by estimating as in (9-61)

$$(9-63) \quad |G_I(k, x_i)| < \exp(-(L - \epsilon)|k - x_i|).$$

For  $k_0 > \frac{4}{5}q_{n-1}$  and  $q_n > 2q_{n-1}$  set

$$I = [j_0 - q_{n-1} + 1, j_0 + q_{n-1} - 1] = [x_1, x_2].$$

Then

$$(9-64) \quad |k - x_i| > \frac{3}{10}q_{n-1}.$$

This implies as before that (9-63) holds for any  $\epsilon > 0$  and sufficiently large  $n$ . This concludes the proof of Lemma 9.4 in all cases.  $\square$

The estimates in the proof of Lemma 9.4 have the following corollary which will be necessary later (when dealing with the resonant case).

LEMMA 9.11. Fix  $\epsilon > 0$ . Assume  $b_n < k \leq \max\{\frac{1}{20}q_n, 50q_{n+1}^{8/9}\}$  and  $k < q_n^C$  for some  $C < \infty$ . Let  $d = \text{dist}(k, \{\ell q_n\}_{\ell \geq 0}) > \frac{1}{10}q_n$ . Let  $\phi = \phi_E$  be a generalized eigenfunction. Then, for sufficiently large  $n$  ( $n > n_0(\epsilon, c, E, C)$ ),

$$|\phi(k)| < \exp(-(L - \epsilon)d/2).$$

*Proof.* Recall that the previous estimates in this section were obtained, under the nonresonance hypothesis  $\text{dist}(k, \{\ell q_n\}_{\ell \geq 0}) > b_n$ , for the range  $b_n < k \leq \max\{\frac{1}{20}q_n, 50q_{n+1}^{8/9}\}$ .

If  $q_n \geq q_{n-1}^{10/9}$ , we have  $s \geq [q_n^{1/10}/10]$ , and the statement follows immediately from (9-4), (9-2) and (9-50), (9-49).

In case  $q_n < q_{n-1}^{10/9}$ , (9-50), (9-49), (9-61), (9-60), (9-63), (9-62) and (9-64) only lead to  $|\phi(k)| < \exp(-(L - \epsilon)cd)$  with certain  $c < 1/2$ . In order to prove the lemma as stated we will need an additional ‘‘patching’’ argument, which is very similar to the one used in Section 9.1.

We will show that in this case

$$(9-65) \quad |\phi(k)| < \exp(-(L - \epsilon)(d - \frac{1}{20}q_{n-1})),$$

from which the statement of the lemma follows. Assume  $\ell q_n < k < (\ell + 1)q_n$ . Using (9-50), (9-49), (9-61),(9-60) and (9-63), (9-62), (9-64) we obtain that for every  $y \in [\ell q_n, (\ell + 1)q_n]$  with  $\text{dist}(y, \{\ell q_n, (\ell + 1)q_n\}) > \frac{1}{20}q_{n-1}$ , there exists an interval  $y \in I(y) = [x_1, x_2] \subset [(\ell - 1)q_n, (\ell + 2)q_n]$  such that

$$(9-66) \quad \text{dist}(y, \partial I(y)) > \frac{1}{40}q_{n-1},$$

$$(9-67) \quad G_{I(y)}(y, x_i) < \exp(-(L - \epsilon)|y - x_i|) \quad \text{for } i = 1, 2$$

(notice that under the condition  $q_n < q_{n-1}^{10/9}$  we have  $b_n = \frac{1}{20}q_{n-1}$ ).

We here denote the boundary of the interval  $I(y)$ , the set  $\{x_1, x_2\}$ , by  $\partial I(y)$ . For  $z \in \partial I(y)$ , we let  $z'$  be the neighbor of  $z$ , (i.e.,  $|z - z'| = 1$ ) not belonging to  $I(y)$ .

If  $x_2 + 1 < (\ell + 1)q_n - \frac{1}{20}q_{n-1}$ , we expand  $\phi(x_2 + 1)$  in (9-4), iterating (9-4) with  $I = I(x_2 + 1)$ , and if  $x_1 - 1 > \ell q_n + \frac{1}{20}q_{n-1}$ , we expand  $\phi(x_1 - 1)$  in (9-4), iterating (9-4) with  $I = I(x_1 - 1)$ . We continue to expand each term of the form  $\phi(z)$  in the same fashion until we arrive to  $z$  such that either  $z + 1 \geq (\ell + 1)q_n - \frac{1}{20}q_{n-1}$  and  $z - 1 \leq \ell q_n + \frac{1}{20}q_{n-1}$ , or the number of  $G_I$  terms in the product becomes  $[40d/q_{n-1}]$ , whichever comes first. We then obtain an expression of the form

$$(9-68) \quad \phi(k) = \sum_{s; z_{i+1} \in \partial I(z'_i)} G_{I(k)}(k, z_1)G_{I(z'_1)}(z'_1, z_2) \cdots G_{I(z'_s)}(z'_s, z_{s+1})\phi(z'_{s+1}).$$

where in each term of the summation we have

$$\ell q_n + \frac{1}{20}q_{n-1} + 1 < z_i < (\ell + 1)q_n - \frac{1}{20}q_{n-1} - 1 \quad \text{for } i = 1, \dots, s,$$

and either  $z_{s+1} \notin [\ell q_n + \frac{1}{20}q_{n-1} + 1, (\ell + 1)q_n - \frac{1}{20}q_{n-1} - 1]$ ,  $s + 1 < [40d/q_{n-1}]$ , or  $s + 1 = [40d/q_{n-1}]$ .

By construction, for each  $z'_i$  with  $i \leq s$ , we have that  $I(z'_i)$  is well defined and satisfies (9-66) and (9-67). We now consider separately the two cases

$$z_{s+1} \notin [\ell q_n + \frac{1}{20}q_{n-1} + 1, (\ell + 1)q_n - \frac{1}{20}q_{n-1} - 1], \quad s + 1 < [40d/q_{n-1}],$$

$$\text{and } s + 1 = [40d/q_{n-1}].$$

Let  $\Gamma$  be the summand in (9-68). In the first case, we have, by (9-67) and (9-2),

$$(9-69) \quad |\Gamma| \leq \exp(-(L - \epsilon)(|k - z_1| + \sum_{i=1}^s |z'_i - z_{i+1}|))(1 + (\ell + 2)q_n)$$

$$\leq \exp(-(L - \epsilon)(|k - z_{s+1}| - (s + 1)))(1 + (\ell + 2)q_n)$$

$$\leq \exp(-(L - \epsilon)(d - \frac{1}{20}q_{n-1} - 40d/q_{n-1}))(1 + q_{n-1}^C).$$

If  $s + 1 = [40d/q_{n-1}]$ , using again (9-2), (9-67), and also (9-66), we obtain

$$|\Gamma| \leq \exp(- (L - \epsilon) \frac{1}{40} q_{n-1} 40d/q_{n-1}) (1 + q_{n-1}^C).$$

In either case,

$$(9-70) \quad |\Gamma| \leq \exp(- (L - 2\epsilon) (d - \frac{1}{20} q_{n-1}))$$

for  $n$  sufficiently large. Finally, we observe that the total number of terms in (9-68) is bounded above by the  $[40d/q_{n-1}]$ -th power of 2. Combining it with (9-68) and (9-70) we obtain that

$$\begin{aligned} |\phi(k)| &\leq 2^{[40d/q_{n-1}]} \exp(- (L - 2\epsilon) (d - \frac{1}{20} q_{n-1})) \\ &< \exp(- (L - 3\epsilon) (d - \frac{1}{20} q_{n-1})) \end{aligned}$$

for large  $n$ . □

9.4. *Resonant case. Proof of Lemma 9.5.* Notice that, under the condition that  $k$  is resonant,  $k \geq \frac{1}{2}q_n$ , which implies  $q_{n+1}^{8/9} \geq \frac{1}{2}q_n$ . This is an implicit hypothesis in the next lemma.

LEMMA 9.12. *For any  $\epsilon > 0$ , for sufficiently large  $n$ , and for any  $b \in [-\frac{13}{8}q_n, -\frac{3}{8}q_n] \cap \mathbb{Z}$ , we have  $\theta + (b + q_n - 1)\alpha \in A_{2q_n-1, 23L/32+\epsilon}$ .*

*Proof.* Let  $b_1 = b - 1$  and  $b_2 = b + 2q_n - 1$ .

Applying Lemma 9.11 we obtain that for  $i = 1, 2$

$$(9-71) \quad |\phi_E(b_i)| < \begin{cases} e^{-(L-\epsilon)(b/2+q_n)} & \text{if } -\frac{13}{8}q_n \leq b \leq -\frac{3}{2}q_n, \\ e^{-(L-\epsilon)|(b+q_n)/2|} & \text{if } -\frac{3}{2}q_n \leq b \leq -\frac{1}{2}q_n, \quad |b + q_n| > \frac{1}{4}q_n, \\ e^{(L-\epsilon)b/2} & \text{if } -\frac{1}{2}q_n \leq b \leq -\frac{3}{8}q_n. \end{cases}$$

Using (9-4) with  $I = [b, b + 2q_n - 2]$ , we get

$$\max(|G_I(0, b)|, |G_I(0, b + 2q_n - 2)|) > \begin{cases} e^{(L-\epsilon)(b/2+q_n)} & \text{if } -\frac{13}{8}q_n \leq b \leq -\frac{3}{2}q_n, \\ e^{(L-\epsilon)|(b+q_n)/2|} & \text{if } -\frac{3}{2}q_n \leq b \leq -\frac{1}{2}q_n, \\ & \text{and } |b + q_n| > \frac{1}{4}q_n, \\ e^{-(L-\epsilon)b/2} & \text{if } -\frac{1}{2}q_n \leq b \leq -\frac{3}{8}q_n, \\ e^{-\epsilon q_n} & \text{if } |b + q_n| < \frac{1}{4}q_n. \end{cases}$$

By (9-6) and (9-7),

$$\begin{aligned} &|Q_{2q_n-1}(\cos 2\pi(\theta + (b + q_n - 1)\alpha))| \\ &= |P_{2q_n-1}(\theta + b\alpha)| \\ &< \min\{|G_I(0, b)|^{-1} e^{(L+\epsilon_1)(b+2q_n-2)}, |G_I(0, b + 2q_n - 2)|^{-1} e^{-(L+\epsilon_1)b}\}. \end{aligned}$$

Therefore, using these last two results, we obtain that  $\theta + (b + q_n - 1)\alpha$  belongs to

$$\begin{cases} A_{2q_n-1, 23L/32+\epsilon} & \text{if } -\frac{13}{8}q_n \leq b \leq -\frac{3}{2}q_n \text{ or } -\frac{1}{2}q_n \leq b \leq -\frac{3}{8}q_n, \\ A_{2q_n-1, 5L/8+\epsilon} & \text{if } -\frac{3}{2}q_n \leq b \leq -\frac{1}{2}q_n \end{cases}$$

for any  $\epsilon > 0$  and for sufficiently large  $n$ . □

Fix  $1 \leq \ell \leq q_{n+1}^{8/9}/q_n$ . Set

$$I_1 = [-\lceil \frac{5}{8}q_n \rceil, \lceil \frac{5}{8}q_n \rceil - 1],$$

$$I_2 = [(\ell - 1)q_n + \lceil \frac{5}{8}q_n \rceil, (\ell + 1)q_n - \lceil \frac{5}{8}q_n \rceil - 1].$$

Set  $\theta_j = \theta + j\alpha$  for  $j \in I_1 \cup I_2$ . The set  $\{\theta_j\}_{j \in I_1 \cup I_2}$  consists of  $2q_n$  elements.

LEMMA 9.13. Assume  $L > \frac{16}{9}\beta$ . There exists an  $\epsilon > 0$  such that for sufficiently large  $n$ , the set  $\{\theta_j : j \in I_1 \cup I_2\}$  is  $(\frac{9}{32}L - \epsilon)$ -uniform.

We will now finish the proof of Lemma 9.5 and prove Lemma 9.13 at the end of the section.

Let  $k$  be resonant. Assume without loss of generality that  $k = \ell q_n + r$ , with  $0 \leq r \leq \max\{q_n^{8/9}, \frac{1}{20}q_{n-1}\}$  and  $1 \leq \ell \leq q_{n+1}^{8/9}/q_n$ .

By Lemmas 9.3, 9.12 and 9.13 there is a  $j_0 \in I_2$  such that  $\theta + j_0\alpha$  does not belong to  $A_{2q_n-1, 23L/32+\epsilon}$ . Set  $I = [j_0 - q_n + 1, j_0 + q_n - 1] = [x_1, x_2]$ . Then  $|G_I(k, x_i)| < e^{(L+\epsilon_1)(2q_n-2-|k-x_i|)-2q_n(23L/32+\epsilon)} < e^{q_n(9L/16+\epsilon)-(L+\epsilon)|k-x_i|}$ .

Since, by a simple computation,  $|k - x_i| > (5/8 - \epsilon - 1/20)q_n$ , we obtain that

$$(9-72) \quad |G_I(k, x_i)| < e^{(-L/46+\epsilon)|k-x_i|},$$

which gives the statement of Lemma 9.4. □

*Proof of Lemma 9.13.* As in the proof of Lemma 9.4 we will first estimate the numerator in (9-8). We have

$$\begin{aligned} (9-73) \quad & \sum_{\substack{j \in I_1 \cup I_2 \\ j \neq i}} \ln |\cos 2\pi a - \cos 2\pi \theta_j| \\ &= \sum_{\substack{j \in I_1 \cup I_2 \\ j \neq i}} \left( \ln \left| \sin 2\pi \frac{a + \theta_j}{2} \right| + \ln \left| \sin 2\pi \frac{a - \theta_j}{2} \right| \right) + (2q_n - 1) \ln 2 \\ &= \Sigma_+ + \Sigma_- + (2q_n - 1) \ln 2. \end{aligned}$$

Both  $\Sigma_+$  and  $\Sigma_-$  consist of 2 terms of the form of (9-27) with  $r = n$ , plus two terms of the form

$$\ln \min_{k=1, \dots, q_n} \left| \sin 2\pi \left( x + \frac{1}{2}(k + \ell_k q_n)\alpha \right) \right|,$$

where  $\ell_k \in \{0, \pm(\ell - 1), \pm\ell\}$  for  $k = 1, \dots, q_n$ , minus  $\ln |\sin 2\pi \frac{1}{2}(a \pm \theta_i)|$ .

Therefore, by (9-27)

$$(9-74) \quad \sum_{\substack{j \in I_1 \cup I_2 \\ j \neq i}} \ln |\cos 2\pi a - \cos 2\pi \theta_j| \leq (2 - 2q_n) \ln 2 + 4 \ln q_n + C \ell \Delta_n q_n \ln q_n.$$

To estimate the denominator of (9-8), we write it in the form (9-73) with  $a = \theta_i$ . Then  $\Sigma_- = \sum_{j \in I_1 \cup I_2, j \neq i} \ln |\sin \pi(i - j)\alpha|$  can be split into two sums of the form (9-27) plus the minimum term. The corresponding minimum term is achieved at  $|i - j_0|$  of the form  $q_n$  or  $\ell q_n$ . Therefore, for any  $\epsilon_1 > 0$  and sufficiently large  $n$

$$(9-75) \quad \begin{aligned} \Sigma_- &> -2q_n \ln 2 + \ln |\sin \pi q_n \alpha| - C \max(\ln q_n, \ell \Delta_n q_n \ln q_n) \\ &> -2q_n \ln 2 - \ln q_{n+1} - C \max(\ln q_n, \ell \Delta_n q_n \ln q_n). \end{aligned}$$

Since

$$\begin{aligned} \sin 2\pi(\theta + \frac{1}{2}(k + i + \ell_k q_n)\alpha) &= \sin 2\pi(\theta + \frac{1}{2}(k + i)\alpha) \cos \pi \ell_k \Delta_n \\ &\quad \pm \cos 2\pi(\theta + \frac{1}{2}(k + i)\alpha) \sin \pi \ell_k \Delta_n \end{aligned}$$

(the  $\pm$  depending on the sign of  $q_n \alpha - p_n$ ) we have by (9-1) that if  $q_{n+1} \geq q_n^{10}$  then

$$\min_{\substack{k, i \in [-q_n, q_n - 1] \\ \ell_k \in \{0, \pm(\ell - 1), \pm \ell\}}} |\sin 2\pi(\theta + \frac{(k + i + \ell_k q_n)\alpha}{2})| > \frac{1}{10} q_n^{-2},$$

and if  $q_{n+1} < q_n^{10}$  then we have the obvious

$$\min_{\substack{k, i \in [-q_n, q_n - 1] \\ \ell_k \in \{0, \pm(\ell - 1), \pm \ell\}}} |\sin 2\pi(\theta + \frac{1}{2}(k + i + \ell_k q_n)\alpha)| > \frac{1}{5} q_{n+1}^{-2} > \frac{1}{5} q_n^{-20}.$$

As before,  $\Sigma_+$  can be split into two sums of the form (9-27) plus two minimum terms minus  $\ln |\sin 2\pi(\theta + i\alpha)|$ . Therefore,

$$(9-76) \quad \Sigma_+ > -2q_n \ln 2 - C \max(\ln q_n, \ell \Delta_n q_n \ln q_n).$$

Combining (9-73), (9-74), (9-75) and (9-76), we obtain

$$(9-77) \quad \max_{j \in I_1 \cup I_2} \prod_{\substack{\ell \in I_1 \cup I_2 \\ \ell \neq j}} \frac{|z - \cos 2\pi \theta_\ell|}{|\cos 2\pi \theta_j - \cos 2\pi \theta_\ell|} < q_n^C e^{(\beta + \epsilon_1)q_n}.$$

For  $\beta < \frac{9}{16}L$  this gives the desired bound. □

### References

[AK06] A. AVILA and R. KRİKORIAN, Reducibility or nonuniform hyperbolicity for quasiperiodic Schrödinger cocycles, *Ann. of Math.* **164** (2006), 911–940. MR 2008h:81044 Zbl 1138.47033



- [AvMS90] J. AVRON, P. H. M. VAN MOUCHE, and B. SIMON, On the measure of the spectrum for the almost Mathieu operator, *Comm. Math. Phys.* **132** (1990), 103–118. MR 92d:39014a Zbl 0724.47002
- [AS83] J. AVRON and B. SIMON, Almost periodic Schrödinger operators. II: The integrated density of states, *Duke Math. J.* **50** (1983), 369–391. MR 85i:34009a Zbl 0544.35030
- [Azb64] M. Y. AZBEL, Energy spectrum of a conduction electron in a magnetic field, *Sov. Phys. JETP* **19** (1964), 634–645.
- [BS82] J. BÉLISSARD and B. SIMON, Cantor spectrum for the almost Mathieu equation, *J. Funct. Anal.* **48** (1982), 408–419. MR 84h:81019 Zbl 0516.47018
- [Ber68] J. M. BEREZANS'KIĬ, *Expansions in Eigenfunctions of Selfadjoint Operators*, *Transl. Math. Monogr.* **17**, Amer. Math. Soc., Providence, R.I., 1968. MR 36 #5768
- [BJ02] J. BOURGAIN and S. JITOMIRSKAYA, Continuity of the Lyapunov exponent for quasiperiodic operators with analytic potential, *J. Statist. Phys.* **108** (2002), 1203–1218. MR 2004c:47073 Zbl 1039.81019
- [CEY90] M. D. CHOI, G. A. ELLIOTT, and N. YUI, Gauss polynomials and the rotation algebra, *Invent. Math.* **99** (1990), 225–246. MR 91b:46067 Zbl 0665.46051
- [DCJ87] C. DE CONCINI and R. A. JOHNSON, The algebraic-geometric AKNS potentials, *Ergodic Theory Dynam. Systems* **7** (1987), 1–24. MR 88f:58065 Zbl 0636.35077
- [Fur97] A. FURMAN, On the multiplicative ergodic theorem for uniquely ergodic systems, *Ann. Inst. H. Poincaré Probab. Statist.* **33** (1997), 797–815. MR 98i:28018 Zbl 0892.60011
- [HS89] B. HELFFER and J. SJÖSTRAND, Semiclassical analysis for Harper's equation. III: Cantor structure of the spectrum, *Mém. Soc. Math. France* (1989), 1–124. MR 92m:35011 Zbl 0699.35273
- [Her83] M.-R. HERMAN, Une méthode pour minorer les exposants de Lyapounov et quelques exemples montrant le caractère local d'un théorème d'Arnol'd et de Moser sur le tore de dimension 2, *Comment. Math. Helv.* **58** (1983), 453–502. MR 85g:58057 Zbl 0554.58034
- [Hör94] L. HÖRMANDER, *Notions of Convexity*, *Progr. Math.* **127**, Birkhäuser, Boston, 1994. MR 95k:00002
- [JKS05] S. JITOMIRSKAYA, D. A. KOSLOVER, and M. S. SCHULTEIS, Localization for a family of one-dimensional quasiperiodic operators of magnetic origin, *Ann. Henri Poincaré* **6** (2005), 103–124. MR 2005j:81057 Zbl 1062.81029
- [JK02] S. Y. JITOMIRSKAYA and I. V. KRASOVSKY, Continuity of the measure of the spectrum for discrete quasiperiodic operators, *Math. Res. Lett.* **9** (2002), 413–421. MR 2004b:81051 Zbl 1020.47002
- [Jit99] S. Y. JITOMIRSKAYA, Metal-insulator transition for the almost Mathieu operator, *Ann. of Math.* **150** (1999), 1159–1175. MR 2000k:81084 Zbl 0946.47018
- [JM82] R. JOHNSON and J. MOSER, The rotation number for almost periodic potentials, *Comm. Math. Phys.* **84** (1982), 403–438. MR 83h:34018 Zbl 0497.35026
- [Las94] Y. LAST, Zero measure spectrum for the almost Mathieu operator, *Comm. Math. Phys.* **164** (1994), 421–432. MR 95f:47096 Zbl 0814.11040
- [Las05] Y. LAST, Spectral theory of Sturm-Liouville operators on infinite intervals: A review of recent developments, in *Sturm-Liouville Theory* (W. O. AMREIN, A. M. HINZ, and D. B. PEARSON, eds.), Birkhäuser, Basel, 2005, pp. 99–120. MR 2145079 Zbl 1098.39011
- [Pui04] J. PUIG, Cantor spectrum for the almost Mathieu operator, *Comm. Math. Phys.* **244** (2004), 297–309. MR 2004k:11129 Zbl 1075.39021

- [Sim83] B. SIMON, Kotani theory for one-dimensional stochastic Jacobi matrices, *Comm. Math. Phys.* **89** (1983), 227–234. MR 85d:60122 Zbl 0534.60057
- [Sim00] ———, Schrödinger operators in the twenty-first century, in *Mathematical Physics 2000* (A. FOKAS, A. GRIGORYAN, T. KIBBLE, and B. ZEGARLINSKI, eds.), Imp. Coll. Press, London, 2000, pp. 283–288. MR 2001g:81071 Zbl 1074.81521
- [Sin87] Y. G. SINAĬ, Anderson localization for one-dimensional difference Schrödinger operator with quasiperiodic potential, *J. Statist. Phys.* **46** (1987), 861–909. MR 88h:82016 Zbl 0682.34023
- [vM89] P. VAN MOUCHE, The coexistence problem for the discrete Mathieu operator, *Comm. Math. Phys.* **122** (1989), 23–33. MR 90e:47027 Zbl 0669.34016

(Received March 19, 2005)

*E-mail address:* artur@math.sunysb.edu

CNRS UMR 7599, LABORATOIRE DE PROBABILITÉS ET MODÈLES ALÉATOIRES,  
UNIVERSITÉ PIERRE ET MARIE CURIE, BOÎTE COURRIER 188, 75252 PARIS CEDEX 05,  
FRANCE

and

IMPA, ESTRADA DONA CASTORINA 110, JARDIM BOTÂNICO, 22460-320 RIO DE JANEIRO, RJ,  
BRAZIL

*E-mail address:* szhitomi@uci.edu

DEPARTMENT OF MATHEMATICS, 340 ROWLAND HALL, UNIVERSITY OF CALIFORNIA,  
IRVINE, CA 92697-3875, UNITED STATES