

ANNALS OF MATHEMATICS

**Nilpotency, almost nonnegative curvature,
and the gradient flow on Alexandrov
spaces**

By VITALI KAPOVITCH, ANTON PETRUNIN, and WILDERICH TUSCHMANN



SECOND SERIES, VOL. 171, NO. 1

January, 2010

ANMAAH

Nilpotency, almost nonnegative curvature, and the gradient flow on Alexandrov spaces

By VITALI KAPOVITCH, ANTON PETRUNIN, and WILDERICH TUSCHMANN

Abstract

We show that almost nonnegatively curved m -manifolds are, up to finite cover, nilpotent spaces in the sense of homotopy theory and have $C(m)$ -nilpotent fundamental groups. We also show that up to a finite cover almost nonnegatively curved manifolds are fiber bundles with simply connected fibers over nilmanifolds.

1. Introduction

Almost nonnegatively curved manifolds were introduced by Gromov in the late 70s [Gro80], with the most significant contributions to their study made by Yamaguchi in [Yam91] and Fukaya and Yamaguchi in [FY92]. Building on their ideas, in the present article we establish several new properties of these manifolds that yield, in particular, new topological obstructions to almost nonnegative curvature. Our techniques also provide simplified proofs of many results from [FY92].

A closed smooth manifold is said to be almost nonnegatively curved if it can Gromov-Hausdorff converge to a single point under a lower curvature bound. By rescaling, this definition is equivalent to the following one, which we will employ throughout this article.

Definition 1.0.1. A closed smooth manifold M is called almost nonnegatively curved if it admits a sequence of Riemannian metrics $\{g_n\}_{n \in \mathbb{N}}$ whose sectional curvatures and diameters satisfy

$$\sec(M, g_n) \geq -1/n \quad \text{and} \quad \text{diam}(M, g_n) \leq 1/n.$$

Almost nonnegatively curved manifolds generalize almost flat as well as nonnegatively curved manifolds. One main source of examples comes from a theorem

V. Kapovitch and A. Petrunin were supported in part by NSF grants #DMS-0204187 and #DMS-0406482, respectively. W. Tuschmann's research was partially supported by a DFG Heisenberg Fellowship.

of Fukaya and Yamaguchi. It states that if $F \rightarrow E \rightarrow B$ is a fiber bundle over an almost nonnegatively curved manifold B whose fiber F is compact and admits a nonnegatively curved metric that is invariant under the structure group, then the total space E is almost nonnegatively curved [FY92]. Further examples are given by closed manifolds that admit cohomogeneity one actions of compact Lie groups (compare [ST04]).

In this work we combine collapsing techniques with a nonsmooth analogue of the gradient flow of concave functions on Alexandrov spaces. This notion is based on the construction of gradient curves of λ -concave functions used in [PP96] and bears many similarities to the Sharafutdinov retraction [Sha78]. In more general settings, the gradient flow was studied in [Lyt05], [Oht09] and [Sav07]. The gradient flow on Alexandrov spaces plays a key role in the proofs of two of the three main results in this paper, and we believe that it should also prove useful for dealing with other problems related to collapsing under a lower curvature bound.

1.1. To put the main theorems of the present work into perspective, let us first briefly recall some previously known results:

Let $M = M^m$ be an almost nonnegatively curved m -manifold.

- Gromov proved in [Gro78] that the minimal number of generators of the fundamental group $\pi_1(M)$ of M can be estimated by a constant $C_1(m)$ depending only on m , and in [Gro81] that the sum of Betti numbers of M with respect to any field of coefficients does not exceed some uniform constant $C_2 = C_2(m)$.
- Yamaguchi showed that, up to a finite cover, M fibers over a flat $b_1(M; \mathbb{R})$ -dimensional torus and M^m is diffeomorphic to a torus if $b_1(M; \mathbb{R}) = m$ [Yam91].
- Fukaya and Yamaguchi proved that $\pi_1(M)$ is almost nilpotent, i.e., contains a nilpotent subgroup of finite index, and also that $\pi_1(M)$ is $C_3(m)$ -solvable, i.e., contains a solvable subgroup of index at most $C_3(m)$ [FY92].
- If a closed manifold has negative Yamabe constant, then it cannot volume collapse with scalar curvature bounded from below; see [Sch89], [LeB01]. In particular, no such manifold can be almost nonnegatively curved.
- The \hat{A} -genus of a closed spin manifold X of almost nonnegative Ricci curvature satisfies the inequality $\hat{A}(X) \leq 2^{\dim(X)/2}$; see [Gro82], [Gal83].

Let us now state the main results of this article.

1.2. Our first result concerns the hitherto unexplored relation between curvature bounds and the actions of the fundamental group on the higher homotopy groups.

An action by automorphisms of a group G on an abelian group V is called nilpotent if V admits a finite sequence of G -invariant subgroups

$$V = V_0 \supset V_1 \supset \cdots \supset V_k = 0$$

such that the induced action of G on V_i/V_{i+1} is trivial for any i . A connected CW-complex X is called *nilpotent* if $\pi_1(X)$ is a nilpotent group that operates nilpotently on $\pi_k(X)$ for every $k \geq 2$.

Nilpotent spaces play an important role in topology since they enjoy some of the best homotopy-theoretic properties of simply connected spaces, like a Whitehead theorem or reasonable Postnikov towers. Furthermore, unlike the category of simply connected spaces, the category of nilpotent ones is closed under many constructions such as the based loop space functor or the formation of function spaces, and group-theoretic functors, like localization and completion, have topological extensions in this category.

THEOREM A (Nilpotency Theorem). *Let M be a closed almost nonnegatively curved manifold. Then a finite cover of M is a nilpotent space.*

It would be interesting to know whether the order of this covering can be estimated solely in terms of the dimension of M .

Example 1.2.1. Let $h : S^3 \times S^3 \rightarrow S^3 \times S^3$ be defined by

$$h : (x, y) \mapsto (xy, yxy).$$

This map is a diffeomorphism with inverse given by

$$h^{-1} : (u, v) \mapsto (u^2v^{-1}, vu^{-1}).$$

The induced map h_* on $\pi_3(S^3 \times S^3)$ is given by the matrix $A_h = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$. Notice that the eigenvalues of A_h are different from 1 in absolute value. Let M be the mapping cylinder of h . Clearly, M has the structure of a fiber bundle $S^3 \times S^3 \rightarrow M \rightarrow S^1$, and the action of $\pi_1(M) \cong \mathbb{Z}$ on $\pi_3(M) \cong \mathbb{Z}^2$ is generated by A_h . In particular, M is not a nilpotent space and hence, by [Theorem A](#), it does not admit almost nonnegative curvature. This fact does not follow from any previously known results.

1.3. Our next main result provides an affirmative answer to a conjecture of Fukaya and Yamaguchi [[FY92](#), Conj. 0.15].

THEOREM B (C-Nilpotency Theorem for π_1). *Let M be an almost nonnegatively curved m -manifold. Then $\pi_1(M)$ is $C(m)$ -nilpotent, i.e., $\pi_1(M)$ contains a nilpotent subgroup of index at most $C(m)$.*

Notice that [Theorem B](#) is new even for manifolds of nonnegative curvature.

Example 1.3.1. For any $C > 0$ there exist prime numbers $p > q > C$ and a finite group G_{pq} of order pq that is solvable but not nilpotent. In particular, G_{pq} does not contain any nilpotent subgroup of index less than or equal to C .

Whereas none of the results mentioned so far excludes G_{pq} from being the fundamental group of some almost nonnegatively curved m -manifold, [Theorem B](#) shows that for $C > C(m)$ none of the groups G_{pq} can be realized as the fundamental group of such a manifold.

1.4. In [\[FY92\]](#), Fukaya and Yamaguchi also conjectured that a finite cover of an almost nonnegatively Ricci curved manifold M fibers over a nilmanifold with a fiber that has nonnegative Ricci curvature and whose fundamental group is finite. This conjecture was later refuted by Anderson [\[And92\]](#).

It is, on the other hand, very natural to consider this conjecture in the context of almost nonnegative *sectional* curvature. In fact, here Yamaguchi's fibration theorem (see [\[Yam91\]](#)) and the results of [\[FY92\]](#) easily imply that a finite cover of an almost nonnegatively curved manifold admits a map onto a nilmanifold whose homotopy fiber is a simply connected closed manifold.

From mere topology, it is, however, not clear whether this homotopy fibration can actually always be made into a genuine fiber bundle. Our next result shows that this is indeed true, and that for manifolds of almost nonnegative sectional curvature Fukaya and Yamaguchi's original conjecture essentially does hold.

THEOREM C (Fibration Theorem). *Let M be an almost nonnegatively curved manifold. Then a finite cover \tilde{M} of M is the total space of a fiber bundle*

$$F \rightarrow \tilde{M} \rightarrow N$$

over a nilmanifold N with a simply connected fiber F . Moreover, the fiber F is almost nonnegatively curved in the sense of the following definition.

Definition 1.4.1. A closed smooth manifold M is called almost nonnegatively curved in the generalized sense if for some nonnegative integer k there exists a sequence of complete Riemannian metrics g_n on $M \times \mathbb{R}^k$ and points $p_n \in M \times \mathbb{R}^k$ such that

- (1) the sectional curvatures of the metric balls of radius n around p_n satisfy

$$\sec(B_n(p_n)) \geq -1/n;$$

- (2) for $n \rightarrow \infty$ the pointed Riemannian manifolds $((M \times \mathbb{R}^k, g_n), p_n)$ converge in the pointed Gromov-Hausdorff distance to $(\mathbb{R}^k, 0)$;
- (3) the regular fibers over 0 are diffeomorphic to M for all large n .

Due to Yamaguchi’s fibration theorem [Yam91], manifolds that are almost nonnegatively curved in the generalized sense play the same central role in collapsing under a lower curvature bound as almost flat manifolds do in the Cheeger–Fukaya–Gromov theory of collapsing with bounded curvature; see [CFG92].

It is not yet known whether all manifolds that are almost nonnegatively curved in the generalized sense are almost nonnegatively curved. Clearly if $k = 0$, this definition reduces to the standard one. Moreover, it is easy to see that all results of the present article, as well as all results about almost nonnegatively curved manifolds mentioned earlier (except possibly for the ones concerning the \hat{A} -genus and Yamabe constant), hold for manifolds that are almost nonnegatively curved in the sense of Definition 1.4.1.

1.5. Let us now describe the structure of the rest of this article.

In Section 2, after providing some necessary background from Alexandrov geometry, we introduce the gradient flow of the square of a distance function. It serves as one of the main technical tools in the proofs of Theorem A and Theorem B.

In Section 3, we prove Theorem A by a direct application of the gradient flow technique.

In Section 4, we prove Theorem B. The proof is also based on the gradient flow, but is more involved and employs further technical tools such as “limit fundamental groups” of Alexandrov spaces.

In Section 5, we prove Theorem C. This section is completely independent from the rest of the article.

In Section 6, we discuss some further open questions related to our results.

2. Alexandrov geometry and the gradient flow

This section provides necessary background in Alexandrov geometry. The results of Sections 2.1–2.3 are mostly duplicated from [PP96], [Pet95] and [Pet07]. See [BGP92] for a general reference on Alexandrov spaces.

2.1. λ -concave functions.

Definition 2.1.1 (for a space without boundary). Let A be an Alexandrov space without boundary. A Lipschitz function $f: A \rightarrow \mathbb{R}$ is called λ -concave if for any unit speed minimizing geodesic γ in A , the function

$$f \circ \gamma(t) - \lambda t^2/2 \quad \text{is concave.}$$

If A is an Alexandrov space with boundary, then its double \tilde{A} is also an Alexandrov space; see [Per91, §5.2]. Let $p: \tilde{A} \rightarrow A$ be the canonical map. Given a function f on A , set $\tilde{f} = f \circ p$.

Definition 2.1.2 (for a space with boundary). Let A be an Alexandrov space with boundary. A Lipschitz function $f: A \rightarrow \mathbb{R}$ is called λ -concave if for any unit speed minimizing geodesic γ in \bar{A} , the function

$$\tilde{f} \circ \gamma(t) - \lambda t^2/2 \quad \text{is concave.}$$

Remark 2.1.3. Notice that the restriction of a linear function on \mathbb{R}^n to a ball is not 0-concave in this sense.

Remark 2.1.4. In the above definitions, the Lipschitz condition is only technical. With some extra work, all results of this section can be extended to continuous functions.

2.2. Tangent cone and differential. Given a point p in an Alexandrov space A , we denote by $T_p = T_p(A)$ the tangent cone at p .

If d denotes the metric of an Alexandrov space A , let us denote by λA the space $(A, \lambda d)$. Let $i_\lambda : \lambda A \rightarrow A$ be the canonical map. The limit of $(\lambda A, p)$ for $\lambda \rightarrow \infty$ is the tangent cone T_p at p ; see [BGP92, §7.8.1].

Definition 2.2.1. For any function $f: A \rightarrow \mathbb{R}$ the function $d_p f: T_p \rightarrow \mathbb{R}$ such that

$$d_p f = \lim_{\lambda \rightarrow \infty} \lambda(f \circ i_\lambda - f(p))$$

is called the differential of f at p .

It is easy to see that for a λ -concave function f , the differential $d_p f$ is defined everywhere, and that $d_p f$ is a 0-concave function on the tangent cone T_p .

Definition 2.2.2. Given a λ -concave function $f: A \rightarrow \mathbb{R}$, a point $p \in A$ is called critical point of f if $d_p f \leq 0$.

2.3. Gradient curves. With a slight abuse of notation we will call elements of the tangent cone T_p the “tangent vectors” at p . The origin of T_p plays the role of the zero vector and is denoted by $o = o_p$. For a tangent vector v at p we define its absolute value $|v|$ as the distance $|ov|$ in T_p . For two tangent vectors u and v at p we can define their “scalar product”

$$\langle u, v \rangle = (|u|^2 + |v|^2 - |uv|^2)/2 = |u| \cdot |v| \cos \alpha,$$

where $\alpha = \angle uov$ in T_p .

For two points $p, q \in A$ we define $\log_p q$ to be a tangent vector v at p such that $|v| = |pq|$ and such that the direction of v coincides with a direction from p to q (if such a direction is not unique, we choose any one of them). Given a curve $\gamma(t)$ in A , we denote by $\gamma^+(t)$ the right and by $\gamma^-(t)$ the left tangent vectors to $\gamma(t)$, where, respectively,

$$\gamma^\pm(t) \in T_{\gamma(t)} \quad \text{and} \quad \gamma^\pm(t) = \lim_{\varepsilon \rightarrow +0} \frac{\log_{\gamma(t)} \gamma(t \pm \varepsilon)}{\varepsilon}.$$

For a real function $f(t)$ with $t \in \mathbb{R}$ we denote by $f^+(t)$ its right derivative and by $-f^-(t)$ its left derivative. Note that our sign convention (which is chosen to agree with the notion of right and left derivatives of curves) is not quite standard. For example,

$$\text{if } f(t) = t, \quad \text{then } f^+(t) \equiv 1 \quad \text{and} \quad f^-(t) \equiv -1.$$

Definition 2.3.1. For f a λ -concave function on A , a vector $g \in T_p(A)$ is called a gradient of f at $p \in A$ (in short: $g = \nabla_p f$) if

- (i) $d_p f(x) \leq \langle g, x \rangle$ for any $x \in T_p$, and
- (ii) $d_p f(g) = \langle g, g \rangle$.

It is easy to see that any λ -concave function has a uniquely defined gradient vector field. Moreover, if $d_p f(x) \leq 0$ for all $x \in T_p$, then $\nabla_p f = o$ (here o denotes the origin of the tangent cone T_p); otherwise

$$\nabla_p f = d_p f(\xi)\xi,$$

where ξ is the (necessarily unique) unit vector for which the function $d_p f$ attains its maximum.

Moreover, for any minimizing geodesic $\gamma : [a, b] \rightarrow U$ parametrized by arc length, the following inequality holds:

$$(2.3.1) \quad \langle \gamma^+(a), \nabla_{\gamma(a)} f \rangle + \langle \gamma^-(b), \nabla_{\gamma(b)} f \rangle \geq -\lambda(b-a).$$

Indeed,

$$\begin{aligned} \langle \gamma^+(a), \nabla_{\gamma(a)} f \rangle + \langle \gamma^-(b), \nabla_{\gamma(b)} f \rangle &\geq d_{\gamma(a)} f(\gamma^+(a)) + d_{\gamma(b)} f(\gamma^-(b)) \\ &= (f \circ \gamma)^+|_a + (f \circ \gamma)^-|_b \geq -\lambda(b-a). \end{aligned}$$

Definition 2.3.2. A curve $\alpha : [a, b] \rightarrow A$ is called an f -gradient curve if

$$\alpha^+(t) = \nabla_{\alpha(t)} f \quad \text{for any } t \in [a, b].$$

PROPOSITION 2.3.3. *Given a λ -concave function $f : A \rightarrow \mathbb{R}$ and a point $p \in A$, there is a unique gradient curve $\alpha : [0, \infty) \rightarrow A$ such that $\alpha(0) = p$.*

Moreover, if α and β are two f -gradient curves, then

$$|\alpha(t_1)\beta(t_1)| \leq |\alpha(t_0)\beta(t_0)| \exp(\lambda(t_1 - t_0)) \quad \text{for all } t_1 \geq t_0.$$

The gradient curve can be constructed as a limit of broken geodesics, made up of short segments with directions close to the gradient. The convergence, uniqueness, as well as the last inequality in [Proposition 2.3.3](#) follow from inequality [\(2.3.1\)](#) above, while [Corollary 2.3.5](#) below guarantees that the limit is indeed a gradient curve, having a unique right tangent vector at each point.

LEMMA 2.3.4. *Let $A_n \xrightarrow{\text{GH}} A$ be a sequence of Alexandrov spaces with curvature $\geq k$ which Gromov-Hausdorff converges to an Alexandrov space A .*

Let $f_n \rightarrow f$, where $f_n : A_n \rightarrow \mathbb{R}$ is a sequence of λ -concave functions converging to $f : A \rightarrow \mathbb{R}$. Let $p_n \rightarrow p$, where $p_n \in A_n$ and $p \in A$. Then

$$|\nabla_p f| \leq \liminf_{n \rightarrow \infty} |\nabla_{p_n} f_n|.$$

COROLLARY 2.3.5. *Given a λ -concave function f on A and a sequence of points $p_n \in A$ such that $p_n \rightarrow p$, we have*

$$|\nabla_p f| \leq \liminf_{n \rightarrow \infty} |\nabla_{p_n} f|.$$

Proof of Lemma 2.3.4. Fix an $\varepsilon > 0$ and choose q near p such that

$$(f(q) - f(p))/|pq| > |\nabla_p f| - \varepsilon.$$

Now choose $q_n \in A_n$ such that $q_n \rightarrow q$. If $|pq|$ is sufficiently small and n is sufficiently large, the λ -concavity of f_n then implies that

$$\liminf_{n \rightarrow \infty} \frac{d_{p_n} f_n(v_n)}{|v_n|} \geq |\nabla_p f| - 2\varepsilon \quad \text{for } v_n = \log_{p_n}(q_n) \in T_{p_n}(A_n).$$

Therefore,

$$\liminf_{n \rightarrow \infty} |\nabla_{p_n} f_n| \geq |\nabla_p f| - 2\varepsilon \quad \text{for any } \varepsilon > 0,$$

i.e.,

$$\liminf_{n \rightarrow \infty} |\nabla_{p_n} f_n| \geq |\nabla_p f|. \quad \square$$

LEMMA 2.3.6. *Let f be a λ -concave function, $\lambda \geq 0$ and $\alpha(t)$ be an f -gradient curve, and let $\bar{\alpha}(s)$ be its reparametrization by arc length. Then $f \circ \bar{\alpha}$ is λ -concave.*

Proof.

$$\begin{aligned} (f \circ \bar{\alpha})^+(s_0) &= |\nabla_{\bar{\alpha}(s_0)} f| \geq \frac{d_{\bar{\alpha}(s_0)} f(\log_{\bar{\alpha}(s_0)}(\bar{\alpha}(s_1)))}{|\bar{\alpha}(s_1) \bar{\alpha}(s_0)|} \\ &\geq \frac{f(\bar{\alpha}(s_1)) - f(\bar{\alpha}(s_0)) - \lambda |\bar{\alpha}(s_1) \bar{\alpha}(s_0)|^2 / 2}{|\bar{\alpha}(s_1) \bar{\alpha}(s_0)|} \\ &\geq \frac{f(\bar{\alpha}(s_1)) - f(\bar{\alpha}(s_0))}{s_1 - s_0} - \lambda |s_1 - s_0| / 2. \end{aligned}$$

Since $|\bar{\alpha}(s_1) \bar{\alpha}(s_0)| / (s_1 - s_0)$ goes to 1 as s_1 goes to s_0+ , it follows that $f \circ \bar{\alpha}$ is λ -concave. □

2.4. *The gradient flow on Alexandrov spaces.* Let f be a λ -concave function on an Alexandrov space A . Consider the map $\Phi_f^T : A \rightarrow A$ defined as follows: $\Phi_f^T(x) = \alpha_x(T)$, where $\alpha_x : [0, \infty) \rightarrow A$ is the f -gradient curve with $\alpha_x(0) = x$. The map Φ_f^T is called the f -gradient flow at time T . From Proposition 2.3.3, it is

clear that Φ_f^T is an $\exp(\lambda T)$ -Lipschitz map. Next we want to prove that this map behaves nicely under Gromov-Hausdorff-convergence.

THEOREM 2.4.1. *Let $A_n \xrightarrow{\text{GH}} A$ be a sequence of Alexandrov spaces with curvature $\geq k$ that converges to an Alexandrov space A .*

Let $f_n \rightarrow f$, where $f_n : A_n \rightarrow \mathbb{R}$ is a sequence of λ -concave functions and $f : A \rightarrow \mathbb{R}$.

Then $\Phi_{f_n}^T \rightarrow \Phi_f^T$.

Theorem 2.4.1 immediately follows from the following lemma:

LEMMA 2.4.2. *Let $A_n \xrightarrow{\text{GH}} A$ be a sequence of Alexandrov spaces with curvature $\geq k$ that converges to an Alexandrov space A .*

Let $f_n \rightarrow f$, where $f_n : A_n \rightarrow \mathbb{R}$ is a sequence of λ -concave functions and $f : A \rightarrow \mathbb{R}$. Let $\alpha_n : [0, \infty) \rightarrow A_n$ be the sequence of f_n -gradient curves with $\alpha_n(0) = p_n$, and let $\alpha : [0, \infty) \rightarrow A$ be the f -gradient curve with $\alpha(0) = p$.

Then $\alpha_n \rightarrow \alpha$.

Proof. We may assume without loss of generality that f has no critical points. (Otherwise consider instead the sequence $A'_n = A_n \times \mathbb{R}$ with $f'_n(a \times x) = f_n(a) + x$.)

Let $\bar{\alpha}_n(s)$ denote the reparametrization of $\alpha_n(t)$ by arc length. Since all $\bar{\alpha}_n$ are 1-Lipschitz, we can choose a converging subsequence from any subsequence of $\bar{\alpha}_n$. Let $\bar{\beta} : [0, \infty) \rightarrow A$ be its limit.

Clearly, $\bar{\beta}$ is also 1-Lipschitz and hence $|\bar{\beta}^+| \leq 1$. Hence by [Lemma 2.3.4](#),

$$\begin{aligned} \lim_{n \rightarrow \infty} f_n \circ \bar{\alpha}_n|_a^b &= \lim_{n \rightarrow \infty} \int_a^b |\nabla_{\bar{\alpha}_n(s)} f_n| ds \\ &\geq \int_a^b |\nabla_{\bar{\beta}(s)} f| \geq \int_a^b d_{\beta(t)} f(\beta^+(t)) = f \circ \beta|_a^b. \end{aligned}$$

On the other hand, since $\bar{\alpha}_n \rightarrow \bar{\beta}$ and $f_n \rightarrow f$, we have $f_n \circ \bar{\alpha}_n|_a^b \rightarrow f \circ \bar{\beta}|_a^b$. Therefore, in both of these inequalities in fact equality holds.

Hence, $|\nabla_{\bar{\beta}(s)} f| = \lim_{n \rightarrow \infty} |\nabla_{\bar{\alpha}_n(s)} f_n|$, $|\bar{\beta}^+(s)| = 1$ and the directions of $\bar{\beta}^+(s)$ and $\nabla_{\bar{\beta}(s)} f$ coincide almost everywhere. This implies that $\bar{\beta}(s)$ is a gradient curve reparametrized by arc length. In other words, if $\bar{\alpha}(s)$ denotes the reparametrization of $\alpha(t)$ by arc length, then $\bar{\beta}(s) = \bar{\alpha}(s)$ for all s . It only remains to show that the original parameter $t_n(s)$ of α_n converges to the original parameter $t(s)$ of α .

Notice that $|\nabla_{\bar{\alpha}_n(s)} f_n| dt_n = ds$ or $dt_n/ds = ds/d(f_n \circ \bar{\alpha}_n)$. Likewise, $dt/ds = ds/d(f \circ \bar{\alpha})$. Then the convergence $t_n \rightarrow t$ follows from the λ -concavity of $f_n \circ \bar{\alpha}_n$ (see [Lemma 2.3.6](#)) and the convergence $f_n \circ \bar{\alpha}_n \rightarrow f \circ \bar{\alpha}$. \square

2.5. *Gradient balls.* Let A be an Alexandrov space, and let $S \subset A$ be a subset of A . A function $f : A \rightarrow \mathbb{R}$ that can be represented as

$$f = \sum_i \theta_i \frac{1}{2} \text{dist}_{a_i}^2, \quad \text{with } \theta_i \geq 0, \quad \sum_i \theta_i = 1 \quad \text{and } a_i \in S$$

will be called a *cocos-function with respect to S* (where ‘‘cocos’’ stands for **convex combination of squares of distance functions**). A broken gradient curve for a collection of such functions will be called *cocos-curve with respect to S* .

For $p \in A$ and $T, r \in \mathbb{R}_+$, let us define $\beta_T^r(p)$, the gradient ball with center p and radius T with respect to $B_r(p)$, as the set of all end points of cocos-curves with respect to $B_r(p)$ that start at p with total time $\leq T$.

LEMMA 2.5.1. (I) *There exists a $T = T(m) \in \mathbb{R}_+$ such that for any m -dimensional Alexandrov space A with curvature ≥ -1 and any $q \in A$ there is a point $p \in A$ such that (i) $|pq| \leq 1$ and (ii) $B_1(p) \subset \beta_T^1(p)$.*

(II) *There exists a $T' = T'(m) \in \mathbb{R}_+$ such that the following holds. Let A be an Alexandrov space that is a quotient $A = \tilde{A}/\Gamma$ of an m -dimensional Alexandrov space \tilde{A} with curvature ≥ -1 by a discrete action of a group of isometries Γ . Let $q \in A$ and $p = p(q) \in A$ be as in part (I) above.*

Then for any lift $\tilde{p} \in \tilde{A}$ of p , one has that $B_1(\tilde{p}) \subset \beta_{T'}^1(\tilde{p})$.

Proof. The proof is similar to the construction of a strained point in an Alexandrov space; see [BGP92].

Set $\delta = 10^{-m}$. Take $a_1 = q$, and take b_1 to be a farthest point from a_1 in the closed ball $\bar{B}_1(a_1)$. Take a_2 to be a midpoint of a_1b_1 , and let b_2 be a farthest point from a_2 such that $|a_1b_2| = |a_1a_2|$ and $|a_2b_2| \leq \delta|a_1b_1|$, etc. On the k -th step we have to take a_k to be a midpoint of $a_{k-1}b_{k-1}$ and b_k to be a farthest point from a_k such that $|a_i b_k| = |a_i a_k|$ for all $i < k$ and $|a_k b_k| \leq \delta|a_{k-1}b_{k-1}|$.

After m steps, take p to be a midpoint of $a_m b_m$. We only have to check that we can find a $T = T(m)$ such that $\beta_T^1(p) \supset B_1(p)$.

Let t_i be the minimal time such that $B_{|a_i b_i|/\delta^m}(p) \subset \beta_{t_i}^1(p)$. Then one can take $T = t_1$. Therefore it is enough to give estimates for t_m and t_{k-1}/t_k only in terms of δ and m . Looking at the ends of broken gradient curves starting at p for the functions $\text{dist}_p^2/2$, $\text{dist}_{a_i}^2/2$ and $\text{dist}_{b_i}^2/2$, we easily see that $t_n \leq 1/\delta^m$. Now, looking at the ends of broken gradient curves starting at $B_{|a_{k-1}b_{k-1}|/\delta^m}(p)$ for the functions $\text{dist}_p^2/2$, $\text{dist}_{a_i}^2/2$ and $\text{dist}_{b_i}^2/2$, we have that $t_{k-1}/t_k \leq 1/\delta^m$. Therefore $t_1 \leq 1/\delta^{m^2} = 10^{-m^3}$. This finishes the proof of part (I).

For part (II), notice that

- (a) for any $r, t > 0$ we have $\beta_t^r(p) \subset B_{re^t}(p)$;
- (b) if $\beta_t^r(p) \supset B_\rho(p)$, then $\beta_{t+\tau}^r(p) \supset B_{\rho e^\tau}(p)$;
- (c) if $\rho = |px|$ and $x \in \beta_t^{r+\rho}(p)$, then $\beta_\tau^r(x) \subset \beta_{t+\tau}^{r+\rho}(p)$.

Take $\varepsilon = e^{-T}/4$ and apply part (I) of the lemma to \tilde{A}/ε to find a point $p' \in \tilde{A}$ such that $|\tilde{p}p'| \leq \varepsilon$ and $B_\varepsilon(p') \subset \beta_T^\varepsilon(p') \subset \tilde{A}$. Then for some deck transformation γ , we have $\gamma p' \in \beta_T^\varepsilon(p) \subset B_{\varepsilon e^T}(p)$. Therefore $\gamma p' \in B_{1/2}(\tilde{p})$. Hence, taking $T' = 2T + 1/\varepsilon = 2T + 4e^T$, we obtain

$$\beta_{T'}^\varepsilon(p) \supset \beta_{T'+1/\varepsilon}^\varepsilon(\gamma p') \supset B_1(\tilde{p}). \quad \square$$

2.6. *Short basis.* We will use the following construction due to Gromov.

Given an Alexandrov space A with a marked point $p \in A$, and a group Γ acting discretely on $A = (A, d)$, one can define a short basis of the action of Γ at p as follows:

For $\gamma \in \Gamma$, define the norm of γ by the formula $|\gamma| = d(p, \gamma(p))$. Choose $\gamma_1 \in \Gamma$ with the minimal norm in Γ . Then choose γ_2 to have minimal norm in $\Gamma \setminus \langle \gamma_1 \rangle$. On the n -th step, choose γ_n to have minimal norm in $\Gamma \setminus \langle \gamma_1, \gamma_2, \dots, \gamma_{n-1} \rangle$. The sequence $\{\gamma_1, \gamma_2, \dots\}$ is called a *short basis* of Γ at p . In general, the number of elements of a short basis can be finite or infinite. In the special case of the action of the fundamental group $\pi_1(A, p)$ on the universal cover of A , one speaks of the short basis of $\pi_1(A, p)$.

It is easy to see that for a short basis $\{\gamma_1, \gamma_2, \dots\}$ of the fundamental group of an Alexandrov space A the following is true:

- (1) If A has diameter d , then $|\gamma_i| \leq 2d$.
- (2) If A is compact, then $\{\gamma_i\}$ is finite.
- (3) For any $i > j$, we have $|\gamma_i| \leq |\gamma_j^{-1}\gamma_i|$.

The third property implies that if $\tilde{p} \in \tilde{A}$ is in the preimage of p in the universal cover \tilde{A} of A and $\tilde{p}_i = \gamma_i(\tilde{p})$, then

$$|\tilde{p}_i \tilde{p}_j| \geq \max\{|\tilde{p} \tilde{p}_i|, |\tilde{p} \tilde{p}_j|\}.$$

As Gromov observed, if A is an Alexandrov space with curvature $\geq \kappa$ and diameter $\leq d$, the last inequality implies that $\angle \tilde{p}_i \tilde{p} \tilde{p}_j > \delta = \delta(\kappa, d) > 0$. This yields an upper bound on the number of elements of a short basis in terms of κ , d and the dimension of A .

3. Nilpotency of almost nonnegatively curved manifolds

In this section we prove [Theorem A](#).

3.1. *Preliminary lemmas.* Suppose M is an almost nonnegatively curved manifold. Let us denote by $M_n = (M, g_n)$ for $n \in \mathbb{N}$ a sequence of Riemannian metrics on M such that $\sec(M_n) \geq -1/n$ and $\text{diam}(M_n) \leq 1/n$. Let us denote by \tilde{M} the universal covering of M , and by $\tilde{M}_n \rightarrow M_n$ the universal Riemannian covering of M_n (i.e., \tilde{M}_n is \tilde{M} equipped with the pullback of the Riemannian metric g_n).

KEY LEMMA 3.1.1. *Given $\varepsilon > 0$ and $r_2 > r_1 > 0$, suppose that $\widetilde{M}_n \supset B_{r_2}(p_n) \supset B_{r_1}(p_n)$. Then, for n sufficiently large, there is a $(1 + \varepsilon)$ -Lipschitz map $\Phi_n : B_{r_2}(p_n) \rightarrow B_{r_1}(p_n)$ that is homotopic to the identity on $B_{r_2}(p_n)$.*

Proof. Fix $R \gg r_2$ (here $R > 1000(1 + 1/\varepsilon)r_2$ will suffice). Note that

$$B_R(p_n) \xrightarrow{\text{GH}} B_R \subset \mathbb{R}^q \quad \text{as } n \rightarrow \infty.$$

Choose a finite $R/1000$ -net $\{a_i\}$ of $\partial B_R \subset \mathbb{R}^q$. Let $a_{i,n} \in M_n$ be sequences such that $a_{i,n} \rightarrow a_n$. Consider the sequence of functions $f_n : M_n \rightarrow \mathbb{R}$ with $f_n = \min_i \text{dist}_{a_{i,n}}^2$.

For large n , the functions f_n are 2-concave in $B_R(p_n)$, so that, in particular, the gradient flows $\Phi_{f_n}^T|_{B_{r_2}(p_n)}$ are e^{2T} -Lipschitz. Moreover, if ξ_x denotes the starting vector of a unit speed shortest geodesic from x to p_n , then for any $x \in B_{r_2}(p_n) \setminus B_{r_1}(p_n)$ we have $\langle \xi_x, \nabla f \rangle \geq R/2$. Therefore,

$$\Phi_{f_n}^T(B_{r_2}(p_n)) \subset B_{r_1}(p_n) \quad \text{if } T = 2r_2/R.$$

Thus $\Phi_n = \Phi_{f_n}^{2r_2/R}$ provides a $4r_2/R$ -Lipschitz map $B_{r_2}(p_n) \rightarrow B_{r_1}(p_n)$, and it is $(1 + \varepsilon)$ -Lipschitz if one chooses R sufficiently large. \square

For $\gamma \in \pi_1(M)$, set $|\gamma|_n = |p\gamma(p)|_{\widetilde{M}_n}$; see [Section 2.6](#).

COROLLARY 3.1.2. *Let M be almost nonnegatively curved manifold. Let*

$$h : \pi_1(M) \rightarrow \text{Aut}(H^*(\widetilde{M}, \mathbb{Z})/\text{tor})$$

be the natural action of $\pi_1(M)$ on $H^(\widetilde{M}, \mathbb{Z})$. Then there is a sequence of norms $\|\cdot\|_n$ on $H^*(\widetilde{M}, \mathbb{Z})/\text{tor}$ such that the following holds. Given any $\varepsilon > 0$, there is $n \in \mathbb{Z}_+$ such that for any $\gamma \in \pi_1(M)$ with $|\gamma|_n \leq 2 \text{diam}(M_n)$, we have $\|h(\gamma)\|_n \leq 1 + \varepsilon$.*

Proof. [[FY92](#), Th. 0.1] and Yamaguchi's fibration theorem [[Yam91](#)] imply that if n is sufficiently large, for any fixed $r \in \mathbb{R}_+$ we have that for any $p_n \in \widetilde{M}_n$ the inclusion map $B_r(p_n) \rightarrow \widetilde{M}_n$ is a homotopy equivalence.

Let $\|\cdot\|_{n,r}$ denote the L_∞ -norm on differential forms on $B_r(p_n) \subset \widetilde{M}_n$.

Fix $r_2 > r_1 > 0$. If ω is a differential form on $B_{r_1}(p_n) \subset M_n$ and n is sufficiently large, [Key Lemma 3.1.1](#) implies that

$$\|\Phi_n^*(\omega)\|_{n,r_2} \leq (1 + \varepsilon)\|\omega\|_{n,r_1} \quad \text{and} \quad 2 \text{diam}(M_n) \leq r_2 - r_1.$$

If now ω is a form on $B_{r_2}(p_n) \in \widetilde{M}_n$ and $\gamma \in \pi_1(M)$ is such that

$$|\gamma|_n = |p_n\gamma(p_n)| \leq 2 \text{diam}(M_n) \leq r_2 - r_1,$$

then $B_{r_1}(p_n) \subset B_{r_2}(\gamma(p_n)) \subset \widetilde{M}_n$, whence

$$\|\Phi_n^*(\gamma^*(\omega))\|_{n,r_2} \leq (1 + \varepsilon)\|\gamma^*(\omega)\|_{n,r_1} \leq (1 + \varepsilon)\|\omega\|_{n,r_2}.$$

Thus, for the induced norms on the de Rham cohomology of \widetilde{M} (and on its integral subspace $H^*(\widetilde{M}, \mathbb{Z})/\text{tor}$), we have

$$\|[\gamma^*(\omega)]\|_{n,r_2} \leq (1 + \varepsilon)\|\omega\|_{n,r_2}.$$

Therefore the sequence of norms $\|*\|_n = \|*\|_{n,r_2}$ satisfies the conditions of the corollary. \square

LEMMA 3.1.3. *There exists a constant $N = N(n, k) \in \mathbb{Z}_+$ such that the following holds. If G is a subgroup of $\text{GL}(n, \mathbb{Z})$ and S is a set of generators of G with $\#(S) \leq k$ such that the eigenvalues of each element of S^N are all equal to 1 in absolute value, then the same is true for the eigenvalues of all elements of G .*

Proof. Let B be the set of all matrices in $\text{GL}(n, \mathbb{Z})$ whose eigenvalues are all equal to 1 in absolute value. Since the characteristic polynomials of such matrices are uniformly bounded and have integer coefficients, there are only finitely many of them. Let \bar{B} be the Zariski closure of B in the set of all real $n \times n$ matrices. By the above, all eigenvalues of elements of \bar{B} also have absolute value 1.

Consider now the space $V = \mathbb{R}^{kn^2}$ of k -tuples of real $n \times n$ matrices.

Consider a collection of matrices $(M_1, M_2, \dots, M_k) \in V$, where $M_i \in \text{GL}(n, \mathbb{R})$. Let F_k be a free group on k generators, generated by $S = \{\gamma_1, \gamma_2, \dots, \gamma_k\}$, and let $h : F_k \rightarrow \text{GL}(n, \mathbb{R})$ be the homomorphism defined by $h(\gamma_i) = M_i$. The property that $h(\gamma)$ is an element of \bar{B} for any $\gamma \in F_k$ then describes an algebraic subset $A_\gamma \subset V$.

The intersection $A = \bigcap_{\gamma \in F_k} A_\gamma$ is also algebraic, and therefore there is a finite number $N = N(n, k)$ such that $A = \bigcap_{\gamma \in S^N} A_\gamma$ for $S^N \subset F_k$. \square

LEMMA 3.1.4. *Let Γ be a subgroup of $\text{GL}(n, \mathbb{Z})$ such that the eigenvalues of each element of Γ are equal to 1 in absolute value. Then Γ contains a subgroup Γ' of finite index whose elements' eigenvalues are all equal to 1.*

Proof. Let G denote the Zariski closure of Γ in $\text{GL}(n, \mathbb{R})$. Then G , being an algebraic group, is a Lie group with finitely many components. Let G_\circ be the identity component of G . By the same argument as in the proof of the previous lemma, the set of all characteristic polynomials of the elements of G is finite. Therefore the characteristic polynomial of any element of G_\circ is identically equal to $(x - 1)^n$.

Thus the subgroup $\Gamma' = \Gamma \cap G_\circ$ satisfies all conditions of the lemma. \square

Remark 3.1.5. As was pointed out to us by Yu. Zarkhin, one can alternatively take Γ' to be the kernel of the composition of the homomorphisms $\Gamma \rightarrow \text{GL}(n, \mathbb{Z}) \rightarrow \text{GL}(n, \mathbb{Z}/3\mathbb{Z})$. In this way one obtains a bound $[\Gamma : \Gamma'] \leq 3^{n^2}$. To see that Γ' satisfies the conclusion of Lemma 3.1.4, one should notice that every element of Γ is a quasi-unipotent matrix, since all its eigenvalues are roots of unity. The desired result then follows from the so-called Minkowski lemma. Apply, for

instance, [SZ96, Th. 7.2] for $n = 3$ and $k = 1$ (so $R(1, 3) = 1$), where we take \mathbb{O} to be the ring of $n \times n$ integer matrices.

3.2. *Proof of Theorem A.* Let M be an almost nonnegatively curved manifold. As usual, denote by $M_n = (M, g_n)$ for $n \in \mathbb{N}$ a sequence of Riemannian metrics on M such that $\sec(M_n) \geq -1/n$ and $\text{diam}(M_n) \leq 1/n$, by \tilde{M} the universal covering of M , and by $\tilde{M}_n \rightarrow M_n$ the universal Riemannian covering of M_n .

After passing to a finite cover of M , by [FY92] we may assume that $\pi_1(M)$ is nilpotent.

Fix $p \in M$, and let $\{\gamma_{i,n}\}$ be a short basis of $\pi_1(M_n, p)$; see Section 2.6. Then, if n is sufficiently large, the short basis $\{\gamma_{i,n}\}$ has at most $k = k(\dim M)$ elements and its elements satisfy $|\gamma_{i,n}|_n \leq 2/n$ for every i . Also, Corollary 3.1.2 implies that given $\varepsilon > 0$, for all large n and every i we have $\|h(\gamma_{i,n})\|_n < 1 + \varepsilon$ and $\|h(\gamma_{i,n}^{-1})\|_n < 1 + \varepsilon$.

Take $N = N(k, m)$ as in Lemma 3.1.3. One can choose $\varepsilon > 0$ so small that if p is a polynomial with integer coefficients whose roots have absolute values lying between $1/(1 + \varepsilon)^N$ and $(1 + \varepsilon)^N$, then all roots of p have absolute values equal to 1. This follows from the fact that the total number of integer polynomials whose roots are all contained in a fixed bounded region is finite.

Set $S_n := \{\gamma_{i,n}\}$. Then for any $\gamma \in S_n^N$ we have

$$\|h(\gamma)\|_n < (1 + \varepsilon)^N \quad \text{and} \quad \|h(\gamma^{-1})\|_n < (1 + \varepsilon)^N.$$

Therefore the absolute values of all eigenvalues lie between $1/(1 + \varepsilon)^N$ and $(1 + \varepsilon)^N$. Since the characteristic polynomial of $h(\gamma)$ has integer coefficients, the absolute values of all the eigenvalues of $h(\gamma)$ are in fact equal to 1.

Apply now Lemma 3.1.3. It follows that the absolute values of all eigenvalues of $h(\gamma)$ are equal to 1 for any $\gamma \in \pi_1(M)$. Then Lemma 3.1.4 implies that after passing to a finite cover M' of M , all eigenvalues of $h(\gamma)$ are equal to 1 for any $\gamma \in \pi_1(M')$. By Engel’s theorem, one can choose an integral basis of $H^*(\tilde{M}, \mathbb{R})$ such that the action of $\pi_1(M)$ on $H^*(\tilde{M}, \mathbb{Z})/\text{tor}$ is given by upper triangular matrices. Therefore, by passing to a finite cover M'' of M' , we can assume that the action of $\pi_1(M'')$ on $H^*(\tilde{M}, \mathbb{Z})$ (and on $H_*(\tilde{M}, \mathbb{Z})$) is nilpotent.

Recall (see, e.g., [HMR75, 2.19]) that a connected CW-complex with nilpotent fundamental group is nilpotent if and only if the action of its fundamental group on the homology of its universal cover is nilpotent. Thus M'' is a nilpotent space, which completes the proof of Theorem A. □

4. C-nilpotency of the fundamental group

4.1. In this section we will prove Theorem B. It will follow from the following somewhat stronger result.

THEOREM 4.1.1. *For any integer m there exist constants $\epsilon(m) > 0$ and $C(m) > 0$ such that the following holds. If M^m is a closed, smooth m -manifold that admits a Riemannian metric g with $\sec(M^m, g) > -\epsilon(m)$ and $\text{diam}(M^m, g) < 1$, then the fundamental group of M^m is $C(m)$ -nilpotent, i.e., $\pi_1(M^m)$ contains a nilpotent subgroup of index $\leq C(m)$.*

Remark 4.1.2. The proofs of Theorems **A** and **C** show that corresponding versions of those results also hold when these theorems are reformulated in a fashion similar to **Theorem 4.1.1**.

By an argument by contradiction, **Theorem 4.1.1** follows from the following statement: Given a sequence of Riemannian m -manifolds (M_n, g_n) with diameters $\text{diam}(M_n, g_n) \leq 1/n$ and sectional curvatures $\sec(g_n) \geq -1/n$, one can find $C \in \mathbb{R}$ such that $\pi_1(M_n)$ is C -nilpotent for all sufficiently large n .

4.2. Algebraic lemmas. Recall that the group of outer automorphisms $\text{Out}(G)$ of a group G is defined as the quotient of its automorphism group $\text{Aut}(G)$ by the subgroup of inner automorphisms $\text{Inn}(G)$.

LEMMA 4.2.1 (a characterization of C -nilpotent groups). *Let*

$$\{1\} = G_\ell \subseteq \cdots \subseteq G_1 \subseteq G_0 = G$$

be a sequence of groups satisfying the following properties: For any i ,

- (i) $G_i \trianglelefteq G$ is normal in G ;
- (ii) *the image of the conjugation homomorphism $h_i : G \rightarrow \text{Out}(G_i/G_{i+1})$ is finite of order at most C_i ;*
- (iii) G_i/G_{i+1} contains an abelian subgroup E_i of index $\leq c_i$.

Then G contains a nilpotent subgroup N of index at most

$$C = C(c_1, \dots, c_\ell, C_1, \dots, C_\ell),$$

where N is of nilpotency class $\leq \ell$.

Proof. First of all, notice that property (i) assures that the objects described in parts (ii) and (iii) of the lemma are well defined.

Set $\Gamma_i := G_i/G_{i+1}$.

Let $H = \bigcap \ker h_i$. Notice that $[G : H] \leq \prod_i C_i$ and that the image of H under the conjugation homomorphism $f_i : G \rightarrow \text{Aut}(\Gamma_i)$ lies in $\text{Inn}(\Gamma_i)$, i.e., $f_i|_H : H \rightarrow \text{Inn}(\Gamma_i)$.

By passing to a subgroup, we can assume $E_i \trianglelefteq \Gamma_i$ is normal of index $\leq C(c_i)$ (we can take $C(c_i) = c_i!$).

By increasing E_i if necessary we can assume E_i contains the center of Γ_i .

Let Z_i be the image of E_i under the projection map $\pi : \Gamma_i \rightarrow \text{Inn}(\Gamma_i)$. Clearly $[\text{Inn}(\Gamma_i) : Z_i] \leq c_i$, and $Z_i \trianglelefteq \text{Inn}(\Gamma_i)$ is normal.

Let $N = H \cap (\bigcap_i f_i^{-1}(Z_i))$ and $N_i = N \cap G_i$. Then

$$[G : N] \leq C = C(c_1, \dots, c_\ell, C_1, \dots, C_\ell)$$

and N satisfies these properties for any i : (i') $N_i \trianglelefteq N$ is normal in N , and (ii') N_i/N_{i+1} is in the center of N/N_{i+1} . That is, N is nilpotent of nilpotency length $\leq \ell$.

Condition (i') is obvious so we only need to check (ii').

To see (ii') observe that by construction the image of the conjugation action $N \rightarrow \text{Aut}(\Gamma_i)$ lies in $\text{Inn}(\Gamma_i)$. In fact, it lies in $\pi(A_i)$ and as such it acts trivially on E_i . Lastly observe that $N_i/N_{i+1} \subset E_i$.

Indeed, by construction, for any $g \in N_i/N_{i+1} \subset \Gamma_i$ there is $a \in E_i$ such that $\pi(g) = \pi(a)$. Therefore, $g = az$ for some z in the center of Γ_i . By our assumption on E_i this means that $g \in E_i$.

Thus N acts trivially on N_i/N_{i+1} , which means that N is nilpotent and G is C -nilpotent. \square

TRIVIAL LEMMA 4.2.2 (a characterization of finite actions). *If S is a finite set of generators of a group G with $S^{-1} = S$, and $h : G \rightarrow H$ is a homomorphism with $|h(S^n)| < n$ for some $n > 0$, then $h(S^n) = h(G)$ and, in particular, $|h(G)| < n$.*

Let now Γ be a group which acts discretely by isometries on an Alexandrov space A with curvature ≥ -1 . Choose a marked point $p \in A$. Assume that $\{\gamma_1, \gamma_2, \dots, \gamma_n\}$ is a finite short basis of Γ at p (see [Section 2.6](#)), and that $\theta \leq |\gamma_i| \leq 1$, where $|\gamma| := |p\gamma(p)|$. Let $\#(R)$ denote the number of elements $\gamma \in \Gamma$ with $|\gamma| \leq R$. The Bishop–Gromov inequality implies that

$$\#(R) \leq v_{-1}^m(R)/v_{-1}^m(\theta),$$

where $m = \dim A$ and $v_{-1}^m(r)$ is the volume of the ball of radius r in the m -dimensional simply connected space form of curvature -1 . Therefore, if $\#\#(L)$ denotes the number of homomorphisms $h : \Gamma \rightarrow \Gamma$ with norm $\leq L$ (i.e., the number of homomorphisms for which $|h(\gamma)| \leq L|\gamma|$ for any $\gamma \in \Gamma$), then

$$(4.2.1) \quad \#\#(L) \leq \#(L)^n \leq \left[\frac{v_{-1}^m(L)}{v_{-1}^m(\theta)} \right]^n.$$

4.3. The blow-up construction. As $n \rightarrow \infty$, the manifolds M_n clearly converge to a point, A_0 .

Set $M_{n,1} := M_n$ and $\vartheta_{n,1} := \text{diam } M_{n,1}$.

Rescale now $M_{n,1}$ by $1/\vartheta_{n,1}$ so that $\text{diam}(M_{n,1}/\vartheta_{n,1}) = 1$. Passing to a subsequence if necessary, one has that the manifolds $(1/\vartheta_{n,1})M_{n,1}$ converge to A_1 , where A_1 is a compact nonnegatively curved Alexandrov space with diameter 1.

Now choose a regular point $\bar{p}_1 \in A_1$, and consider distance coordinates around $\bar{p}_1 \in U_1 \rightarrow \mathbb{R}^{k_1}$, where k_1 is the dimension of A_1 . The distance functions can be lifted to $U_{n,1} \subset (1/\vartheta_{n,1})M_{n,1}$.

Let $M_{n,2}$ be the level set of $U_{n,1} \rightarrow \mathbb{R}^{k_1}$ that corresponds to \bar{p}_1 . Clearly, $M_{n,2}$ is a compact submanifold of codimension k_1 . Set $\vartheta_{n,2} := \text{diam } M_{n,2}$. Passing again to a subsequence if necessary, one has that the sequence $(1/\vartheta_{n,2})M_{n,2}$ converges to some Alexandrov space A_2 . As before, A_2 is a compact nonnegatively curved Alexandrov space with diameter 1. Define $k_2 := k_1 + \text{dim } A_2$. If one now chooses a marked point in $M_{n,2}$, then, as $n \rightarrow \infty$, $M_n/\vartheta_{n,2}$ converges to $A_2 \times \mathbb{R}^{k_1}$, which is of some dimension $k_2 > k_1$.

We repeat this procedure until, at some step, $k_\ell = m$.

As a result one obtains a sequence $\{A_i\}$ of compact nonnegatively curved Alexandrov spaces with diameter 1 that satisfies

$$\dim A_i = k_i - k_{i-1}, \quad \text{so that} \quad \sum_{i=1}^{\ell} \dim A_i = m.$$

We also obtain a sequence of rescaling factors $\vartheta_{n,i} = \text{diam } M_{n,i}$, and a nested sequence of submanifolds

$$\{p_n\} = M_{n,\ell} \subset \dots \subset M_{n,2} \subset M_{n,1} = M_n,$$

which in turn induces a sequence of homomorphisms

$$\{1\} = \pi_1(M_{n,\ell}) \xrightarrow{l} \dots \xrightarrow{l} \pi_1(M_{n,2}) \xrightarrow{l} \pi_1(M_{n,1}) = \pi_1(M_n).$$

Let $G_i := G_i(n) := l^i \pi_1(M_{n,i})$.

For n sufficiently large, the subgroups $G_i(n)$ are those that are generated by elements of norm $\leq 3\vartheta_{n,i}$. Equivalently, if one takes a short basis $\{\gamma_i\}$ of $G(n)$, then G_i is the subgroup generated by all elements γ_i such that $|\gamma_i| \leq 3\vartheta_{n,i}$.

4.4. *Limit fundamental groups of Alexandrov spaces.* We will now define the limit, or L -fundamental groups, of the Alexandrov spaces A_i constructed above. This notion is similar to the notion of the fundamental group of an orbifold. However, we note in advance that the construction of the L -fundamental group does not only depend on the spaces A_i , but also on the chosen rescaled subsequence of M_n . In fact, the following construction shows that the limit fundamental group $\pi_1^L(A_i)$ of A_i is isomorphic to $\pi_1(M_{n,i}, M_{n,i+1})$ for all large enough n . But, unlike $\pi_1(M_{n,i})$, the groups $\pi_1^L(A_i)$ will not depend on n .

The limit fundamental groups of A_i . Consider the converging sequence

$$(M_n/\vartheta_{n,i}, p_n) \xrightarrow{\text{GH}} (A_i \times \mathbb{R}^{k_i-1}, \bar{p}_i \times 0)$$

(here the interesting case is collapsing). Recall that $\bar{p}_i \in A_i$ is a regular point. Fix $\varepsilon > 0$ such that $\text{dist}_{\bar{p}_i}$ on A_i does not have critical values in $(0, 2\varepsilon)$. Take a sequence R_n that converges very slowly to infinity (here we will need $R_n \vartheta_{n,i} / \vartheta_{n,i-1} \rightarrow 0$ and $R_n \rightarrow \infty$).

Consider then a sequence of Riemannian coverings

$$\Pi : (\tilde{B}_n, \tilde{p}_n) \rightarrow (B_{R_n}(p_n), p_n) \quad \text{of } B_{R_n}(p_n) \subset M_n / \vartheta_{n,i}$$

with $\pi_1(\tilde{B}_n, \tilde{p}_n) = \pi_1(B_\varepsilon(p_n), p_n)$, where $B_\varepsilon(p_n) \subset M_n / \vartheta_{n,i}$.

After passing to a subsequence if necessary, the sequence $(\tilde{B}_n, \tilde{p}_n)$ converges to a nonnegatively curved Alexandrov space $\tilde{A}_i \times \mathbb{R}^{k_i-1}$, where the space \tilde{A}_i has the same dimension as A_i . Indeed, by construction it contains an isometric copy of $B_\varepsilon(p_{n,i})$, and therefore

$$\dim \tilde{A}_i + k_{i-1} = \dim \lim_{i \rightarrow \infty} B_\varepsilon(p_{n,i}) = \dim A_i + k_{i-1}.$$

Let us show that for all sufficiently large n ,

$$\iota(\pi_1(M_{n,i+1})) \trianglelefteq \pi_1(M_{n,i}).$$

Assume that $\Pi(\tilde{q}_n) = \tilde{p}_n$ and that $\tilde{q}_n \rightarrow \bar{q}_n \in \tilde{A}_i$. Connect \bar{p}_n and \bar{q}_n by a geodesic, which, by [Pet98], only passes through regular points. Note that in a small neighborhood of this geodesic in M_n we have two copies of $M_{n,i+1}$, near \tilde{p}_n and \tilde{q}_n . Therefore, applying Yamaguchi’s fibration theorem in a small neighborhood of this geodesic, we can construct a diffeomorphism from $M_{n,i+1}$ to itself. This implies that for any loop γ that after lifting connects \tilde{p}_n and \tilde{q}_n , we have $\gamma^{-1} \iota \pi_1(M_{n,i+1}) \gamma \subset \iota \pi_1(M_{n,i+1})$, i.e., $\iota \pi_1(M_{n,i+1}) \triangleleft \pi_1(M_{n,i})$ (for an alternative argument see also [FY92]).

This easily yields that $A_i = \tilde{A}_i / \Gamma_i$, where Γ_i is a group of isometries that acts discretely on \tilde{A}_i . The group Γ_i is denoted by $\pi_1^L(A_i)$ (the *limit* or *L-fundamental group* of A_i). This group is clearly isomorphic to

$$\pi_1(M_{n,i}, M_{n,i+1}) = \pi_1(M_{n,i}) / \iota(\pi_1(M_{n,i+1}))$$

for all sufficiently large n , and the space \tilde{A}_i will be called the *universal covering* of A_i .

Because \tilde{A}_i is nonnegatively curved and $A_i = \tilde{A}_i / \pi_1^L(A_i)$ is compact, Toponogov’s splitting theorem says \tilde{A}_i isometrically splits as $\tilde{A}_i = K_i \times \mathbb{R}^{s_i}$, where K_i is a compact Alexandrov space with $\text{curv} \geq 0$. Since $\pi_1^L(A_i)$ is a group of isometries that acts discretely on \tilde{A}_i , it follows that $\pi_1^L(A_i)$ is a virtually abelian group.

4.5. *Final steps.* Consider now the corresponding series

$$\{1\} = G_\ell(n) \subset \cdots \subset G_1(n) \subset G_0(n) = \pi_1(M_n).$$

The theorem then follows from this lemma:

LEMMA 4.5.1. *For all sufficiently large n , the series*

$$\{1\} = G_\ell(n) \subset \cdots \subset G_1(n) \subset G_0(n)$$

constructed above satisfies the assumptions of Lemma 4.2.1 for numbers C_i and c_i not depending on n .

We first prove the following.

SUBLEMMA 4.5.2. *Each subgroup $G_i(n)$ is normal in $G(n)$.*

Proof. We will show by reverse induction on k that $G_i(n) \trianglelefteq G_k(n)$ for any $k \leq i$. Let us assume that we already know that $G_i(n) \trianglelefteq G_{k+1}(n)$. Since

$$\iota\pi_1(M_{n,k+1}) \trianglelefteq \pi_1(M_{n,k}),$$

we know that $G_{k+1}(n) \trianglelefteq G_k(n)$. Consider the covering

$$\Pi_{k+1} : (\tilde{M}_{n,k+1}, \tilde{p}_{n,k+1}) \rightarrow (M_n, p_n)$$

with covering group $\Gamma_{k+1}(n)$.

Clearly $(\tilde{M}_{n,k+1}, \tilde{p}_{n,k+1}) \xrightarrow{\text{GH}} \mathbb{R}^{s_i}$ for some integer s_i . From Lemma 2.5.1, it follows that for any $a \in G$ with $|a| < 1$ there is a cocos-curve γ in $\tilde{M}_{n,k+1}$ with total time T connecting \tilde{p}_n and $a(\tilde{p}_n)$ in $\tilde{M}_{n,k+1}$. Then clearly $\gamma \sim ga$ for some $g \in G_{k+1}(n)$. Let us denote by $\Phi^T : \tilde{M}_{n,i} \rightarrow \tilde{M}_{n,i}$ the gradient flow corresponding to γ .

Let γ_j be a loop from the short basis of $G_i(n)$. As was mentioned in Section 4.3, if n is large, then $\text{length}(\gamma_j) \leq 3\vartheta_{n,i}$. Let us denote by $\tilde{\gamma}_j$ a lift of γ_j to $\tilde{M}_{n,i}$. Let $\tilde{p}_{n,j} \in \tilde{M}_{n,i}$ be its starting point. Since $[\gamma_j] \in G_i(n)$, we have that $\tilde{\gamma}_j$ is a loop in $\tilde{M}_{n,i}$. Consider then the loop $\gamma'_j = \Pi \circ \Phi^T \circ \tilde{\gamma}_j$. Clearly,

$$[\gamma_j] = a^{-1}g^{-1}[\gamma'_j]ga, \quad \text{or} \quad [\gamma'_j] = ga[\gamma_j]a^{-1}g^{-1}.$$

Proposition 2.3.3 implies that $\text{length}(\gamma'_j) \leq \exp(2T) \text{length}(\gamma_j)$. Therefore, $ga[\gamma_j]a^{-1}g^{-1} \in G_i(n)$ for sufficiently large n , and since $g \in G_i(n) \trianglelefteq G_{k+1}(n)$ it follows that $a[\gamma_j]a^{-1} \in G_i(n)$, i.e., $G_i(n) \trianglelefteq G_k(n)$. \square

Proof of Lemma 4.5.1. The group

$$\pi_1^L(A_i) = \pi_1(M_{n,i}, M_{n,i+1}) = \pi_1(M_{n,i}) / \iota(\pi_1(M_{n,i}))$$

is virtually abelian. Let d_i be the minimal index of an abelian subgroup of $\pi_1^L(A_i)$. The epimorphism $\iota^i : \pi_1(M_{n,i}) \rightarrow G_i$ induces an epimorphism $\pi_1^L(A_i) \rightarrow G_i(n) / G_{i+1}(n)$. Hence $G_i(n) / G_{i+1}(n)$ is d_i -abelian for all large n .

Consider the covering $\Pi_i : \tilde{M}_{n,i} \rightarrow M_n$ with covering group $G_i(n)$, and let $\tilde{p}_{n,i}$ be a preimage of p_n . Clearly

$$(\tilde{M}_{n,i}, \tilde{p}_{n,i}) \xrightarrow{\text{GH}} \mathbb{R}^{s_i} \quad \text{for some integer } s_i.$$

Applying [Lemma 2.5.1](#), it follows that for any $a \in G(n)$ with $|a| < 1$ there is a cocos-curve γ in $\widetilde{M}_{n,i}$ that connects p and $a(p)$. Then clearly $\gamma \sim ga$ for some $g \in G_i(n)$. Let us denote by $\Phi^T : \widetilde{M}_{n,i} \rightarrow \widetilde{M}_{n,i}$ the gradient flow corresponding to γ .

Let $b \in G_i(n)$ and β be a loop representing b . Let us denote by $\tilde{\beta}$ a lift of β to $\widetilde{M}_{n,i}$. Let $\tilde{p}_{n,i} \in \widetilde{M}_{n,i}$ be its starting point. Since $[\beta] \in G_i(n)$, we have that $\tilde{\beta}$ is a loop in $\widetilde{M}_{n,i}$.

Consider now the loop $\beta' = \Pi \circ \Phi^T \circ \tilde{\beta}$. Clearly,

$$b = [\beta] = a^{-1}g^{-1}[\beta']ga, \quad \text{or} \quad [\beta'] = gaba^{-1}g^{-1}.$$

[Proposition 2.3.3](#) then implies that $\text{length}(\beta') \leq \exp(2T) \text{length}(\beta)$. Therefore, if

$$h_a : G_i(n)/G_{i+1}(n) \rightarrow G_i(n)/G_{i+1}(n)$$

is induced by the conjugation $b \rightarrow aba^{-1}$, then for any $a \in G(n)$ there is a $g \in G_i(n)$ such that $|h_{ga}| \leq \exp(2T)$.

Let now δ_i be the minimal norm of the elements of $\pi_1^L(A_i)$, where $\pi_1^L(A_i)$ acts on \tilde{A}_i . Then [\(4.2.1\)](#) implies that the image of the action of $G(n)$ by conjugation in $\text{Out}(G_i(n)/G_{i+1}(n))$ is C_i -finite, where C_i depends only on c_i, T , and δ_i . \square

4.6. *Remark on nonfree actions.* [Theorem 4.1.1](#) can be reformulated:

There exists a constant $\epsilon(m) > 0$ such that if N^m is a Riemannian manifold that admits a free discrete isometric action by a group Γ such that

$$\text{sec}(N) > -\epsilon(m) \quad \text{and} \quad \text{diam}(N/\Gamma) < 1,$$

then Γ is $C(m)$ -nilpotent.

B. Wilking pointed out to us that one can actually easily remove from this reformulation of [Theorem 4.1.1](#) the assumption that the Γ action be free.

COROLLARY 4.6.1. *There exists a constant $\epsilon(m) > 0$ such that if N^m is a Riemannian manifold that admits a discrete isometric action by a group Γ such that $\text{sec}(N) > -\epsilon(m)$ and $\text{diam}(N/\Gamma) < 1$, then Γ is $C(m)$ -nilpotent.*

Proof. Let $\epsilon = \epsilon(m)$ be as provided by [Theorem 4.1.1](#), and suppose N satisfies the assumptions of the corollary for this ϵ . Let F be the frame bundle of N . Then the action of Γ on N lifts to a free isometric action on F . As was observed in [\[FY92\]](#), using Cheeger’s rescaling trick, F can be equipped with a Γ -invariant metric satisfying $\text{sec}(F) > -\epsilon(m)$ and $\text{diam}(F/\Gamma) < 1$. Since the induced action of Γ on F is free, the claim now follows from [Theorem 4.1.1](#). \square

5. Proof of the fibration theorem

5.1. Let M be an almost nonnegatively curved manifold. Let us denote by $M_n = (M, g_n)$ a sequence of Riemannian metrics on M such that $\sec(M_n) \geq -1/n$ and $\text{diam}(M_n) \leq 1/n$.

Let us denote by \tilde{M} the universal cover of M and by $\tilde{M}_n \rightarrow M_n$ the universal Riemannian covering of M_n (i.e., \tilde{M} equipped with the pull back of the metric g_n on M).

By [FY92], passing to a finite cover we may assume that $\Gamma = \pi_1(M)$ is a nilpotent group without torsion. Hence, to prove the topological part of [Theorem C](#), it is enough to show the following:

THEOREM 5.1.1. *Suppose M is a closed almost nonnegatively curved m -manifold such that $\Gamma = \pi_1(M)$ is a nilpotent group without torsion. Then M is the total space of a fiber bundle*

$$F \rightarrow M \rightarrow N,$$

where the base N is a nilmanifold and the fiber F is simply connected.

The assumption on Γ implies that we can fix a series

$$\Gamma = \Gamma_0 \triangleright \Gamma_1 \triangleright \Gamma_2 \triangleright \cdots \triangleright \Gamma_\ell = \{1\}$$

such that Γ_i is normal in Γ and $\Gamma_i/\Gamma_{i+1} \cong \mathbb{Z}$.

Let us first give an informal proof.

5.2. *An informal proof of [Theorem 5.1.1](#).* We use induction to construct the bundles

$$F_i \longrightarrow M \xrightarrow{f_i} N_i,$$

where each N_i is a nilmanifold with $\pi_1(N_i) = \Gamma/\Gamma_i$ and $\pi_1(F_i) \cong \Gamma_i$. Since the base of induction is trivial, we are only interested in the induction step.

Fix $p \in N_i$, and let $F_i(p)$ be the fiber over p . For any sufficiently large n , choose a subgroup $G_i = G_i(n)$ such that $\Gamma_i \triangleleft G_i \triangleleft \Gamma_{i+1}$ and $[\Gamma_i : G_i]$ is finite, but large enough so that the cover $\bar{F}_i(p)$ of $F_i(p)$ corresponding to G_i is Hausdorff close to a unit circle S^1 .

Construct now a bundle map $\varphi_p : \bar{F}_i(p) \rightarrow S^1$ by lifting distance functions from S^1 . (This can be done by a slight generalization of a construction in [FY92] and [BGP92].) Let $\omega_p = d\varphi_p$.

Then ω_p is a closed integral nondegenerate one-form on $F_i(p)$. Since deck transformations are isometries, after averaging by \mathbb{Z}_a , where $a = [\Gamma_i : G_i]$, we can assume that ω_p is \mathbb{Z}_a -invariant. Thus ω_p descends to a form on $F_i(p)$, which when integrated gives a bundle map $F_i(p)$ onto a small S^1 .

Note that although this bundle is defined only up to rotations of S^1 , its fibers are well defined.

Since Γ_{i+1} is normal in Γ , the choice of the covering $\bar{F}_i(p)$ of $F_i(p)$ is unambiguous for all $p \in N_i$. By using a partition of unity on N_i we can glue the forms ω_p into a global 1-form on M which satisfies the properties that

- (a) $\omega|_{F(p)}$ is closed and integral for any p ;
- (b) $\omega|_{F(p)}$ is nondegenerate.

Integrating ω over the various $F(p)$, we construct a continuous family of bundles $F_p \rightarrow S^1$. The level sets partition each $F(p)$ and hence the whole M into fibers of a fiber bundle, whose quotient space is then a circle bundle N_{i+1} over N_i . $\approx \square$

This gives a good idea of the proof. However, to make it precise some extra work has to be done. In particular, one has to be careful with the construction of ω . To make this construction possible we have to keep track of how $F(p)$ was obtained. Namely, we have to use that the fiber $F(p)$ was obtained by a construction as in Yamaguchi’s fibration theorem; see [Yam91] or [BGP92]. This makes the induction proof quite technical.

We now proceed with the rigorous proof of [Theorem 5.1.1](#).

5.3. Proof of [Theorem 5.1.1](#). Let us denote by $\tilde{M}_{n,i}$ the Riemannian covering of M_n with respect to Γ_i .

For any choice of marked points p_n we have

$$(\tilde{M}_{n,i}, p_n, \pi_1(M)) \xrightarrow{\text{GH}} (\mathbb{R}^i, 0, \mathbb{R}^i)$$

in equivariant Gromov-Hausdorff convergence, where \mathbb{R}^i acts on itself by translations. Indeed, the limit space must be a nonnegatively curved simply connected Alexandrov space, and since $\text{diam } M_n \rightarrow 0$, it possesses a transitive group action by a nilpotent group. Then Euclidean space, acting as a group of translations, is here the only choice, and it is easy to see that the dimension of the limit must be equal to i .

Therefore $(\tilde{M}_n, p_n, \pi_1(M)) \xrightarrow{\text{GH}} (\mathbb{R}^\ell, 0, \mathbb{R}^\ell)$, and we may also assume that

$$(\tilde{M}_n, p_n, \Gamma_i) \xrightarrow{\text{GH}} (\mathbb{R}^\ell, 0, \mathbb{R}^{\ell-i}) \quad \text{for each } i,$$

where $\mathbb{R}^{\ell-i}$ is the coordinate subspace of \mathbb{R}^ℓ that corresponds to the first $\ell - i$ elements of the standard basis.

Now, let us give a technical definition:

If R is a Riemannian manifold, let us denote by $\widetilde{\text{dist}}_p$ the average of a distance function over a small ball around p . This enables us to work with the C^1 function $\widetilde{\text{dist}}_p$ instead of the Lipschitz function dist_p .

Definition 5.3.1. Suppose $R_n \xrightarrow{\text{GH}} R$ is a sequence of Riemannian m -manifolds with curvature $\geq \kappa$ that Gromov-Hausdorff converges to a Riemannian m' -manifold R , where $m' \leq m$. A sequence of forms ω_n on R_n is said to ε -approximate a form ω on R , if

- (i) for any point $p \in R$ there is a neighborhood $U \ni p$ that admits a distance chart $f : U \rightarrow \mathbb{R}^{m'}$,

$$f(x) = (\text{dist}_{a_1}(x), \text{dist}_{a_2}(x), \dots, \text{dist}_{a_{m'}}(x))$$

that is a smooth regular map, and

- (ii) smooth lifts of f to $U_n \subset R_n$ give, for n large enough, regular maps

$$f_n(x) = (\widetilde{\text{dist}}_{a_{1,n}}(x), \widetilde{\text{dist}}_{a_{2,n}}(x), \dots, \widetilde{\text{dist}}_{a_{m',n}}(x))$$

with $a_{i,n} \in M_n$, $a_{i,n} \rightarrow a_n$ such that

$$|(f_n \circ f^{-1})^*(\omega) - \omega_n|_{C^0} < \varepsilon \quad \text{for all sufficiently large } n.$$

Theorem 5.1.1 now easily follows from the following lemma:

LEMMA 5.3.2. *Given $\varepsilon > 0$, there is a sequence of one-forms*

$$\{\omega_{1,n}, \omega_{2,n}, \dots, \omega_{k,n}\}$$

on \widetilde{M}_n with the following properties:

- (i) For each i , $\omega_{i,n}$ is a $\pi_1(M)$ -invariant form on \widetilde{M}_n .
- (ii) The forms $\omega_{i,n}$ ε -approximate the coordinate forms dx_i on \mathbb{R}^k . In particular, the forms $\{\omega_{i,n}\}$ are nowhere zero and almost orthonormal at each point.
- (iii) If $\omega_{j,n}(X) = \omega_{j,n}(Y) = 0$ for all $j < i$, then $d\omega_{i,n}(X, Y) = 0$. In particular, for each i and all sufficiently large n , the distribution corresponding to the system of equations

$$\omega_{j,n}(X) = 0 \quad \text{for all } j \leq i$$

defines on \widetilde{M}_n a foliation $\mathcal{F}_{i,n}$.

- (iv) If $\widetilde{F}_{i,n}(x) \subset \widetilde{M}_n$ denotes the fiber of the foliation $\mathcal{F}_{i,n}$ through the point $x \in \widetilde{M}_n$, then each $\widetilde{F}_{i,n}(x)$ is Γ_i -invariant, i.e., for any $\gamma \in \Gamma_i$ one has that $\widetilde{F}_{i,n}(x) = \widetilde{F}_{i,n}(\gamma x)$. Moreover, the action of Γ_i on $\widetilde{F}_{i,n}(x)$ is cocompact for each i . In particular, $\mathcal{F}_{i,n}$ induces on M_n the structure of a fiber bundle.

Proof. We will construct these forms by induction. Assume that we have already constructed one-forms $\omega_1, \omega_2, \dots, \omega_{i-1}$ with all the required properties. They give a $\pi_1(M)$ -invariant fibration of \widetilde{M}_n by submanifolds $\widetilde{F}_{i-1,n}(x)$ through each point $x \in \widetilde{M}_n$, with tangent spaces defined by the equations $\omega_j(X) = 0$ for $j = 1, \dots, i - 1$.

Denote by $\theta : \mathbb{R} \rightarrow [0, 1]$ a smooth monotone function that is equal to 1 before 0 and to 0 after 1. Choose numbers $\delta_n > 0$ slowly converging to 0, and let $\Theta_{i,n} : \widetilde{M}_n \rightarrow \mathbb{R}_+$ be the function defined by

$$\Theta_{i,n}(x) = \min_{y \in F_{i-1,n}(x)} \theta(|p_n y| / \delta_n).$$

Clearly $\Theta_{i,n}$ is a continuous Γ_{i-1} -invariant function, which is constant on each $F_{i-1,n}(x)$. Moreover, for large n , $\Theta_{i,n}$ has support in some $C_i\delta_n$ -neighborhood of $F_{i-1,n}(p_n)$, and is equal to 1 in some $c_i\delta_n$ -neighborhood of $F_{i-1,n}(p_n)$.

Now let $\varphi : \mathbb{R} \rightarrow [0, 1]$ be a smooth nondecreasing function that is 0 before $1/2$ and 1 after $3/2$. Consider the form

$$\omega'_{i,n} = \Theta_{i,n} \cdot d(\varphi \circ \widetilde{\text{dist}}_{\Gamma_i a_{i,n}}),$$

where $a_{i,n} \in \widetilde{M}_n$ is a sequence of points converging to $-e_i \in \mathbb{R}^\ell$, and $\widetilde{\text{dist}}_{\Gamma_i a_{i,n}}$ is the average of $\text{dist}_{\Gamma_i x}$ for x in a small ball around $a_{i,n}$. The support of ω'_i has two components, one that contains p_n (notice here that $p_n \rightarrow 0 \in \mathbb{R}^\ell$), and another that does not. (It follows from the construction that the limit of $F_{i-1,n}(p_n)$ is a coordinate plane in \mathbb{R}^ℓ .)

Set $\omega''_{i,n} := \omega'_{i,n}$ on the component of p_n , and let this form be 0 otherwise. Clearly, $\omega''_{i,n}$ is then a continuous Γ_i -invariant form whose restriction to $\widetilde{F}_{i-1,n}(x)$ is exact. Moreover, each level set of its integral over $\widetilde{F}_{i-1,n}(x)$ is Γ_i -invariant.

By construction, the form $\omega''/|\omega''|$ is now (in the sense of [Definition 5.3.1](#)) close to dx_i at the points where $|\omega''| \neq 0$. Take

$$\omega_{i,n} = c \sum_{\gamma \in \Gamma/\Gamma_i} \gamma \omega',$$

where the coefficient c is chosen in such a way that $|\omega_{i,n}(p_n)| = 1$. As δ_n is a sequence slowly converging to zero, we may assume that $\text{diam}(M_n)/\delta_n \rightarrow 0$. Therefore, $\omega_{i,n}$ is the form we need. □

Notice that the proof actually shows that the fibers in [Theorem 5.1.1](#) are almost nonnegatively curved manifolds in the generalized sense with $k = \ell$. Therefore, the proof of [Theorem C](#) is complete. □

6. Open questions

We would like to conclude this work by posing a number of related open questions.

6.1. *Is the torsion contained in the center?* As was noted earlier, [Theorem B](#) is new even for manifolds of nonnegative curvature. For such manifolds it is known that their fundamental groups are almost abelian, and Fukaya and Yamaguchi conjectured the following:

CONJECTURE 6.1.1 (see [\[FY92\]](#)). *The fundamental group of a nonnegatively curved m -manifold is $C(m)$ -abelian.*

In this regard we would like to pose the following two conjectures:

MAIN CONJECTURE 6.1.2. *There is a $C = C(m)$ such that if M^m is almost nonnegatively curved, then there is a nilpotent subgroup $N \subset \pi_1(M)$ of index $\leq C$ whose torsion is contained in its center (or, at least, whose torsion is commutative).*

CONJECTURE 6.1.3. *If M^m is almost nonnegatively curved, then the action of $\pi_1(M)$ on $\pi_2(M)$ is almost trivial (or maybe even $C(m)$ -trivial), i.e., there exists a finite index subgroup of $\pi_1(M)$ (or, respectively, a subgroup of index $\leq C(m)$) that acts trivially on $\pi_2(M)$.*

Main Conjecture 6.1.2 implies in particular that the fundamental groups of closed positively curved m -manifolds are $C(m)$ -abelian.

In fact, as was pointed out to us by B. Wilking, the truth of **Main Conjecture 6.1.2** would also imply a positive answer to **Conjecture 6.1.1**. Indeed, if $\sec(M) \geq 0$, then the universal cover \tilde{M} of M is isometric to the product $\mathbb{R}^n \times K$, where K is a compact Riemannian manifold and the $\pi_1(M)$ action on $\mathbb{R}^n \times K$ is diagonal. It follows from [Wil00, Cor. 6.3] that one can deform the metric on M so that its universal cover is still isometric to $\mathbb{R}^n \times K$ and the induced action on K is finite. By passing, as in the proof of **Corollary 4.6.1**, to the induced action on the frame bundle of K , one reduces the statement to **Main Conjecture 6.1.2**.

We tried to prove these conjectures by studying successive blow-ups of the collapsing sequence M_n as done in **Section 4.3**.

We can prove **Conjectures 6.1.2** and **6.1.3** in the case where all spaces A_i that appear in the construction in **Section 4.3** are closed Riemannian manifolds. Moreover, we believe we have an argument to prove it if all the A_i are Alexandrov spaces without boundary.

It seems that if we would have just a slightly better understanding of collapsing to a ray, then we could prove the conjectures. Here is the simplest related question we cannot solve:

Question 6.1.4. Let $M_n = (S^2 \times \mathbb{R}^2, g_n)$ be a sequence of complete Riemannian manifolds with $\sec(M_n) \geq -\varepsilon_n$, where $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$. Assume $(M_n, p_n) \xrightarrow{\text{GH}} (\mathbb{R}_+, 0)$ for a sequence of points $p_n \in M_n$. Let $q_n \in M_n$ be a sequence of points such that $|p_n q_n| = 1$ and such that there is a sequence of rescalings $\lambda_n \rightarrow \infty$ such that $(\lambda_n M_n, q_n) \xrightarrow{\text{GH}} S^2 \times S^1 \times \mathbb{R}$, where the latter space is equipped with the product of the canonical metrics.

- (i) Can it happen that $(\lambda_n M_n, p_n) \xrightarrow{\text{GH}} (\mathbb{R}_+, 0)$?
- (ii) Is the Gromov-Hausdorff limit of $(\lambda_n M_n, p_n)$ at least 3-dimensional?
- (iii) What are the possible limits of $(\lambda_n M_n, p_n)$?

In the case where all A_i are Riemannian manifolds, crucial to the proof is the following topological statement:

THEOREM 6.1.5. *Given manifolds F_1, F_2, \dots, F_n such that each F_i is either S^1 or is simply connected, if E is the total space of a tower of fiber bundles*

$$E = E_n \xrightarrow{F_n} E_{n-1} \xrightarrow{F_{n-1}} \dots \xrightarrow{F_1} E_0 = \{\text{point}\}$$

and each of the bundles $E_k \xrightarrow{F_k} E_{k-1}$ are homotopically trivial over the 1-skeleton, then $\pi_1(E)$ contains a nilpotent subgroup N such that

$$[\pi_1(E) : N] \leq C(F_1, F_2, \dots, F_n) \quad \text{and} \quad \text{Tor}(N) \subset Z(N).$$

Our current proof of this theorem is surprisingly nontrivial,¹ and finding an easier proof will probably help to prove our conjecture in full generality.

Conjectures 6.1.1 and 6.1.2 are also related to the following conjecture of Rong (cf. [Ron96a], [Ron96b]):

CONJECTURE 6.1.6 (Rong). *Positively curved m -manifolds have $C(m)$ -cyclic fundamental groups.*

This conjecture has been proved by Rong [Ron96a] under the additional assumption of a uniform upper curvature bound. We also believe that if one could carry out the above program for proving **Main Conjecture 6.1.2**, one would have a good shot at handling Rong’s conjecture as well.

6.2. The simply connected case. So far we have only discussed manifolds with nontrivial fundamental groups. However, some of our arguments also work in a more general setting. We hope that it might be possible to use them to obtain new restrictions on simply connected almost nonnegatively curved manifolds as well as on collapsing with a lower curvature bound.

Let us indicate one possible approach to do so.

Let us denote by $\mathcal{M}(F)$ the space of self homotopy equivalences of a manifold F . Assume now that F is simply connected and that $\tilde{f} : S^k \times F \rightarrow F$ is a map such that $\tilde{f}_u : F \rightarrow F$ is homotopic to the identity for some (and therefore any) $u \in S^k$. Then \tilde{f} represents an element $\alpha = [\tilde{f}] \in \pi_k(\mathcal{M}(F))$. Let g be a Riemannian metric on F . Define

$$\text{dil}_g(\tilde{f}) = \max_{u \in S^k} \text{dil}_g(\tilde{f}_u),$$

where $\text{dil}_g(\tilde{f}_u)$ is the optimal Lipschitz constant of \tilde{f}_u with respect to g . For any $\alpha \in \pi_k(\mathcal{M}(F))$ define

$$\text{dil}_g(\alpha) = \inf_{[h]=\alpha} \text{dil}_g(h).$$

Finally, define

$$\text{DIL}(\alpha) = \inf_g \text{dil}_g(\alpha)$$

¹The note <http://www.math.psu.edu/petrinin/papers/almpos/TinZ.pdf> gives a sketch of the proof of **Theorem 6.1.5**.

over all Riemannian metrics on F and

$$\text{DIL}_+(\alpha) = \inf_g \text{dil}_g(\alpha)$$

over all Riemannian metrics on F with $\text{diam} \leq 1$ and $\text{sec} \geq -1$.

Clearly, both $\text{DIL}(\alpha)$ and $\text{DIL}_+(\alpha)$ are homotopy invariants of α .

Now suppose that $M_n \xrightarrow{\text{GH}} S^{k+1}$ is a sequence of Riemannian manifolds collapsing to a round sphere with $\text{sec}(M_n) \geq k$. By Yamaguchi's fibration theorem, M_n is a fiber bundle over S^{k+1} with almost nonnegatively curved fiber F_n . This bundle is classified by an element α of $\pi_k(\text{Aut}(F_n))$, and by using our gradient flow technique we can estimate $\text{DIL}_+(\alpha)$ (and hence $\text{DIL}(\alpha)$) from above.

Therefore, if one could find examples of a simply connected F and an α with arbitrary big $\text{DIL}_+(\alpha)$, one would obtain new restrictions on collapsing to a sphere with curvature bounded from below, and probably more restrictions for the topological type of manifolds with lower curvature and upper diameter bounds in general. In fact, F need not be simply connected as long as the total space of the bundle $F \rightarrow M \rightarrow S^{k+1}$ is.

While we believe that finding examples with arbitrary large $\text{DIL}(\alpha)$ is very difficult (and might even be impossible), we have several candidates to produce large $\text{DIL}_+(\alpha)$.

On the other hand, the problem of finding α with $\text{DIL}(\alpha) > 1$ seems quite interesting in its own right and might have other applications unrelated to collapsing.

Let us next describe some possible sources of examples with $\text{DIL}_+(\alpha) > 1$:

Example 6.2.1. Obviously, if $\text{dil}_g(h) = 1$, then h_u is a homotopy of isometries of (F, g) . Let G be the isometry group of F . Then G can be viewed as a subset of $\mathcal{M}(F)$. Therefore, if $[h] \neq 0$ in $\pi_k(\mathcal{M}(F))$, then $[h_u] \neq 0$ in $\pi_k(G)$. Now G is a compact Lie group, in particular, $\pi_2(G) = 0$ (and even more generally $\pi_{2n}(G)$ is finite). On the other hand, there are spaces F such that the space $\mathcal{M}(F)$ might have nontrivial second homotopy; for example, one can take $F = \text{SU}(6)/\text{SU}(3) \times \text{SU}(3)$, for which the canonical metric on F has nonnegative curvature, and it follows from [TO97, Ch. 5, Ex. 4.14] that $\pi_2(\mathcal{M}(F)) \otimes \mathbb{Q}$ is nontrivial. Therefore, there is an $\alpha \in \pi_2(\mathcal{M}(F))$ such that $\text{dil}_g(\alpha) > 1$ for any metric g on F ; we believe it should be true that $\text{DIL}_+(\alpha) > 1$. Still, it might happen that $\text{DIL}(\alpha) = 1$.

Another possible source of such manifolds is provided by the following example due to D. Sullivan.

Example 6.2.2. Let N^7 be the total space of an S^3 bundle over S^4 with zero Euler class and nontrivial p_1 . Clearly N^7 is rationally equivalent to $S^4 \times S^3$. In particular, its minimal model has no nontrivial derivations of degree -1 . Therefore, by [Sul77, 13.3], there exists a diffeomorphism $f : N \rightarrow N$ that is homotopic to the identity but whose obstruction to being diffeotopic to the identity is a nonzero

element of $H^3(N, \mathbb{Z}) \cong \mathbb{Z}$. Let M^8 be the mapping cylinder of f . Clearly M is homotopy equivalent to $N \times S^1$, and hence it is spin with signature zero. On the other hand, by construction, $p_1^2(M) \neq 0$. Since the signature of M is zero we must necessarily have that $p_2(M) \neq 0$, and hence $\hat{A}(M) \neq 0$. In particular, by the Atiyah-Hirzebruch theorem, M does not admit an S^1 action, and hence the corresponding element $\alpha \in \pi_1(\mathcal{M}(M))$ has $\text{dil}_g(\alpha) > 1$ for any metric g on M .

Remark 6.2.3. As mentioned in the introduction, it is known that a spin manifold X of almost nonnegative Ricci curvature has $\hat{A}(X) \leq 2^{\dim(X)/2}$; see [Gro82, p. 41] and [Gal83]. Clearly, a finite cover of the manifold M constructed above violates this restriction and therefore M does not admit almost nonnegative Ricci curvature. However, it could possibly be almost nonnegatively curved in the generalized sense.

6.3. *Further questions.* Recall that a simply connected space C is called *rationally elliptic* if it is homotopy equivalent to a finite CW-complex and

$$\dim \pi_*(C, \mathbb{Q}) < \infty.$$

A conjecture of Grove and Halperin [GH82] says that simply connected nonnegatively curved manifolds are rationally elliptic. This conjecture was extended by Grove to include almost nonnegatively curved manifolds [Gro02]. Later, Totaro has proposed the following definition of rationally elliptic spaces, which covers manifolds with infinite fundamental groups:

A connected topological space X is *rationally elliptic* if it is homotopy equivalent to a finite CW-complex, it has a finite covering that is a nilpotent space, and its universal cover is rationally elliptic in the ordinary sense.

With this definition one can extend Grove's conjecture to manifolds that are not simply connected, as follows:

CONJECTURE 6.3.1. *Any almost nonnegatively curved manifold in the generalized sense is rationally elliptic.*

Theorem A reduces this conjecture to the simply connected case, which is undoubtedly the most difficult part of the problem.

It has been shown in [PP06] that if M is a nilpotent closed manifold that admits a Riemannian metric with zero topological entropy, then its universal cover \tilde{M} is rationally elliptic. Coupled with **Theorem A**, this means that to prove **Conjecture 6.3.1** it would be sufficient to show that a manifold with almost nonnegative curvature in the generalized sense admits a metric with zero topological entropy. However, we think that this might be wrong in general.

As pointed out in the discussion in **Section 1** before **Theorem C**, it already follows from Yamaguchi's fibration theorem and [FY92] that a finite cover of an

almost nonnegatively curved manifold maps onto a nilmanifold with homotopy fiber a simply connected closed manifold. While this is formally weaker than the statement of [Theorem C](#), it would be interesting to have an answer to the following, purely topological, question:

Question 6.3.2. Let $M \xrightarrow{f} N$ be a map from a closed manifold M to a nilmanifold N such that the homotopy fiber of f is a simply connected closed manifold. Is it true that after passing to a finite cover, the map f becomes homotopic to a fiber bundle projection?

Question 6.3.3. Is it true that manifolds that are almost nonnegatively curved in the generalized sense are almost nonnegatively curved?

In view of [Theorems A](#) and [B](#) it is also reasonable to pose the following question:

Question 6.3.4. Is it true that almost nonnegatively curved m -manifolds M^m are $C(m)$ -nilpotent spaces?

It is clear from the proof of [Theorems A](#) and [B](#) that this is true if the universal cover of M^m has torsion free integral cohomology.

In view of [Theorem B](#) it is also natural to raise the following question:

Question 6.3.5. Can one give an explicit bound on $C(m)$ in [Theorem B](#)?

Acknowledgments

We would like to express our thanks to the following people for helpful conversations during the preparation of this work: I. Belegradek, V. Gorbunov, E. Formanek, I. Kapovitch, A. Lytchak, R. Matveyev, D. Robinson, D. Sullivan, B. Wilking, and Yu. Zarkhin.

We are also especially grateful to Yunhui Wu for noticing a mistake in the proof of [Theorem 4.1.1](#) in the original version of this article.

References

- [And92] M. T. ANDERSON, [Hausdorff perturbations of Ricci-flat manifolds and the splitting theorem](#), *Duke Math. J.* **68** (1992), 67–82. [MR 94k:53054](#) [Zbl 0767.53029](#)
- [BGP92] Y. BURAGO, M. GROMOV, and G. PEREL'MAN, [A. D. Aleksandrov spaces with curvatures bounded below](#), *Uspekhi Mat. Nauk* **47** (1992), 3–51, 222, In Russian; translated in *Russian Math. Surveys* **47** (1992), 1–58. [MR 93m:53035](#)
- [CFG92] J. CHEEGER, K. FUKAYA, and M. GROMOV, [Nilpotent structures and invariant metrics on collapsed manifolds](#), *J. Amer. Math. Soc.* **5** (1992), 327–372. [MR 93a:53036](#) [Zbl 0758.53022](#)
- [FY92] K. FUKAYA and T. YAMAGUCHI, [The fundamental groups of almost non-negatively curved manifolds](#), *Ann. of Math.* **136** (1992), 253–333. [MR 93h:53041](#) [Zbl 0770.53028](#)

- [Gal83] S. GALLOT, Inégalités isopérimétriques, courbure de Ricci et invariants géométriques, II, *C. R. Acad. Sci. Paris Sér. I Math.* **296** (1983), 365–368. MR 85c:53069b Zbl 0535.53035
- [Gro78] M. GROMOV, *Almost flat manifolds*, *J. Differential Geom.* **13** (1978), 231–241. MR 80h:53041 Zbl 0432.53020
- [Gro80] ———, Synthetic geometry in Riemannian manifolds, in *Proc. Internat. Congress of Mathematicians* (Helsinki, 1978) (O. LEHTO, ed.), Acad. Sci. Fennica, 1980, pp. 415–419. MR 81g:53029 Zbl 0427.53018
- [Gro81] ———, *Curvature, diameter and Betti numbers*, *Comment. Math. Helv.* **56** (1981), 179–195. MR 82k:53062 Zbl 0467.53021
- [Gro82] ———, *Volume and bounded cohomology*, *Inst. Hautes Études Sci. Publ. Math.* (1982), 5–99. MR 84h:53053 Zbl 0516.53046
- [Gro02] K. GROVE, Geometry of, and via, symmetries, in *Conformal, Riemannian and Lagrangian Geometry* (Knoxville, TN, 2000), *Univ. Lecture Ser.* **27**, Amer. Math. Soc., Providence, RI, 2002, pp. 31–53. MR 2003g:53046 Zbl 1019.53001
- [GH82] K. GROVE and S. HALPERIN, *Contributions of rational homotopy theory to global problems in geometry*, *Inst. Hautes Études Sci. Publ. Math.* (1982), 171–177. MR 84b:58030 Zbl 0508.55013
- [HMR75] P. HILTON, G. MISLIN, and J. ROITBERG, *Localization of Nilpotent Groups and Spaces*, *Mathematics Studies* **15**, North-Holland, Amsterdam, 1975. MR 57 #17635 Zbl 0323.55016
- [LeB01] C. LEBRUN, *Ricci curvature, minimal volumes, and Seiberg-Witten theory*, *Invent. Math.* **145** (2001), 279–316. MR 2002h:53061 Zbl 0999.53027
- [Lyt05] A. LYTCHAK, *Open map theorem for metric spaces*, *Algebra i Analiz* **17** (2005), 139–159, In Russian; translated in *St. Petersburg Math. J.* **17** (2006), 477–491. MR 2006e:53075 Zbl 1152.53033
- [Oht09] S.-I. OHTA, *Gradient flows on Wasserstein spaces over compact Alexandrov spaces*, *Amer. J. Math.* **131** (2009), 475–516. MR 2503990 Zbl 1169.53053
- [PP06] G. P. PATERNAIN and J. PETEAN, *Zero entropy and bounded topology*, *Comment. Math. Helv.* **81** (2006), 287–304. MR 2007g:37023 Zbl 1093.53090
- [Per91] G. PERELMAN, *Alexandrov spaces with curvatures bounded from below*, II, preprint, 1991.
- [PP96] G. PERELMAN and A. PETRUNIN, *Quasigeodesics and gradient curves in Alexandrov spaces*, preprint, 1996. Available at http://www.math.psu.edu/petrunin/papers/qg_ams.pdf
- [Pet95] A. PETRUNIN, *Quasigeodesics in multidimensional Alexandrov spaces*, thesis, University of Illinois at Urbana Champaign, 1995.
- [Pet98] ———, *Parallel transportation for Alexandrov space with curvature bounded below*, *Geom. Funct. Anal.* **8** (1998), 123–148. MR 98j:53048 Zbl 0903.53045
- [Pet07] ———, *Semiconcave functions in Alexandrov’s geometry*, to appear in *Surveys in Comparison Geometry*, 2007. Zbl 1166.53001
- [Ron96a] X. RONG, *The almost cyclicity of the fundamental groups of positively curved manifolds*, *Invent. Math.* **126** (1996), 47–64. MR 97g:53045 Zbl 0859.53022
- [Ron96b] ———, *On the fundamental groups of manifolds of positive sectional curvature*, *Ann. of Math.* **143** (1996), 397–411. MR 97a:53067 Zbl 0974.53029
- [Sav07] G. SAVARÉ, *Gradient flows and diffusion semigroups in metric spaces under lower curvature bounds*, *C. R. Math. Acad. Sci. Paris* **345** (2007), 151–154. MR 2008j:49113 Zbl 1125.53064

- [Sch89] R. M. SCHOEN, Variational theory for the total scalar curvature functional for Riemannian metrics and related topics, in *Topics in Calculus of Variations* (Montecatini Terme, 1987) (M. GIAQUINTA, ed.), *Lecture Notes in Math.* **1365**, Springer-Verlag, New York, 1989, pp. 120–154. [MR 90g:58023](#) [Zbl 0702.49038](#)
- [ST04] L. J. SCHWACHHÖFER and W. TUSCHMANN, Metrics of positive Ricci curvature on quotient spaces, *Math. Ann.* **330** (2004), 59–91. [MR 2005e:53053](#) [Zbl 1062.53027](#)
- [Sha78] V. A. SHARAFUTDINOV, The Pogorelov-Klingenberg theorem for manifolds homeomorphic to \mathbb{R}^n , *Sib. Math. J.* **18** (1978), 649–657. [Zbl 0411.53031](#)
- [SZ96] A. SILVERBERG and Y. G. ZARHIN, Variations on a theme of Minkowski and Serre, *J. Pure Appl. Algebra* **111** (1996), 285–302. [MR 97c:11065](#) [Zbl 0885.14006](#)
- [Sul77] D. SULLIVAN, Infinitesimal computations in topology, *Inst. Hautes Études Sci. Publ. Math.* (1977), 269–331 (1978). [MR 58 #31119](#) [Zbl 0374.57002](#)
- [TO97] A. TRALLE and J. OPREA, *Symplectic Manifolds with no Kähler Structure*, *Lecture Notes in Mathematics* **1661**, Springer-Verlag, New York, 1997. [MR 98k:53038](#) [Zbl 0891.53001](#)
- [Wil00] B. WILKING, On fundamental groups of manifolds of nonnegative curvature, *Differential Geom. Appl.* **13** (2000), 129–165. [MR 2001g:53076](#) [Zbl 0993.53018](#)
- [Yam91] T. YAMAGUCHI, Collapsing and pinching under a lower curvature bound, *Ann. of Math.* **133** (1991), 317–357. [MR 92b:53067](#) [Zbl 0737.53041](#)

(Received June 14, 2005)

(Revised March 26, 2007)

E-mail address: vtk@math.toronto.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, ONTARIO M5S 2E4,
CANADA

E-mail address: petrunin@math.psu.edu

DEPARTMENT OF MATHEMATICS, PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK,
STATE COLLEGE, PA 16802, UNITED STATES

E-mail address: tusch@math.uni-kiel.de

MATHEMATISCHES SEMINAR, CHRISTIAN-ALBRECHTS-UNIVERSITÄT ZU KIEL,
LUDEWIG-MEYN-STRASSE 4-8, D-24118 KIEL, GERMANY