

# Monodromy and irreducibility of leaves

By CHING-LI CHAI and FRANS OORT

## 1. Introduction

Let  $p$  be a prime number, fixed throughout this article. Let  $n, d$  be positive integers with  $\gcd(n, dp) = 1$ . Denote by  $\mathcal{A}_{g,d,n}$  the moduli space over  $\mathbb{F}_p$  of  $g$ -dimensional abelian varieties with a polarization of degree  $d^2$  and with a symplectic level- $n$  structure in characteristic  $p$ . The moduli space  $\mathcal{A}_{g,d,n}$  has a stratification

$$\{\mathcal{W}_\xi^0(\mathcal{A}_{g,d,n}) \mid \xi\}$$

which is indexed by symmetric Newton polygons  $\xi$  of height  $2g$ . Every stratum  $\mathcal{W}_\xi^0(\mathcal{A}_{g,d,n})$  is a reduced locally closed subscheme of  $\mathcal{A}_{g,d,n}$  whose geometric points correspond to  $g$ -dimensional principally polarized abelian varieties with Newton polygon equal to  $\xi$ ; see [Section 2.5](#). The smallest among the Newton polygon strata is the supersingular stratum  $\mathcal{W}_\sigma^0(\mathcal{A}_{g,d,n})$ , corresponding to supersingular polarized abelian varieties, i.e. those with Newton polygon isoclinic of slope  $1/2$ , or equivalently, geometrically isogenous to a product of supersingular elliptic curves.

In this paper we prove three global properties of nonsupersingular Newton polygon strata and leaves in  $\mathcal{A}_{g,d,n}$ .

**THEOREM A.** *For  $\xi \neq \sigma$ , the Newton stratum  $\mathcal{W}_\xi^0 := \mathcal{W}_\xi^0(\mathcal{A}_{g,1,n})$  is geometrically irreducible. See [Theorem 3.1](#).*

**THEOREM B.** *For any geometric point  $x \in \mathcal{A}_{g,d,n}(k)$  not contained in the supersingular stratum  $\mathcal{W}_\sigma^0(\mathcal{A}_{g,d,n}) = \mathcal{W}_\sigma(\mathcal{A}_{g,d,n})$ , the leaf  $\mathcal{C}(x)$  in  $\mathcal{A}_{g,d,n}$  is irreducible. See [Theorem 4.1](#).*

**THEOREM C.** *For any geometric point  $x \in \mathcal{A}_{g,d,n}(k)$  not contained in the supersingular stratum  $\mathcal{W}_\sigma(\mathcal{A}_{g,d,n})$ , the  $p$ -adic monodromy for the leaf  $\mathcal{C}(x)$  is maximal. See [Theorem 5.6](#).*

The definition of the leaf  $\mathcal{C}(x)$  in  $\mathcal{A}_{g,d,n}$  passing through  $x$  is recalled in [Section 2.2](#). Let  $\bar{\eta}_{\mathcal{C}(x)}$  be a geometric generic point of the leaf  $\mathcal{C}(x)$ . In

**Theorem C** the  $p$ -adic monodromy for  $\mathcal{C}(x)$  refers to the action of the Galois group of the function field of  $\mathcal{C}(x)$  on the polarized  $p$ -divisible group  $(A_{\bar{\eta}_{\mathcal{C}(x)}}[p^\infty], \lambda_{\eta_{\mathcal{C}(x)}}[p^\infty])$  attached to the polarized abelian variety  $(A_{\bar{\eta}_{\mathcal{C}(x)}}, \lambda_{\eta_{\mathcal{C}(x)}})$  over the generic point of  $\mathcal{C}(x)$ ; see [Definition 5.1](#) and [Section 5.4](#) for the definition. The assertion in [Theorem C](#) is that the image of the monodromy action, regarded as a subgroup of the group  $\text{Aut}(A_{\bar{\eta}_{\mathcal{C}(x)}}[p^\infty], \lambda_{\eta_{\mathcal{C}(x)}}[p^\infty])$  of automorphisms of the polarized  $p$ -divisible group  $(A_{\bar{\eta}_{\mathcal{C}(x)}}[p^\infty], \lambda_{\eta_{\mathcal{C}(x)}}[p^\infty])$ , is equal to the whole group  $\text{Aut}(A_{\bar{\eta}_{\mathcal{C}(x)}}[p^\infty], \lambda_{\eta_{\mathcal{C}(x)}}[p^\infty])$ . Note that the number of irreducible components of  $\mathcal{W}_\sigma(\mathcal{A}_{g,d,n})$  is a suitable class number, which is large when  $p$  is large. Similarly, every leaf in the supersingular locus is 0-dimensional whose cardinality is a class number. So the nonsupersingularity assumptions in [Theorems A, B, and C](#) are necessary.

Note that the conclusion of [Theorem A](#) is not correct for  $\mathcal{W}_\xi^0(\mathcal{A}_{g,d,n})$  for arbitrary  $d$ . However in [Theorems B and C](#) we need not restrict to the principally polarized case. Note that *all*  $p$ -adic invariants are constant on  $\mathcal{C}(x)$ , in the sense that the geometric isomorphism class of polarized  $p$ -divisible groups  $(A_y, \lambda_y)[p^\infty]$  attached to points  $y$  on  $\mathcal{C}(x)$  is fixed.

Questions on global topological properties of subvarieties of moduli spaces in characteristic  $p$  are often challenging. Subvarieties such as  $\mathcal{W}_\xi^0(\mathcal{A}_{g,1,n}) \subset \mathcal{A}_{g,1,n}$  and  $\mathcal{C}(x) \subset \mathcal{A}_{g,d,n}$  are stable under all prime-to- $pd$  Hecke correspondences. One knows from [\[4\]](#) that such a subvariety  $Z \subset \mathcal{A}_{g,d,n}$  is irreducible if  $Z$  is not contained in the supersingular locus and the prime-to- $pd$  Hecke correspondences operate transitively on the set of geometrically irreducible components of  $Z$ ; this is the first “reduction step” in the proof of [A and B](#). Although this reduction step offers a general strategy for proving irreducibility statements of subvarieties of modular varieties of PEL-type, some special properties are used in each example that has been worked out.

The reader will find another reduction method that allows us to deduce [B](#) from the irreducibility statement in [A](#). That method is based on the notion of *hypersymmetric points* introduced in [\[3\]](#) and [\[6\]](#). These points in  $\mathcal{A}_{g,d,n}$  correspond to abelian varieties whose endomorphism rings are “as large as allowed by their geometric  $p$ -divisible groups”, or equivalently, “maximal under the given slope constraint”; for hypersymmetric abelian varieties, see [Section 2.7](#). The idea of the proof of [B](#) is to use the irreducibility in [A](#) to produce hypersymmetric points on every irreducible components of the leaf  $\mathcal{C}(x)$ , and then connect these points by suitable prime-to- $p$  Hecke correspondences with the help from group theory.

The existence of hypersymmetric points on the leaf  $\mathcal{C}(x)$  allows us to apply the third method in [\[5\]](#) and deduce the maximality statement in [C](#). For technical reasons it is more convenient to deduce [Theorems B and C](#) by proving

first the case when  $d$  is a power of  $p$ . To make the logical structure of the proof as clear as possible, we first prove [Theorem 5.6](#) in the special case when the  $p$ -divisible group  $A_x[p^\infty]$  is *completely slope divisible*; see [Theorem 5.9](#). The general case is proved by the same method in the proof of [5.9](#), but with some additional technical components; see [5.13](#). In [5.14](#) we indicate how [Theorem 5.6](#) can be deduced from its special case, [Theorem 5.9](#).

Methods in this article are developed for the Hecke orbit conjecture, first formulated in [\[25\]](#) in the case of Siegel modular varieties; see also [\[3\]](#) for a sketch of a proof. Implicit in the proof sketched in [\[3\]](#) is a proof of the irreducibility statement in [Theorem B](#). That proof depends on the irreducibility of central leaves in Hilbert modular varieties proved by C.-F. Yu in [\[34\]](#), as well as the ‘‘Hilbert trick’’ and the trick of ‘‘splitting at supersingular points’’. The older proof of [Theorem B](#) is more involved in terms of the complexity of the logical structure than the proof here.

*Acknowledgment.* We thank J-P. Serre for a helpful discussion on weak approximation.

## 2. Prerequisites

2.1. *Some notation to be used below.* Let  $p$  be a prime number, fixed in this article. All abelian varieties and  $p$ -divisible groups are defined over a field or a base scheme of characteristic  $p$ . We write  $\mathbb{F}$  for an algebraic closure of  $\mathbb{F}_p$ . For an abelian variety  $A$  we write  $X = A[p^\infty]$  for its  $p$ -divisible group.

We write  $k$  and  $\Omega$  for an algebraically closed field of characteristic  $p$ . All base fields and base schemes considered will be in characteristic  $p$ .

For a group scheme  $G$  over a field  $K$  we write  $a(G)$  for the dimension of the  $L$ -vector space  $\text{Hom}(\alpha_{p,L}, G_L)$ , where  $L \supset K$  is a perfect field. Note that  $\text{End}(\alpha_{p,L}) = L$ ; hence the right- $L$ -module  $\text{Hom}(\alpha_{p,L}, G_L)$  is a vector space over  $L$ . Suppose  $L \subset L'$  are both perfect fields containing  $K$ ; then the natural map

$$\text{Hom}(\alpha_{p,L}, G_L) \otimes_L L' \xrightarrow{\sim} \text{Hom}(\alpha_{p,L'}, G_{L'})$$

is an isomorphism. In particular, the number  $a(G)$  is independent of the choice of the perfect field  $L$  containing  $K$ .

For a scheme  $W$  over a field  $K$  we write  $\Pi_0(W)$  for the set of geometrically irreducible components of  $W$ ; i.e., we choose  $K \subset k$ , and we consider the set of irreducible components of  $W \times_{\text{Spec}(K)} \text{Spec}(k)$ .

We write  $\sigma = \sigma_g$  for the supersingular Newton polygon (i.e. all slopes are equal to  $1/2$ ).

We write  $\mathcal{A}_{g,d,n}$  for the moduli space of polarized abelian varieties in characteristic  $p$ , with degree of polarization equal to  $d^2$ , and with symplectic level- $n$  structure; here  $g, d, n \in \mathbb{Z}_{>0}$  and  $\text{gcd}(n, dp) = 1$ . This moduli space should be

denoted by  $\mathcal{A}_{g,d,n} \times_{\text{Spec}(\mathbb{Z})} \text{Spec}(\mathbb{F}_p)$ , but we abbreviate notation. In most cases we assume  $n \geq 3$ . We write  $\mathcal{A}_{g,d}$  as an abbreviation for  $\mathcal{A}_{g,d,1}$ . When working over a field  $K \supset \mathbb{F}_p$  we still write  $\mathcal{A}_{g,d}$  instead of  $\mathcal{A}_{g,d} \times_{\text{Spec}(\mathbb{F}_p)} \text{Spec}(K)$ .

We write  $\ell$  for a prime number different from  $p$ , sometimes also not dividing  $d$ .

2.2. *Central leaves* (main reference: [25]). Work over an algebraically closed field  $k \supset \mathbb{F}_p$ . Recall that for each geometric point  $x = [(A, \lambda, \iota_A)] \in \mathcal{A}_{g,d,n}(k)$ , there is a locally closed, reduced  $k$ -subscheme  $\mathcal{C}(x)$  of  $\mathcal{A}_{g,d,n}$ , called the *central leaf* passing through  $x$ , characterized by the property that

$$\begin{aligned} \mathcal{C}(x)(\Omega) &= \left\{ (B, \mu, \iota_B) \in \mathcal{A}_{g,d,n}(\Omega) \mid (B, \mu)[p^\infty] \right. \\ &\quad \left. \cong (A, \lambda)[p^\infty] \otimes \Omega, \text{type}^{(p)}(\mu) = \text{type}^{(p)}(\lambda) \right\} \end{aligned}$$

for every algebraically closed field  $\Omega \supset k$ ; see [25]. Here  $\iota_A$  is a symplectic level- $n$  structure of the polarized abelian variety  $(A, \lambda)$ , where  $(A, \lambda)[p^\infty]$  is the quasi-polarized  $p$ -divisible group attached to  $(A, \lambda)$ , and  $\text{type}^{(p)}(\lambda)$  is the sequence of elementary divisors of the alternating pairing on  $\prod_{\ell \neq p} H_1(A, \mathbb{Z}_\ell)$  induced by the polarization  $\lambda$ ; similarly for  $(B, \mu, \iota_B)$ . For  $n \geq 3$  we have that  $\mathcal{C}(x)$  is smooth over  $k$ . The sequence  $\text{type}^{(p)}(\lambda)$  of elementary divisors has the form  $(d_1, \dots, d_g)$ , where  $d_1, \dots, d_g$  are positive integers such that  $d_1 \mid d_2 \mid \dots \mid d_g$  and  $d = \prod_{i=1}^g d_i$ . It is known that the leaf  $\mathcal{C}(x)$  is closed in the open Newton stratum  $W_\xi^0(\mathcal{A}_{g,1,n})$  containing  $\mathcal{C}(x)$ ; see [25, Th. 3.3]. See Section 2.5 for the notation and properties of Newton strata. We will shorten ‘‘central leaf’’ to ‘‘leaf’’ if confusion (with ‘‘isogeny leaf’’) is unlikely.

The above definition of the leaf  $\mathcal{C}(x)$  in  $\mathcal{A}_{g,d,n}$  is slightly different from the definition in [25] because of the extra condition that we require here that the discrete invariant  $\text{type}^{(p)}(\mu)$  is constant on  $\mathcal{C}(x)$ . This condition enables us to announce Theorems B and C in the present short form.

2.3. *Newton polygons*. Newton polygons with slopes between 0 and 1 will be denoted by a symbol like  $\zeta$  or  $\xi$ . By  $\zeta = \sum_i (m_i, n_i)$ , we intend to say that the (lower convex) Newton polygon starting from the origin of the plane, such that the multiplicity of a slope  $\nu$  in  $\zeta$  is equal to  $\sum_{\nu=m_i/(m_i+n_i)} (m_i + n_i)$ . In the above, it is understood that  $m_i, n_i \in \mathbb{Z}_{\geq 0}$  and  $\text{gcd}(m_i, n_i) = 1$  for all  $i$ ; ‘‘lower convex’’ means that  $\zeta$  is the graph of a convex piecewise linear function defined on  $[0, h]$ , where  $h = \sum_i (m_i + n_i)$ . A Newton polygon  $\zeta = \sum_i (m_i, n_i)$  can be specified by its *slope sequence*  $(\nu_1, \dots, \nu_h)$ , where  $0 \leq \nu_1 \leq \dots \leq \nu_h \leq 1$ , and for every  $\nu \in \mathbb{Q} \cap [0, 1]$ , the multiplicity of  $\nu$  in the slope sequence above, defined as  $\text{Card}\{j \mid \mu_j = \nu, 1 \leq j \leq h\}$ , is equal to the multiplicity of  $\nu$  in  $\zeta$ .

A Newton polygon  $\xi$  is said to be *symmetric* if the multiplicity of  $\nu$  in  $\xi$  is equal to the multiplicity of  $1 - \nu$  in  $\xi$  for every slope  $\nu$ . We say that

two Newton polygons are *disjoint* if they have no slopes in common. Every symmetric Newton polygon  $\xi$  can be written as a sum of disjoint symmetric Newton polygons, each having at most two slopes.

Every symmetric Newton polygon  $\xi$  can be written in a unique way in the following *standard form*

$$\xi = (\gamma_0 \cdot ((1, 0) + (0, 1))) + \left( \sum_{1 \leq i \leq t} \gamma_i \cdot ((m_i, n_i) + (n_i, m_i)) \right) + (\gamma_{t+1} \cdot (1, 1)),$$

where  $f = \gamma_0 \in \mathbb{Z}_{\geq 0}$ ,  $t \in \mathbb{Z}_{\geq 0}$ ,  $\gamma_1, \dots, \gamma_t, \gamma_{t+1} \in \mathbb{Z}_{>0}$  and  $m_i > n_i \geq 0$  for  $1 \leq i \leq t$ , and  $(m_i, n_i) \neq (m_j, n_j)$  if  $1 \leq i \neq j \leq t$ . The coefficients  $\gamma_0, \gamma_1, \dots, \gamma_{t+1}$  are called the *multiplicities of the simple parts* of  $\xi$ . Define  $g(\xi) = \gamma_0 + \sum_{1 \leq i \leq t} \gamma_i \cdot (m_i + n_i) + \gamma_{t+1}$ .

2.4. According to the Dieudonné-Manin classification of  $p$ -divisible groups over an algebraically closed field (see [15, p. 35]), every  $p$ -divisible group  $X$  over an algebraically closed field  $k \supset \mathbb{F}_p$  is isogenous to a direct product of isoclinic  $p$ -divisible groups  $G_{m,n}$ , with  $m, n \in \mathbb{Z}_{\geq 0}$  and  $\gcd(m, n) = 1$ , with  $\dim(G_{m,n}) = m$ ; in this case  $G_{m,n}$  has height  $m + n$  and is isoclinic of slope  $m/(m + n)$ . The Newton polygon of a  $p$ -divisible group  $X$  isogenous to  $\prod_i G_{m_i, n_i}$  is

$$\sum_i (m_i, n_i) =: \mathcal{N}(X).$$

For an abelian variety  $A$  over a field  $K \supset \mathbb{F}_p$ , the Newton polygon attached to  $A[p^\infty]$  is a symmetric Newton polygon  $\mathcal{N}(A)$ , and it can be written in standard form as above. Then we have  $\dim(A) = g(\xi)$ . We hope there will be no confusion caused by the formal sum expressing  $\xi$  and the summation as in the formula for  $g$ .

Working over a field  $K \supset \mathbb{F}_p$ , if no confusion can arise, we write  $G_{m,n}$  instead of  $G_{m,n} \otimes_{\mathbb{F}_p} K$ . The result is that there is a bijection between the set of  $k$ -isogeny classes of  $p$ -divisible groups over  $k$  and the set of Newton polygons:

THEOREM (Dieudonné and Manin; see [15, p. 35, ‘‘Classification theorem’’]).

$$\{X\} / \sim_k \xrightarrow{\sim} \{\text{Newton polygon}\}.$$

2.5. *Newton polygon strata.* The set of Newton polygons is partially ordered; we write  $\gamma \prec \beta$  if no point of  $\gamma$  is below  $\beta$ :

$$\gamma \prec \beta \iff \gamma \text{ is ‘‘above’’ } \beta.$$

If  $(\nu_1, \dots, \nu_h)$  and  $(\mu_1, \dots, \mu_{h'})$  are the slope sequence of  $\gamma$  and  $\beta$  respectively, with  $0 \leq \nu_1 \leq \dots \leq \nu_h \leq 1$  and  $0 \leq \mu_1 \leq \dots \leq \mu_{h'} \leq 1$ , then  $\gamma \prec \beta$  if and only if

$$h = h', \sum_{j=1}^m \nu_j \geq \sum_{j=1}^m \mu_j \quad \text{for } m = 1, \dots, h - 1 \quad \text{and} \quad \sum_{j=1}^h \nu_j = \sum_{j=1}^h \mu_j.$$

An abelian variety  $A$  is isogenous with its dual  $A^t$ . Using the duality theorem (see [20, 19.1]), we conclude that  $X \sim X^t$ ; hence  $\mathcal{N}(A) =: \xi$  is symmetric.

For a symmetric Newton polygon  $\xi$  we write:

$$\mathcal{W}_\xi(\mathcal{A}_{g,d,n}) := \{[(A, \lambda)] \mid \mathcal{N}(A) \prec \xi\},$$

$$\mathcal{W}_\xi^0(\mathcal{A}_{g,d,n}) := \{[(A, \lambda)] \mid \mathcal{N}(A) = \xi\}.$$

Grothendieck-Katz:

$$\mathcal{W}_\xi(\mathcal{A}_{g,d,n}) \subset \mathcal{A}_{g,d,n} \text{ is closed}$$

(see [12, pp. 149–159], [13, Th. 2.3.1, p. 143]); hence

$$\mathcal{W}_\xi^0(\mathcal{A}_{g,d,n}) \subset \mathcal{A}_{g,d,n} \text{ is locally closed.}$$

These are called the *Newton polygon strata*. We write

$$W_\xi = \mathcal{W}_\xi(\mathcal{A}_{g,1}), \quad W_\xi^0 = \mathcal{W}_\xi^0(\mathcal{A}_{g,1}).$$

2.6. THEOREM (Oort; see [25, Th. 3.3]). *Every leaf*

$$\mathcal{C}(y) \subset \mathcal{W}_\xi^0(\mathcal{A}_{g,d,n})$$

*is a closed subset of the open Newton polygon stratum  $\mathcal{W}_\xi^0(\mathcal{A}_{g,d,n})$ .*

2.7. *Hypersymmetric abelian varieties* (main reference: [6]).

*Definition.* Let  $B$  be an abelian variety over a field  $K \supset \mathbb{F}_p$ . We say that  $B$  is *hypersymmetric* if one (or all) of the following equivalent conditions are satisfied:

(1) The natural map

$$\text{End} \left( B \times_{\text{Spec}(K)} \text{Spec}(\overline{K}) \right) \otimes_{\mathbb{Z}} \mathbb{Z}_p \xrightarrow{\sim} \text{End} \left( B[p^\infty] \times_{\text{Spec}(K)} \text{Spec}(\overline{K}) \right)$$

is an isomorphism.

(2) The natural map

$$\text{End} \left( B \times_{\text{Spec}(K)} \text{Spec}(\overline{K}) \right) \otimes_{\mathbb{Z}} \mathbb{Q}_p \xrightarrow{\sim} \text{End} \left( B[p^\infty] \times_{\text{Spec}(K)} \text{Spec}(\overline{K}) \right) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$$

is an isomorphism.

(3) There exist an abelian variety  $A$  defined over  $\mathbb{F}$  and an isogeny

$$B \times_{\text{Spec}(K)} \text{Spec}(\overline{K}) \sim A \times_{\text{Spec}(\mathbb{F})} \text{Spec}(\overline{K}),$$

such that the natural map

$$\text{End}(A) \otimes_{\mathbb{Z}} \mathbb{Z}_p \xrightarrow{\sim} \text{End}(A[p^\infty])$$

is an isomorphism.

*Remark.* Tate proved that for an abelian variety  $A$  defined over a finite field  $\mathbb{F}_q$ , the natural homomorphism

$$\text{End}(A) \otimes_{\mathbb{Z}} \mathbb{Z}_p \xrightarrow{\sim} \text{End}(A[p^\infty])$$

is an isomorphism; see [33, Th. 1, p. 60]. This shows that  $\text{End}(A) \otimes_{\mathbb{Z}} \mathbb{Z}_p$  is identified with the  $\text{Gal}(\mathbb{F}/\mathbb{F}_q)$ -invariant endomorphisms of

$$\text{End}\left(A[p^\infty] \times_{\text{Spec}(\mathbb{F}_q)} \text{Spec}(\mathbb{F})\right).$$

Suppose that

$$\text{End}(A) \xrightarrow{\sim} \text{End}\left(A \times_{\text{Spec}(\mathbb{F}_q)} \text{Spec}(\mathbb{F})\right).$$

Then  $A$  is hypersymmetric if and only if every element of the ring of endomorphisms of the  $p$ -divisible group  $A[p^\infty] \times_{\text{Spec}(\mathbb{F}_q)} \text{Spec}(\mathbb{F})$  is invariant under every element of  $\text{Gal}(\mathbb{F}/\mathbb{F}_q)$ . From this we see that there are “many” abelian varieties over a finite field which are not hypersymmetric. Also we see that for a hypersymmetric abelian variety this Galois action is “in diagonal form” for every isoclinic part of  $A[p^\infty]$ . This can be made precise as follows:

*Let  $K$  be a finite field and let  $B$  be an abelian variety over  $K$ . Then  $B$  is hypersymmetric if and only if there exists a positive integer  $n$  such that the  $n$ -th power of the Frobenius  $\pi_B$  of  $B$  lies in the center of  $\text{End}^0\left(B[p^\infty] \times_{\text{Spec}(K)} \text{Spec}(\mathbb{F})\right)$ . In other words, any two eigenvalues of the action of  $\pi_B$  on the Dieudonné module of  $B[p^\infty] \times_{\text{Spec}(K)} \text{Spec}(\mathbb{F})$  which have the same  $p$ -adic absolute value, differ by a root of unity.*

*Remark.* In [6] we showed that for any symmetric Newton polygon  $\xi$  there exists a hypersymmetric point in the Newton stratum  $W_\xi^0$ . This statement does *not* hold for general modular varieties of PEL-type, where a point is said to be hypersymmetric if the underlying abelian variety with prescribed endomorphism ring is hypersymmetric in the sense of [6, Def. 6.4]. Therefore the methods in this article do not apply to all leaves in a modular variety of PEL-type. Explicit conditions for the existence of hypersymmetric points on a Newton stratum can be found in [36].

2.8. *Hecke orbits.* Suppose we are given a field  $K$ , and a polarized abelian variety  $(A, \lambda)$  over  $K$ . We define the *Hecke orbit* of the moduli point  $x := [(A, \lambda)]$  to be the set of points  $y = [(B, \mu)]$  over some field  $L$  such that there exist a field  $\Omega$  containing  $K$  and  $L$ , and a quasi-isogeny  $\varphi : A \rightarrow B$  such that

$$\varphi^*(\mu) = \lambda.$$

*Notation.* We write  $y \in \mathcal{H}(x)$ . The set  $\mathcal{H}(x)$  is called the *Hecke orbit* of  $x$ .

Recall that a *quasi-isogeny*  $\varphi : A \rightarrow B$  is an equivalence class of diagrams  $A \xleftarrow{u} C \xrightarrow{v} B$ , where  $u$  and  $v$  are isogenies between abelian varieties. We

write  $\varphi = v \circ u^{-1}$  when  $\varphi$  is represented by such a diagram. By definition, two diagrams

$$A \xleftarrow{u} C \xrightarrow{v} B \quad \text{and} \quad A \xleftarrow{u'} C' \xrightarrow{v'} B$$

are equivalent if there exists an abelian variety  $D$  and isogenies  $w: D \rightarrow C$  and  $w': D \rightarrow C'$  such that  $u \circ w = u' \circ w'$  and  $v \circ w = v' \circ w'$ . Let  $N$  be a non-zero integer and let  $\ell$  be a prime number different from  $p$ . A quasi-isogeny  $\varphi$  is said to be relatively prime to  $N$  (resp.  $\ell$ -primary) if  $\varphi$  can be represented as  $v \circ u^{-1}$  such that the degrees of the isogenies  $u$  and  $v$  are both relatively prime to  $N$  (resp. the degrees of  $u$  and  $v$  are powers of  $\ell$ ).

*Hecke-prime-to- $N$ -orbits.* Let  $N \in \mathbb{Z}_{>0}$ . If in the definition of the Hecke orbit of a moduli point  $x = [(A, \lambda)]$  the isogeny  $\varphi$  is relatively prime to  $N$ , we say that  $[(B, \mu)] = y$  is in the Hecke-prime-to- $N$ -orbit of  $x$ .

*Notation.*  $y \in \mathcal{H}^{(N)}(x)$ .

*Hecke- $\ell$ -orbits.* Fix a prime number  $\ell$  different from  $p$ . We say that  $[(B, \mu)] = y$  is in the Hecke- $\ell$ -orbit of  $x$  if the quasi-isogeny  $\varphi$  is  $\ell$ -primary in the previous definition.

*Notation.*  $y \in \mathcal{H}_\ell(x)$ .

*Remark.* We have given the definition of the so-called  $\mathrm{Sp}_{2g}(\mathbb{A}_f)$ -Hecke orbit, because, for every positive integer multiple  $N$  of  $pd$ , the prime-to- $N$  Hecke orbit  $\mathcal{H}^{(N)}(x)$  of a geometric point  $x \in \mathcal{A}_{g,d,n}(k)$  can be described as follows. Consider the tower of moduli spaces  $(\mathcal{A}_{g,d,m})$ , where  $m$  runs through all multiples of  $n$  which are prime to  $pd$ . The group  $\mathrm{Sp}_{2g}(\mathbb{A}_f^{(N)})$  operates on the tower, where  $\mathbb{A}_f^{(N)} = \prod'_{\mathrm{gcd}(\ell, dp)=1} \mathbb{Q}_\ell$ . Let  $\tilde{x}$  be a  $k$ -point of the projective limit  $\varprojlim \mathcal{A}_{g,d,m}$  lying above  $x$ . Then  $\mathcal{H}^{(N)}(x)$  is the image in  $\mathcal{A}_{g,d,n}$  of  $\mathrm{Sp}_{2g}(\mathbb{A}_f^{(N)}) \cdot \tilde{x}$ , the orbit of  $\tilde{x}$  under the action of  $\mathrm{Sp}_{2g}(\mathbb{A}_f^{(N)})$ . See [7, §1] for a further discussion.

*Hecke stable.* We say a set  $T \subset \mathcal{A}_g$  is  $\mathcal{H}_\ell$ -stable if for every  $x := [(A, \lambda)] \in T$  we have  $\mathcal{H}_\ell(x) \subset T$ , and analogously for  $\mathcal{H}^{(p)}$ -stable. Note that Newton polygon strata are Hecke-stable, and EO-strata and central leaves are  $\mathcal{H}^{(p)}$ -stable.

*The  $\mathcal{H}_\alpha$ -orbit.* By considering Hecke correspondences where all isogenies involved are geometrically successive extensions of  $\alpha_p$ , we define the Hecke- $\alpha$ -orbit of a moduli point. If in the notation just used the kernels of  $\varphi_1$  and of  $\varphi_2$  are successive extensions of  $\alpha_p$  over an algebraically closed field, we write  $[(B, \mu)] = y \in \mathcal{H}_\alpha(x)$ .

We define  $I(y) \subset \mathcal{A}_g$ , a maximal  $\mathcal{H}_\alpha$ -set, by taking the union of all irreducible components of the  $\mathcal{H}_\alpha$ -orbit of  $y$  containing  $y$ . This is called the *isogeny leaf* passing through  $y = [(B, \mu)]$ . We give this the induced reduced scheme structure.



The completion of  $\mathcal{A}_{g,d,n}$  over  $k$  along an isogeny leaf, in the sense of [25], is étale locally isomorphic to the reduction modulo  $p$  of a suitable Rapoport-Zink space; see [31, Th. 6.2.3] for a more precise statement. This implies that the formal completion  $I(y)^{/y}$  of an isogeny leaf at a  $k$ -point  $y \in \mathcal{A}_{g,d,n}(k)$  is isomorphic to the formal completion at a suitable  $k$ -point of the reduction modulo  $p$  of a Rapoport-Zink space with reduced structure.

2.9. THEOREM (Oort) (“central leaves and isogeny leaves almost give a product structure on an irreducible component of a Newton polygon stratum”); see [25, Th. 5.3]). *Work over an algebraically closed field  $k$ . Choose a symmetric Newton polygon  $\xi$ , an irreducible component  $W$  of  $\mathcal{W}_\xi^0(\mathcal{A}_g)$ , an irreducible component  $C$  of a central leaf, and an irreducible component  $I$  of an isogeny leaf. There exist finite surjective morphisms  $f: T \rightarrow C$ ,  $g: J \rightarrow I$ , a finite surjective morphism*

$$\Phi: T \times J \rightarrow W,$$

and a polarization-preserving quasi-isogeny

$$\Theta: f^*(A, \lambda) \rightarrow \Phi^*(B, \mu)$$

such that for every  $u \in J$ ,

$$\Phi(T \times \{u\}) \text{ is a component of a central leaf}$$

and for every  $t \in T$ ,

$$\Phi(\{t\} \times J) \text{ is a component of an isogeny leaf.}$$

2.10. *Minimal  $p$ -divisible groups* (main reference: [26]; also see [8]). For a pair  $(m, n)$  of coprime nonnegative integers we define  $H_{m,n}$  to be a  $p$ -divisible group of dimension  $m$ , of height  $m + n$ , of slope  $m/(m + n)$  such that over  $k$  the endomorphism ring  $\text{End}(H_{m,n})$  is the maximal order in  $\text{End}^0(H_{m,n}) := \text{End}(H_{m,n}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$ . This defines this group uniquely up to isomorphism over an algebraically closed field; the group can be constructed over  $\mathbb{F}_p$ . For  $\zeta = \sum_i (m_i, n_i)$  we write  $H(\zeta) = \prod_i H_{m_i, n_i}$ . This  $p$ -divisible group is defined over  $\mathbb{F}_p$ ; if no confusion can arise we write  $H(\zeta)$  over any field  $K \supset \mathbb{F}_p$ , meaning  $H(\zeta) \otimes_{\mathbb{F}_p} K$ .

*Definition.* We say a  $p$ -divisible group  $X$  is *minimal* if there exists an isomorphism  $X_\Omega \cong H(\zeta)$ .

2.11. THEOREM (Oort; see [26, Th. 1.2]). *Work over an algebraically closed field  $k$ . Let  $X$  be a  $p$ -divisible group, and let  $\zeta$  be some Newton polygon. Suppose there exists an isomorphism  $X[p] \cong H(\zeta)[p]$ . Then there exists an isomorphism  $X \cong H(\zeta)$ .*

Suppose  $[(A, \lambda)] = x$ , where  $A[p^\infty]$  is *minimal* and  $\lambda$  is a *principal* polarization. In this case the central leaf  $\mathcal{C}(x)$  is denoted by  $\mathcal{Z}_\xi = \mathcal{C}(x)$ , where

$\xi = \mathcal{N}(A[p^\infty])$ , and it is called the *central stream* inside  $W_\xi^0$ . Note that a principal quasi-polarization on the minimal  $p$ -divisible  $A[p^\infty]$  is unique up to pre-composing the given principal quasi-polarization  $\lambda[p^\infty]$  with an element of  $\text{Aut}(A[p^\infty])$  which is fixed under the Rosati involution  $*_{\lambda[p^\infty]}$  attached  $\lambda[p^\infty]$ ; see [25, 3.7].

*Results on  $\ell$  and prime-to- $p$  monodromy and irreducibility* (main reference: [4]).

2.12. THEOREM (Chai). *Work over an algebraically closed field  $k$ . Suppose  $n \in \mathbb{Z}_{\geq 3}$  is relatively prime with  $pd$ . Let  $Z \subset \mathcal{A}_{g,d,n}$  be a locally closed nonsingular subset not contained in the supersingular locus. Suppose  $Z$  is  $\mathcal{H}_{\text{Sp}}^{(pd)}$ -stable. Assume that  $\mathcal{H}_{\text{Sp}}^{(pd)}$  acts transitively on  $\Pi_0(Z)$ . Then  $Z$  is irreducible and the prime-to- $pd$  monodromy representation on  $Z$  equals*

$$\text{Sp}_{2g}(\widehat{\mathbb{Z}}^{(pd)})(n) := \{\gamma \in \text{Sp}_{2g}(\widehat{\mathbb{Z}}^{(pd)}) \mid \gamma \equiv 1 \pmod{n}\}.$$

See [4, Prop. 4.5.4] for a proof. More precisely, in [4] the degree of the polarization is assumed to be 1, but the proof there gives a proof of [Theorem 2.12](#).

Note that [Theorem 2.12](#) implies that the image of the  $\ell$ -adic monodromy homomorphism for any leaf not contained in the supersingular stratum  $\mathcal{A}_{g,d,n}$  is maximal for  $\ell$  not dividing  $pdn$ .

We will need a slightly more general version of the irreducibility part in [Theorem 2.12](#).

2.13. THEOREM (Chai). *Let  $k \supset \mathbb{F}_p$  be an algebraically closed field. Let  $n, d$  be positive integers with  $\text{gcd}(n, pd) = 1$  and  $n \geq 3$ . Let  $Z \subset \mathcal{A}_{g,d,n}$  be a locally closed subscheme over  $k$  smooth over  $k$ , not contained in the supersingular locus and stable under all prime-to- $pd$  Hecke correspondences. Let  $W \rightarrow Z$  be a finite étale scheme over  $Z$ , and let*

$$\widetilde{W} := (W \times_{\mathcal{A}_{g,d,n}} \mathcal{A}_{g,d,m} : n|m, \text{gcd}(m, pd) = 1)$$

be the fiber product of  $Z$  with the prime-to- $pd$  tower

$$\widetilde{\mathcal{A}}_{g,d} = (\mathcal{A}_{g,d,m} : n|m, \text{gcd}(m, pd) = 1)$$

above  $\mathcal{A}_{g,d,n}$ , indexed by positive integers  $m$  which are relatively prime to  $pd$ . Assume that the prime-to- $pd$  Hecke correspondences lift to  $W$  in the sense that the action of the group  $\text{Sp}_{2g}(\mathbb{A}_f^{(pd)})$  on the restriction to  $Z$  of the prime-to- $pd$  tower  $\widetilde{\mathcal{A}}_{g,d}$  lifts to an action on the tower  $\widetilde{W}$ . Assume, moreover, that the prime-to- $pd$  Hecke correspondences on  $W$  induced by the Hecke action on the tower  $\widetilde{W}$  operate transitively on the set  $\Pi_0(W)$  of irreducible components of  $W$ . Then  $W$  is irreducible. Moreover, the prime-to- $pdn$  monodromy of  $Z$  is maximal.

The situation of [Theorem 2.13](#) is different from that of [4, Prop. 4.4]. In [4], the variety in question is a subvariety of the moduli space  $\mathcal{A}_{g,1,n}$ , while in [Theorem 2.13](#) the variety in question is a cover of subvariety of  $\mathcal{A}_{g,d,n}$ . However the proof of [4, Prop. 4.4] gives a proof of [Theorem 2.13](#); see also the proof of Proposition 5.5 in [5]. The key point of this argument is the fact that the group  $\mathrm{Sp}(\mathrm{H}_1(A_0, \mathbb{A}_f^{(pd)}), \langle, \rangle)$  has no proper subgroup of finite index; see [4, Lemma 3.1]. The assumption in [Theorem 2.13](#) implies that  $\Pi_0(Z)$  is a surjective image of a quotient of the group  $\mathrm{Sp}(\mathrm{H}_1(A_0, \mathbb{A}_f^{(pd)}), \langle, \rangle)$  by a subgroup of finite index. Since every such quotient of  $\mathrm{Sp}(\mathrm{H}_1(A_0, \mathbb{A}_f^{(pd)}), \langle, \rangle)$  is a one-point set,  $\Pi_0(Z)$  has only one element; i.e.,  $Z$  is irreducible.

2.14. *Unwinding a finite level over a leaf* (main reference: [25, Th. 1.3]). We say that a  $p$ -divisible group  $X \rightarrow S$  is *geometrically fiberwise constant*, abbreviated gfc, if fibers are mutually geometrically isomorphic; see [25, 1.1] for more details. In [25, Th. 1.3], we show that *if  $S$  satisfies reasonable finiteness conditions,  $X \rightarrow S$  is gfc, and  $i \in \mathbb{Z}_{>0}$ , then there exists a finite surjective morphism  $T_i = T \twoheadrightarrow S$  such that  $X[p^i] \times_S T$  is constant over  $T$* . We need a part of the proof of this theorem which is contained in [29]. And actually we need the part of the proof which goes back to Hasse and Witt, to Dieudonné, and which was described by Zink in [35, §2]; also see [29, 1.6].

2.15. *Completely slope divisible  $p$ -divisible groups*. Let  $t \geq r_1 > r_2 > \dots > r_m \geq 0$  be integers. Let  $q = p^t$ , and let  $S$  be a scheme over  $\mathbb{F}_q$ . A  $p$ -divisible group  $Y \rightarrow S$  is said to be *completely slope divisible* with respect to the numbers  $t \geq r_1 > r_2 > \dots > r_m \geq 0$  if there exist a filtration

$$0 = Y_0 \subset Y_1 \subset \dots \subset Y_m = Y$$

of  $Y$  by  $p$ -divisible subgroups over  $S$  such that

- each  $Y_i/Y_{i-1}$  is a  $p$ -divisible group for  $i = 1, \dots, m$ ,
- the quasi-isogenies  $p^{-r_i} \mathrm{Fr}^t : Y_i \rightarrow Y_i^{(p^t)}$  are isogenies for  $i = 1, \dots, m$ ,
- the induced homomorphisms  $p^{-r_i} \mathrm{Fr}^t : (Y_i/Y_{i-1}) \rightarrow (Y_i/Y_{i-1})^{(p^t)}$  are isomorphisms for  $i = 1, \dots, m$ .

See [29, 1.1].

Let  $0 = Y_0 \subset Y_1 \subset \dots \subset Y_m = Y$  be the slope filtration of a completely slope divisible  $p$ -divisible group  $Y \rightarrow S$  as above. Let  $G_i = Y_i/Y_{i-1}$  for  $i = 1, \dots, m$ . Let  $q = p^t$ . By the theory of  $\Phi$ -étale part of a Frobenius module in [35, §1], for each  $b \geq 1$  there exist fppf sheaves of commutative  $\mathbb{F}_q$ -algebras  $\Gamma_{i,b} = \Gamma_{i,b,q}$  on  $S$  which are representable by finite étale schemes over  $S$ , and natural isomorphisms  $\mathcal{O}_S \otimes_{\mathbb{F}_q} \Gamma_{i,b} \xrightarrow{\sim} \mathcal{O}_{G_i[p^b]}$ . The sheaf  $\Gamma_{i,b}$  is subsheaf of the structure sheaf of  $G_i[p^b]$  for the fppf topology fixed under the  $\mathrm{Frob}_S^t$ -linear map  $\Phi_i : \mathcal{O}_{G_i[p^b]} \rightarrow \mathcal{O}_{G_i[p^b]}$  induced by the composition of the isomorphism

$p^{-r_i} \text{Fr}^t : G_i[p^b] \rightarrow G_i[p^b]^{(p^t)}$  and the map  $W_s : G_i[p^b]^{(p^t)} \rightarrow G_i[p^b]$  coming from the base change by the absolute Frobenius  $\text{Frob}_S^t : S \rightarrow S$ . The group law on  $G_i[p^b]$  induces a comultiplication on  $\Gamma_{i,b}$ , making  $\Gamma_{i,b}$  a sheaf of commutative cocommutative bialgebras over  $\mathbb{F}_q$ . The inclusion maps  $G_i[p^b] \hookrightarrow G_i[p^{b+1}]$  induces surjective homomorphism  $\Gamma_{i,b+1} \rightarrow \Gamma_{i,b}$  between sheaves of bialgebras over  $\mathbb{F}_q$ .

2.16. *Remarks.* (1) A  $p$ -divisible group  $Y$  over a field  $K \supset \mathbb{F}_p$  is completely slope divisible if and only if  $Y \times_{\text{Spec}(K)} \text{Spec}(K^a)$  is.

(2) Let  $Y$  be a  $p$ -divisible group over a perfect  $K \supset \mathbb{F}_p$ . Then  $Y$  is completely slope divisible if and only if there exists a finite field  $\mathbb{F}_q \subset k$  and isoclinic  $p$ -divisible groups  $Z_1, \dots, Z_m$  over  $\mathbb{F}_q$  with distinct slopes such that  $Y$  is isomorphic to

$$Z_1 \times_{\text{Spec}(\mathbb{F}_q)} \cdots \times_{\text{Spec}(\mathbb{F}_q)} Z_m \times_{\text{Spec}(\mathbb{F}_q)} \text{Spec}(k).$$

2.17. *Remark.* Let  $Y$  be a  $p$ -divisible group over a field  $K \supset \mathbb{F}_p$ . By [35, Cor. 13],  $Y$  has a slope filtration; i.e., there exist  $p$ -divisible subgroups  $0 \subset Y_1 \subset Y_2 \subset \cdots \subset Y_m = Y$  over  $K$  such that each quotient  $Z_i := Y_i/Y_{i-1}$  is an isoclinic  $p$ -divisible group of slope  $\mu_i$  over  $K$  and  $\mu_1 > \mu_2 > \cdots > \mu_m$ . We have a natural inclusion

$$\text{End}(Y) \hookrightarrow \text{End}(Z_1) \times \cdots \times \text{End}(Z_m) = \text{End}(Z_1 \times \cdots \times Z_m).$$

This inclusion is an equality if and only if  $Y$  is isomorphic to  $Z_1 \times \cdots \times Z_m$ . The “if” direction is obvious. For the “only if” part, it suffices to note that the  $\mathbb{Z}_p$ -span of

$$(\mathbb{Z}_p^\times \cdot \text{Id}_{Z_1}) \times \cdots \times (\mathbb{Z}_p^\times \cdot \text{Id}_{Z_m})$$

in the algebra  $\text{End}(Z_1) \times \cdots \times \text{End}(Z_m)$  is equal to  $(\mathbb{Z}_p \cdot \text{Id}_{Z_1}) \times \cdots \times (\mathbb{Z}_p \cdot \text{Id}_{Z_m})$ . In particular this  $\mathbb{Z}_p$ -span contains the  $m$  idempotents corresponding to the  $m$  factors in the product  $Z_1 \times \cdots \times Z_m$ .

Several notations and prerequisites, perhaps not sufficiently well explained here, are to be found in [7] and [27].

### 3. Irreducibility of Newton polygon strata

In order to start the proof of irreducibility of leaves, we first show the irreducibility of Newton polygon strata in the principally polarized case. Note however that *supersingular* Newton polygon strata are reducible (for  $p$  large). For the case of elliptic curves in characteristic  $p$  this is classical. By Hasse, Deuring, and Igusa we know that

$$\sum_{j(E) \text{ is ss}} \frac{1}{\#(\text{Aut}(E))} = \frac{p-1}{24};$$

this implies that the number of supersingular  $j$ -invariants is asymptotically  $p/12$  for  $p \rightarrow \infty$ .

For fixed  $g$  and large  $p$  the supersingular locus has “many components” (see [14, 4.9]):

$$\begin{aligned} \#(\Pi_0(W_\sigma)) &= H_g(p, 1) \quad \text{if } g \text{ is odd,} \\ \#(\Pi_0(W_\sigma)) &= H_g(1, p) \quad \text{if } g \text{ is even.} \end{aligned}$$

Note that for  $g$  fixed and  $p \rightarrow \infty$ , indeed  $\#(\Pi_0(W_\sigma)) \rightarrow \infty$ .

3.1. THEOREM. *Let  $\xi$  be a symmetric Newton polygon which is not supersingular. The stratum  $W_\xi := \mathcal{W}_\xi(\mathcal{A}_{g,1})$  is geometrically irreducible.*

A sketch of the proof below can be found in [19].

Step 1. *The moduli scheme  $\mathcal{A}_{g,1} \otimes \mathbb{F}_p$  is geometrically irreducible.* This was proved by Faltings; see [10, Kor., p. 364]. In [1], it was proved by Chai for the case  $p > 2$ . See also [11, Chap. IV, Cor. 6.8]. For a pure characteristic  $p$  proof, see [24, Cor. 1.4].

Step 2. *Deformation to  $a \leq 1$ .* Let  $(A, \lambda)$  be a principally polarized abelian variety. There exists a deformation to a principally polarized abelian variety  $(B, \mu)$  with  $\mathcal{N}(A) = \mathcal{N}(B)$  and  $a(B) \leq 1$ .

This is a difficult theorem. For the proof we use deformation to  $a \leq 1$  for simple  $p$ -divisible groups and purity; see [8, Cor. 5.12]. Once this is established the general result follows by considering deformation of filtered  $p$ -divisible groups; see [23, §3].

COROLLARY.  $W_\xi^0(a \leq 1)$  is dense in  $W_\xi^0$ .

Step 3. *The Cayley-Hamilton theorem.* In [22] we study deformations of (polarized)  $p$ -divisible groups with  $a$ -number equal to 1. As a corollary, using the corollary in Step 2, we show that the strata  $\{W_\xi^0(a \leq 1) \mid \xi\}$  are ordered by inclusion-in-the-boundary exactly as prescribed by the NP-graph. Moreover, we compute the dimension of these strata; see [22, Th. 3.4].

Step 4. COROLLARY. *For symmetric Newton polygons  $\xi_1 \prec \xi_2$  and for an irreducible component  $P_1$  of  $W_{\xi_1}^0$ , there exists an irreducible component  $P_2$  of  $W_{\xi_2}^0$  such that  $P_1 \subset (P_2)^{\text{Zar}}$  and  $P_2$  is unique.*

COROLLARY.  $W_\xi^0(a \leq 1)$  is dense in  $W_\xi$ .

COROLLARY.  $\cup_{\xi' \preceq \xi} W_{\xi'}^0$  is contained in  $(W_\xi^0)^{\text{Zar}}$ .

Step 5. Using the facts

- every nonsupersingular Hecke- $\ell$  orbit is nonfinite (see [2, Prop. 1, p. 448]),
- EO-strata are Hecke- $\ell$  stable,

- EO-strata are quasi-affine (see [24, Th. 1.2]),

we show

*for every symmetric nonsupersingular Newton polygon  $\xi_2$  and every irreducible component  $P_2$  of  $W_{\xi_2}^0$  there exists  $\xi_1 \not\preceq \xi_2$  such that  $W_{\xi_1}^0 \cap (P_2)^{\text{Zar}} \neq \emptyset$ .*

*Proof.* Fix  $P_2 \subset W_{\xi_2}^0$ . Let  $\mathcal{P}$  be a union of irreducible components of  $W_{\xi_2}^0$ , the union taken over the Hecke- $\ell$ -orbit of  $[P_2]$  in  $\Pi_0(W_{\xi_2}^0)$ .

Consider all elementary sequences, in the sense of [25], which appear on  $P_2$ . This is a finite set  $\Gamma(P_2)$ . On the set of elementary sequences we have a partial ordering, denoted by  $\subset$  in [25, 14.3]; i.e., we write  $\varphi' \subset \varphi$  if and only if  $S_{\varphi'}$  is contained in the closure of  $S_{\varphi}$ . Indeed, this defines a partial ordering by [25, 12.5]. Let  $\varphi$  be a *minimal element* in  $\Gamma(P_2)$ . Let  $x \in W_{\xi_2}^0 \cap S_{\varphi}$ .

CLAIM.  $S_{\varphi} \cap P_2$  is not a finite set.

Consider  $\mathcal{H}_{\ell}$ . Note that  $\mathcal{H}_{\ell}(x) \subset \mathcal{P}$ . This follows because any Hecke- $\ell$  correspondence is étale and finite-to-finite. As  $x$  is not contained in the supersingular locus, it follows from [2, Prop. 1, p. 448] that this Hecke orbit  $\mathcal{H}_{\ell}(x)$  is nonfinite. Note that  $S_{\varphi}$  and  $\mathcal{P}$  are Hecke- $\ell$ -stable. Hence  $\mathcal{H}_{\ell}(x) \subset (S_{\varphi} \cap \mathcal{P})$ . The Zariski closure  $\mathcal{H}$  of  $\mathcal{H}_{\ell}(x)$  is not finite. It contains an irreducible component of positive dimension. As Hecke- $\ell$  correspondences are étale and finite-to-finite, this shows that there is an irreducible component  $H \subset (S_{\varphi} \cap P_2)$  of positive dimension containing  $x$ .

Consider the stratum  $T_{\varphi} \subset \mathcal{A}_g^*$  as defined [24, 6.1]; this consists of  $S_{\varphi}$  and all boundary points in  $\mathcal{A}_g^*$  corresponding with the elementary sequence  $\varphi$ . The stratum  $T_{\varphi}$ , in the minimal compactification  $\mathcal{A}_g^*$ , is quasi-affine (see [24, 6.5]) and  $\varphi$  is also minimal on  $T_{\varphi}$ ; see [24, 6.3]. Hence there is a point  $y'$  in the closure of  $T_{\varphi}$  not contained in  $T_{\varphi}$  itself. If  $y' \in \mathcal{A}_g$ , we write  $y = y'$ . If  $y' \notin \mathcal{A}_g$ , using [24, 6.3], we conclude there exists a point  $y \in \mathcal{A}_{g,1}$  with  $y \notin T_{\varphi}$ . This shows that the Zariski closure of  $H \subset \mathcal{A}_{g,1}$  considered above contains a point  $y \in \mathcal{A}_{g,1}$  with  $y \notin T_{\varphi}$ ; hence  $y \notin S_{\varphi}$ . Let  $\varphi'$  be the elementary sequence defined by  $y$ . Let  $\xi_1$  be the Newton polygon defined by  $y$ . As  $y \notin S_{\varphi}$  we have  $\varphi' \not\subseteq \varphi$ . Because the minimality assumption on  $\varphi$ , we see that  $y \notin W_{\xi_1}^0$  and  $\xi_1 \not\preceq \xi_2$ . This proves Step 5:

$$W_{\xi_1}^0 \cap (P_2)^{\text{Zar}} \neq \emptyset. \quad \square$$

*Remark.* Instead of the argument above, a variant of a method indicated in [2] can be used. If  $\xi$  is ordinary, then the conclusion of Step 5 follows from the fact that the ordinary locus is dense: either use that  $\mathcal{A}_{g,1}$  is irreducible, or use [17]. Suppose  $\xi$  is not ordinary (and not supersingular). Choose a point  $x$  with minimal EO-type on  $\mathcal{P}$  as above, which, moreover, is defined over a

finite field. Consider a Hilbert modular variety  $\mathcal{M}$  such that its image in  $\mathcal{A}_g$  contains  $x$ . On a Hilbert modular variety the Hecke orbit of a nonsupersingular is nonfinite. Using [24] we see that EO-strata on  $\mathcal{M}$  are quasi-affine. Moreover, any nonordinary Newton polygon stratum does not contain a boundary point in the minimal compactification of  $\mathcal{M}$ . Hence the Zariski closure of  $H \subset \mathcal{A}_{g,1}$  considered above contains a point  $y \in \mathcal{A}_{g,1}$  with  $y \notin S_\varphi$ , and we conclude as above.

*Step 6.* For  $\xi_1 \prec \xi_2$  define  $i_{\xi_2}^{\xi_1} : \Pi_0(W_{\xi_1}) \rightarrow \Pi_0(W_{\xi_2})$  by

$$i_{\xi_2}^{\xi_1}(P_1) = P_2 \iff P_1 \subset (P_2)^{\text{Zar}}.$$

Using Steps 3, 4, and 5 we conclude:

**COROLLARY.** *This map is well-defined.*

**COROLLARY.** *The map  $i_\xi^\sigma : \Pi_0(W_\sigma) \twoheadrightarrow \Pi_0(W_\xi)$  is surjective for every  $\xi$ .*

Note that this map is  $\mathcal{H}_\ell^{\text{Sp}}$ -equivariant.

**COROLLARY.** *The map  $i_{\xi_2}^{\xi_1}$  is surjective.*

*Step 7. Notation.* For  $g \in \mathbb{Z}_{>1}$  and  $j \in \mathbb{Z}_{\geq 0}$  we write  $\Lambda_{g,j}$  for the set of isomorphism classes of polarizations  $\mu$  on the superspecial abelian variety  $A = E^g \times_{\text{Spec}(\mathbb{F}_p)} \text{Spec}(k)$  such that  $\text{Ker}(\mu) = A[F^j]$ ; here  $E$  is a supersingular elliptic curve defined over  $\mathbb{F}_p$ . Note that  $\Lambda_{g,j} \xrightarrow{\sim} \Lambda_{g,j+2}$  under  $\mu \mapsto F^t \cdot \mu \cdot F$ .

*Fact. Characterization of components of  $W_\sigma$ .* There is a canonical bijective map

$$\Pi_0(W_\sigma) \xrightarrow{\sim} \Lambda_{g,g-1}.$$

See [14, 3.6 and 4.2]; this uses [18, 2.2 and 3.1].

*Step 8. Transitivity.* The action of  $\mathcal{H}_\ell^{\text{Sp}}$  on  $\Lambda_{g,g-1}$  is transitive.

*Proof.* Consider  $[(A_1, \mu_1)], [(A_2, \mu_2)] \in \Lambda_{g,g-1}$ . Let  $G_i$  be the unitary group over  $\mathbb{Q}$  defined by the semisimple algebra with involution  $(\text{End}^0(A_i), *_i)$ , where  $*_i$  is the Rosati involution attached to  $\mu_i$ . In other words,

$$G_i(R) = \{y \in (\text{End}^0(A_i) \otimes_{\mathbb{Q}} R)^\times \mid y \cdot *_i(y) = 1 = *_i(y) \cdot y\}$$

for every commutative  $\mathbb{Q}$ -algebra  $R$ . Denote by  $\mathcal{T}$  the affine scheme of finite type over  $\mathbb{Q}$  such that

$$\mathcal{T}(R) = \{\phi \in \text{Hom}(A_2, A_1) \otimes_{\mathbb{Z}} R \mid \phi^*(\lambda_1) = \lambda_2 \text{ in } \text{Hom}(A_1, A_1^t) \otimes_{\mathbb{Z}} R\}$$

for all commutative  $\mathbb{Q}$ -algebra  $R$ . It is clear that  $\mathcal{T}$  has a natural structure as a left  $G_1$ -torsor and also a right  $G_2$ -torsor; moreover, the left  $G_1$ -action commutes with the right  $G_2$ -action. We claim that this torsor  $\mathcal{T}$  is trivial; i.e., it has a  $\mathbb{Q}$ -rational point.

We use the left  $G_1$ -torsor structure of  $\mathcal{T}$ . It is clear that  $\mathcal{T}(\mathbb{Q}_\ell) \neq \emptyset$  for all prime numbers  $\ell \neq p$ . We see that  $\mathcal{T}(\mathbb{R}) \neq \emptyset$  because the Rosati involutions  $*_1$  and  $*_2$  are positive definite. The set  $\mathcal{T}(\mathbb{Q}_p)$  of  $\mathbb{Q}_p$ -points of  $\mathcal{T}$  is not empty because  $(A_1, \lambda_1)[p^\infty] \cong (A_2, \lambda_2)[p^\infty]$  by assumption. Since the group  $G_1$  is connected and simply connected, the Hasse principle applies; see [30, Th. 6.6]. We conclude that  $\mathcal{T}(\mathbb{Q}) \neq \emptyset$ .

As the torsor  $\mathcal{T}$  over  $\mathbb{Q}$  is trivial, the weak approximation theorem for semisimple simply connected algebraic groups applies; see [30, Th. 7.8, p. 415], or Lemma 4.6 for a direct proof of weak approximation for unitary groups. In particular, there exists a quasi-isogeny  $\phi$  from  $A_2$  to  $A_1$  which induces an isomorphism from  $(A_2, \lambda_2)[p^\infty]$  to  $(A_1, \lambda_1)[p^\infty]$ . The is prime-to- $p$  quasi-isogeny  $\phi$  defines a prime-to- $p$  Hecke correspondence connecting the two elements  $[(A_1, \mu_1)], [(A_2, \mu_2)]$  of  $\Lambda_{g,g-1}$ . □

**COROLLARY.** *The action of  $\mathcal{H}_\ell^{\text{Sp}}$  on  $\Pi_0(W_\sigma)$  is transitive.*

This follows using Step 7.

*Step 9. End of the proof of Theorem 3.1.* We have seen that the map  $i_\xi^\sigma : \Pi_0(W_\sigma) \rightarrow \Pi_0(W_\xi)$  is surjective and  $\mathcal{H}_\ell^{\text{Sp}}$  equivariant for every  $\xi$ ; by the previous step this implies that the action of  $\mathcal{H}_\ell^{\text{Sp}}$  on  $\Pi_0(W_\xi)$  is transitive. By Theorem 2.12 this implies that  $W_\xi$  is geometrically irreducible for  $\xi \neq \sigma$ . This finishes a proof of Theorem 3.1. □

*Remark.* The same proof shows that  $W_\xi(\mathcal{A}_{g,1,n})$  is geometrically irreducible for every  $n \neq p$ .

**3.2. Remark.** We have seen that the strata  $W_\xi^0 := W_\xi^0(\mathcal{A}_{g,1})$ , in the (principally polarized case), form a stratification of  $\mathcal{A}_{g,1}$ ; in particular for  $\xi_1 \succcurlyeq \xi_2$  the stratum  $W_{\xi_1}^0$  is contained in the closure of  $W_{\xi_2}^0$ .

However there are many cases where  $\xi_1 \succcurlyeq \xi_2$  such that a component of  $W_{\xi_1}^0(\mathcal{A}_{g,d})$  is not contained in the closure of  $W_{\xi_2}^0(\mathcal{A}_{g,d})$  for an appropriate choice of  $d$ ; this has been worked out in [28, §3].

Note the definition of  $W_\xi(\mathcal{A}_{g,d})$  is not given as the closure of  $W_\xi^0(\mathcal{A}_{g,d})$ .

We know that the dimension of a central leaf is determined by the related Newton polygon: all leaves in one Newton polygon stratum have the same dimension, independent of the degrees of the polarizations considered. However, different irreducible components of  $W_\xi^0(\mathcal{A}_{g,d})$  for varying  $d$  can have different dimension. All this has been computed and discussed in [28].

### 4. Irreducibility of leaves

**4.1. THEOREM.** *Let  $g, n$ , and  $d$  be positive integers with  $n \geq 3$  and  $\gcd(pd, n) = 1$ . Let  $k \supset \mathbb{F}_p$  be an algebraically closed field. Let  $x = [(A, \lambda, \iota)] \in \mathcal{A}_{g,d,n}(k)$  be a geometric point of  $\mathcal{A}_{g,d,n}$  not contained in the supersingular stratum. The central leaf  $\mathcal{C}(x) \subset \mathcal{A}_{g,d,n}$  passing through  $x$  is irreducible.*



Write  $d = d' \cdot m$ , where  $p$  does not divide  $m$  and  $d'$  is a power of  $p$ . Suppose that this theorem has been proved with polarizations of degree  $(d')^2$  with  $\gcd(pd', n) = 1$ . The theorem follows using [Proposition 2.12](#) and the remark following that proposition. Consider the finite étale morphism  $\pi_{g,d,n;d'} : \mathcal{A}_{g,d,n} \rightarrow \mathcal{A}_{g,d',n}$ . Let  $y := \pi_{g,d,n;d'}(x)$  and let  $\mathcal{C}(y)$  be the leaf in  $\mathcal{A}_{g,d',n}$  passing through  $y$ . The morphism  $\pi_{g,d,n;d'}$  induces a finite étale morphism  $f : \mathcal{C}(x) \rightarrow \mathcal{C}(y)$ , which is dominated by the profinite étale Galois cover over  $\mathcal{C}(y)$  given by  $m^\infty$ -level structures. If  $\mathcal{C}(y)$  has been shown to be irreducible, then it follows from [Proposition 2.12](#) that  $\mathcal{C}(x)$  is irreducible as well. *From now on in this section we suppose that  $d$  is a power of  $p$ .*

The starting point of our proof of [Theorem 4.1](#) is that the irreducibility nonsupersingular Newton polygon strata in  $\mathcal{A}_{g,1,n}$  implies the existence of points in any irreducible component of a central leaf  $C$  in  $\mathcal{A}_{g,1,n}$  which belongs to a specified isogeny class in the Newton polygon stratum containing  $C$ .

**4.2. PROPOSITION.** *Let  $g, n, d$  be positive integers as in [Theorem 4.1](#). Let  $k \supset \mathbb{F}_p$  be an algebraically closed field. Let  $C$  be a central leaf in  $\mathcal{A}_{g,d,n}$  over  $k$ , and let  $\xi$  be the symmetric Newton polygon attached to  $C$ . Let  $W_\xi^0$  be the open Newton polygon stratum in  $\mathcal{A}_{g,1,n}$  with symmetric Newton polygon  $\xi$ . Let  $C'$  be an irreducible component of  $C$ . For any point  $y = [(B_y, \mu_y, \iota_y)] \in W_\xi^0(k)$ , there exists a point  $x = [(A_x, \lambda_x, \iota_x)] \in C'(k)$  and a quasi-isogeny  $\theta \in \text{Hom}(A_x, B_y) \otimes_{\mathbb{Z}} \mathbb{Q}$  such that  $\theta^*(\mu_y) = \lambda_x$ .*

*Proof.* If  $C$  is contained in the supersingular locus, this statement is clear. For the rest of the proof we assume  $C$  is not contained in the supersingular locus. By the almost product structure explained in [Theorem 2.9](#), we obtain finite surjective morphisms  $f : T \rightarrow C$  and  $g : J \rightarrow I$  over the central leaf  $C$  and the isogeny leaf  $I$  respectively, a finite surjective morphism  $\Phi : T \times J \rightarrow W_\xi^0$ , and a quasi-isogeny

$$\Theta \in \text{Hom} \left( A_{T \times J}, B_{W_\xi^0 \times_{W_\xi^0, \Phi} (T \times J)} \right) \otimes_{\mathbb{Z}} \mathbb{Q}$$

such that  $\Theta^*(\lambda_{A_{T \times J}}) = \mu_{B_{T \times J}}$ . Here  $(A_{T \times J}, \lambda_{T \times J})$  is the pull-back to  $T \times J$  of the universal polarized abelian scheme over the central leaf  $C$  in  $\mathcal{A}_{g,d,n}$  and  $(B_{W_\xi^0 \times_{W_\xi^0, \Phi} (T \times J)}, \mu_{B_{T \times J}})$  is the pull-back by  $\Phi$  of the universal principally polarized abelian scheme over the open Newton polygon stratum  $W_\xi^0$  in  $\mathcal{A}_{g,1,n}$ .

Let  $T' = f^{-1}(C')$ , an open subscheme of  $T$ . The image  $\Phi(T' \times J)$  of  $T' \times J$  under  $\Phi$  is a union of irreducible components of  $W_\xi^0$ , hence equal to  $W_\xi^0$  because  $W_\xi^0$  is irreducible by [Theorem 3.1](#). Let  $(t, u) \in (T' \times J)(k)$  be a  $k$ -point of the pre-image  $f^{-1}(y)$  of  $y$  under  $\Phi$  and let  $x = f(t) = (A_x, \lambda_x, \iota_x)$  be the image of  $t$  in  $C'$ . Let  $\theta = \Theta_t$ , the fiber of the quasi-isogeny  $\Theta$  at  $t$ , which is a quasi-isogeny from  $A_x$  to  $B_y$  such that  $\theta^*(\mu_y) = \lambda_x$ . □

*Remark.* A special case of [Proposition 4.2](#) is the following claim in the introduction: For any hypersymmetric point  $y = (B_y, \mu_y, \iota_y)$  in an open Newton polygon  $W_\xi^0$  and any irreducible component  $C$  of a leaf contained in  $W_\xi^0$ , there exists a hypersymmetric point  $x = (A_x, \lambda_x, \iota_x)$  such that  $A_x$  is isogenous to  $B_y$ .

**4.3. PROPOSITION.** *Let  $g, d, n$  be positive integers as in [Theorem 4.1](#). Let  $C$  be a leaf in  $\mathcal{A}_{g,d,n}$  over an algebraically closed field  $k \supset \mathbb{F}_p$ , and let  $x_1 = [(B_1, \mu_1, \iota_1)]$ ,  $x_2 = [(B_2, \mu_2, \iota_2)] \in C(k)$  be two hypersymmetric  $k$ -points  $C$  such that there exists a quasi-isogeny  $\theta$  from  $B_2$  to  $B_1$  with  $\theta^*(\mu_2) = \mu_1$ . Then  $x_2$  belongs to the prime-to- $p$  Hecke orbit of  $x_1$ ; i.e., there exists a quasi-isogeny  $\psi \in \text{Hom}(B_2, B_1) \otimes_{\mathbb{Z}} \mathbb{Q}$  such that  $\psi^*(\mu_1) = \mu_2$  and the homomorphism  $\psi[p^\infty] : B_2[p^\infty] \rightarrow B_1[p^\infty]$  induced by  $\psi$  is an isomorphism.*

*Reduction of [Theorem 4.1](#) to [Proposition 4.3](#).* Let  $\xi$  be the symmetric Newton polygon attached to  $C$ . By [[6](#), Prop. 4.1], there exists a hypersymmetric point  $y \in W_\xi(k)$ . Apply [Proposition 4.2](#) to a hypersymmetric point  $y \in W_\xi(k)$ . One sees that every irreducible component of  $C$  contains a  $k$ -point  $x$  such that there exists a quasi-isogeny  $\theta : B_x \rightarrow A_y$  with  $\theta^*(\lambda_y) = \mu_x$ . [Proposition 4.3](#) tells us that the prime-to- $p$  Hecke correspondences operate transitively on the set of irreducible components of  $C$ . By [Theorem 2.12](#) this implies [Theorem 4.1](#).

**4.4.** We set up notation for a proof of [Proposition 4.3](#) by group theory. Suppose we have a polarized abelian variety  $(A, \lambda)$  over a field  $K$ . Let  $*_\lambda : \text{End}^0(A) \rightarrow \text{End}^0(A)$  be the Rosati involution attached to the polarization  $\lambda$ . We define a linear algebraic group  $G = G_{(A)} = G_{(A, \lambda)}$  over  $\mathbb{Q}$  by

$$G(R) = \text{U}(\text{End}(A) \otimes R, *) := \{x \mid x \cdot *(x) = 1 = *(x) \cdot x\}$$

for every commutative  $\mathbb{Q}$ -algebra  $R$ . In other words,  $G_{(A, \lambda)}$  is the unitary group attached to the semisimple  $\mathbb{Q}$ -algebra with involution  $(\text{End}^0(A), *_\lambda)$ . Clearly a quasi-isogeny  $\beta : B \rightarrow A$  induces an isomorphism between  $G_{(A, \lambda_A)}$  and  $G_{(B, \beta^* \lambda_A)}$ .

**4.5. LEMMA.** *Let  $(A, \lambda)$  be a polarized abelian variety over an algebraically closed field  $k \supset \mathbb{F}$ . Assume that there is an abelian variety  $B$  over  $\mathbb{F}$  such that  $B \times_{\text{Spec}(\mathbb{F})} \text{Spec}(k)$  is isogenous to  $A$ . The group  $G = G^{(A, \lambda)}$  is a connected reductive linear algebraic group, isomorphic to a product  $\prod_{i=1}^a G_i$ , where each  $G_i$  is a reductive algebraic group over  $\mathbb{Q}$  such that  $G_i \times_{\text{Spec}(\mathbb{Q})} \text{Spec}(\overline{\mathbb{Q}})$  is isomorphic to  $\text{Sp}_{2m}^b$  or  $\text{GL}_N^b$  for suitable integers  $b, m, N$ .*

*Proof.* Let  $\alpha$  be a quasi-isogeny from  $B \times_{\text{Spec}(\mathbb{F})} \text{Spec}(k)$  to  $A$  and let  $\mu$  be a polarization on  $B$  which is a multiple of  $\alpha^*(\lambda)$ . Then  $\alpha$  induces an isomorphism between the two semisimple algebras with involution,  $(\text{End}^0(A), *_\lambda)$

and  $(\text{End}^0(B), *_{\mu})$ . So we may and do assume that  $k = \mathbb{F}$ . Changing  $A$  by an isogeny, we may assume that  $A \cong \prod_{1 \leq j \leq r} C_j^{m_j}$ , where each  $C_j$  is a simple abelian variety over  $\mathbb{F}$ , each  $m_j$  is a positive integer, and  $\text{Hom}(C_j, C_{j'}) = (0)$  if  $j \neq j'$ . There exist polarizations  $\lambda_1, \lambda_r$  on  $C_1, \dots, C_r$  such that  $\lambda$  is equal to the product polarization  $\lambda_1 \times \dots \times \lambda_r$  on  $A = C_1^{m_1} \times \dots \times C_r^{m_r}$ . Let  $*_j$  be the Rosati involution on  $\text{End}^0(C_j)$  attached to  $\lambda_j$ . We have a product decomposition  $(\text{End}^0(A), *_{\lambda}) = \prod_{1 \leq j \leq r} (\text{End}^0(C_j^{m_j}), *_{\lambda_j})$  of the semisimple algebra with involution  $\text{End}^0(A), *_{\lambda}$ , which induces a product decomposition  $G_{(A, \lambda)} = \prod_{1 \leq j \leq r} G_{(C_j^{m_j}, \lambda_j)}$ . So we may and do assume that  $A \cong C^m$  for a simple abelian variety  $C$  over  $\mathbb{F}$ . The group  $G$  is isomorphic to the unitary group attached to the semisimple algebra with positive involution  $(M_m(D), *)$ , where  $D = \text{End}^0(C)$  is a division algebra which has a positive definite involution. A  $\mathbb{Q}$ -algebra of finite rank with a positive definite involution is called an *Albert algebra*. For the classification by Albert, for more information and notation, and for references see [16, §21], [21], and [7, 10.12–10.14].

According to Albert’s classification, there are four possible types for the division algebra  $D = \text{End}^0(A)$ , the endomorphism algebra of an abelian variety  $A$ , simple over a field  $K$ . In every characteristic all four types do occur for some abelian variety over some field (Albert, Shimura, Gerritzen). However, as Tate proved, over a *finite field* only two types can appear:

- (III)  $D$  is a totally definite quaternion division algebra over a totally real number field  $F$ , unramified at all finite places of  $F$  away from  $p$ .
- (IV)  $D$  is a central division algebra over a totally imaginary quadratic extension  $L$  of a totally real number field  $F$ ,  $D$  is unramified at every place of  $F$  not lying above  $p$ , and  $K/F$  is split at all places of  $F$  above  $p$ .

See [32]. For further results and references see [27, §5].

Write  $G_{\overline{\mathbb{Q}}} := G \times_{\text{Spec } \mathbb{Q}} \text{Spec } \overline{\mathbb{Q}}$ . We list some of the basic properties of  $G$  below. The proofs are straight forward.

- In case (III) above,  $G_{\overline{\mathbb{Q}}}$  is isomorphic to  $\text{Sp}_{2m}^{[F:\mathbb{Q}]}$ . So  $G$  is connected and simply connected.
- In case (IV) above, the group  $G_{\overline{\mathbb{Q}}}$  is isomorphic to  $\text{GL}_N^{[F:\mathbb{Q}]}$ , where  $N = m \cdot \sqrt{\dim_L(D)}$ . The algebraic group  $G$  over  $\mathbb{Q}$  is connected; its center is the kernel of the norm homomorphism  $\text{Nm}_{L/F} : \text{Res}_{L/\mathbb{Q}} \mathbb{G}_m \rightarrow \text{Res}_{F/\mathbb{Q}} \mathbb{G}_m$ . The derived group  $G^{\text{der}}$  of  $G$  is simply connected. The quotient torus  $G/G^{\text{der}}$  is isomorphic to the torus  $\text{Ker}(\text{Nm}_{L/F})$  above.

We have shown that the group  $G$  in Lemma 4.5 is a product of a finite number of algebraic groups over  $\mathbb{Q}$  of type (III) or (IV) above. Clearly  $G$  is connected.  $\square$

We need the weak approximation property for the reductive group  $G$  over  $\mathbb{Q}$  for the prime  $p$ .

4.6. LEMMA. *Let  $G$  be the unitary group attached to a semisimple algebra with positive definite involution  $(S, *)$  over  $\mathbb{Q}$ . Suppose that  $(S, *) = \prod_{i=1}^a (S_i, *_i)$  and each  $S_i$  is a matrix algebra over a division algebra  $D_i$  of type (III) or (IV) as in the proof of Lemma 4.5. Then the connected reductive linear algebraic group  $G$  is a  $\mathbb{Q}$ -rational variety. Especially weak approximation holds for  $G$ .*

*Proof.* Let  $S^-$  be the  $\mathbb{Q}$ -vector subspace of  $S$  consisting of all elements  $x \in S$  satisfying  $x + *(x) = 0$ ; it is naturally identified with the Lie algebra of  $G$ . Let  $\underline{S}$  be the ring scheme over  $\mathbb{Q}$  attached to  $S$ ; we see that  $\underline{S}$  is isomorphic to  $\mathbb{G}_a^{[S:\mathbb{Q}]}$  as a vector scheme over  $\mathbb{Q}$  and  $\underline{S}(\mathbb{Q})$  is naturally identified with  $S$ . Moreover,  $G$  is a closed subscheme of  $\underline{S}$ . Let  $\underline{S}^-$  be the vector subscheme of  $\underline{S}$  over  $\mathbb{Q}$  such that  $\underline{S}^-(\mathbb{Q}) = S^-$ . The standard formulas

$$u = (1 - a) \cdot (1 + a)^{-1}, a = (1 - u) \cdot (1 + u)^{-1}a \in S^{-1}, u \in G$$

for the Cayley transform define a  $\mathbb{Q}$ -birational map  $f: \underline{S}^- \dashrightarrow G$ ; i.e., the Cayley transform induces an isomorphism between an open dense subscheme of  $\underline{S}^-$  and an open dense subscheme of  $G$ . We have shown that  $G$  is  $\mathbb{Q}$ -rational. It follows that weak approximation holds for  $G$ ; see [30, Prop. 7.3]. □

4.7. We explain the definition of a quasi-bitorsor to be used in the proof of Proposition 4.3. Let  $(A, \lambda)$  and  $(B, \mu)$  be polarized abelian varieties over a field  $K$  (eventually we will assume all endomorphisms are defined over  $K$ , the polarizations are principal, and the polarized abelian varieties are in the same Hecke orbit).

For a commutative  $\mathbb{Q}$ -algebra  $R$ , a symplectic  $R$ -isogeny from  $(A, \lambda)$  to  $(B, \mu)$  is, by definition, an element of  $\text{Hom}(A, B) \otimes_{\mathbb{Z}} R$  which has an inverse in  $\text{Hom}(B, A) \otimes_{\mathbb{Z}} R$  and which respects the polarizations. Define a functor  $\mathcal{T} = \mathcal{T}_{(A,B)} = \mathcal{T}_{((A,\lambda),(B,\mu))}$  on the category of all commutative  $\mathbb{Q}$ -algebras by requiring that  $\mathcal{T}(R)$  is the set of symplectic  $R$ -isogenies from  $(A, \lambda)$  to  $(B, \mu)$ , for every commutative  $\mathbb{Q}$ -algebra  $R$ . (We should not confuse the notions of an  $R$ -isogeny and of an isogeny defined over  $R$ .) The linear group  $G_{(A,\lambda)}$  operates naturally on  $\mathcal{T}$  on the right by composing arrows in the following pattern  $A \rightarrow A \rightarrow B$ . Similarly, the linear group  $G_{(B,\mu)}$  operates on  $\mathcal{T}$  on the left by composing arrows in the pattern  $A \rightarrow B \rightarrow B$ , compatible with the previous right action by  $G_{(A,\lambda)}$ . This functor  $\mathcal{T}$  is representable, either by the empty scheme, or by a  $(G_{(B,\mu)}, G_{(A,\lambda)})$ -bitorsor over  $\mathbb{Q}$ .

*Proof of Proposition 4.3.* Let  $G_1 = G_{(B_1,\mu_1)}$ ,  $G_2 = G_{(B_2,\mu_2)}$ , and  $\mathcal{T} = \mathcal{T}_{((B_2,\mu_2),(B_1,\mu_1))}$ . We know from Lemma 4.5 that  $G_1$  and  $G_2$  are connected reductive linear algebraic groups. We also know that  $\mathcal{T}(\mathbb{Q}) \neq \emptyset$  by assumption, so  $\mathcal{T}$  is a trivial  $(G_1, G_2)$ -bi-torsor over  $\mathbb{Q}$ . By Lemma 4.6,  $G_1$  satisfies weak approximation, so the  $G_1$ -torsor  $\mathcal{T}$  also satisfies weak approximation. In other

words, the image of  $\mathcal{T}(\mathbb{Q})$  in  $\mathcal{T}(\mathbb{Q}_p)$  is dense for the  $p$ -adic topology. Since  $x_1$  and  $x_2$  are on the same leaf  $C$ , there exists an isomorphism  $\psi_p : (B_2, \mu_2)[p^\infty] \xrightarrow{\sim} (B_1, \mu_1)[p^\infty]$ . Because  $B_1$  and  $B_2$  are hypersymmetric,  $\text{Hom}(B_2, B_1) \otimes_{\mathbb{Z}} \mathbb{Q}_p \xrightarrow{\sim} \text{Hom}(B_2[p^\infty], B_1[p^\infty])$  and  $\psi_p$  corresponds to a  $\mathbb{Q}_p$ -point  $t_p$  of  $\mathcal{T}$  under the above isomorphism. Let  $U_p$  be the open subset of  $\mathcal{T}(\mathbb{Q}_p)$  consisting of all elements  $t'_p \in \mathcal{T}(\mathbb{Q}_p)$  such that  $t'_p$  induces an isomorphism from  $(B_2, \mu_2)[p^\infty]$  to  $(B_1, \mu_1)[p^\infty]$ ; it is not empty because  $t_p \in U_p$ . By the weak approximation property of  $\mathcal{T}$ , there exists an element  $t \in \mathcal{T}(\mathbb{Q}) \cap U$  under the natural injection  $\mathcal{T}(\mathbb{Q}) \hookrightarrow \mathcal{T}(\mathbb{Q}_p)$ . The quasi-isogeny  $\psi$  from  $B_2$  to  $B_1$  corresponding to  $t$  satisfies the required properties. We have proved [Proposition 4.3](#) and [Theorem 4.1](#).  $\square$

*Remark.* The assumption that  $x_1, x_2$  are hypersymmetric is used in the proof above to guarantee that  $U_p$  is not empty.

The next theorem is a special case of [Theorem 4.1](#). We state this special case explicitly because the proof below is completely different from the proof of [Theorem 4.1](#) given above.

4.8. THEOREM. *Let  $(A, \lambda)$  be a principally polarized abelian variety which is not supersingular. Suppose  $A[p^\infty]$  is a minimal  $p$ -divisible group. The central stream  $\mathcal{Z}_\xi$  passing through  $[(A, \lambda)] =: x \in \mathcal{A}_{g,1}$  is geometrically irreducible.*

The proof of [Theorem 4.8](#) is divided into three steps.

*Step 1.* For every Newton polygon  $\xi$  the central stream  $\mathcal{Z}_\xi \subset \mathcal{A}_{g,1}$  is an EO-stratum:  $\mathcal{Z}_\xi := \mathcal{C}(x) = S_\varphi$ , where  $\varphi := ((A, \lambda)[p] \bmod \cong)$ .

*Proof of Step 1.* The central stream is defined by the fact that it is the central leaf where the  $p$ -divisible groups are minimal. For a minimal  $p$ -divisible group  $X$  over  $k$ , we know that  $X[p]$  determines the isomorphism class of  $X$ ; see [26, 1.2] and [Theorem 2.11](#). Moreover, a principal quasi-polarization on a minimal  $p$ -divisible group is unique; see [25, 3.7]. This proves that the usual inclusion  $\mathcal{C}(x) \subset S_\varphi$  in this case is an equality.  $\square$

*Step 2.* Use notation as in [24]. Let  $\varphi = \{\varphi(1), \dots, \varphi(g)\}$  be an elementary sequence and let  $S_\varphi \subset \mathcal{A}_{g,1}$  be the corresponding EO-stratum. Write  $g = 2r$ , respectively  $g = 2r - 1$ ; i.e.,  $r = \lceil g/2 \rceil$ . Then

$$\varphi(r) = 0 \iff S_\varphi \subset W_\sigma.$$

*Proof of Step 2.* Let  $N_1 \subset \dots \subset N_{2g} = N = X[p]$  be a final filtration. Suppose  $\varphi(r) = 0$ ; we see that  $N_{g+r}/N_r$  is annihilated by  $F$  and by  $V$ . Hence  $X/N_r$  is superspecial. Thus  $X$  is supersingular. We conclude  $S_\varphi \subset W_\sigma$ .

Define  $u(\xi)$  to be the elementary sequence of the minimal  $p$ -divisible group with Newton polygon equal to  $\xi$ . Suppose  $\xi$  is ‘‘almost supersingular; i.e., either  $\xi = (r, r - 1) + (1, 1) + (r - 1, 1)$  or  $\xi = (r, r - 1) + (r - 1, r)$ . Direct computation

shows that

$$u(\xi) := \text{ES}(H(\xi)[p]) = \{0, \dots, \varphi(r-1) = 0, \varphi(r) = 1, \dots, 1\}.$$

In this case  $S_{u(\xi)} \not\subset W_\sigma$ . For every  $\varphi$  with  $\varphi(r) \neq 0$  we have  $\varphi \succ \text{ES}(H(\xi)[p])$ . Using [24, Th. 1.3], we conclude

$$\varphi(r) \neq 0 \implies S_{u(\xi)} \subset S_\varphi; \quad \text{hence } S_\varphi \not\subset W_\sigma. \quad \square$$

*Step 3. End of the proof of Theorem 4.8.* In [9, Th. 11.5], we see that certain EO-strata are geometrically irreducible. By Step 2 we see that these are exactly the EO-strata not contained in the supersingular locus. Moreover, using Step 1 we conclude that  $\mathcal{Z}_\xi = S_{u(\xi)}$  is geometrically irreducible for every  $\xi \neq \sigma$  and  $u(\xi) := \text{ES}(H(\xi)[p])$ . This proves Theorem 4.8.  $\square$

### 5. Monodromy

In this section we show that the  $p$ -adic monodromy of a leaf is maximal using the method in [5, §5] as indicated in [5, 6.3].

Maximality of monodromy is an irreducibility statement relative to the base. We have already seen that every nonsupersingular leaf  $C$  is irreducible, so the maximality of  $p$ -adic monodromy means that certain profinite étale cover  $\mathcal{S}$  of  $C$  is irreducible, or equivalently every finite quotient cover  $\mathcal{S}_b$  of  $C$  is irreducible. The covers  $\mathcal{S}_b$  are described in Theorem 5.13. The point of the proof is to show that the fiber in  $\mathcal{S}_b$  above a hypersymmetric point of the leaf  $C$  belongs to the same prime-to- $pd$  Hecke orbit, so  $\mathcal{S}_b$  is irreducible by Theorem 2.13.

This phenomenon that the  $p$ -adic monodromy is related to the prime-to- $p$  Hecke correspondences is reminiscent of the product formula as envisioned by H. Hida.

5.1. *Definition.* Let  $K$  be a field of characteristic  $p > 0$ , let  $\overline{K}$  be an algebraic closure of  $K$ , and let  $K^{\text{perf}}$  be the perfection of  $K$  in  $\overline{K}$ . Let  $\text{Gal}_K := \text{Gal}(\overline{K}/K^{\text{perf}})$ , naturally isomorphic to  $\text{Gal}(K^{\text{sep}}/K)$ . Let  $X$  be a  $p$ -divisible group over  $K$ . Let  $X_0$  be a  $p$ -divisible group over a finite subfield  $\mathbb{F}_q \subset K$  such that

- (i) There exists a quasi-isogeny

$$\psi_0 : X_0 \times_{\text{Spec}(\mathbb{F}_q)} \text{Spec}(\overline{K}) \longrightarrow X \times_{\text{Spec}(K)} \text{Spec}(\overline{K}).$$

- (ii) The natural map  $\text{can}_q$  in the diagram

$$\text{End}^0(X_0) \xrightarrow{\text{can}_q} \text{End}^0(X_0 \times_{\text{Spec}(\mathbb{F}_q)} \text{Spec}(\mathbb{F})) \xrightarrow{\sim} \text{End}^0(X_0 \times_{\text{Spec}(\mathbb{F}_q)} \text{Spec}(\overline{K}))$$

is an isomorphism.

Consider the set

$$\text{QIsog}[X_0, X] := \text{QIsog}\left(X_0 \times_{\text{Spec}(\mathbb{F}_q)} \text{Spec}(\overline{K}), X \times_{\text{Spec}(K)} \text{Spec}(\overline{K})\right)$$

of quasi-isogenies, which has the following properties:

- $\text{QIsog}[X_0, X]$  has a natural structure as a right  $\text{End}^0(X_0)^\times$ -torsor and a natural continuous left action by  $\text{Gal}_K$ .
- The right action of  $\text{End}^0(X_0)^\times$  on  $\text{QIsog}[X_0, X]$  is compatible with the left Galois action, i.e.  $\sigma(\psi \cdot u) = (\sigma\psi) \cdot u$  for all  $\sigma \in \text{Gal}_K$ , all  $\psi \in \text{QIsog}[X_0, X]$ , and all  $u \in \text{End}^0(X_0)^\times$ .

Define a continuous homomorphism  $\rho_{X, \psi_0} : \text{Gal}_K \rightarrow \text{End}^0(X_0)^\times$  by

$$\sigma\psi_0 = \psi_0 \cdot \rho_{X, \psi_0}(\sigma) \forall \sigma \in \text{Gal}_K.$$

We call  $\rho_{X, \psi_0}$  the *p-adic monodromy homomorphism* of  $X$  with respect to  $\psi_0$ .

5.2. *Remarks.* (1) Suppose that  $t$  is a positive integer which is a common multiple of the denominators of the slopes of  $X$ . Then for  $q = p^t$ , there exists a *decent*  $p$ -divisible group  $X_0$  over the finite field  $\mathbb{F}_q$  which satisfies the requirements (i) and (ii); see [Remark 5.3](#).

(2) It is not difficult to see that if  $X_0, X_1$  are two  $p$ -divisible groups over  $\mathbb{F}_q$  satisfying (i) and (ii), then  $X_1$  is up to isogeny a Galois twist of  $X_0$  via a continuous homomorphism

$$\chi : \text{Gal}(\mathbb{F}/\mathbb{F}_q) \rightarrow Z\left(\text{End}^0(X_0)\right)^\times,$$

where  $Z\left(\text{End}^0(X_0)\right)$  is the center of  $\text{End}^0(X_0)$ . In other words the Galois group  $\text{Gal}(\mathbb{F}/\mathbb{F}_q)$  acts on  $\text{QIsog}[X_0, X_1]$  via a character  $\chi$  as above. If, in addition,  $X_0$  and  $X_1$  are decent, then the character  $\chi$  has finite image; see [Remark 5.3](#).

(3) Suppose that  $\mu$  is a polarization on  $X$  and  $\mu_0$  is a polarization on  $X_0$ , and  $\psi_0^*(\mu) = \mu_0$  as polarizations on  $X_0 \times_{\text{Spec}(\mathbb{F}_q)} \text{Spec}(\overline{K})$ . Denote by  $U$  the subgroup of  $\text{End}^0(X_0)^\times$  consisting of all quasi-isogenies from  $X_0$  to itself which preserve the polarization  $\mu_0$ . Let

$$\text{UQIsog}[X_0, X] := \text{UQIsog}\left(X_0 \times_{\text{Spec}(\mathbb{F}_q)} \text{Spec}(\overline{K}), X \times_{\text{Spec}(K)} \text{Spec}(\overline{K})\right)$$

be the set of all elements  $\psi$  in  $\text{QIsog}(X_0 \times_{\text{Spec}(\mathbb{F}_q)} \text{Spec}(\overline{K}), X \times_{\text{Spec}(K)} \text{Spec}(\overline{K}))$  such that  $\psi^*(\mu) = \mu_0$ . Then  $\text{UQIsog}[X_0, X]$  is stable under both the left action by  $\text{Gal}_K$  and the right action by the subgroup  $U$  of  $\text{End}^0(X_0)^\times$ . So the  $p$ -adic monodromy homomorphism  $\rho_{X, \psi_0} : \text{Gal}_K \rightarrow \text{End}^0(X_0)^\times$  factors through the subgroup  $U \subset \text{End}^0(X_0)^\times$ .

(4) If  $\psi_0$  is an isomorphism

$$\psi_0 : X_0 \times_{\text{Spec}(\mathbb{F}_q)} \text{Spec}(\overline{K}) \xrightarrow{\sim} X \times_{\text{Spec}(K)} \text{Spec}(\overline{K}),$$

then

$$\rho_{X, \psi_0}(\text{Gal}_K) \subseteq \text{Aut}(X_0).$$

Similarly to the situation of (2) above, if  $\psi_0$  is an isomorphism

$$\psi_0 : (X_0, \mu_0) \times_{\text{Spec}(\mathbb{F}_q)} \text{Spec}(\overline{K}) \xrightarrow{\sim} (X, \mu) \times_{\text{Spec}(K)} \text{Spec}(\overline{K}),$$

then  $\rho_{X, \psi_0}(\text{Gal}_K) \subseteq \text{Aut}(X_0, \mu_0)$ .

(5) Suppose that  $X_2$  is a  $p$ -divisible group over  $\mathbb{F}_q$ ,  $\psi_2$  is an element of  $\text{QIsog}[X_2, X]$ , and  $\delta$  is the quasi-isogeny from

$$X_2 \times_{\text{Spec}(\mathbb{F}_q)} \text{Spec}(\mathbb{F}) \text{ to } X_0 \times_{\text{Spec}(\mathbb{F}_q)} \text{Spec}(\mathbb{F})$$

such that  $\psi_0 \circ \delta = \psi_2$ . Then

$$(\dagger) \quad \rho_{X, \psi_2}(\sigma) = \delta^{-1} \circ \rho_{X, \psi_0}(\sigma) \circ \sigma \delta \quad \forall \sigma \in \text{Gal}_K,$$

where  $\psi_2 = \psi_0 \circ \delta$ . In particular, if  $\psi_1 = \psi \circ \gamma$  for an element  $\gamma \in \text{End}^0(X_0)^\times$ , then

$$\rho_{X, \psi_1}(\sigma) = \gamma^{-1} \cdot \rho_{X, \psi_0}(\sigma) \cdot \gamma \quad \forall \sigma \in \text{Gal}_K.$$

The formula  $(\dagger)$  also shows that if  $K \supset \mathbb{F}$ , then the  $p$ -adic monodromy homomorphism for a  $p$ -divisible group  $X$  is unique up to conjugation.

(6) Suppose that  $\beta : X \rightarrow X'$  is a quasi-isogeny of  $p$ -divisible groups over  $K$  and that  $\psi' = \beta \circ \psi$ . Then  $\rho_{X', \psi'} = \rho_{X, \psi}$ .

(7) From (1) and (4) above, we see that the  $p$ -adic monodromy homomorphism  $\rho_X$  attached to a  $p$ -divisible group is unique up to

- (a) conjugation, and
- (b) multiplication by a continuous homomorphism

$$\chi : \text{Gal}_K \longrightarrow Z(\text{End}^0(X))^\times.$$

5.3. *Remarks.* (1) Recall that a  $p$ -divisible group  $Y$  over a perfect field  $L \supset \mathbb{F}_p$  is said to be *decent* for if its associated isocrystal is generated by elements  $v_i$  satisfying an equation of the form  $F^t v = p^{r_i} v$  with  $t, r_i \in \mathbb{Z}_{>0}$ ; see [31, 2.13]. The group  $G_{m,n}$  in Section 2.4 is a decent  $p$ -divisible group over  $\mathbb{F}_p$ ; its base change to  $\mathbb{F}_{p^{m+n}}$  satisfies requirement (i) in Definition 5.1.

(2) If  $Y_1, Y_2$  are decent  $p$ -divisible groups over a perfect field  $K \supset \mathbb{F}_p$ , then there exists a finite extension field  $L$  of  $K$  such that the canonical map

$$\begin{aligned} \text{Hom}(Y_1 \times_{\text{Spec}(K)} \text{Spec}(L), Y_2 \times_{\text{Spec}(K)} \text{Spec}(L)) \\ \xrightarrow{\sim} \text{Hom}(Y_1 \times_{\text{Spec}(K)} \text{Spec}(\overline{K}), Y_2 \times_{\text{Spec}(K)} \text{Spec}(\overline{K})) \end{aligned}$$

is an isomorphism.

(3) From (2) above and Remarks 5.2(1) and (4), we see that if we require the “base quasi-isogeny”  $\psi_0$  in Definition 5.1 to be decent, then the  $p$ -adic monodromy homomorphism attached to a  $p$ -divisible group  $X$  is unique up to

- (a) conjugation, and
- (b) multiplication by a homomorphism  $\chi : \text{Gal}_K \longrightarrow Z(\text{End}^0(X))^\times$  with finite image.



5.4. We discuss a variant of [Definition 5.1](#) on  $p$ -adic monodromy in the case of a completely slope divisible  $p$ -divisible group over an integral base  $\mathbb{F}_p$ -scheme and also in the case of a  $p$ -divisible group with constant Newton polygon over a noetherian normal integral  $\mathbb{F}_p$ -scheme.

Let  $0 = Y_0 \subset Y_1 \subset \dots \subset Y_m = Y$  be the slope filtration of a completely slope divisible  $p$ -divisible group  $Y \rightarrow S$  with respect to natural numbers  $t \geq r_1 > r_2 > \dots > r_m \geq 0$  as in [Section 2.15](#), where  $S$  is an integral  $\mathbb{F}_q$  scheme,  $q = p^t$ . We will show that

*The  $p$ -adic monodromy homomorphism for  $Y \rightarrow S$  factors through the fundamental group  $\pi_1(S)$  of  $S$  and can be computed via a projective system of finite étale coverings of  $S$ .*

We follow the notation in [Section 2.15](#).

For every geometric point  $\bar{s}$  of  $S$ , we get homomorphisms

$$\rho_{G_i, b, \bar{s}}: \pi_1(S, \bar{s}) \longrightarrow \text{Aut} \left( (\Gamma_{i, b})_{\bar{s}} \right)$$

from the fundamental group of  $S$  to the group of automorphisms of the commutative bialgebra  $(\Gamma_{i, b})_{\bar{s}}$  over  $\mathbb{F}_q$ . The homomorphisms  $\rho_{G_i, b, \bar{s}}$  are compatible if we fix  $i$  and let  $b$  vary. Taking the limit, we get homomorphisms

$$\rho_{G_i, \bar{s}}: \pi_1(S, \bar{s}) \longrightarrow \varprojlim_b \text{Aut} \left( (\Gamma_{i, b})_{\bar{s}} \right).$$

Notice that the target  $\varprojlim_b \text{Aut}((\Gamma_{i, b})_{\bar{s}})$  of the homomorphism  $\rho_{G_i, \bar{s}}$  is naturally isomorphic to the opposite of the group of automorphisms of  $G_i \times_S \bar{s}$ . Let

$$\rho_{Y, \bar{s}}: \pi_1(S, \bar{s}) \longrightarrow \prod_{i=1}^m \varprojlim_b \text{Aut} \left( (\Gamma_{i, b})_{\bar{s}} \right)$$

be the product of the homomorphisms  $\rho_{G_i, \bar{s}}$ . The composition of  $\rho_{Y, \bar{s}}$  with the inverse map on  $\pi_1(S, \bar{s})$  can be regarded as a homomorphism from  $\pi_1(S, \bar{s})$  to the group of automorphisms  $\text{Aut}(Y_{\bar{s}})$  of the geometric fiber  $Y_{\bar{s}}$  of  $Y$ ; see [Remarks 2.16](#) and [2.17](#). We denote the composition also by  $\rho_{Y, \bar{s}}$  if no confusion is possible.

Suppose that  $X \rightarrow S$  is a  $p$ -divisible group with constant Newton polygon over a noetherian normal integral  $\mathbb{F}_p$ -scheme  $S$ . By [[29](#), Th. 2.1], there exists a completely slope divisible  $p$ -divisible group  $Y \rightarrow S$  and a quasi-isogeny  $\phi: Y \rightarrow X$  over  $S$ . Let  $\bar{s}$  be a geometric point of  $S$ . The quasi-isogeny  $\phi_{\bar{s}}$  induces an injection from  $j_{\phi, \bar{s}}: \text{Aut}(Y_{\bar{s}})$  to  $\text{End}^0(X_{\bar{s}})^\times$ . Let  $\rho_{X, \phi, \bar{s}}$  be the composition of  $\rho_{Y, \bar{s}}$  with  $j_{\phi, \bar{s}}$ ; we call it the  *$p$ -adic monodromy* attached to  $Y \xrightarrow{\phi} X \rightarrow S$  with base point  $\bar{s}$ . It is not difficult to check that the homomorphism

$$\rho_{X, \phi, \bar{s}}: \pi_1(X, \bar{s}) \rightarrow \text{End}^0(X_{\bar{s}})^\times$$

is independent of the choice of quasi-isogeny  $\phi$ , so the subscript “ $\phi$ ” can be dropped from the notation  $\rho_{X, \phi, \bar{s}}$ .

5.5. *Remarks.* (1) The construction/definition of  $\rho_{Y, \bar{s}}$  is independent of the choice of  $q = p^t$  in the following sense. Suppose that  $\mathcal{O}_S$  contains  $\mathbb{F}_{q^a}$  for some positive integer  $a$ . Then  $C_{i,b,q^a} = C_{i,b,q} \otimes_{\mathbb{F}_q} \mathbb{F}_{q^a}$  for all  $i = 1, \dots, m$  and all  $b \geq 1$ . Consequently the homomorphisms  $\rho_{Y,q,\bar{s}}$  and  $\rho_{Y,q^a,\bar{s}}$  defined using  $q$  and  $q^a$  respectively are equal, as homomorphisms from  $\pi_1(S, \bar{s})$  to  $\text{Aut}(Y_{\bar{s}})$ .

(2) The construction/definition of the  $p$ -adic monodromy homomorphism in Section 5.4 coincides with Definition 5.1 up to conjugation. Let  $S$  be an integral scheme with function field  $K$ . Let  $\eta$  be the generic point of  $S$  and let  $\bar{\eta}$  be the geometric generic point of  $S$ . Let  $Y \rightarrow S$  be a completely  $p$ -divisible group as in Section 5.4. For  $i = 1, \dots, m$ , let  $X_i$  be the  $p$ -divisible group over  $\mathbb{F}_q$  with  $X_i[p^b] = \text{Spec}((\Gamma_{i,b})_{\bar{\eta}})$  over  $\mathbb{F}_q$ ; each  $X_i$  is decent by construction. Let  $X = X_1 \times_{\text{Spec}(\mathbb{F}_q)} \cdots \times_{\text{Spec}(\mathbb{F}_q)} X_m$ . The theory of  $\Phi$ -étale part in [35] defines an isomorphism

$$\psi: X \times_{\text{Spec} \mathbb{F}_q} \bar{\eta} \xrightarrow{\sim} Y_{\bar{\eta}}.$$

The  $p$ -adic monodromy homomorphism

$$\rho_{Y_{\eta}, \psi_0}: \text{Gal}(K^a/K^{\text{perf}}) \longrightarrow \text{End}^0(X)^\times = \text{End}^0(X_1)^\times \times \cdots \times \text{End}^0(X_m)^\times$$

defined in Definition 5.1 is compatible with the homomorphism

$$\rho_{Y_{\eta}, \psi}: \pi_1(S, \bar{\eta}) \rightarrow \text{Aut}(X) = \text{Aut}(X_1) \times \cdots \times \text{Aut}(X_m)$$

via the natural surjection  $\text{Gal}(K^a/K^{\text{perf}}) \rightarrow \pi_1(S, \bar{\eta})$  and the inclusion  $\text{Aut}(X) \hookrightarrow \text{End}^0(X)^\times$ . To see this, it suffices to check for the case when  $m = 1$ , because  $Y_{\bar{\eta}}$  splits canonically into a product  $(Y_1)_{\bar{\eta}} \times (Y_2/Y_1)_{\bar{\eta}} \cdots \times (Y_m/Y_{m-1})_{\bar{\eta}}$ , which induces a splitting of  $\psi$  as a product  $\psi_1 \times \cdots \times \psi_m$ . Furthermore, the  $\rho_{Y_{\eta}, \psi}$  is the product of the  $p$ -adic monodromy homomorphisms  $\rho_{(Y_i/Y_{i-1})_{\eta}, \psi_i}$ . Assume that  $Y$  is isoclinic and completely  $p$ -divisible; let  $C(b)$  be the étale sheaf commutative  $\mathbb{F}_q$ -bialgebras over  $S$  attached to  $Y[p^b]$  as before. The theory of  $\Phi$ -étale part shows that the projective system  $\varprojlim_b \text{Jsom}(C(b), C(b)_{\bar{\eta}} \times S)$  of sheaves on  $S_{\text{fppf}}$  is isomorphic to the projective system  $\varprojlim_b \text{Jsom}(X[p^b] \times_{\text{Spec}(\mathbb{F}_q)} S, Y[p^b])$  of sheaves. The claim on compatibility follows.

5.6. **THEOREM.** *Let  $k \supset \mathbb{F}_p$  be an algebraically closed field. Let  $n$  be a positive integers with  $n \geq 3$  with  $\text{gcd}(n, pd) = 1$ . Let  $C$  be a leaf over  $k$  in  $\mathcal{A}_{g,d,n}$ , and let  $x = [(A_x, \lambda_x, \eta_x)]$  be a  $k$ -point on  $C$ . Then the  $p$ -adic monodromy homomorphism*

$$\rho_{(A,\lambda)[p^\infty]/C,x}: \pi_1(C, x) \longrightarrow \text{Aut}((A_x, \lambda_x)[p^\infty])$$

for the polarized  $p$ -divisible group attached to the universal polarized abelian scheme  $(A, \lambda)$  over  $C$  is surjective.

Write  $d = d' \cdot m$ , where  $p$  does not divide  $m$  and  $d'$  is a power of  $p$ . Suppose that this theorem has been proved with polarizations of degree  $(d')^2$ . Then

the theorem follows using [Proposition 2.13](#) and the remarks following that proposition, as in the paragraph after the statement of [Theorem 4.1](#). From now on we suppose that  $d$  is a power of  $p$ .

*Remark.* It suffices to prove [Theorem 5.6](#) for one  $k$ -point  $x \in C(k)$ . In the following we will take a hypersymmetric point  $x_0 \in C(k)$  and prove [Theorem 5.6](#) in the case  $x = x_0$ .

We will first prove [Theorem 5.6](#) in the special case [Theorem 5.9](#), when  $A_0[p^\infty]$  is completely slope divisible. The basic argument is more transparent and not burdened with extra technical complications such as [Proposition 5.11](#).

5.7. Let  $k$  be an algebraically closed field as before. Let  $C \subset \mathcal{A}_{g,d,n}$  be a leaf over  $k$  such that the restriction to  $C$  of the  $p$ -divisible group attached to the universal abelian scheme is completely slope divisible, or equivalently there exists a point  $x = [(A_x, \lambda_x, \iota_x)]$  on  $C$  such that  $A_x[p^\infty]$  is completely slope divisible. Assume in addition that  $C$  is not contained in the supersingular locus. By [Theorem 4.1](#), the leaf  $C$  contains a  $k$ -point  $x_0 = [(A_0, \lambda_0, \iota_0)]$  which is hypersymmetric. Moreover,  $C$  is smooth and irreducible. It is convenient to use the definition of  $p$ -adic monodromy explained in [Section 5.4](#), because the  $p$ -divisible group  $A[p^\infty] \rightarrow C$  is completely slope divisible. Let

$$0 = Y_0 \subset Y_1 \subset \dots \subset Y_m = A[p^\infty]$$

be the slope filtration of  $A[p^\infty]$  over  $C$ . Let

$$\rho_{A[p^\infty]/C, x_0} : \pi_1(C, x_0) \longrightarrow \text{Aut}((A_0, \lambda_0)[p^\infty])$$

be the  $p$ -adic monodromy homomorphism explained in [Section 5.4](#). Here the polarization  $\lambda$  of the abelian scheme  $A \rightarrow C$  induces a polarization of the  $p$ -divisible group  $A[p^\infty] \rightarrow C$ , again denoted by  $\lambda$ , which is compatible with the slope filtrations on  $A[p^\infty] \rightarrow C$  and on the dual of  $A[p^\infty] \rightarrow C$ . Hence the  $p$ -adic monodromy homomorphism factors through the subgroup  $\text{Aut}((A_0, \lambda_0)[p^\infty])$  of  $\text{Aut}(A_0[p^\infty])$ .

Let  $L_{\mathbb{Z}_p} := \text{Aut}((A_0, \lambda_0)[p^\infty])$  and denote by  $L_{\mathbb{Z}/p^b\mathbb{Z}}$  the finite image of  $L_{\mathbb{Z}_p}$  in the quotient group  $\text{Aut}((A_0, \lambda_0)[p^b])$  for every positive integer  $b$ . Note that the group  $\text{Aut}((A_0, \lambda_0)[p^b])$  can be infinite, while  $L_{\mathbb{Z}_p/p^b\mathbb{Z}}$  is finite for all  $b$ . Clearly we have

$$L_{\mathbb{Z}_p} = \varprojlim_b L_{\mathbb{Z}/p^b\mathbb{Z}},$$

the projective limit of the finite groups  $L_{\mathbb{Z}/p^b\mathbb{Z}}$ , where the transition maps in the projective system are all surjective. The  $p$ -adic monodromy homomorphism

$$\rho_{A[p^\infty]/C, x_0} : \pi_1(C, x_0) \longrightarrow \text{Aut}((A_0, \lambda_0)[p^\infty])$$

comes from a right étale  $L_{\mathbb{Z}_p}$ -torsor  $\mathcal{T}$  over  $C$ , constructed from the projective system

$$\left( \prod_{i=1}^m \Gamma_{i,b} \right)_b$$

of étale covers of  $C$ , analogous to the “reduction of the structural group” procedure. Stated differently, we have a projective system of finite étale  $L_{\mathbb{Z}/p^b\mathbb{Z}}$ -torsors  $\mathcal{T}_b$  over  $C$  such that  $\mathcal{T} = \varprojlim_b \mathcal{T}_b$  and  $\prod_{i=1}^m \Gamma_{i,b}$  is the contraction product  $\mathcal{T}_b \times^{L_{\mathbb{Z}/p^b\mathbb{Z}}} (\Gamma_{i,b})_{x_0}$ , i.e., the finite étale cover of  $C$  attached to the  $L_{\mathbb{Z}/p^b\mathbb{Z}}$ -torsor  $\mathcal{T}_b$  and the natural action of  $\mathcal{T}_b$  on the fiber  $(\Gamma_{i,b})_{x_0}$  of  $\Gamma_{i,b}$  at  $x_0$ . By construction, for each  $b \geq 1$  the composition of the homomorphism  $\pi_1(C, x_0) \rightarrow L_{\mathbb{Z}/p^b\mathbb{Z}}$  associated to the  $L_{\mathbb{Z}/p^b\mathbb{Z}}$  and the inclusions

$$L_{\mathbb{Z}/p^b\mathbb{Z}} \hookrightarrow \text{Aut}(A_0[p^b]) = \prod_{i=1}^m \text{Aut}((\Gamma_{i,b})_{x_0})$$

is equal to the homomorphism from  $\pi_1(C, x_0)$  to  $\prod_{i=1}^m \text{Aut}((\Gamma_{i,b})_{x_0})$  attached to the family of étale covers  $(\Gamma_{i,b})_{i=1, \dots, m}$ . The inverse limit of the image in  $\text{Aut}(A_0[p^b])$  of this homomorphism  $\pi_1(C, x_0) \rightarrow \text{Aut}(A_0[p^b])$ , as  $b$  goes to  $\infty$ , is equal to the image of the  $p$ -adic monodromy homomorphism  $\rho_{(A,\lambda)[p^\infty]/C,x_0}$ . The natural projections  $\mathcal{T}_{b+1} \rightarrow \mathcal{T}_b$  are surjective and  $\mathcal{T}$  is the projective limit of the  $\mathcal{T}_b$ 's.

The set of  $k$ -points of the  $L_{\mathbb{Z}_p}$ -torsor  $\mathcal{T}$  can be canonically identified with the set of isomorphism classes of quadruples  $(B, \lambda_B, \iota, \psi)$ , where  $[(B, \lambda_B, \psi)]$  is a  $k$ -point of  $C$ , in particular  $\iota$  is a symplectic level- $n$  structure of the principally polarized abelian variety  $(B, \lambda)$ , and

$$\psi: (A_0, \lambda_0)[p^\infty] \xrightarrow{\sim} (B, \lambda_B)[p^\infty]$$

is an isomorphism. The set of  $k$ -points of the  $L_{\mathbb{Z}/p^b\mathbb{Z}}$ -torsor  $\mathcal{T}_b$  is the set of isomorphism classes of quadruples  $[(B, \lambda_B, \iota, \psi_b)]$ , where  $[(B, \lambda_B, \iota)]$  is a  $k$ -point of  $C$  and  $\psi_b: (A_0, \lambda_0)[p^b] \xrightarrow{\sim} (B, \lambda_B)[p^b]$  is an isomorphism which can be lifted to an isomorphism  $\psi: (A_0, \lambda_0)[p^\infty] \xrightarrow{\sim} (B, \lambda_B)[p^\infty]$  of polarized  $p$ -divisible groups.

More generally, for a  $k$ -scheme  $S$ , the set of  $S$ -points of the  $L_{\mathbb{Z}/p^b\mathbb{Z}}$ -torsor  $\mathcal{T}_b$  can be described similarly. It is the set of isomorphism classes of quadruples  $[(B, \lambda_B, \iota, \psi_b)]$ , where  $[(B, \lambda_B, \iota)]$  is an  $S$ -point of  $C$  as before, so that  $B \rightarrow S$  is a  $g$ -dimensional abelian scheme,  $\lambda_B$  is a polarization on  $B \rightarrow S$  of degree  $d$ ,  $\iota$  is a level- $n$  structure for  $B \rightarrow S$ , and all geometric fibers of  $(B, \lambda_B)[p^\infty] \rightarrow S$  are isomorphic to  $(A_0, \lambda_0)[p^\infty]$  after suitable extension of base fields. The fourth element  $\psi_b$  in the above quadruple is an isomorphism

$$\psi_b: \prod_{i=1}^m (X_i/X_{i-1})[p^b] \times_{\text{Spec}(k)} S \xrightarrow{\sim} \prod_{i=1}^m (Y_i/Y_{i-1})[p^b] \times_C S,$$

which respects the polarizations induced by  $\lambda_0$  and  $\lambda_B$  respectively, such that for each  $N \geq b$  the isomorphism  $\psi_b$  over  $S$  can be lifted étale locally to a polarization-preserving isomorphism

$$\psi_N: \prod_{i=1}^m (X_i/X_{i-1})[p^N] \times_{\text{Spec}(k)} S \xrightarrow{\sim} \prod_{i=1}^m (Y_i/Y_{i-1})[p^N] \times_C S.$$

In the above,  $0 \subset X_1 \subset \dots \subset X_m = A_0[p^\infty]$  denotes the slope filtration on  $A_0[p^\infty]$  and  $0 \subset Y_1 \subset \dots \subset Y_m = A[p^\infty]$  is the slope filtration on the completely slope divisible  $p$ -divisible group  $A[p^\infty] \rightarrow C$ . Notice that the base change to  $S$  of the slope filtration for  $A[p^\infty] \rightarrow C$  is the slope filtration for  $B[p^\infty] \rightarrow S$ .

Our goal is to prove a strong version of the maximality of the  $p$ -adic monodromy homomorphism  $\rho_{A[p^\infty]/C, x_0}$  which asserts that its image is equal to  $\text{Aut}((A_0, \lambda_0)[p^\infty])$ . This is equivalent to the statement that the  $L_{\mathbb{Z}/p^b\mathbb{Z}}$ -torsor  $\mathcal{T}_b$  is geometrically irreducible for every  $b \geq 1$ .

5.8. (notation as in §5.7). To prove that  $L_{\mathbb{Z}/p^b\mathbb{Z}}$ -torsor  $\mathcal{T}_b$  is irreducible, we will use the prime-to- $p$  Hecke correspondences on  $L_{\mathbb{Z}/p^b}$ . These correspondences come from the action of the locally compact group  $\text{Sp}_{2g}(\mathbb{A}_f^{(p)})$  on the prime-to- $p$  tower  $(\mathcal{T}_{b;c})_c$  defined below, where  $c$  runs through all positive multiples of  $n$  such that  $\text{gcd}(c, p) = 1$ . For each positive integer multiple  $c$  of  $n$  not dividing  $p$ , let  $C_c := C \times_{\mathcal{A}_{g,d,n}} \mathcal{A}_{g,d,c}$  and let  $\mathcal{T}_{b;c} := \mathcal{T}_b \times_C C_c$ . Thus  $\mathcal{T}_{b;c}$  is a torsor over  $\mathcal{T}_b$  for the group  $\text{Sp}_{2g}(\mathbb{Z}/c\mathbb{Z})(n)$ , the group of all elements of  $\text{Sp}_{2g}(\mathbb{Z}/c\mathbb{Z})$  which are congruent to  $\text{Id}$  modulo  $n$ . Let  $\mathcal{T}_b^\sim$  be the projective limit of the tower  $(\mathcal{T}_{b;c})_c$ .

The action of the group  $\text{Sp}_{2g}(\mathbb{A}_f^{(p)})$  on the prime-to- $p$  tower

$$C^\sim := (C \times_{\mathcal{A}_{g,d,n}} \mathcal{A}_{g,d,c})_c$$

lifts to an action on  $\mathcal{T}_b^\sim$ , described below.

First we describe the set of  $S$ -valued points of  $\mathcal{T}_b^\sim$  for any  $k$ -scheme  $S$ . It consists of isomorphism classes of all quadruples  $(B, \lambda_B, \tilde{\iota}, \psi_b)$ , where

- $(B, \lambda_B) \rightarrow S$  is a  $g$ -dimensional polarized abelian scheme over  $S$  up to prime-to- $p$  quasi-isogenies,  $\text{deg}(\lambda_B) = d^2$ ;
- $\tilde{\iota}: \text{H}_1(A_0, \mathbb{A}_f^{(p)}) \times S \xrightarrow{\sim} \underline{\text{H}}_1(B/S, \mathbb{A}_f^{(p)})$  is a symplectic isomorphism such that the image of  $[(B, \lambda_B, \tilde{\iota})]$  in  $\mathcal{A}_{g,d,n}$  factors through  $C \hookrightarrow \mathcal{A}_{g,d,n}$ ; and
- $\psi_b = (\psi_{b,i})_{1 \leq i \leq m}$  is a family of isomorphisms  $\psi_{b,i}$ ,  $i = 1, \dots, m$  as in Section 5.7, so that each  $\psi_{b,i}$  is an isomorphism  $(X_i/X_{i-1})[p^b] \times_{\text{Spec}(k)} S \xrightarrow{\sim} (Y_i/Y_{i-1})[p^b] \times_C S$  such that for each  $N \geq b$   $\psi_{b,i}$  can be extend étale locally to an isomorphism

$$(X_i/X_{i-1})[p^N] \times_{\text{Spec}(k)} S \xrightarrow{\sim} (Y_i/Y_{i-1})[p^N] \times_C S.$$

An isomorphism between two quadruples  $(B, \lambda_B, \tilde{\iota}, \psi_b)$  and  $(B', \lambda_{B'}, \tilde{\iota}', \psi'_b)$  is a prime-to- $p$  isogeny  $\beta$  from  $B$  to  $B'$  which sends  $\tilde{\iota}$  to  $\tilde{\iota}'$  and  $\psi_b$  to  $\psi'_b$

Under the above description of points of the projective limit  $\mathcal{T}_b^\sim$ , the symplectic group  $\mathrm{Sp}(\mathrm{H}_1(A_0, \mathbb{A}_f^{(p)}), \langle, \rangle)$  operates on the right of  $\mathcal{T}_b^\sim$ . Suppose that  $\gamma$  is an element of the symplectic group  $\mathrm{Sp}(\mathrm{H}_1(A_0, \mathbb{A}_f^{(p)}), \langle, \rangle)$  and  $[(B, \lambda_B, \iota^\sim, \psi_b)]$  is a point of  $\mathcal{T}_b^\sim$ ; then

$$[(B, \lambda_B, \tilde{\iota}, \psi_b)] \cdot \gamma = [B, \lambda_B, \tilde{\iota} \circ \gamma, \psi_b].$$

As usual, the action of the group  $\mathrm{Sp}(\mathrm{H}_1(A_0, \mathbb{A}_f^{(p)}), \langle, \rangle)$  on  $\mathcal{T}_b^\sim$  induces a family of algebraic correspondences on  $\mathcal{T}_b$ . Recall that  $A_0$  is a hypersymmetric abelian variety; hence the natural map  $\mathrm{End}(A_0) \otimes_{\mathbb{Z}} \mathbb{Z}_p \rightarrow \mathrm{End}(A_0[p^\infty])$  is an isomorphism. Let  $*_0$  be the Rosati involution attached to the polarization  $\lambda_0$  of  $A_0$ . Let  $H$  be the unitary group over  $\mathbb{Q}$  attached to the semisimple algebra  $\mathrm{End}^0(A_0)$  with involution  $*_0$ ; it is the linear algebra group over  $\mathbb{Q}$  such that for every commutative  $\mathbb{Q}$ -algebra  $R$  the set  $H(R)$  of  $R$ -valued points of  $H$  consists of all elements  $x \in (\mathrm{End}^0(A_0) \otimes_{\mathbb{Q}} R)^\times$  such that  $*_0(x) \cdot x = x \cdot *_0(x) = 1$  in  $(\mathrm{End}^0(A_0) \otimes_{\mathbb{Q}} R)^\times$ . In other words  $H(R)$  is the group of symplectic  $R$ -isogenies from  $A_0$  to  $A_0$ . We abuse the notation and denote by  $H(\mathbb{Z}_p)$  the subgroup of  $H(\mathbb{Q}_p)$  consisting of symplectic  $\mathbb{Q}_p$ -isogenies which are automorphisms of  $A_0[p^\infty]$ . Note that  $H(\mathbb{Z}_p)$  is a compact open subgroup of  $H(\mathbb{Q}_p)$ . Since  $A_0$  is hypersymmetric, the group  $H(\mathbb{Z}_p)$  is canonically isomorphic to the group  $\mathrm{Aut}((A_0, \lambda)[p^\infty])$  of automorphisms of the polarized  $p$ -divisible group  $(A_0, \lambda_0)[p^\infty]$ .

5.9. THEOREM (notation as in §§5.7 and 5.8). *In particular,  $k \supset \mathbb{F}_p$  is an algebraically closed field,  $\gcd(n, p) = 1$ ,  $n \geq 3$ , and  $C$  is a leaf in  $\mathcal{A}_{g,d,n}$  over  $k$  such that the  $p$ -divisible of the universal abelian scheme  $A \rightarrow C$  is completely slope divisible.*

- (i) *For any two  $k$ -points  $y_1, y_2$  of  $\mathcal{T}_b$  above  $x_0$ , there exists an element  $\delta$  in the group  $\mathrm{Sp}(\mathrm{H}_1(A_0, \mathbb{A}_f^{(p)}), \langle, \rangle)$  such that  $y_1$  belongs to the image of  $y_2$  under the algebraic correspondence induced by  $\delta$ . In particular, the Hecke correspondences induced by elements of  $\mathrm{Sp}(\mathrm{H}_1(A_0, \mathbb{A}_f^{(p)}), \langle, \rangle)$  operates transitively on the set  $\Pi_0(\mathcal{T}_b)$  of irreducible components of  $\mathcal{T}_b$  for every positive integer  $b$ .*
- (ii) *The  $L_{\mathbb{Z}/p^n\mathbb{Z}}$ -torsor  $\mathcal{T}_b$  over  $k$  is irreducible for every  $b \geq 1$ . Therefore the  $p$ -adic monodromy homomorphism*

$$\rho_{(A,\lambda)[p^\infty]/C,x_0} : \pi_1(C, x_0) \longrightarrow \mathrm{Aut}((A_0, \lambda_0)[p^\infty])$$

*attached to the leaf  $C$  in  $\mathcal{A}_{g,d,n}$  over  $k$  is surjective.*

*Proof.* The first assertion in statement (ii) follows from (i) as in [Theorem 2.13](#) by the argument of [[4](#), Prop. 4.4]; see also the proof of Proposition 5.5 in [[5](#)]. The second part of (ii) follows from the first part.

As indicated in [[5](#), 6.3.5], the methods in [[5](#), §5] can be used to give a proof of statement (i). Write  $y_1 = [(A_0, \lambda_0, \iota, \psi_b)]$   $y_2 = [(A_0, \lambda_0, \iota, \psi'_b)]$  in the notation of [Section 5.7](#), where the  $\iota$  and  $\iota'$  are symplectic level- $n$  structures on  $(A_0, \lambda_0)$  and  $\psi_b, \psi'_b$  are automorphisms of  $(A_0, \lambda_0)[p^b]$  induced by automorphisms of  $(A_0, \lambda_0)[p^\infty]$ . Pick a point  $z = [(A_0, \lambda_0, \tilde{\iota}, \psi_b)]$  of  $\mathcal{T}^\sim$  above  $y_1$ . Then  $z' = [(A_0, \lambda_0, \tilde{\iota}, \psi'_b)]$  is a point above  $y_2$ . Denote by  $H(\mathbb{Z}_{(p)})$  the intersection  $H(\mathbb{Q}) \cap H(\mathbb{Z}_p)$  in  $H(\mathbb{Q}_p)$ . Each element  $\gamma \in H(\mathbb{Z}_{(p)})$  can be regarded as a prime-to- $p$  symplectic isogeny from  $A_0$  to itself. Moreover,  $\gamma$  induces an isomorphism from  $(A_0, \lambda_0, \tilde{\iota}, \psi_b)$  to  $(A_0, \lambda_0, \gamma^{(p)} \circ \tilde{\iota}, \gamma_{p,b} \circ \psi_b)$ , where  $\gamma^{(p)}$  is the automorphism of  $H_1(A_0, \mathbb{A}_f^{(p)})$  induced by  $\gamma$  and  $\gamma_{p,b}$  is the automorphism of  $(A_0, \lambda_0)[p^b]$  induced by  $\gamma$ . The weak approximation theorem holds for the reductive group  $H$  over  $\mathbb{Q}$ ; see [Lemma 4.6](#). In particular there exists an element  $\xi \in H(\mathbb{Z}_{(p)})$  such that  $\xi_{p,b} \circ \psi_b = \psi'_b$ . Let  $\delta$  be the element of  $\text{Sp}(H_1(A_0, \mathbb{A}_f^{(p)}), \langle, \rangle)$  such that  $\xi^{(p)} \circ \tilde{\iota} = \tilde{\iota} \circ \delta$ . Then we have

$$z = [(A_0, \lambda_0, \xi^{(p)} \circ \tilde{\iota}, \psi'_b)] = z' \cdot \delta,$$

so  $y_1$  is an image of  $y_2$  under the prime-to- $p$  Hecke correspondence induced by  $\delta$ . We have proved that the image of the  $p$ -adic monodromy for  $C$  is equal to  $L_{\mathbb{Z}}$ . □

We formulate [Proposition 5.11](#) and the related [Lemma 5.10](#), convenient for the exposition of the proof of [Theorem 5.6](#).

5.10. LEMMA (Zink). *Let  $Y$  be a  $p$ -divisible group over a field  $K \supset \mathbb{F}_p$  and let  $L$  be a field containing  $K$ . Then there exist isogenies  $\alpha : X \rightarrow Y$  and  $\beta : Y \rightarrow Z$  with  $X, Z$  completely slope divisible, with the following properties:*

- (1) *Every isogeny  $\alpha' : X' \rightarrow Y$  of  $p$ -divisible groups with  $X'$  completely slope divisible factors through  $\alpha$ ; i.e., there exist an isogeny  $\delta : X' \rightarrow X$  such that  $\alpha' = \alpha \circ \delta$ . Similarly, every isogeny  $\beta' : Y \rightarrow Z'$  with  $Z'$  completely slope divisible factors through  $\beta$ ; i.e., there exists an isogeny  $\epsilon : Z \rightarrow Z'$  such that  $\beta' = \epsilon \circ \beta$ . These properties characterize  $\alpha$  and  $\beta$  respectively. Moreover, these properties hold after extension of base fields  $K \rightarrow L$ .*
- (2) *The isogeny  $\alpha$  induces an inclusion*

$$\text{End}(X \times_{\text{Spec}(K)} \text{Spec}(L)) \supset \text{End}(Y \times_{\text{Spec}(K)} \text{Spec}(L));$$

*the isogeny  $\beta$  induces an inclusion*

$$\text{End}(Y \times_{\text{Spec}(K)} \text{Spec}(L)) \subset \text{End}(Z \times_{\text{Spec}(K)} \text{Spec}(L)).$$

*Proof.* The construction of  $\beta : Y \rightarrow X$  is implicit in the proof of [[35](#), Th. 7]; it is given by an inductive procedure. We recall briefly the construction. In

Lemma 9 of [35], a “formula” for  $\beta$  is given in terms of Dieudonné modules when the base field  $K$  is perfect and  $Y$  is isoclinic. This construction in terms of Dieudonné modules can be reformulated in terms of the “small image” of a suitable homomorphism between  $p$ -divisible groups, which settles the case when  $Y$  is isoclinic. An induction on the length of the slope filtration finishes the proof. See [35] for details.  $\square$

5.11. PROPOSITION (Zink) (notation as in Theorem 5.6). *There exists an abelian scheme  $B$  over the leaf  $C$  and a  $p$ -primary isogeny  $\beta: A \rightarrow B$  over  $C$  with the following properties:*

- (i) *The  $p$ -divisible group  $B[p^\infty] \rightarrow C$  is completely slope divisible.*
- (ii) *For every geometric point  $x \in C(\Omega)$ , the isogeny  $\beta_x: A_x \rightarrow B_x$  induces an inclusion*

$$\text{End}(A_x[p^\infty]) \subseteq \text{End}(B_x[p^\infty]).$$

*In particular,  $\beta_x$  induces an inclusion  $\text{End}(A_x) \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)} \subseteq \text{End}(B_x) \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}$ .*

*Proof.* Clearly it suffices to establish the existence of an isogeny

$$\beta[p^\infty]: A[p^\infty] \rightarrow B[p^\infty]$$

between  $p$ -divisible groups satisfying (i) and (ii). Applying Lemma 5.10 to the generic fiber of  $A[p^\infty] \rightarrow C$ , one obtains the desired  $p$ -primary isogeny  $\beta_{K(C)}: A[p^\infty]_{K(C)} \rightarrow B[p^\infty]_{K(C)}$  over the spectrum of the function field  $K(C)$  of  $C$ , which satisfies the properties in Lemma 5.10. Proposition 14 in [35] and the proof of Theorem 7 in [35] show that the  $p$ -divisible group  $B[p^\infty]$  over  $K(C)$  extends to a completely slope divisible  $p$ -divisible group  $B[p^\infty]$  over  $C$  and the isogeny over  $K(C)$  extends to  $C$ .

We know that the  $p$ -divisible groups  $A[p^\infty]$  over  $C$  is geometrically fiberwise constant by assumption, while the  $p$ -divisible group  $B[p^\infty]$  over  $C$  is geometrically fiberwise constant because it is completely slope divisible. Because  $C$  is irreducible, we conclude that any isogeny between  $A[p^\infty]$  and  $B[p^\infty]$  is also geometrically fiberwise constant. This implies that the same procedure, which produced the isogeny  $\beta_{K(C)}$  over the function field  $K(C)$  from the  $p$ -divisible group  $A[p^\infty]$  over  $K(C)$ , also produces the fiber  $\beta_y$  of  $\beta$  at  $y$  when applied to the fiber  $A[p^\infty]_y$  of  $A[p^\infty]$ , for every geometric point  $y \in C(k)$ .  $\square$

5.12. (choose notation as in §§5.7 and 5.8). Let  $\beta: A \rightarrow B$  be as in Proposition 5.11 and let  $\beta_0: A_0 \rightarrow B_0$  be the fiber of  $\beta$  over  $x_0$ . Let  $\mu$  be a polarization of  $B$  over  $C$  such that  $\beta^*(\mu)$  is a multiple of  $\lambda$ . Let  $\mu_0$  be the fiber of  $\mu$  at  $x_0$ , a polarization of  $B_0$ , and let  $\iota'_0$  be the level- $n$  structure on  $B_0$  induced by  $\beta_0$ . Write  $x_0 = (A_0, \lambda_0, \iota_0)$ ,  $y_0 = (B_0, \lambda'_0, \iota'_0)$ . Let

$$\rho_{(B, \lambda') [p^\infty]}: \pi_1(C, x_0) \rightarrow \text{Aut}((B_0, \lambda'_0)[p^\infty])$$

be the  $p$ -adic monodromy homomorphism attached to  $(B, \lambda') \rightarrow C$ .



We use the definition of  $p$ -adic monodromy in [Section 5.4](#). By the property of  $\beta$  stated in [Proposition 5.11](#), the isogeny  $\beta_0$  identifies  $\text{Aut}((A_0, \lambda_0)[p^\infty])$  with a subgroup of  $\text{Aut}(B_0, \mu_0)$ . We know that the image of  $\rho_{(B, \mu)}$  is contained in the subgroup  $\text{Aut}((A_0, \lambda_0)[p^\infty])$  of  $\text{Aut}(B_0, \mu_0)$ . We must show that image of  $\rho_{(B, \mu)}$  is equal to  $\text{Aut}((A_0, \lambda_0)[p^\infty])$ .

Let  $H$  be the unitary group over  $\mathbb{Q}$  attached to

$$(\text{End}^0(A_0), *_{\lambda_0}) = (\text{End}^0(B_0), *_{\mu_0}).$$

Let

$$L_{\mathbb{Z}_p} = L(A_0, \lambda_0)_{\mathbb{Z}_p} := \text{Aut}((A_0, \lambda_0)[p^\infty])$$

and

$$L'_{\mathbb{Z}_p} := L(B_0, \mu_0)_{\mathbb{Z}_p} := \text{Aut}((B_0, \mu_0)[p^\infty]).$$

Let  $L'_{\mathbb{Z}/b\mathbb{Z}}$  be the image of  $L'_{\mathbb{Z}_p}$  in  $\text{Aut}((B_0, \mu)[p^b])$  and let  $L_{\mathbb{Z}/b\mathbb{Z}}$  be the image of  $L_{\mathbb{Z}_p}$  in  $\text{Aut}((A_0, \lambda)[p^b])$ . So  $L_{\mathbb{Z}_p}$  and  $L'_{\mathbb{Z}_p}$  are both compact open subgroups of  $H(\mathbb{Q}_p)$ .

5.13. The  $p$ -adic monodromy representation attached to the polarized  $p$ -divisible group  $(B[p^\infty], \mu) \rightarrow C$  comes from a right  $L'_{\mathbb{Z}_p}$ -torsor  $\mathcal{T}$  over  $C$  defined the same way as before. As before, the right  $L'_{\mathbb{Z}_p}$ -torsor  $\mathcal{T} \rightarrow C$  is the projective of a tower of right  $L'_{\mathbb{Z}/b\mathbb{Z}}$ -torsors  $\mathcal{T}_b$  and the transition morphisms  $\mathcal{T}_{b_1} \rightarrow \mathcal{T}_b$  are finite étale surjective if  $b|b_1$ .

The set of  $k$ -points  $\mathcal{T}$  can be described as follows. It consists of all isomorphism classes of quadruples  $(A_x, \lambda_x, \iota_x, \psi)$ , where  $[(A_x, \lambda_x, \iota_x)] =: x$  is a  $k$ -point of  $C$  and

$$\psi: (B_0, \mu_0)[p^\infty] \xrightarrow{\sim} (B_x, \mu_x)[p^\infty]$$

is an isomorphism. In the above  $B_x$  (resp.  $\mu_x$ ) denotes the fiber of  $B$  (resp.  $\mu$ ) above the point  $x \in C$ . The set of  $k$ -points of  $\mathcal{T}_b$  and the set of  $S$ -points of  $\mathcal{T}_b$  of a  $k$ -scheme  $S$  can be described explicitly in a similar way as in [Sections 5.7](#) and [5.8](#).

We know that the image of the  $p$ -adic monodromy representation is contained in  $L_{\mathbb{Z}_p}$ , so the right  $L'_{\mathbb{Z}_p}$ -torsor  $\mathcal{T}$  over  $C$  is of the form  $\mathcal{S} \times^{L_{\mathbb{Z}_p}} L'_{\mathbb{Z}_p}$ , the push-forward via  $L_{\mathbb{Z}_p} \rightarrow L'_{\mathbb{Z}_p}$  of a right  $L_{\mathbb{Z}_p}$ -torsor  $\mathcal{S}$  over  $C$ . As before, the  $L_{\mathbb{Z}_p}$ -torsor  $\mathcal{S} \rightarrow C$  is a projective limit of  $L_{\mathbb{Z}/b\mathbb{Z}}$ -torsors  $\mathcal{S}_b \rightarrow C$ . Below are explicit descriptions of  $\mathcal{S}$ .

(I)  $k$ -points of  $\mathcal{S}$ .

The set of  $k$ -points of the  $L_{\mathbb{Z}_p}$ -torsor  $\mathcal{S} \rightarrow C$  is the set of isomorphism classes of quadruples  $(A_x, \lambda_x, \iota_x, \psi)$ , where

- $[(A_x, \lambda_x, \iota_x)] =: x$  is a  $k$ -point of  $C$ , and
- $\psi: (B_0, \mu_0)[p^\infty] \xrightarrow{\sim} (B_x, \mu_x)[p^\infty]$  is an isomorphism of polarized  $p$ -divisible groups which induces an isomorphism  $(A_0, \lambda_0)[p^\infty] \xrightarrow{\sim} (A_x, \lambda_x)[p^\infty]$ .

(II)  $D$ -points of  $\mathcal{S}$  for a  $k$ -scheme  $D$ .

It is more convenient to describe the set  $\mathcal{S}_b(D)$  of  $D$ -points for members  $\mathcal{S}_b$  in the tower  $(\mathcal{S}_b)_{b \geq 1}$  whose limit is  $\mathcal{S}$ . The set  $\mathcal{S}_b(D)$  consists of all isomorphism classes of quadruples

$$(A_D \rightarrow D, \lambda_D, \iota_D, \psi_b),$$

where  $(A \rightarrow D, \lambda_D, \iota_D)$  defines a  $D$ -point of the leaf  $C$  and

$$\psi_b : \prod_{i=1}^m (X_i/X_{i-1})[p^b] \times_{\text{Spec}(\mathbb{F}_q)} D \xrightarrow{\sim} \prod_{i=1}^m (Z_i/Z_{i-1})[p^b] \times_C D$$

is an isomorphism over  $D$  satisfying the following properties:

- (1) The isomorphism  $\psi_b$  respects the polarizations induced by  $\mu_0$  and  $\mu_D$  respectively.
- (2) For every geometric point  $y \in D$ , there exist isomorphisms

$$\psi_y : (B_0, \mu_0)[p^\infty] \xrightarrow{\sim} (B_y, \mu_y)[p^\infty] \quad \text{and} \quad \phi_y : (A_0, \lambda_0)[p^\infty] \xrightarrow{\sim} (A_y, \mu_y)[p^\infty]$$

such that  $\psi_y \circ \beta_0 = \beta_y \circ \phi_y$  and the fiber of  $\psi_b$  at  $y$  is induced by  $\psi_y$ .

- (3) For every integer  $N > b$ , there exists an isomorphism

$$\psi_N : \prod_{i=1}^m (X_i/X_{i-1})[p^N] \times_{\text{Spec}(\mathbb{F}_q)} D \xrightarrow{\sim} \prod_{i=1}^m (Z_i/Z_{i-1})[p^N] \times_C D$$

such that properties (1) and (2) hold with  $b$  replaced by  $N$ .

The following notation is used in the above.

- $\beta : A \rightarrow B$  is the isogeny over the leaf  $C$  in [Proposition 5.11](#),
- $(B_D, \mu_D)$  is the pull-back to  $D$  of the polarized abelian scheme  $(B, \mu)$  as in [Proposition 5.11](#),
- $0 = Z_0 \subset Z_1 \subset \dots \subset Z_m = B[p^\infty]$  is the slope filtration of the completely slope divisible  $p$ -divisible group  $B[p^\infty]$  over  $C$ ,
- $0 = X_0 \subset X_1 \subset \dots \subset X_m = B_0[p^\infty]$  is the slope filtration of the completely slope divisible  $p$ -divisible group  $B_0[p^\infty]$ .

Write  $L'_{\mathbb{Z}_p} = \coprod_{i=1}^m (L_{\mathbb{Z}_p} \cdot \zeta_i)$ , where  $\zeta_1 = 1, \dots, \zeta_m$  is a system of representatives of  $L_{\mathbb{Z}_p} \setminus L'_{\mathbb{Z}_p}$ , with  $\zeta_1 = 1$ . Then

$$L'_{\mathbb{Z}_p} = \coprod_{i=1}^m \mathcal{S} \times^{L_{\mathbb{Z}_p}} (L_{\mathbb{Z}_p} \cdot \zeta_i),$$

a disjoint union of contraction products, and  $\mathcal{S}$  is equal to  $\mathcal{S} \times^{L_{\mathbb{Z}_p}} (L_{\mathbb{Z}_p} \cdot \zeta_1)$ .

*Proof of Theorem 5.6.* We need to show that the  $L_{\mathbb{Z}_p}$ -torsor  $\mathcal{S}$  is irreducible, or equivalently each of the  $L_{\mathbb{Z}/b\mathbb{Z}}$ -torsors  $\mathcal{S}_b \rightarrow C$  is irreducible.

The group  $\text{Sp}(\mathbb{H}_1(A_0, \mathbb{A}_f^{(p)}), \langle, \rangle) \cong \text{Sp}(\mathbb{H}_1(B_0, \mathbb{A}_f^{(p)}), \langle, \rangle)$  operates on the torsor  $\mathcal{S} = \varprojlim \mathcal{S}_b$ , inducing prime-to- $p$  Hecke correspondences on each  $\mathcal{S}_b$ .

The argument of [Theorem 5.9](#) applies to the present situation and shows that any two points  $y_1, y_2$  of  $\mathcal{S}_b$  lying above the hypersymmetric point  $x_0$  belong to the same prime-to- $p$  Hecke orbit. Suppose that  $y_i = [(A_0, \lambda_0, \iota_0, \psi_{i,b})]$ ,  $i = 1, 2$ , where  $\psi_{i,b}: (B_0, \mu_0)[p^b] \xrightarrow{\sim} (B_0, \mu_0)[p^b]$  is the image of an element  $\psi_i \in L_{\mathbb{Z}_p}$ , and  $\psi_2 \in L_{\mathbb{Z}_p} \cdot \psi_1$ . Let  $\tilde{\iota}$  be a prime-to- $p$  symplectic level structure of  $(A_0, \lambda_0)$  above  $\iota_0$ . Let  $U_b$  be the open subset of  $L_{\mathbb{Z}_p}$  consisting of all elements of  $\gamma_p \in L_{\mathbb{Z}_p}$  such that the image of  $\gamma_p \circ \psi_1$  in  $L'_{\mathbb{Z}/b\mathbb{Z}}$  is  $\psi_{2,b}$ . By the weak approximation theorem applied to the connected  $\mathbb{Q}$ -group  $H$ , there exists an element  $\gamma \in H(\mathbb{Q})$  such that the image of  $\gamma \in H(\mathbb{Q}_p)$  belongs to  $U_b$ . Let  $\delta$  be the element of  $\text{Sp}(H_1(A_0, \mathbb{A}_f^{(p)}), \langle, \rangle)$  such that  $\gamma^{(p)} \circ \tilde{\iota} = \tilde{\iota} \circ \delta$ . Then  $y_1$  is an image of  $y_2$  under the prime-to- $p$  Hecke correspondence induced by  $\delta$ . By [Theorem 2.13](#), we conclude that  $\mathcal{S}_b$  is irreducible for every  $b \geq 1$ . Therefore  $\mathcal{S}$  is irreducible and the image of the  $p$ -adic monodromy homomorphism  $\rho_{(A,\lambda)[p^\infty]/C,x_0}$  is equal to  $\text{Aut}((A_0, \lambda_0)[p^\infty])$ . This finishes the proof of [Theorem 5.6](#).  $\square$

5.14. *Remark.* As an alternative proof of [Theorem 5.6](#), we indicate how [5.6](#) can be deduced from its special case, [Theorem 5.9](#). Let  $\alpha: B \rightarrow A$  be a  $p$ -primary isogeny which is a multiple of the quasi-isogeny  $\beta^{-1}$  and let  $\mu' := \alpha^*(\lambda)$ . Let  $d' > 0$  be the square root of the degree of  $\mu'$ . Let  $\mu'_0$  be the fiber of  $\mu'$  at  $x_0 \in C$  and let  $\iota'_0$  be the symplectic level- $n$  structure on  $B_0$  corresponding to the level- $n$  structure on  $A_0$  under the  $p$ -primary isogeny  $\alpha$ . Let  $y_0$  be the  $k$ -point  $[(B_0, \mu'_0, \iota'_0)]$  in  $\mathcal{A}_{g,d',n}$ , and let  $C' = \mathcal{C}(y_0)$  be the leaf in  $\mathcal{A}_{g,d',n}$  passing through  $y_0$ . Let  $(B', \nu) \rightarrow C'$  be the universal polarized abelian scheme over the leaf  $C'$  in  $\mathcal{A}_{g,d',n}$  and let  $(B', \nu)[p^\infty] \rightarrow C'$  be the associated polarized  $p$ -divisible group. Let  $C'_{\text{perf}}$  be the perfection of  $C'$ .

We know from [Theorem 4.1](#) that the leaf  $C'$  is (geometrically) irreducible. By [[29](#), Prop. 1.3], the completely slope divisible group  $B'[p^\infty] \times_{C'} C'_{\text{perf}} \rightarrow C'_{\text{perf}}$  over  $C'_{\text{perf}}$  splits into a direct sum of isoclinic completely slope divisible groups with distinct slopes. From [Theorem 5.9](#) we know that the  $p$ -adic monodromy for  $(B', \mu')[p^\infty] \rightarrow C'$  is equal to  $L'_{\mathbb{Z}_p} = \text{Aut}((B'_0, \mu'_0)[p^\infty])$ . According to the theory of  $\Phi$ -étale part, there exists an *irreducible* profinite étale Galois cover  $\mathcal{Z}$  of  $C'_{\text{perf}}$  with group  $L'_{\mathbb{Z}_p}$  such that  $(B', \mu')[p^\infty] \times_{C'_{\text{perf}}} \mathcal{Z}$  is a constant polarized  $p$ -divisible group over  $\mathcal{Z}$ .

By [Proposition 5.11](#), the kernel of the isogeny  $\alpha_0: B_0 \rightarrow A_0$  is stable under the subgroup  $L_{\mathbb{Z}_p} = \text{Aut}((A_0, \lambda_0)[p^\infty])$  of  $L_{\mathbb{Z}_p}$ . Let  $D = \mathcal{Z}/L_{\mathbb{Z}_p}$  be the finite étale cover of  $C'_{\text{perf}}$  corresponding to the subgroup  $L_{\mathbb{Z}_p}$ . On  $D$  we have a  $p$ -primary isogeny  $\alpha': B' \times_{C'} D \rightarrow A'$  such that the fiber  $\alpha'_{y_0}$  of  $\alpha'$  at  $y_0$  is equal to  $\alpha_{x_0}$ . Moreover, the polarization  $\nu$  on  $B' \times_{C'} D \rightarrow D$  descends to a polarization  $\lambda'$  on  $A' \rightarrow D$  such that  $(\alpha')^*(\lambda') = \nu$  and the fiber of  $\lambda'$  at  $y_0$  is equal to  $\lambda'_0$ .

The polarized abelian scheme  $(A', \lambda') \rightarrow D$  defines a morphism  $D \rightarrow \mathcal{A}_{g,d,n}$  which factorizes as  $D \rightarrow C \hookrightarrow \mathcal{A}_{g,d,n}$  by construction, such that  $(A', \lambda')$  is the pull-back to  $D$  of the universal polarized abelian scheme over  $C$ . By functoriality of the fundamental group, the  $p$ -adic monodromy for the leaf  $C$  contains the  $p$ -adic monodromy attached to the polarized  $p$ -divisible group  $(A', \lambda')[p^\infty]$  over  $D$ . Since the latter is equal to  $L_{\mathbb{Z}_p}$  by construction, the image of the  $p$ -adic monodromy representation for  $C$  contains  $L_{\mathbb{Z}_p}$ , hence is equal  $L_{\mathbb{Z}_p}$ .

## References

- [1] C.-L. CHAI, *Compactification of Siegel Moduli Schemes*, *London Math. Soc. Lect. Note Series* **107**, Cambridge Univ. Press, Cambridge, 1985. [MR 0853543](#). [Zbl 0578.14009](#).
- [2] ———, Every ordinary symplectic isogeny class in positive characteristic is dense in the moduli, *Invent. Math.* **121** (1995), 439–479. [MR 1353306](#). [Zbl 0990.11039](#). [doi: 10.1007/BF01884309](#).
- [3] ———, Hecke orbits on Siegel modular varieties, in *Geometric Methods in Algebra and Number Theory*, *Progr. Math.* **235**, Birkhäuser, Boston, MA, 2005, pp. 71–107. [MR 2159378](#). [Zbl 1143.14021](#). [doi: 10.1007/0-8176-4417-2\\_4](#).
- [4] ———, Monodromy of Hecke-invariant subvarieties, *Pure Appl. Math. Q.* **1** (2005), 291–303. [MR 2194726](#). [Zbl 1140.11031](#).
- [5] ———, Methods for  $p$ -adic monodromy, *J. Inst. Math. Jussieu* **7** (2008), 247–268. [MR 2400722](#). [Zbl 1140.14019](#). [doi: 10.1017/S1474748007000199](#).
- [6] C.-L. CHAI and F. OORT, Hypersymmetric abelian varieties, *Pure Appl. Math. Q.* **2** (2006), 1–27. [MR 2217565](#). [Zbl 1156.14320](#).
- [7] ———, Moduli of abelian varieties and  $p$ -divisible groups, in *Arithmetic Geometry*, *Clay Math. Proc.* **8**, Amer. Math. Soc., Providence, RI, 2009, pp. 441–536. [MR 2498069](#). [Zbl 1188.14028](#).
- [8] A. J. DE JONG and F. OORT, Purity of the stratification by Newton polygons, *J. Amer. Math. Soc.* **13** (2000), 209–241. [MR 1703336](#). [Zbl 0954.14007](#). [doi: 10.1090/S0894-0347-99-00322-7](#).
- [9] T. EKEDAHL and G. VAN DER GEER, Cycle classes of the E-O stratification on the moduli of abelian varieties, in *Algebra, Arithmetic, and Geometry: in Honor of Yu. I. Manin. Vol. I*, *Progr. Math.* **269**, Birkhäuser, Boston, MA, 2009, pp. 567–636. [MR 2641181](#). [Zbl 1200.14089](#). [doi: 10.1007/978-0-8176-4745-2\\_13](#).
- [10] G. FALTINGS, Arithmetische Kompaktifizierung des Modulraums der abelschen Varietäten, in *Workshop Bonn 1984*, *Lecture Notes in Math.* **1111**, Springer-Verlag, New York, 1985, pp. 321–383. [MR 0797429](#). [doi: 10.1007/BFb0084598](#).
- [11] G. FALTINGS and C.-L. CHAI, *Degeneration of Abelian Varieties*, *Ergeb. Math. Grenzgeb.* **22**, Springer-Verlag, New York, 1990. [MR 1083353](#). [Zbl 0744.14031](#).
- [12] A. GROTHENDIECK, *Groupes de Barsotti-Tate et Cristaux de Dieudonné*, Les Presses de l'Université de Montréal, Montreal, Que., 1974, Séminaire de Mathématiques Supérieures, No. 45 (Été, 1970). [MR 0417192](#). [Zbl 0331.14021](#).

- [13] N. M. KATZ, Slope filtration of  $F$ -crystals, in *Journées de Géométrie Algébrique de Rennes, Vol. I, Astérisque* **63**, Soc. Math. France, Paris, 1979, pp. 113–163. [MR 0563463](#). [Zbl 0426.14007](#).
- [14] K.-Z. LI and F. OORT, *Moduli of Supersingular Abelian Varieties, Lecture Notes in Math.* **1680**, Springer-Verlag, New York, 1998. [MR 1611305](#). [Zbl 0920.14021](#).
- [15] J. I. MANIN, Theory of commutative formal groups over fields of finite characteristic, *Uspehi Mat. Nauk* **18** (1963), 3–90. [MR 0157972](#). [Zbl 0128.15603](#).
- [16] D. MUMFORD, *Abelian Varieties, Tata Inst. Fund. Res. Stud. Math.* **5**, Published for the Tata Institute of Fundamental Research, Bombay, 1970. [MR 0282985](#). [Zbl 0223.14022](#).
- [17] P. NORMAN and F. OORT, Moduli of abelian varieties, *Ann. of Math.* **112** (1980), 413–439. [MR 0595202](#). [Zbl 0483.14010](#). doi: [10.2307/1971152](#).
- [18] T. ODA and F. OORT, Supersingular abelian varieties, in *Proceedings of the International Symposium on Algebraic Geometry* (Kyoto Univ., Kyoto, 1977), Kinokuniya Book Store, Tokyo, 1978, pp. 595–621. [MR 0578876](#). [Zbl 0402.14016](#).
- [19] F. OORT, Monodromy, Hecke orbits and Newton polygon strata, Manuscript, Seminar Algebraic Geometry, (Bonn, 24 - II - 2003); 9 pp. Available at <http://www.math.uu.nl/~oort0109/>.
- [20] ———, *Commutative Group Schemes, Lecture Notes in Math.* **15**, Springer-Verlag, New York, 1966. [MR 0213365](#). [Zbl 0216.05603](#).
- [21] ———, Endomorphism algebras of abelian varieties, in *Algebraic Geometry and Commutative Algebra, Vol. II*, Kinokuniya, Tokyo, 1988, pp. 469–502. [MR 0977774](#). [Zbl 0697.14029](#).
- [22] ———, Newton polygons and formal groups: conjectures by Manin and Grothendieck, *Ann. of Math.* **152** (2000), 183–206. [MR 1792294](#). [Zbl 0991.14016](#). doi: [10.2307/2661381](#).
- [23] ———, Newton polygon strata in the moduli space of abelian varieties, in *Moduli of Abelian Varieties, Progr. Math.* **195**, Birkhäuser, Basel, 2001, pp. 417–440. [MR 1827028](#). [Zbl 1086.14037](#).
- [24] ———, A stratification of a moduli space of abelian varieties, in *Moduli of Abelian Varieties, Progr. Math.* **195**, Birkhäuser, Basel, 2001, pp. 345–416. [MR 1827027](#). [Zbl 1052.14047](#).
- [25] ———, Foliations in moduli spaces of abelian varieties, *J. Amer. Math. Soc.* **17** (2004), 267–296. [MR 2051612](#). [Zbl 1041.14018](#). doi: [10.1090/S0894-0347-04-00449-7](#).
- [26] ———, Minimal  $p$ -divisible groups, *Ann. of Math.* **161** (2005), 1021–1036. [MR 2153405](#). [Zbl 1081.14065](#). doi: [10.4007/annals.2005.161.1021](#).
- [27] ———, Abelian varieties over finite fields, in *Higher-Dimensional Geometry over Finite Fields, NATO Sci. Peace Secur. Ser. D Inf. Commun. Secur.* **16**, IOS, Amsterdam, 2008, pp. 123–188. [MR 2484079](#). [Zbl 1181.14029](#).
- [28] ———, Foliations in moduli spaces of abelian varieties and dimension of leaves, in *Algebra, Arithmetic, and Geometry: In Honor of Yu. I. Manin. Vol. II, Progr. Math.* **270**, Birkhäuser, Boston, MA, 2009, pp. 465–501. [MR 2641199](#). [Zbl 05778264](#). doi: [10.1007/978-0-8176-4747-6\\_15](#).

- [29] F. OORT and T. ZINK, Families of  $p$ -divisible groups with constant Newton polygon, *Doc. Math.* **7** (2002), 183–201. [MR 1938119](#). [Zbl 1022.14013](#).
- [30] V. PLATONOV and A. RAPINCHUK, *Algebraic Groups and Number Theory*, *Pure Appl. Math.* **139**, Academic Press, Boston, MA, 1994. [MR 1278263](#). [Zbl 0841.20046](#).
- [31] M. RAPOPORT and T. ZINK, *Period Spaces for  $p$ -Divisible Groups*, *Ann. of Math. Stud.* **141**, Princeton Univ. Press, Princeton, NJ, 1996. [MR 1393439](#). [Zbl 0873.14039](#).
- [32] J. TATE, Classes d'isogénie de variétés abéliennes sur un corps fini (d'après Z. Honda), in *Sém. Bourbaki* **21**, 1968/69, Exp. 352, *Lecture Notes Math.* **179**, 1971, pp. 95–110. [Zbl 0212.25702](#).
- [33] W. C. WATERHOUSE and J. S. MILNE, Abelian varieties over finite fields, in 1969 *Number Theory Institute*, Amer. Math. Soc., Providence, R.I., 1971, pp. 53–64. [MR 0314847](#). [Zbl 0216.33102](#).
- [34] C.-F. YU, Discrete Hecke orbit problem on Hilbert-Blumenthal modular varieties, preprint.
- [35] T. ZINK, On the slope filtration, *Duke Math. J.* **109** (2001), 79–95. [MR 1844205](#). [Zbl 1061.14045](#). doi: [10.1215/S0012-7094-01-10913-7](https://doi.org/10.1215/S0012-7094-01-10913-7).
- [36] Y. ZONG, On hypersymmetric abelian varieties, Ph.D. dissertation, Univ. of Pennsylvania, 2008. Available at <http://www.math.upenn.edu/grad/dissertations/YingZongThesis.pdf>.

(Received: February 22, 2008)

(Revised: July 16, 2009)

UNIVERSITY OF PENNSYLVANIA, PHILADELPHIA, PA

*E-mail*: [chai@math.upenn.edu](mailto:chai@math.upenn.edu)

<http://www.math.upenn.edu/~chai/>

UNIVERSITY OF UTRECHT, UTRECHT (STAD), NETHERLANDS

*E-mail*: [F.Oort@uu.nl](mailto:F.Oort@uu.nl)

<http://www.math.uu.nl/~oort0109/>