

On a problem posed by Steve Smale

By PETER BÜRGISSER and FELIPE CUCKER

Abstract

The 17th of the problems proposed by Steve Smale for the 21st century asks for the existence of a deterministic algorithm computing an approximate solution of a system of n complex polynomials in n unknowns in time polynomial, on the average, in the size N of the input system. A partial solution to this problem was given by Carlos Beltrán and Luis Miguel Pardo who exhibited a randomized algorithm doing so. In this paper we further extend this result in several directions. Firstly, we exhibit a linear homotopy algorithm that efficiently implements a nonconstructive idea of Mike Shub. This algorithm is then used in a randomized algorithm, call it LV, à la Beltrán-Pardo. Secondly, we perform a smoothed analysis (in the sense of Spielman and Teng) of algorithm LV and prove that its smoothed complexity is polynomial in the input size and σ^{-1} , where σ controls the size of the random perturbation of the input systems. Thirdly, we perform a condition-based analysis of LV. That is, we give a bound, for each system f , of the expected running time of LV with input f . In addition to its dependence on N this bound also depends on the condition of f . Fourthly, and to conclude, we return to Smale's 17th problem as originally formulated for deterministic algorithms. We exhibit such an algorithm and show that its average complexity is $N^{O(\log \log N)}$. This is nearly a solution to Smale's 17th problem.

Contents

1. Introduction	1786
Acknowledgments	1791
2. Preliminaries	1791
2.1. Setting and notation	1791
2.2. Newton's method	1793
2.3. Condition numbers	1793
2.4. Gaussian distributions	1794

P.B. was partially supported by DFG grant BU 1371/2-1 and Paderborn Institute for Scientific Computation (PaSCo). F. C. was partially supported by GRF grant CityU 100810.

An extended abstract of this work was presented at STOC 2010 under the title "Solving Polynomial Equations in Smoothed Polynomial Time and a Near Solution to Smale's 17th Problem".

3.	Statement of main results	1794
3.1.	The homotopy continuation routine ALH	1794
3.2.	Randomization and complexity: the algorithm LV	1795
3.3.	Average analysis of LV	1796
3.4.	Smoothed analysis of LV	1797
3.5.	The main technical result	1798
3.6.	Condition-based analysis of LV	1798
3.7.	A near solution of Smale's 17th problem	1798
4.	Complexity analysis of ALH	1799
5.	A useful change of variables	1802
6.	Analysis of LV	1804
6.1.	Average-case analysis of LV (proof)	1804
6.2.	Smoothed analysis of LV (proof)	1806
7.	Smoothed analysis of a matrix condition number	1806
8.	Smoothed analysis of the mean square condition number	1810
8.1.	Orthogonal decompositions of \mathcal{H}_d	1810
8.2.	Outline of proof of Theorem 3.6	1811
8.3.	Coarea formula	1813
8.4.	Normal Jacobians	1814
8.5.	Induced probability distributions	1818
8.6.	Expected number of real zeros	1820
9.	Effective sampling in the solution variety	1822
10.	Homotopies with a fixed extremity	1823
10.1.	Condition-based analysis of LV (proof)	1826
10.2.	The complexity of a deterministic homotopy continuation	1826
11.	A near solution to Smale's 17th problem	1828
11.1.	The case $D \leq n$	1828
11.2.	The case $D > n$	1830
11.3.	Proof of Lemma 11.3	1833
	Note added in proof	1834
	References	1834

1. Introduction

In 2000, Steve Smale published a list of mathematical problems for the 21st century [29]. The 17th problem in the list reads as follows:

Can a zero of n complex polynomial equations in n unknowns be found approximately, on the average, in polynomial time with a uniform algorithm?

Smale pointed out that “it is reasonable” to homogenize the polynomial equations by adding a new variable and to work in projective space after which he made precise the different notions intervening in the question above. We

provide these definitions in full detail in Section 2. Before doing so, in the remainder of this section, we briefly describe the recent history of Smale's 17th problem and the particular contribution of the present paper. The following summary of notations should suffice for this purpose.

We denote by $\mathcal{H}_{\mathbf{d}}$ the linear space of complex homogeneous polynomial systems in $n + 1$ variables, with a fixed degree pattern $\mathbf{d} = (d_1, \dots, d_n)$. We let $D = \max_i d_i$, $N = \dim_{\mathbb{C}} \mathcal{H}_{\mathbf{d}}$, and $\mathcal{D} = \prod_i d_i$. We endow this space with the unitarily invariant Bombieri-Weyl Hermitian product and consider the unit sphere $S(\mathcal{H}_{\mathbf{d}})$ with respect to the norm induced by this product. We then make this sphere a probability space by considering the uniform measure on it. The expression "on the average" refers to expectation on this probability space. Also, the expression "approximate zero" refers to a point for which Newton's method, starting at it, converges immediately, quadratically fast.

This is the setting underlying the series of papers [22], [23], [24], [26], [25] — commonly referred to as "the Bézout series" — written by Shub and Smale during the first half of the 1990s, a collection of ideas, methods, and results that pervade all the research done in Smale's 17th problem since this was proposed. The overall idea in the Bézout series is to use a linear homotopy. That is, one starts with a system g and a zero ζ of g and considers the segment $E_{f,g}$ with extremities f and g . Here f is the system whose zero we want to compute. Almost surely, when one moves from g to f , the zero ζ of g follows a curve in projective space to end in a zero of f . The homotopy method consists of dividing the segment $E_{f,g}$ in a number, say k , of subsegments E_i small enough to ensure that an approximate zero x_i of the system at the origin of E_i can be made into an approximate zero x_{i+1} of the system at its end (via one step of Newton's method). The difficulty of this overall idea lies in the following issues:

- (1) How does one choose the initial pair (g, ζ) ?
- (2) How does one choose the subsegments E_i ? In particular, how large should k be?

The state of the art at the end of the Bézout series, i.e., in [25], showed an incomplete picture. For (2), the rule consisted of taking a regular subdivision of $E_{f,g}$ for a given k , executing the path-following procedure, and repeating with k replaced by $2k$ if the final point could not be shown to be an approximate zero of f . (Shub and Smale provided criteria for checking this.) Concerning (1), Shub and Smale proved that good initial pairs (g, ζ) (in the sense that the average number of iterations for the rule above was polynomial in the size of f) existed for each degree pattern \mathbf{d} , but they could not exhibit a procedure to generate one such pair.

The next breakthrough took a decade to come. Beltrán and Pardo proposed in [4], [5] that the initial pair (g, ζ) should be randomly chosen. The consideration of randomized algorithms departs from the formulation of Smale's

17th problem¹ but it is widely accepted that, in practical terms, such algorithms are as good as their deterministic siblings. And in the case at hand this departure turned out to pay off. The average (over f) of the expected (over (g, ζ)) number of iterations of the algorithm proposed in [5] is $\mathcal{O}(n^5 N^2 D^3 \log \mathcal{D})$. One of the most notable features of the ideas introduced by Beltrán and Pardo is the use of a measure on the space of pairs (g, ζ) which is friendly enough to perform a probabilistic analysis while, at the same time, does allow for efficient sampling.

Shortly after the publication of [4], [5] Shub wrote a short paper of great importance [21]. Complexity bounds in both the Bézout series and the Beltrán-Pardo results rely on condition numbers. Shub and Smale had introduced a measure of condition $\mu_{\text{norm}}(f, \zeta)$ for $f \in \mathcal{H}_{\mathbf{d}}$ and $\zeta \in \mathbb{C}^{n+1}$ which, in case ζ is a zero of f , quantifies how much ζ varies when f is slightly perturbed. Using this measure they defined the *condition number* of a system f by taking

$$(1.1) \quad \mu_{\max}(f) := \max_{\zeta|f(\zeta)=0} \mu_{\text{norm}}(f, \zeta).$$

The bounds mentioned above make use of an estimate for the worst-conditioned system along the segment $E_{f,g}$, that is, of the quantity

$$(1.2) \quad \max_{q \in E_{f,g}} \mu_{\max}(q).$$

The main result in [21] shows that there exists a partition of $E_{f,g}$ which successfully computes an approximate zero of f whose number k of pieces satisfies

$$(1.3) \quad k \leq CD^{3/2} \int_{q \in E_{f,g}} \mu_2^2(q) dq,$$

where C is a constant and $\mu_2(q)$ is the *mean square condition number* of q given by

$$(1.4) \quad \mu_2^2(q) := \frac{1}{\mathcal{D}} \sum_{\zeta|q(\zeta)=0} \mu_{\text{norm}}^2(q, \zeta).$$

This partition is explicitly described in [21], but no constructive procedure to compute the partition is given there.

¹In his description of Problem 17 Smale writes “Time is measured by the number of arithmetic operations and comparisons, \leq , using real machines (as in Problem 3)” and in the latter he points that, “In [Blum-Shub-Smale,1989] a satisfactory definition [of these machines] is proposed.” The paper [9] quoted by Smale deals exclusively with deterministic machines. Furthermore, Smale adds that “a probability measure must be put on the space of all such f , for each $\mathbf{d} = (d_1, \dots, d_n)$, and the time of an algorithm is averaged over the space of f .” Hence, the expression ‘average time’ refers to expectation over the input data only.

In an oversight of this nonconstructibility, Beltrán and Pardo [6] provided a new version of their randomized algorithm² with an improved complexity of $\mathcal{O}(D^{3/2}nN)$.

A first goal of this paper is to validate Beltrán and Pardo's analysis in [6] by exhibiting an efficiently constructible partition of $E_{f,g}$ which satisfies a bound like (1.3). Our way of doing so owes much to the ideas in [21]. The path-following procedure ALH relying on this partition is described in detail in Section 3.1 together with a result, Theorem 3.1, bounding its complexity as in (1.3).

The second goal of this paper is to perform a smoothed analysis of a randomized algorithm (essentially Beltrán-Pardo randomization plus ALH) computing a zero of f , which we call LV. What smoothed analysis is, is succinctly explained in the citation of the Gödel prize 2008 awarded to its creators, Daniel Spielman and Teng Shang-Hua.³

Smoothed Analysis is a novel approach to the analysis of algorithms. It bridges the gap between worst-case and average case behavior by considering the performance of algorithms under a small perturbation of the input. As a result, it provides a new rigorous framework for explaining the practical success of algorithms and heuristics that could not be well understood through traditional algorithm analysis methods.

In a nutshell, smoothed analysis is a probabilistic analysis which replaces the 'evenly spread' measures underlying the usual average-case analysis (uniform measures, standard normals, ...) by a measure centered at the input data. That is, it replaces the 'average data input' (an unlikely input in actual computations) by a small random perturbation of a worst-case data and substitutes the typical quantity studied in the average-case context,

$$\mathbb{E}_{f \sim \mathcal{R}} \varphi(f),$$

by

$$\sup_{\bar{f}} \mathbb{E}_{f \sim \mathcal{C}(\bar{f}, r)} \varphi(f).$$

²The algorithm in [6] explicitly calls as a subroutine "the homotopy algorithm of [21]" without noticing that the partition in [21] is nonalgorithmic. Actually, the word 'algorithm' is never used in [21]. The main goal of [21], as stated in the abstract, is to motivate "the study of short paths or geodesics in the condition metric" —the proof of (1.3) does not require the homotopy to be linear and one may wonder whether other paths in $\mathcal{H}_{\mathbf{a}}$ may substantially decrease the integral in the right-hand side. This goal has been addressed, but not attained, in [7]. As of today it remains a fascinating open problem.

³See <http://www.fmi.uni-stuttgart.de/ti/personen/Diekert/citation08.pdf> for the whole citation.

Here $\varphi(f)$ is the function of f one is interested in (e.g., the complexity of an algorithm over input f), \mathcal{R} is the ‘evenly spread’ measure mentioned above and $\mathcal{C}(\bar{f}, r)$ is an isotropic measure centered at \bar{f} with a dispersion (e.g., variance) given by a (small) parameter $r > 0$.

An immediate advantage of smoothed analysis is its robustness with respect to the measure \mathcal{C} (see §3.4 below). This is in contrast with the most common critique to average-case analysis: “A bound on the performance of an algorithm under one distribution says little about its performance under another distribution, and may say little about the inputs that occur in practice” [31].

The precise details of the smoothed analysis we perform for zero finding are in Section 3.4.

To describe the third goal of this paper we recall Smale’s ideas of complexity analysis as exposed in [28]. In this program-setting paper Smale writes that he sees “much of the complexity theory [...] of numerical analysis conveniently represented by a two-part scheme.” The first part amounts to obtain, for the running time $\text{time}(f)$ of an algorithm on input f , an estimate of the form

$$(1.5) \quad \text{time}(f) \leq K(\text{size}(f) + \mu(f))^c,$$

where K and c are positive constants and $\mu(f)$ is a condition number for f . The second takes the form

$$(1.6) \quad \text{Prob}\{\mu(f) \geq T\} \leq T^{-c},$$

“where a probability measure has been put on the space of inputs.” The first part of this scheme provides understanding on the behavior of the algorithm for specific inputs f (in terms of their condition as measured by $\mu(f)$). The second, combined with the first, allows one to obtain probability bounds for $\text{time}(f)$ in terms of $\text{size}(f)$ only. But these bounds say little about $\text{time}(f)$ for actual input data f .

Part one of Smale’s program is missing in the work related with his 17th problem. All estimates on the running time of path-following procedures for a given f occurring in both the Bézout series and the work by Beltrán and Pardo are expressed in terms of the quantity in (1.2) or the integral in (1.3), not purely in terms of the condition of f . We fill this gap by showing for the expected running time of LV a bound like (1.5) with $\mu(f) = \mu_{\max}(f)$. The precise statement, Theorem 3.7, is in Section 3.6 below.

Last but not least, to close this introduction, we return to its opening theme: Smale’s 17th problem. Even though randomized algorithms are efficient in theory and reliable in practice, they do not offer an answer to the question of the existence of a deterministic algorithm computing approximate

zeros of complex polynomial systems in average polynomial time. The situation is akin to the development of primality testing. It was precisely with this problem that randomized algorithms became a means to deal with apparently intractable problems [30], [17]. Yet, the eventual display of a deterministic polynomial-time algorithm [1] was justly welcomed as a major achievement. The fourth main result in this paper exhibits a deterministic algorithm computing approximate zeros in average time $N^{\mathcal{O}(\log \log N)}$. To do so we design and analyze a deterministic homotopy algorithm, call it MD, whose average complexity is polynomial in n and N and exponential in D . This already yields a polynomial-time algorithm when one restricts the degree D to be at most $n^{1-\varepsilon}$ for any fixed $\varepsilon > 0$ (and, in particular, when D is fixed as in a system of quadratic or cubic equations). Algorithm MD is fast when D is small. We complement it with an algorithm that uses a procedure proposed by Jim Renegar [18] and which computes approximate zeros similarly fast when D is large.

In order to prove the results described above we have relied on a number of ideas and techniques. Some of them —e.g., the use of the coarea formula or of the Bombieri-Weyl Hermitian inner product— are taken from the Bézout series and are pervasive in the literature on the subject. Some others —notably the use of the Gaussian distribution and its truncations in Euclidean space instead of the uniform distribution on a sphere or a projective space— are less common. The blending of these ideas has allowed us a development which unifies the treatment of the several situations we consider for zero finding in this paper.

Acknowledgments. Thanks go to Carlos Beltrán and Jean-Pierre Dedieu for helpful comments. We are very grateful to Mike Shub for constructive criticism and insightful comments that helped to improve the paper considerably. This work was finalized during the special semester on Foundations of Computational Mathematics in the fall of 2009. We thank the Fields Institute in Toronto for hospitality and financial support.

2. Preliminaries

2.1. *Setting and notation.* For $d \in \mathbb{N}$ we denote by \mathcal{H}_d the subspace of $\mathbb{C}[X_0, \dots, X_n]$ of homogeneous polynomials of degree d . For $f \in \mathcal{H}_d$, we write

$$f(x) = \sum_{\alpha} \binom{d}{\alpha}^{1/2} a_{\alpha} X^{\alpha},$$

where $\alpha = (\alpha_0, \dots, \alpha_n)$ is assumed to range over all multi-indices such that $|\alpha| = \sum_{k=0}^n \alpha_k = d$, $\binom{d}{\alpha}$ denotes the multinomial coefficient, and $X^{\alpha} := X_0^{\alpha_0} X_1^{\alpha_1} \cdots X_n^{\alpha_n}$. That is, we take for basis of the linear space \mathcal{H}_d the *Bombieri-Weyl* basis consisting of the monomials $\binom{d}{\alpha}^{1/2} X^{\alpha}$. A reason to do so is that the

Hermitian inner product associated to this basis is unitarily invariant. That is, if $g \in \mathcal{H}_d$ is given by $g(x) = \sum_{\alpha} \binom{d}{\alpha}^{1/2} b_{\alpha} X^{\alpha}$, then the canonical Hermitian inner product

$$\langle f, g \rangle = \sum_{|\alpha|=d} a_{\alpha} \overline{b_{\alpha}}$$

satisfies, for all element ν in the unitary group $\mathcal{U}(n + 1)$, that

$$\langle f, g \rangle = \langle f \circ \nu, g \circ \nu \rangle.$$

Fix $d_1, \dots, d_n \in \mathbb{N} \setminus \{0\}$ and let $\mathcal{H}_{\mathbf{d}} = \mathcal{H}_{d_1} \times \dots \times \mathcal{H}_{d_n}$ be the vector space of polynomial systems $f = (f_1, \dots, f_n)$ with $f_i \in \mathbb{C}[X_0, \dots, X_n]$ homogeneous of degree d_i . The space $\mathcal{H}_{\mathbf{d}}$ is naturally endowed with a Hermitian inner product $\langle f, g \rangle = \sum_{i=1}^n \langle f_i, g_i \rangle$. We denote by $\|f\|$ the corresponding norm of $f \in \mathcal{H}_{\mathbf{d}}$.

Recall that $N = \dim_{\mathbb{C}} \mathcal{H}_{\mathbf{d}}$ and $D = \max_i d_i$. Also, in the rest of this paper, we assume $D \geq 2$ (the case $D = 1$ being solvable with elementary linear algebra).

Let $\mathbb{P}^n := \mathbb{P}(\mathbb{C}^{n+1})$ denote the complex projective space associated to \mathbb{C}^{n+1} and $S(\mathcal{H}_{\mathbf{d}})$ the unit sphere of $\mathcal{H}_{\mathbf{d}}$. These are smooth manifolds that naturally carry the structure of a Riemannian manifold (for \mathbb{P}^n the metric is called Fubini-Study metric). We will denote by $d_{\mathbb{P}}$ and $d_{\mathbb{S}}$ their Riemannian distances which, in both cases, amount to the angle between the arguments. Specifically, for $x, y \in \mathbb{P}^n$, one has

$$(2.1) \quad \cos d_{\mathbb{P}}(x, y) = \frac{|\langle x, y \rangle|}{\|x\| \|y\|}.$$

Ocasionally, for $f, g \in \mathcal{H}_{\mathbf{d}} \setminus \{0\}$, we will abuse language and write $d_{\mathbb{S}}(f, g)$ to denote this angle, that is, the distance $d_{\mathbb{S}}\left(\frac{f}{\|f\|}, \frac{g}{\|g\|}\right)$.

We define the *solution variety* to be

$$V_{\mathbb{P}} := \{(f, \zeta) \in \mathcal{H}_{\mathbf{d}} \times \mathbb{P}^n \mid f \neq 0 \text{ and } f(\zeta) = 0\}.$$

This is a smooth submanifold of $\mathcal{H}_{\mathbf{d}} \times \mathbb{P}^n$ and hence also carries a Riemannian structure. We denote by $V_{\mathbb{P}}(f)$ the zero set of $f \in \mathcal{H}_{\mathbf{d}}$ in \mathbb{P}^n . By Bézout’s theorem, it contains \mathcal{D} points for almost all f . Let $Df(\zeta)|_{T_{\zeta}}$ denote the restriction of the derivative of $f: \mathbb{C}^{n+1} \rightarrow \mathbb{C}^n$ at ζ to the tangent space $T_{\zeta} := \{v \in \mathbb{C}^{n+1} \mid \langle v, \zeta \rangle = 0\}$ of \mathbb{P}^n at ζ . The *subvariety of ill-posed pairs* is defined as

$$\Sigma'_{\mathbb{P}} := \{(f, \zeta) \in V_{\mathbb{P}} \mid \text{rank } Df(\zeta)|_{T_{\zeta}} < n\}.$$

Note that $(f, \zeta) \notin \Sigma'_{\mathbb{P}}$ means that ζ is a simple zero of f . In this case, by the implicit function theorem, the projection $V_{\mathbb{P}} \rightarrow \mathcal{H}_{\mathbf{d}}, (g, x) \mapsto g$ can be locally inverted around (f, ζ) . The image Σ of $\Sigma'_{\mathbb{P}}$ under the projection $V_{\mathbb{P}} \rightarrow \mathcal{H}_{\mathbf{d}}$ is called the *discriminant variety*.

2.2. *Newton's method.* In [20], Mike Shub introduced the following projective version of Newton's method. We associate to $f \in \mathcal{H}_d$ (with $Df(x)$ of rank n for some x) a map $N_f : \mathbb{C}^{n+1} \setminus \{0\} \rightarrow \mathbb{C}^{n+1} \setminus \{0\}$ defined (almost everywhere) by

$$N_f(x) = x - Df(x)|_{T_x}^{-1} f(x).$$

Note that $N_f(x)$ is homogeneous of degree 0 in f and of degree 1 in x so that N_f induces a rational map from \mathbb{P}^n to \mathbb{P}^n (which we will still denote by N_f), and this map is invariant under multiplication of f by constants.

We note that $N_f(x)$ can be computed from f and x very efficiently: since the Jacobian $Df(x)$ can be evaluated with $\mathcal{O}(N)$ arithmetic operations [3], one can do with a total of $\mathcal{O}(N + n^3)$ arithmetic operations.

It is well known that when x is sufficiently close to a simple zero ζ of f , the sequence of Newton iterates beginning at x will converge quadratically fast to ζ . This property led Steve Smale to define the following intrinsic notion of approximate zero.

Definition 2.1. By an *approximate zero* of $f \in \mathcal{H}_d$ associated with a zero $\zeta \in \mathbb{P}^n$ of f , we understand a point $x \in \mathbb{P}^n$ such that the sequence of Newton iterates (adapted to projective space)

$$x_{i+1} := N_f(x_i)$$

with initial point $x_0 := x$ converges immediately quadratically to ζ , i.e.,

$$d_{\mathbb{P}}(x_i, \zeta) \leq \left(\frac{1}{2}\right)^{2^i - 1} d_{\mathbb{P}}(x_0, \zeta)$$

for all $i \in \mathbb{N}$.

2.3. *Condition numbers.* How close need x be from ζ to be an approximate zero? This depends on how well conditioned the zero ζ is.

For $f \in \mathcal{H}_d$ and $x \in \mathbb{C}^{n+1} \setminus \{0\}$, we define the (*normalized*) *condition number* $\mu_{\text{norm}}(f, x)$ by

$$\mu_{\text{norm}}(f, x) := \|f\| \left\| \left(Df(x)|_{T_x} \right)^{-1} \text{diag}(\sqrt{d_1} \|x\|^{d_1-1}, \dots, \sqrt{d_n} \|x\|^{d_n-1}) \right\|,$$

where T_x denotes the Hermitian complement of $\mathbb{C}x$, the right-hand side norm is the spectral norm, and $\text{diag}(a_i)$ denotes the diagonal matrix with entries a_i . Note that $\mu_{\text{norm}}(f, x)$ is homogeneous of degree 0 in both arguments; hence it is well defined for $(f, x) \in \mathcal{H}_d \times \mathbb{P}^n$. If x is a simple zero of f , then $\ker Df(x) = \mathbb{C}x$ and hence $(Df(x)|_{T_x})^{-1}$ can be identified with the Moore-Penrose inverse $Df(x)^\dagger$ of $Df(x)$. We have $\mu_{\text{norm}}(f, x) \geq 1$; cf. [8, §12.4, Cor. 3].

The following result (essentially, a γ -Theorem in Smale's theory of estimates for Newton's method [27]) quantifies our claim above.

THEOREM 2.2. *Assume $f(\zeta) = 0$ and $d_{\mathbb{P}}(x, \zeta) \leq \frac{u_0}{D^{3/2} \mu_{\text{norm}}(f, \zeta)}$, where $u_0 := 3 - \sqrt{7} \approx 0.3542$. Then x is an approximate zero of f associated with ζ .*

Proof. This is an immediate consequence of the projective γ -Theorem in [8, p. 263, Th. 1] combined with the higher derivative estimate [8, p. 267, Th. 2]. \square

2.4. *Gaussian distributions.* The distribution of input data will be modelled with Gaussians. Let $\bar{x} \in \mathbb{R}^n$ and $\sigma > 0$. We recall that the Gaussian distribution $N(\bar{x}, \sigma^2 \mathbf{I})$ on \mathbb{R}^n with mean \bar{x} and covariance matrix $\sigma^2 \mathbf{I}$ is given by the density

$$\rho(x) = \left(\frac{1}{\sigma\sqrt{2\pi}} \right)^n \exp\left(-\frac{\|x - \bar{x}\|^2}{2\sigma^2} \right).$$

3. Statement of main results

3.1. *The homotopy continuation routine ALH.* Suppose that we are given an input system $f \in \mathcal{H}_{\mathbf{d}}$ and an initial pair (g, ζ) in the solution variety $V_{\mathbb{P}}$ such that f and g are \mathbb{R} -linearly independent. Let $\alpha = d_{\mathbb{S}}(f, g)$. Consider the line segment $E_{f,g}$ in $\mathcal{H}_{\mathbf{d}}$ with endpoints f and g . We parametrize this segment by writing

$$E_{f,g} = \{q_{\tau} \in \mathcal{H}_{\mathbf{d}} \mid \tau \in [0, 1]\},$$

with q_{τ} being the only point in $E_{f,g}$ such that $d_{\mathbb{S}}(g, q_{\tau}) = \tau\alpha$ (see Figure 1). Explicitly, we have $q_{\tau} = tf + (1-t)g$, where $t = t(\tau)$ is given by equation (5.4) below. If $E_{f,g}$ does not intersect the discriminant variety Σ , there is a unique continuous map $[0, 1] \rightarrow V_{\mathbb{P}}, \tau \mapsto (q_{\tau}, \zeta_{\tau})$ such that $(q_0, \zeta_0) = (g, \zeta)$, called the *lifting* of $E_{f,g}$ with origin (g, ζ) . In order to find an approximation of the zero ζ_1 of $f = q_1$, we may start with the zero $\zeta = \zeta_0$ of $g = q_0$ and numerically follow the path (q_{τ}, ζ_{τ}) by subdividing $[0, 1]$ with points $0 = \tau_0 < \tau_1 < \dots < \tau_k = 1$ and by successively computing approximations x_i of ζ_{τ_i} by Newton's method.

More precisely, we consider the following algorithm ALH (Adaptive Linear Homotopy) with the stepsize parameter $\lambda = 6.67 \cdot 10^{-3}$.

Algorithm ALH

input $f, g \in \mathcal{H}_{\mathbf{d}}$ and $\zeta \in \mathbb{P}^n$ such that $g(\zeta) = 0$

$\alpha := d_{\mathbb{S}}(f, g)$, $r := \|f\|$, $s := \|g\|$

$\tau := 0$, $q := g$, $x := \zeta$

repeat

$\Delta\tau := \frac{\lambda}{\alpha D^{3/2} \mu_{\text{norm}}^2(q, x)}$

$\tau := \min\{1, \tau + \Delta\tau\}$

$t := \frac{s}{r \sin \alpha \cot(\tau\alpha) - r \cos \alpha + s}$

$q := tf + (1-t)g$

$x := N_q(x)$

until $\tau = 1$

RETURN x

Our main result for this algorithm, which we will prove in Section 4, is the following.

THEOREM 3.1. *The algorithm ALH stops after at most k steps with*

$$k \leq 245 D^{3/2} d_{\mathbb{S}}(f, g) \int_0^1 \mu_{\text{norm}}^2(q_{\tau}, \zeta_{\tau}) d\tau.$$

The returned point x is an approximate zero of f with associated zero ζ_1 .

Remark 3.2. 1. The bound in Theorem 3.1 is optimal up to a constant factor. This easily follows by an inspection of its proof given in Section 4.

2. Algorithm ALH requires the computation of μ_{norm} which, in turn, requires the computation of the operator norm of a matrix. This cannot be done exactly with rational operations and square roots only. We can do, however, with a sufficiently good approximation of $\mu_{\text{norm}}^2(q, x)$, and there exist several numerical methods efficiently computing such an approximation. We will therefore neglect this issue pointing, however, for the skeptical reader that another course of action is possible. Indeed, one may replace the operator by the Frobenius norm in the definition of μ_{norm} and use the bounds $\|M\| \leq \|M\|_F \leq \sqrt{\text{rank}(M)}\|M\|$ to show that this change preserves the correctness of ALH and adds a multiplicative factor n in the right-hand side of Theorem 3.1. A similar comment applies to the computation of α and $\cot(\tau\alpha)$ in algorithm ALH which cannot be done exactly with rational operations.

3.2. *Randomization and complexity: the algorithm LV.* ALH will serve as the basic routine for a number of algorithms computing zeros of polynomial systems in different contexts. In these contexts both the input system f and the origin (g, ζ) of the homotopy may be randomly chosen: in the case of (g, ζ) as a computational technique and in the case of f in order to perform a probabilistic analysis of the algorithm's running time.

In both cases, a probability measure is needed: one for f and one for the pair (g, ζ) . The measure for f will depend on the kind of probabilistic analysis (standard average-case or smoothed analysis) we perform. In contrast, we will consider only one measure on $V_{\mathbb{P}}$ — which we denote by ρ_{st} — for the initial pair (g, ζ) . It consists of drawing g from $\mathcal{H}_{\mathbf{d}}$ from the standard Gaussian distribution (defined via the isomorphism $\mathcal{H}_{\mathbf{d}} \simeq \mathbb{R}^{2N}$ given by the Bombieri-Weyl basis) and then choosing one of the (almost surely) \mathcal{D} zeros of g from the uniform distribution on $\{1, \dots, \mathcal{D}\}$. The formula for the density of ρ_{st} will be derived later; see Lemma 8.8(5). The above procedure is clearly nonconstructive as computing a zero of a system is the problem we wanted to solve in the first place. One of the major contributions in [4] was to show that this drawback can be repaired. The following result (a detailed version of the

effective sampling in [6]) will be proved in Section 9 as a special case of more general results we will need in our development.

PROPOSITION 3.3. *We can compute a random pair $(g, \zeta) \in V_{\mathbb{P}}$ according to the density ρ_{st} with $\mathcal{O}(N)$ choices of random real numbers from the standard Gaussian distribution and $\mathcal{O}(DnN + n^3)$ arithmetic operations (including square roots of positive numbers).*

Algorithms using randomly drawn data are called probabilistic (or randomized). Those that always return a correct output are said to be of type *Las Vegas*. The following algorithm (which uses Proposition 3.3) belongs to this class:

Algorithm LV
input $f \in \mathcal{H}_{\mathbf{d}}$
 draw $(g, \zeta) \in V_{\mathbb{P}}$ **from** ρ_{st}
 run ALH **on input** (f, g, ζ)

For an input $f \in \mathcal{H}_{\mathbf{d}}$ algorithm LV either outputs an approximate zero x of f or loops forever. By the *running time* $t(f, g, \zeta)$ we will understand the number of elementary operations (i.e., arithmetic operations, evaluations of the elementary functions sin, cos, cot, square root, and comparisons) performed by LV on input f with initial pair (g, ζ) . For fixed f , this is a random variable and its expectation $t(f) := \mathbb{E}_{(g, \zeta) \sim \rho_{\text{st}}}(t(f, g, \zeta))$ is said to be the *expected running time* of LV on input f .

For all f, g, ζ , the running time $t(f, g, \zeta)$ is given by the *number of iterations* $K(f, g, \zeta)$ of ALH with input this triple times the cost of an iteration, the latter being dominated by that of computing one Newton iterate (which is $\mathcal{O}(N + n^3)$ independently of the triple (f, g, ζ) ; see §2.2). It therefore follows that analyzing the expected running times of LV amounts to do so for the expected value — over $(g, \zeta) \in V_{\mathbb{P}}$ drawn from ρ_{st} — of $K(f, g, \zeta)$. We denote this expectation by

$$K(f) := \mathbb{E}_{(g, \zeta) \sim \rho_{\text{st}}}(K(f, g, \zeta)).$$

3.3. Average analysis of LV. To talk about average complexity of LV requires specifying a measure for the set of inputs. The most natural choice is the standard Gaussian distribution on $\mathcal{H}_{\mathbf{d}}$. Since $K(f)$ is invariant under scaling, we may equivalently assume that f is chosen in the unit sphere $S(\mathcal{H}_{\mathbf{d}})$ from the uniform distribution. With this choice, we say a Las Vegas algorithm is *average polynomial time* when the average — over $f \in S(\mathcal{H}_{\mathbf{d}})$ — of its expected running time is polynomially bounded in the size N of f . The following result shows that LV is average polynomial time. It is essentially the main result in [6] (modulo the existence of ALH and with specific constants).

THEOREM 3.4. *The average of the expected number of iterations of Algorithm LV is bounded as ($n \geq 4$)*

$$\mathbb{E}_{f \in \mathcal{S}(\mathcal{H}_d)} K(f) \leq 4185 D^{3/2} N(n+1).$$

3.4. Smoothed analysis of LV. A smoothed analysis of an algorithm consists of bounding, for all possible input data \bar{f} , the average of its running time (its expected running time if it is a Las Vegas algorithm) over small perturbations of \bar{f} . To perform such an analysis, a family of measures (parametrized by a parameter r controlling the size of the perturbation) is considered with the following characteristics:

- (1) The density of an element f depends only on the distance $\|f - \bar{f}\|$.
- (2) The value of r is closely related to the variance of $\|f - \bar{f}\|$.

Then, the average above is estimated as a function of the data size N and the parameter r , and a satisfying result, which is described by the expression *smoothed polynomial time*, demands that this function is polynomially bounded in r^{-1} and N . Possible choices for the measures' family are the Gaussians $N(\bar{f}, \sigma^2 \mathbf{I})$ (used, for instance, in [14], [19], [32], [33]) and the uniform measure on disks $B(\bar{f}, r)$ (used in [2], [11], [12]). Other families may also be used and an emerging impression is that smoothed analysis is robust in the sense that its dependence on the chosen family of measures is low. This tenet was argued for in [15] where a uniform measure is replaced by an adversarial measure (one having a pole at \bar{f}) without a significant loss in the estimated averages.

In this paper, for reasons of technical simplicity and consistency with the rest of the exposition, we will work with truncated Gaussians defined as follows. For $\bar{f} \in \mathcal{H}_d$ and $\sigma > 0$, we shall denote by $N(\bar{f}, \sigma^2 \mathbf{I})$ the Gaussian distribution on $\mathcal{H}_d \simeq \mathbb{R}^{2N}$ (defined with respect to the Bombieri-Weyl basis) with mean \bar{f} and covariance matrix $\sigma^2 \mathbf{I}$. Further, for $A > 0$, let $P_{A,\sigma} := \text{Prob}\{\|f\| \leq A \mid f \sim N(0, \sigma^2 \mathbf{I})\}$. We define the *truncated Gaussian* $N_A(\bar{f}, \sigma^2 \mathbf{I})$ with center $\bar{f} \in \mathcal{H}_d$ as the probability measure on \mathcal{H}_d with density

$$(3.1) \quad \rho(f) = \begin{cases} \frac{\rho_{\bar{f},\sigma}(f)}{P_{A,\sigma}} & \text{if } \|f - \bar{f}\| \leq A \\ 0 & \text{otherwise,} \end{cases}$$

where $\rho_{\bar{f},\sigma}$ denotes the density of $N(\bar{f}, \sigma^2 \mathbf{I})$. Note that $N_A(\bar{f}, \sigma^2 \mathbf{I})$ is isotropic around its mean \bar{f} .

For our smoothed analysis we will take $A = \sqrt{2N}$. In this case, we have $P_{A,\sigma} \geq \frac{1}{2}$ for all $\sigma \leq 1$ (Lemma 6.1). Note also that $\text{Var}(\|f - \bar{f}\|) \leq \sigma^2$, so that any upper bound polynomial in σ^{-2} is also an upper bound polynomial in $\text{Var}(\|f - \bar{f}\|)^{-1}$.

We can now state our smoothed analysis result for LV.

THEOREM 3.5. *For any $0 < \sigma \leq 1$, Algorithm LV satisfies*

$$\sup_{\bar{f} \in S(\mathcal{H}_d)} \mathbb{E}_{f \sim N_A(\bar{f}, \sigma^2 \mathbf{I})} K(f) \leq 4185 D^{3/2} (N + 2^{-1/2} \sqrt{N}) (n + 1) \frac{1}{\sigma}.$$

3.5. *The main technical result.* The technical heart of the proof of the mentioned results on the average and smoothed analysis of LV is the following smoothed analysis of the mean square condition number.

THEOREM 3.6. *Let $\bar{q} \in \mathcal{H}_d$ and $\sigma > 0$. For $q \in \mathcal{H}_d$ drawn from $N(\bar{q}, \sigma^2 \mathbf{I})$, we have*

$$\mathbb{E}_{\mathcal{H}_d} \left(\frac{\mu_2^2(q)}{\|q\|^2} \right) \leq \frac{e(n + 1)}{2\sigma^2}.$$

We note that no bound on the norm of \bar{q} is required here. Indeed, using $\mu_2(\lambda q) = \mu_2(q)$, it is easy to see that the assertion for \bar{q}, σ implies the assertion for $\lambda \bar{q}, \lambda \sigma$, for any $\lambda > 0$.

3.6. *Condition-based analysis of LV.* We are here interested in estimating $K(f)$ for a fixed input system $f \in S(\mathcal{H}_d)$. Such an estimate will have to depend on, besides N, n , and D , the condition of f . We measure the latter using Shub and Smale’s [22] $\mu_{\max}(f)$ defined in (1.1). Our condition-based analysis of LV is summarized in the following statement.

THEOREM 3.7. *The expected number of iterations of Algorithm LV with input $f \in S(\mathcal{H}_d) \setminus \Sigma$ is bounded as*

$$K(f) \leq 200411 D^3 N (n + 1) \mu_{\max}^2(f).$$

3.7. *A near solution of Smale’s 17th problem.* We finally want to consider deterministic algorithms finding zeros of polynomial systems. Our goal is to exhibit one such algorithm working in nearly-polynomial average time, more precisely in average time $N^{\mathcal{O}(\log \log N)}$. A first ingredient to do so is a deterministic homotopy algorithm which is fast when D is small. This consists of algorithm ALH plus the initial pair (\bar{U}, \mathbf{z}_1) , where $\bar{U} = (\bar{U}_1, \dots, \bar{U}_n) \in S(\mathcal{H}_d)$ with $\bar{U}_i = \frac{1}{\sqrt{2n}}(X_0^{d_i} - X_i^{d_i})$ and $\mathbf{z}_1 = (1 : 1 : \dots : 1)$.

We consider the following algorithm MD (Moderate Degree):

Algorithm MD
input $f \in \mathcal{H}_d$
 run ALH on input $(f, \bar{U}, \mathbf{z}_1)$

We write $K_{\bar{U}}(f) := K(f, \bar{U}, \mathbf{z}_1)$ for the number of iterations of algorithm MD with input f . We are interested in computing the average over f of $K_{\bar{U}}(f)$ for f randomly chosen in $S(\mathcal{H}_d)$ from the uniform distribution.

The complexity of MD is bounded as follows.

THEOREM 3.8. *The average number of iterations of Algorithm MD is bounded as*

$$\mathbb{E}_{f \in S(\mathcal{H}_d)} K_{\overline{U}}(f) \leq 400821 D^3 N(n+1)^{D+1}.$$

Algorithm MD is efficient when D is small, say, when $D \leq n$. For $D > n$ we use another approach, namely, a real number algorithm designed by Jim Renegar [18] which in this case has a performance similar to that of MD when $D \leq n$. Putting both pieces together we will reach our last main result.

THEOREM 3.9. *There is a deterministic real number algorithm that on input $f \in \mathcal{H}_d$ computes an approximate zero of f in average time $N^{\mathcal{O}(\log \log N)}$, where $N = \dim \mathcal{H}_d$ measures the size of the input f . Moreover, if we restrict data to polynomials satisfying*

$$D \leq n^{\frac{1}{1+\varepsilon}} \quad \text{or} \quad D \geq n^{1+\varepsilon}$$

for some fixed $\varepsilon > 0$, then the average time of the algorithm is polynomial in the input size N .

4. Complexity analysis of ALH

The goal of this section is to prove Theorem 3.1. An essential component in this proof is an estimate of how much $\mu_{\text{norm}}(f, \zeta)$ changes when f or ζ (or both) are slightly perturbed. The following result gives upper and lower bounds on this variation. It is a precise version, with explicit constants, of Theorem 1 of [21].

PROPOSITION 4.1. *Assume $D \geq 2$. Let $0 < \varepsilon \leq 0.13$ be arbitrary and $C \leq \frac{\varepsilon}{5.2}$. For all $f, g \in S(\mathcal{H}_d)$ and all $x, \zeta \in \mathbb{P}^n$, if $d(f, g) \leq \frac{C}{D^{1/2} \mu_{\text{norm}}(f, \zeta)}$ and $d(\zeta, x) \leq \frac{C}{D^{3/2} \mu_{\text{norm}}(f, \zeta)}$, then*

$$\frac{1}{1+\varepsilon} \mu_{\text{norm}}(g, x) \leq \mu_{\text{norm}}(f, \zeta) \leq (1+\varepsilon) \mu_{\text{norm}}(g, x).$$

In what follows, we will fix the constants as $\varepsilon = 0.13$ and $C = \frac{\varepsilon}{5.2} = 0.025$.

Remark 4.2. The constants C and ε implicitly occur in the statement of Theorem 3.1 since the 245 therein is a function of these numbers. But their role is not limited to this since they also occur in the algorithm ALH in the parameter $\lambda = \frac{C(1-\varepsilon)}{2(1+\varepsilon)^4}$ controlling the update $\tau + \Delta\tau$ of τ . We note that for the former we could do without precise values by using the big Oh notation. In contrast, we cannot talk of a constructive procedure unless all of its steps are precisely given.

Proof of Theorem 3.1. Let $0 = \tau_0 < \tau_1 < \dots < \tau_k = 1$ and $\zeta_0 = x_0, x_1, \dots, x_k$ be the sequences of τ -values and points in \mathbb{P}^n generated by the

algorithm ALH. To simplify notation we write q_i instead of q_{τ_i} and ζ_i instead of ζ_{τ_i} .

We claim that, for $i = 0, \dots, k - 1$, the following inequalities are true:

- (a) $d_{\mathbb{P}}(x_i, \zeta_i) \leq \frac{C}{D^{3/2}\mu_{\text{norm}}(q_i, \zeta_i)}$;
- (b) $\frac{\mu_{\text{norm}}(q_i, x_i)}{(1 + \varepsilon)} \leq \mu_{\text{norm}}(q_i, \zeta_i) \leq (1 + \varepsilon)\mu_{\text{norm}}(q_i, x_i)$;
- (c) $d_{\mathbb{S}}(q_i, q_{i+1}) \leq \frac{C}{D^{3/2}\mu_{\text{norm}}(q_i, \zeta_i)}$;
- (d) $d_{\mathbb{P}}(\zeta_i, \zeta_{i+1}) \leq \frac{C}{D^{3/2}\mu_{\text{norm}}(q_i, \zeta_i)} \frac{(1 - \varepsilon)}{(1 + \varepsilon)}$;
- (e) $d_{\mathbb{P}}(x_i, \zeta_{i+1}) \leq \frac{2C}{(1 + \varepsilon)D^{3/2}\mu_{\text{norm}}(q_i, \zeta_i)}$.

We proceed by induction showing that

$$(\mathbf{a}, i) \Rightarrow (\mathbf{b}, i) \Rightarrow ((\mathbf{c}, i) \text{ and } (\mathbf{d}, i)) \Rightarrow (\mathbf{e}, i) \Rightarrow (\mathbf{a}, i + 1).$$

Inequality (a) for $i = 0$ is trivial.

Assume now that (a) holds for some $i \leq k - 1$. Then, Proposition 4.1 (with $f = g = q_i$) implies

$$\frac{\mu_{\text{norm}}(q_i, x_i)}{(1 + \varepsilon)} \leq \mu_{\text{norm}}(q_i, \zeta_i) \leq (1 + \varepsilon)\mu_{\text{norm}}(q_i, x_i)$$

and thus (b). We now show (c) and (d). To do so, put $p_\tau := \frac{q_\tau}{\|q_\tau\|}$ and let $\tau_* > \tau_i$ be such that $\int_{\tau_i}^{\tau_*} (\|\dot{p}_\tau\| + \|\dot{\zeta}_\tau\|)d\tau = \frac{C}{D^{3/2}\mu_{\text{norm}}(q_i, \zeta_i)} \frac{(1 - \varepsilon)}{(1 + \varepsilon)}$ or $\tau_* = 1$, whichever the smallest. Then, for all $t \in [\tau_i, \tau_*]$,

$$\begin{aligned} d_{\mathbb{P}}(\zeta_i, \zeta_t) &= \int_{\tau_i}^t \|\dot{\zeta}_\tau\| d\tau \leq \int_{\tau_i}^{\tau_*} (\|\dot{p}_\tau\| + \|\dot{\zeta}_\tau\|)d\tau \\ &\leq \frac{C}{D^{3/2}\mu_{\text{norm}}(q_i, \zeta_i)} \frac{(1 - \varepsilon)}{(1 + \varepsilon)} \end{aligned}$$

and, similarly,

$$d_{\mathbb{S}}(q_i, q_t) \leq \frac{C}{D^{3/2}\mu_{\text{norm}}(q_i, \zeta_i)} \frac{(1 - \varepsilon)}{(1 + \varepsilon)} \leq \frac{C}{D^{3/2}\mu_{\text{norm}}(q_i, \zeta_i)}.$$

It is therefore enough to show that $\tau_{i+1} \leq \tau_*$. This is trivial if $\tau_* = 1$. We therefore assume $\tau_* < 1$. The two bounds above allow us to apply Proposition 4.1 and to deduce, for all $\tau \in [\tau_i, \tau_*]$,

$$\mu_{\text{norm}}(q_\tau, \zeta_\tau) \leq (1 + \varepsilon)\mu_{\text{norm}}(q_i, \zeta_i).$$

From $\|\dot{\zeta}_\tau\| \leq \mu_{\text{norm}}(q_\tau, \zeta_\tau) \|\dot{p}_\tau\|$ (cf. [8, §12.3-12.4]) and $\mu_{\text{norm}}(q_\tau, \zeta_\tau) \geq 1$ it follows that

$$\begin{aligned} \frac{C}{D^{3/2}\mu_{\text{norm}}(q_i, \zeta_i)} \frac{(1 - \varepsilon)}{(1 + \varepsilon)} &= \int_{\tau_i}^{\tau_*} (\|\dot{p}_\tau\| + \|\dot{\zeta}_\tau\|) d\tau \leq \int_{\tau_i}^{\tau_*} 2\mu_{\text{norm}}(q_\tau, \zeta_\tau) \|\dot{p}_\tau\| d\tau \\ &\leq 2(1 + \varepsilon)\mu_{\text{norm}}(q_i, \zeta_i) \int_{\tau_i}^{\tau_*} \|\dot{p}_\tau\| d\tau \leq 2d_{\mathbb{S}}(q_i, q_{\tau_*})(1 + \varepsilon)\mu_{\text{norm}}(q_i, \zeta_i). \end{aligned}$$

Consequently, using (b), we obtain

$$d_{\mathbb{S}}(q_i, q_{\tau_*}) \geq \frac{C(1 - \varepsilon)}{2(1 + \varepsilon)^2 D^{3/2} \mu_{\text{norm}}^2(q_i, \zeta_i)} \geq \frac{C(1 - \varepsilon)}{2(1 + \varepsilon)^4 D^{3/2} \mu_{\text{norm}}^2(q_i, x_i)}.$$

The parameter λ in ALH is chosen as $\frac{C(1-\varepsilon)}{2(1+\varepsilon)^4}$ (or slightly less). By the definition of $\tau_{i+1} - \tau_i$ in ALH we have $\alpha(\tau_{i+1} - \tau_i) = \frac{\lambda}{D^{3/2} \mu_{\text{norm}}^2(q_i, x_i)}$. So we obtain

$$d_{\mathbb{S}}(q_i, q_{\tau_*}) \geq \alpha(\tau_{i+1} - \tau_i) = d_{\mathbb{S}}(q_i, q_{i+1}).$$

This implies $\tau_{i+1} \leq \tau_*$ as claimed and hence, inequalities (c) and (d). With them, we may apply Proposition 4.1 to deduce, for all $\tau \in [\tau_i, \tau_{i+1}]$,

$$(4.1) \quad \frac{\mu_{\text{norm}}(q_i, \zeta_i)}{1 + \varepsilon} \leq \mu_{\text{norm}}(q_\tau, \zeta_\tau) \leq (1 + \varepsilon)\mu_{\text{norm}}(q_i, \zeta_i).$$

Next we use the triangle inequality, (a), and (d), to obtain

$$\begin{aligned} d_{\mathbb{P}}(x_i, \zeta_{i+1}) &\leq d_{\mathbb{P}}(x_i, \zeta_i) + d_{\mathbb{P}}(\zeta_i, \zeta_{i+1}) \\ &\leq \frac{C}{D^{3/2}\mu_{\text{norm}}(q_i, \zeta_i)} + \frac{C}{D^{3/2}\mu_{\text{norm}}(q_i, \zeta_i)} \frac{(1 - \varepsilon)}{(1 + \varepsilon)} \\ &= \frac{2C}{(1 + \varepsilon)D^{3/2}\mu_{\text{norm}}(q_i, \zeta_i)}, \end{aligned}$$

which proves (e). Theorem 2.2 yields that x_i is an approximate zero of q_{i+1} associated with its zero ζ_{i+1} . Indeed, by our choice of C and ε , we have $2C \leq u_0(1 + \varepsilon)$ and hence $d_{\mathbb{P}}(x_i, \zeta_{i+1}) \leq \frac{u_0}{D^{3/2}\mu_{\text{norm}}(q_i, \zeta_i)}$. Therefore, $x_{i+1} = N_{q_{i+1}}(x_i)$ satisfies

$$d_{\mathbb{P}}(x_{i+1}, \zeta_{i+1}) \leq \frac{1}{2} d_{\mathbb{P}}(x_i, \zeta_{i+1}).$$

Using (e) and the right-hand inequality in (4.1) with $t = t_{i+1}$, we obtain

$$d_{\mathbb{P}}(x_{i+1}, \zeta_{i+1}) \leq \frac{C}{(1 + \varepsilon)D^{3/2}\mu_{\text{norm}}(q_i, \zeta_i)} \leq \frac{C}{D^{3/2}\mu_{\text{norm}}(q_{i+1}, \zeta_{i+1})},$$

which proves (a) for $i + 1$. The claim is thus proved.

The estimate $d_{\mathbb{P}}(x_k, \zeta_k) \leq \frac{C}{D^{3/2}\mu_{\text{norm}}(q_k, \zeta_k)}$ just shown for $i = k - 1$ implies by Theorem 2.2 that the returned point x_k is an approximate zero of $q_k = f$ with associated zero ζ_1 .

Consider now any $i \in \{0, \dots, k - 1\}$. Using (4.1) and (b) we obtain

$$\begin{aligned} \int_{\tau_i}^{\tau_{i+1}} \mu_{\text{norm}}^2(q_\tau, \zeta_\tau) d\tau &\geq \int_{\tau_i}^{\tau_{i+1}} \frac{\mu_{\text{norm}}^2(q_i, \zeta_i)}{(1 + \varepsilon)^2} d\tau = \frac{\mu_{\text{norm}}^2(q_i, \zeta_i)}{(1 + \varepsilon)^2} (\tau_{i+1} - \tau_i) \\ &\geq \frac{\mu_{\text{norm}}^2(q_i, x_i)}{(1 + \varepsilon)^4} (\tau_{i+1} - \tau_i) \\ &= \frac{\mu_{\text{norm}}^2(q_i, x_i)}{(1 + \varepsilon)^4} \frac{\lambda}{\alpha D^{3/2} \mu_{\text{norm}}^2(q_i, x_i)} \\ &= \frac{\lambda}{(1 + \varepsilon)^4 \alpha D^{3/2}} \geq \frac{1}{245} \frac{1}{\alpha D^{3/2}}. \end{aligned}$$

This implies

$$\int_0^1 \mu_{\text{norm}}^2(q_\tau, \zeta_\tau) d\tau \geq \frac{k}{245} \frac{1}{\alpha D^{3/2}},$$

which proves the stated bound on k . □

5. A useful change of variables

We first draw a conclusion of Theorem 3.1, that we will need several times. Recall the definition (1.4) of the mean square condition number $\mu_2(q)$.

PROPOSITION 5.1. *The expected number of iterations of ALH on input $f \in \mathcal{H}_{\mathbf{d}} \setminus \Sigma$ is bounded as*

$$K(f) \leq 245 D^{3/2} \mathbb{E}_{g \in S(\mathcal{H}_{\mathbf{d}})} \left(d_{\mathbb{S}}(f, g) \int_0^1 \mu_2^2(q_\tau) d\tau \right).$$

Proof. Fix $g \in \mathcal{H}_{\mathbf{d}}$ such that the segment $E_{f,g}$ does not intersect the discriminant variety Σ (which is the case for almost all g , as $f \notin \Sigma$). To each of the zeros $\zeta^{(i)}$ of g there corresponds a lifting $[0, 1] \rightarrow V, \tau \mapsto (q_\tau, \zeta_\tau^{(i)})$ of $E_{f,g}$ such that $\zeta_0^{(i)} = \zeta^{(i)}$. Theorem 3.1 states that

$$K(f, g, \zeta^{(i)}) \leq 245 D^{3/2} d_{\mathbb{S}}(f, g) \int_0^1 \mu_{\text{norm}}^2(q_\tau, \zeta_\tau^{(i)}) d\tau.$$

Since $\zeta_\tau^{(1)}, \dots, \zeta_\tau^{(\mathcal{D})}$ are the zeros of q_τ , we have by the definition (1.4) of the mean square condition number

$$(5.1) \quad \frac{1}{\mathcal{D}} \sum_{i=1}^{\mathcal{D}} K(f, g, \zeta^{(i)}) \leq 245 D^{3/2} d_{\mathbb{S}}(f, g) \int_0^1 \mu_2^2(q_\tau) d\tau.$$

The assertion follows now from (compare the forthcoming Lemma 8.8)

$$K(f) = \mathbb{E}_{(g, \zeta) \sim \rho_{\text{st}}} (K(f, g, \zeta)) = \mathbb{E}_{g \in S(\mathcal{H}_{\mathbf{d}})} \left(\frac{1}{\mathcal{D}} \sum_{i=1}^{\mathcal{D}} K(f, g, \zeta^{(i)}) \right). \quad \square$$

The remaining of this article is devoted to prove Theorems 3.4– 3.9. All of them involve expectations — over random f and/or g — of the integral $\int_0^1 \mu_2^2(q_\tau) d\tau$. In all cases, we will eventually deal with such an expectation with f and g Gaussian. Since a linear combination (with fixed coefficients) of two such Gaussian systems is Gaussian as well, it is convenient to parametrize the interval $E_{f,g}$ by a parameter $t \in [0, 1]$ representing a ratio of Euclidean distances (instead of a ratio of angles as τ does). Thus we write, abusing notation, $q_t = tf + (1 - t)g$. For fixed t , as noted before, q_t follows a Gaussian law. For this new parametrization we have the following result.

PROPOSITION 5.2. *Let $f, g \in \mathcal{H}_d$ be \mathbb{R} -linearly independent and $\tau_0 \in [0, 1]$. Then*

$$d_{\mathbb{S}}(f, g) \int_{\tau_0}^1 \mu_2^2(q_\tau) d\tau \leq \int_{t_0}^1 \|f\| \|g\| \frac{\mu_2^2(q_t)}{\|q_t\|^2} dt,$$

where

$$t_0 = \frac{\|g\|}{\|g\| + \|f\|(\sin \alpha \cot(\tau_0 \alpha) - \cos \alpha)}$$

is the fraction of the Euclidean distance $\|f - g\|$ corresponding to the fraction τ_0 of the angle $\alpha = d_{\mathbb{S}}(f, g)$.

Proof. For $t \in [0, 1]$, abusing notation, we let $q_t = tf + (1 - t)g$ and $\tau(t) \in [0, 1]$ be such that $\tau(t)\alpha$ is the angle between g and q_t . This defines a bijective map $[t_0, 1] \rightarrow [\tau_0, 1], t \mapsto \tau(t)$. We denote its inverse by $\tau \mapsto t(\tau)$. We claim that

$$(5.2) \quad \frac{d\tau}{dt} = \frac{\sin \alpha}{\alpha} \frac{\|f\| \cdot \|g\|}{\|q_t\|^2}.$$

Note that the stated inequality easily follows from this claim by the transformation formula for integrals together with the bound $\sin \alpha \leq 1$.

To prove Claim (5.2), denote $r = \|f\|$ and $s = \|g\|$. We will explicitly compute $t(\tau)$ by some elementary geometry. For this, we introduce cartesian coordinates in the plane spanned by f and g and assume that g has the coordinates $(s, 0)$ and f has the coordinates $(r \cos \alpha, r \sin \alpha)$; see Figure 1.

Then, the lines determining q_τ have the equations

$$x = y \frac{\cos(\tau\alpha)}{\sin(\tau\alpha)} \quad \text{and} \quad x = y \frac{r \cos \alpha - s}{r \sin \alpha} + s$$

from where it follows that the coordinate y of q_τ is

$$(5.3) \quad y = \frac{rs \sin \alpha \sin(\tau\alpha)}{r \sin \alpha \cos(\tau\alpha) - r \cos \alpha \sin(\tau\alpha) + s \sin(\tau\alpha)}.$$

Since $t(\tau) = \frac{y}{r \sin \alpha}$ it follows that

$$(5.4) \quad t(\tau) = \frac{s}{r \sin \alpha \cot(\tau\alpha) - r \cos \alpha + s}.$$




Figure 1. Computing $t(\tau)$.

This implies the stated formula for $t_0 = t(\tau_0)$. Differentiating with respect to τ , using (5.3) and $\sin(\tau\alpha) = \frac{y}{\|q_\tau\|}$, we obtain from (5.4)

$$\begin{aligned} \frac{dt}{d\tau} &= \frac{\alpha r s \sin \alpha}{(r \sin \alpha \cos(\tau\alpha) - r \cos \alpha \sin(\tau\alpha) + s \sin(\tau\alpha))^2} \\ &= \frac{\alpha y^2}{r s \sin^2(\tau\alpha) \sin \alpha} = \frac{\alpha \|q_{t(\tau)}\|^2}{r s \sin \alpha}. \end{aligned}$$

This finishes the proof of Claim (5.2). □

In all the cases we will deal with, the factor $\|f\| \|g\|$ will be easily bounded and factored out the expectation. We will ultimately face the problem of estimating expectations of $\frac{\mu_2^2(q_t)}{\|q_t\|^2}$ for different choices of \bar{q}_t and σ_t . This is achieved by Theorem 3.6 stated in Section 3.5.

6. Analysis of LV

We derive here from Theorem 3.6 our main results on the average and smoothed analysis of LV stated in Section 3. The proof of Theorem 3.6 is postponed to Sections 7–8.

6.1. *Average-case analysis of LV (proof)*. To warm up, we first prove Theorem 3.4, which illustrates the blending of the previous results in a simpler setting.

In the following we set $A := \sqrt{2N}$ and write $P_{A,\sigma} = \text{Prob}\{\|f\| \leq A \mid f \sim N(0, \sigma^2 \mathbf{I})\}$ for $\sigma > 0$.

LEMMA 6.1. *We have $P_{A,\sigma} \geq \frac{1}{2}$ for all $0 < \sigma \leq 1$.*

Proof. Clearly it suffices to assume $\sigma = 1$. The random variable $\|f\|^2$ is chi-square distributed with $2N$ degrees of freedom. Its mean equals $2N$. In [13, Cor. 6] it is shown that the median of a chi-square distribution is always less than its mean. □

Proof of Theorem 3.4. We use Proposition 5.1 to obtain

$$\begin{aligned} \mathbb{E}_{f \in S(\mathcal{H}_d)} K(f) &\leq 245 D^{3/2} \mathbb{E}_{f \in S(\mathcal{H}_d)} \mathbb{E}_{g \in S(\mathcal{H}_d)} \left(d_{\mathbb{S}}(f, g) \int_0^1 \mu_2^2(q_\tau) d\tau \right) \\ &= 245 D^{3/2} \mathbb{E}_{f \sim N_A(0, I)} \mathbb{E}_{g \sim N_A(0, I)} \left(d_{\mathbb{S}}(f, g) \int_0^1 \mu_2^2(q_\tau) d\tau \right). \end{aligned}$$

The equality follows from the fact that, since both $d_{\mathbb{S}}(f, g)$ and $\mu_2^2(q_\tau)$ are homogeneous of degree 0 in both f and g , we may replace the uniform distribution on $S(\mathcal{H}_d)$ by any rotationally invariant distribution on \mathcal{H}_d , in particular by the centered truncated Gaussian $N_A(0, I)$ defined in (3.1). Now we use Proposition 5.2 (with $\tau_0 = 0$) to get

$$(6.1) \quad \mathbb{E}_{f \in S(\mathcal{H}_d)} K(f) \leq 245 D^{3/2} A^2 \mathbb{E}_{f \sim N_A(0, I)} \mathbb{E}_{g \sim N_A(0, I)} \left(\int_0^1 \frac{\mu_2^2(q_t)}{\|q_t\|^2} dt \right).$$

Denoting by $\rho_{0,1}$ the density of $N(0, I)$, the right-hand side of (6.1) equals

$$\begin{aligned} &245 D^{3/2} \frac{A^2}{P_{A,1}^2} \int_{\|f\| \leq A} \int_{\|g\| \leq A} \left(\int_0^1 \frac{\mu_2^2(q_t)}{\|q_t\|^2} dt \right) \rho_{0,1}(g) \rho_{0,1}(f) dg df \\ &\leq 245 D^{3/2} \frac{A^2}{P_{A,1}^2} \mathbb{E}_{f \sim N(0, I)} \mathbb{E}_{g \sim N(0, I)} \left(\int_0^1 \frac{\mu_2^2(q_t)}{\|q_t\|^2} dt \right) \\ &= 245 D^{3/2} \frac{A^2}{P_{A,1}^2} \int_0^1 \mathbb{E}_{q_t \sim N(0, (t^2 + (1-t)^2)I)} \left(\frac{\mu_2^2(q_t)}{\|q_t\|^2} \right) dt, \end{aligned}$$

where the last equality follows from the fact that, for fixed t , the random polynomial system $q_t = tf + (1 - t)g$ has a Gaussian distribution with law $N(0, \sigma_t^2 I)$, where $\sigma_t^2 := t^2 + (1 - t)^2$. Note that we deal with nonnegative integrands, so the interchange of integrals is justified by Tonelli’s theorem. By Lemma 6.1 we have $\frac{A^2}{P_{A,1}^2} \leq 8N$.

We now apply Theorem 3.6 to deduce that

$$\int_0^1 \mathbb{E}_{q_t \sim N(0, \sigma_t^2 I)} \left(\frac{\mu_2^2(q_t)}{\|q_t\|^2} \right) dt \leq \frac{e(n + 1)}{2} \int_0^1 \frac{dt}{t^2 + (1 - t)^2} = \frac{e\pi(n + 1)}{4}.$$

Consequently,

$$\mathbb{E}_{f \in S(\mathcal{H}_d)} K(f) \leq 245 D^{3/2} \cdot 8N \cdot \frac{e\pi(n + 1)}{4} \leq 4185 D^{3/2} N(n + 1). \quad \square$$

Remark 6.2. The proof (modulo the existence of ALH) for the average complexity of LV given by Beltrán and Pardo in [6] differs from the one above. It relies on the fact (elegantly shown by using integral geometry arguments) that, for all $\tau \in [0, 1]$, when f and g are uniformly drawn from the sphere, so is $q_\tau / \|q_\tau\|$. The extension of this argument to more general situations appears to be considerably more involved. In contrast, as we shall shortly see, the

argument based on Gaussians in the proof above carries over, *mutatis mutandis*, to the smoothed analysis context.

6.2. *Smoothed analysis of LV (proof).* The smoothed analysis of LV is shown similarly to its average-case analysis.

Proof of Theorem 3.5. Fix $\bar{f} \in S(\mathcal{H}_d)$. Reasoning as in the proof of Theorem 3.4 and using $\|f\| \leq \|\bar{f}\| + \|f - \bar{f}\| \leq 1 + A$, we show that

$$\begin{aligned} \mathbb{E}_{f \sim N_A(\bar{f}, \sigma^2 \mathbf{I})} K(f) &\leq 245 D^{3/2} \frac{(A+1)A}{P_{A,\sigma} P_{A,1}} \mathbb{E}_{f \sim N(\bar{f}, \sigma^2 \mathbf{I})} \mathbb{E}_{g \sim N(0, \mathbf{I})} \left(\int_0^1 \frac{\mu_2^2(q_t)}{\|q_t\|} dt \right) \\ &= 245 D^{3/2} \frac{(A+1)A}{P_{A,\sigma} P_{A,1}} \int_0^1 \mathbb{E}_{q_t \sim N(\bar{q}_t, \sigma_t^2 \mathbf{I})} \left(\frac{\mu_2^2(q_t)}{\|q_t\|} \right) dt \end{aligned}$$

with $\bar{q}_t = t\bar{f}$ and $\sigma_t^2 = (1-t)^2 + \sigma^2 t^2$. We now apply Theorem 3.6 to deduce

$$\int_0^1 \mathbb{E}_{q_t \sim N(\bar{q}_t, \sigma_t^2 \mathbf{I})} \left(\frac{\mu_2^2(q_t)}{\|q_t\|^2} \right) dt \leq \frac{e(n+1)}{2} \int_0^1 \frac{dt}{(1-t)^2 + \sigma^2 t^2} = \frac{e\pi(n+1)}{4\sigma}.$$

Consequently, using Lemma 6.1, we get

$$\mathbb{E}_{f \sim N_A(\bar{f}, \sigma^2 \mathbf{I})} K(f) \leq 245 D^{3/2} \cdot 4 \cdot (2N + \sqrt{2N}) \frac{e\pi(n+1)}{4\sigma}$$

which proves the assertion. □

The next two sections are devoted to the proof of Theorem 3.6. First, in Section 7, we give a particular smoothed analysis of a matrix condition number (Proposition 7.1). Then, in Section 8, we reduce Theorem 3.6 to this smoothed analysis of matrix condition numbers.

7. Smoothed analysis of a matrix condition number

In the following we fix $\bar{A} \in \mathbb{C}^{n \times n}$, $\sigma > 0$ and denote by ρ the Gaussian density of $N(\bar{A}, \sigma^2 \mathbf{I})$ on $\mathbb{C}^{n \times n}$. Moreover, we consider the related density

$$(7.1) \quad \tilde{\rho}(A) = c^{-1} |\det A|^2 \rho(A) \quad \text{where } c := \mathbb{E}_{A \sim \rho} (|\det A|^2).$$

The following result is akin to a smoothed analysis of the matrix condition number $\kappa(A) = \|A\| \cdot \|A^{-1}\|$, with respect to the probability densities $\tilde{\rho}$ that are not Gaussian, but closely related to Gaussians.

PROPOSITION 7.1. *We have $\mathbb{E}_{A \sim \tilde{\rho}} (\|A^{-1}\|^2) \leq \frac{e(n+1)}{2\sigma^2}$.*

The proof is based on ideas in Sankar et al. [19, §3]; see also [10]. We will actually prove tail bounds from which the stated bound on the expectation easily follows.

We denote by $\mathbb{S}^{n-1} := \{\zeta \in \mathbb{C}^n \mid \|\zeta\| = 1\}$ the unit sphere in \mathbb{C}^n .

LEMMA 7.2. For any $v \in \mathbb{S}^{n-1}$ and any $t > 0$, we have

$$\text{Prob}_{A \sim \tilde{\rho}} \left\{ \|A^{-1}v\| \geq t \right\} \leq \frac{1}{4\sigma^4 t^4}.$$

Proof. We first claim that, because of unitary invariance, we may assume that $v = e_n := (0, \dots, 0, 1)$. To see this, take $S \in U(n)$ such that $v = Se_n$. Consider the isometric map $A \mapsto B = S^{-1}A$ which transforms the density $\tilde{\rho}(A)$ to a density of the same form, namely

$$\tilde{\rho}'(B) = \tilde{\rho}(A) = c^{-1} |\det A|^2 \rho(A) = c^{-1} |\det B|^2 \rho'(B),$$

where $\rho'(B)$ denotes the density of $N(S^{-1}\bar{A}, \sigma^2\mathbf{I})$ and $c = \mathbb{E}_\rho(|\det A|^2) = \mathbb{E}_{\rho'}(|\det B|^2)$. Thus the assertion for e_n and random B (chosen from any isotropic Gaussian distribution) implies the assertion for v and A , noting that $A^{-1}v = B^{-1}e_n$. This proves the claim.

Let a_i denote the i th row of A . Almost surely, the rows a_1, \dots, a_{n-1} are linearly independent. We are going to characterize $\|A^{-1}e_n\|$ in a geometric way. Let $S_n := \text{span}\{a_1, \dots, a_{n-1}\}$ and denote by a_n^\perp the orthogonal projection of a_n onto S_n^\perp . Consider $w := A^{-1}e_n$, which is the n th column of A^{-1} . Since $AA^{-1} = \mathbf{I}$ we have $\langle w, a_i \rangle = 0$ for $i = 1, \dots, n-1$ and hence $w \in S_n^\perp$. Moreover, $\langle w, a_n \rangle = 1$, so $\|w\| \|a_n^\perp\| = 1$ and we arrive at

$$(7.2) \quad \|A^{-1}e_n\| = \frac{1}{\|a_n^\perp\|}.$$

Let $A_n \in \mathbb{C}^{(n-1) \times n}$ denote the matrix obtained from A by omitting a_n . We shall write $\text{vol}(A_n) = \det(AA^*)^{1/2}$ for the $(n-1)$ -dimensional volume of the parallelepiped spanned by the rows of A_n . Similarly, $|\det A|$ can be interpreted as the n -dimensional volume of the parallelepiped spanned by the rows of A .

Now we write $\rho(A) = \rho_1(A_n)\rho_2(a_n)$, where ρ_1 and ρ_2 are the density functions of $N(\bar{A}_n, \sigma^2\mathbf{I})$ and $N(\bar{a}_n, \sigma^2\mathbf{I})$, respectively (the meaning of \bar{A}_n and \bar{a}_n being clear). Moreover, note that

$$\text{vol}(A)^2 = \text{vol}(A_n)^2 \|a_n^\perp\|^2.$$

Fubini's theorem combined with (7.2) yields for $t > 0$

$$(7.3) \quad \int_{\|A^{-1}e_n\| \geq t} \text{vol}(A)^2 \rho(A) dA = \int_{A_n \in \mathbb{C}^{(n-1) \times n}} \text{vol}(A_n)^2 \rho_1(A_n) \cdot \left(\int_{\|a_n^\perp\| \leq 1/t} \|a_n^\perp\|^2 \rho_2(a_n) da_n \right) dA_n.$$

We next show that for fixed, linearly independent a_1, \dots, a_{n-1} and $\lambda > 0$

$$(7.4) \quad \int_{\|a_n^\perp\| \leq \lambda} \|a_n^\perp\|^2 \rho_2(a_n) da_n \leq \frac{\lambda^4}{2\sigma^2}.$$

For this, note that $a_n^\perp \sim N(\bar{a}_n^\perp, \sigma^2 \mathbf{I})$ in $S_n^\perp \simeq \mathbb{C}$, where \bar{a}_n^\perp is the orthogonal projection of \bar{a}_n onto S_n^\perp . Thus, proving (7.4) amounts to showing

$$\int_{|z| \leq \lambda} |z|^2 \rho_{\bar{z}}(z) dz \leq \frac{\lambda^4}{2\sigma^2}$$

for the Gaussian density $\rho_{\bar{z}}(z) = \frac{1}{2\pi\sigma^2} e^{-\frac{1}{2\sigma^2}|z-\bar{z}|^2}$ of $z \in \mathbb{C}$, where $\bar{z} \in \mathbb{C}$. Clearly, it is enough to show that

$$\int_{|z| \leq \lambda} \rho_{\bar{z}}(z) dz \leq \frac{\lambda^2}{2\sigma^2}.$$

Without loss of generality we may assume that $\bar{z} = 0$, since the integral in the left-hand side is maximized at this value of \bar{z} . The substitution $z = \sigma w$ yields $dz = \sigma^2 dw$ (dz denotes the Lebesgue measure on \mathbb{R}^2) and we get

$$\begin{aligned} \int_{|z| \leq \lambda} \rho_0(z) dz &= \int_{|w| \leq \frac{\lambda}{\sigma}} \frac{1}{2\pi} e^{-\frac{1}{2}|w|^2} dw = \int_0^{\frac{\lambda}{\sigma}} \frac{1}{2\pi} e^{-\frac{1}{2}r^2} 2\pi r dr \\ &= -e^{-\frac{1}{2}r^2} \Big|_0^{\frac{\lambda}{\sigma}} = 1 - e^{-\frac{\lambda^2}{2\sigma^2}} \leq \frac{\lambda^2}{2\sigma^2}, \end{aligned}$$

which proves inequality (7.4).

A similar argument shows that

$$(7.5) \quad 2\sigma^2 \leq \int |z|^2 \rho_{\bar{z}}(z) dz = \int \|a_n^\perp\|^2 \rho_2(a_n) da_n.$$

Plugging in this inequality into (7.3) (with $t = 0$) we conclude that

$$(7.6) \quad 2\sigma^2 \mathbb{E}_{\rho_1}(\text{vol}(A_n)^2) \leq \mathbb{E}_\rho(\text{vol}(A)^2).$$

On the other hand, plugging in (7.4) with $\lambda = \frac{1}{t}$ into (7.3), we obtain

$$\int_{\|A^{-1}e_n\| \geq t} \text{vol}(A)^2 \rho(A) dA \leq \frac{1}{2\sigma^2 t^4} \mathbb{E}_{\rho_1}(\text{vol}(A_n)^2).$$

Combined with (7.6) this yields

$$\int_{\|A^{-1}e_n\| \geq t} \text{vol}(A)^2 \rho(A) dA \leq \frac{1}{4\sigma^4 t^4} \mathbb{E}_\rho(\text{vol}(A)^2).$$

By the definition of the density $\tilde{\rho}$, this means that

$$\text{Prob}_{A \sim \tilde{\rho}} \{ \|A^{-1}e_n\| \geq t \} \leq \frac{1}{4\sigma^4 t^4},$$

which was to be shown. □

LEMMA 7.3. *For fixed $u \in \mathbb{S}^{n-1}$, $0 \leq s \leq 1$, and random v uniformly chosen in \mathbb{S}^{n-1} , we have*

$$\text{Prob}_v \{ |u^T v| \geq s \} = (1 - s^2)^{n-1}.$$

Proof. Recall the Riemannian distance $d_{\mathbb{P}}$ in $\mathbb{P}^{n-1} := \mathbb{P}(\mathbb{C}^n)$ from (2.1). Accordingly, for $0 \leq \theta \leq \pi/2$, we have

$$\text{Prob}_v \left\{ |u^T v| \geq \cos \theta \right\} = \frac{\text{vol} \left\{ [v] \in \mathbb{P}^{n-1} \mid d_{\mathbb{P}}([u], [v]) \leq \theta \right\}}{\text{vol } \mathbb{P}^{n-1}} = (\sin \theta)^{2(n-1)},$$

where the last equality is due to [11, Lemma 2.1]. □

LEMMA 7.4. *For any $t > 0$, we have*

$$\text{Prob}_{A \sim \tilde{\rho}} \left\{ \|A^{-1}\| \geq t \right\} \leq \frac{e^2(n+1)^2}{16\sigma^4} \frac{1}{t^4}.$$

Proof. We use an idea in Sankar et al. [19, §3]. For any invertible $A \in \mathbb{C}^{n \times n}$ there exists $u \in \mathbb{S}^{n-1}$ such that $\|A^{-1}u\| = \|A^{-1}\|$. For almost all A , the vector u is uniquely determined up to a scaling factor θ of modulus 1. We shall denote by u_A a representative of such u .

The following is an easy consequence of the singular value decomposition of $\|A^{-1}\|$: for any $v \in \mathbb{S}^{n-1}$, we have

$$(7.7) \quad \|A^{-1}v\| \geq \|A^{-1}\| \cdot |u_A^T v|.$$

We choose now a random pair (A, v) with A following the law $\tilde{\rho}$ and, independently, $v \in \mathbb{S}^{n-1}$ from the uniform distribution. Lemma 7.2 implies that

$$\text{Prob}_{A,v} \left\{ \|A^{-1}v\| \geq t \sqrt{\frac{2}{n+1}} \right\} \leq \frac{(n+1)^2}{16\sigma^4 t^4}.$$

On the other hand, by (7.7) we have

$$\begin{aligned} & \text{Prob}_{A,v} \left\{ \|A^{-1}v\| \geq t \sqrt{2/(n+1)} \right\} \\ & \geq \text{Prob}_{A,v} \left\{ \|A^{-1}\| \geq t \text{ and } |u_A^T v| \geq \sqrt{2/(n+1)} \right\} \\ & \geq \text{Prob}_A \left\{ \|A^{-1}\| \geq t \right\} \text{Prob}_{A,v} \left\{ |u_A^T v| \geq \sqrt{2/(n+1)} \mid \|A^{-1}\| \geq t \right\}. \end{aligned}$$

Lemma 7.3 tells us that for any fixed $u \in \mathbb{S}^{n-1}$, we have

$$\text{Prob}_v \left\{ |u^T v| \geq \sqrt{2/(n+1)} \right\} = (1 - 2/(n+1))^{n-1} \geq e^{-2},$$

the last inequality as $(\frac{n+1}{n-1})^{n-1} = (1 + \frac{2}{n-1})^{n-1} \leq e^2$. We thus obtain

$$\text{Prob}_A \left\{ \|A^{-1}\| \geq t \right\} \leq e^2 \text{Prob}_{A,v} \left\{ \|A^{-1}v\| \geq t \sqrt{\frac{2}{n+1}} \right\} \leq \frac{e^2(n+1)^2}{16\sigma^4 t^4},$$

as claimed. □

Proof of Proposition 7.1. By Lemma 7.4 we obtain, for any $T_0 > 0$,

$$\begin{aligned} \mathbb{E}(\|A^{-1}\|^2) &= \int_0^\infty \text{Prob}\{\|A^{-1}\|^2 \geq T\} dT \\ &\leq T_0 + \int_{T_0}^\infty \text{Prob}\{\|A^{-1}\|^2 \geq T\} dT \leq T_0 + \frac{\epsilon^2(n+1)^2}{16\sigma^4} \frac{1}{T_0}, \end{aligned}$$

using $\int_{T_0}^\infty T^{-2} dT = T_0^{-1}$. Now choose $T_0 = \frac{\epsilon(n+1)}{4\sigma^2}$. □

8. Smoothed analysis of the mean square condition number

The goal of this section is to accomplish the proof of Theorem 3.6.

8.1. *Orthogonal decompositions of \mathcal{H}_d .* For reasons to become clear soon we have to distinguish points in \mathbb{P}^n from their representatives ζ in the sphere $\mathbb{S}^n = \{\zeta \in \mathbb{C}^{n+1} \mid \|\zeta\| = 1\}$.

For $\zeta \in \mathbb{S}^n$ we consider the subspace R_ζ of \mathcal{H}_d consisting of all systems h that vanish at ζ of higher order:

$$R_\zeta := \{h \in \mathcal{H}_d \mid h(\zeta) = 0, Dh(\zeta) = 0\}.$$

We further decompose the orthogonal complement R_ζ^\perp of R_ζ in \mathcal{H}_d (defined with respect to the Bombieri-Weyl Hermitian inner product). Let L_ζ denote the subspace of R_ζ^\perp consisting of the systems vanishing at ζ and let C_ζ denote its orthogonal complement in R_ζ^\perp . Then we have an orthogonal decomposition

$$(8.1) \quad \mathcal{H}_d = C_\zeta \oplus L_\zeta \oplus R_\zeta$$

parametrized by $\zeta \in \mathbb{S}^n$.

LEMMA 8.1. *The space C_ζ consists of the systems $(c_i \langle X, \zeta \rangle^{d_i})$ with $c_i \in \mathbb{C}$. The space L_ζ consists of the systems*

$$g = (\sqrt{d_i} \langle X, \zeta \rangle^{d_i-1} \ell_i),$$

where ℓ_i is a linear form vanishing at ζ . Moreover, if $\ell_i = \sum_{j=0}^n m_{ij} X_j$ with $M = (m_{ij})$, then $\|g\| = \|M\|_F$.

Proof. By unitary invariance it suffices to verify the assertions in the case $\zeta = (1, 0, \dots, 0)$. In this case this follows easily from the definition of the Bombieri-Weyl inner product. □

The Bombieri-Weyl inner product on \mathcal{H}_d and the standard metric on the sphere \mathbb{S}^n define a Riemannian metric on $\mathcal{H}_d \times \mathbb{S}^n$ on which the unitary group $\mathcal{U}(n+1)$ operates isometrically. The “lifting”

$$V := \{(q, \zeta) \in \mathcal{H}_d \times \mathbb{S}^n \mid q(\zeta) = 0\}$$

of the solution variety $V_{\mathbb{P}}$ is easily seen to be a $\mathcal{U}(n+1)$ -invariant Riemannian submanifold of $\mathcal{H}_d \times \mathbb{S}^n$.

The projection $\pi_2: V \rightarrow \mathbb{S}^n, (q, \zeta) \mapsto \zeta$ defines a vector bundle with fibers $V_\zeta := \pi_2^{-1}(\zeta)$. In fact, (8.1) can be interpreted as an orthogonal decomposition of the trivial Hermitian vector bundle $\mathcal{H}_d \times \mathbb{S}^n \rightarrow \mathbb{S}^n$ into subbundles $C, L,$ and R over \mathbb{S}^n . Moreover, the vector bundle V is the orthogonal sum of L and R : we have $V_\zeta = L_\zeta \oplus R_\zeta$ for all ζ .

In the special case where all the degrees d_i are one, \mathcal{H}_d can be identified with the space $\mathcal{M} := \mathbb{C}^{n \times (n+1)}$ of matrices and the solution manifold V specializes to the manifold

$$W := \{(M, \zeta) \in \mathcal{M} \times \mathbb{S}^n \mid M\zeta = 0\}.$$

The map π_2 specializes to the vector bundle $p_2: W \rightarrow \mathbb{S}^n, (M, \zeta) \mapsto \zeta$ with the fibers

$$W_\zeta := \{M \in \mathcal{M} \mid M\zeta = 0\}.$$

Lemma 8.1 tells us that for each ζ we have *isometrical* linear maps

$$(8.2) \quad W_\zeta \rightarrow L_\zeta, \quad M \mapsto g_{M,\zeta} := (\sqrt{d_i} \langle X, \zeta \rangle^{d_i-1} \sum_j m_{ij} X_j).$$

In other words, the Hermitian vector bundles W and L over \mathbb{S}^n are isometric. The fact that the map (8.2) depends on the choice of the representative of ζ forces us to work over \mathbb{S}^n instead over \mathbb{P}^n . (All other notions introduced so far only depend on the base point in \mathbb{P}^n .)

We compose the orthogonal bundle projection $V_\zeta = L_\zeta \oplus R_\zeta \rightarrow L_\zeta$ with the bundle isometry $L_\zeta \simeq W_\zeta$ obtaining the map of vector bundles

$$(8.3) \quad \Psi: V \rightarrow W, \quad (g_{M,\zeta} + h, \zeta) \mapsto (M, \zeta)$$

with fibers $\Psi^{-1}(M, \zeta)$ isometric to R_ζ .

LEMMA 8.2. *We have $\Psi(q, \zeta) = (\Delta^{-1}Dq(\zeta), \zeta)$, where $\Delta := \text{diag}(\sqrt{d_i})$.*

Proof. Let $(q, \zeta) \in V$ and $(M, \zeta) := \Psi(q, \zeta)$. Then we have the decomposition $q = 0 + g_{M,\zeta} + h \in C_\zeta \oplus L_\zeta \oplus R_\zeta$. It is easily checked that $Dg_{M,\zeta}(\zeta) = \Delta M$. Since $Dq(\zeta) = Dg_{M,\zeta}(\zeta)$ we obtain $M = \Delta^{-1}Dq(\zeta)$. \square

The lemma shows that the condition number $\mu_{\text{norm}}(q, \zeta)$ (cf. §2.3) can be described in terms of Ψ as follows:

$$(8.4) \quad \frac{\mu_{\text{norm}}(q, \zeta)}{\|q\|} = \|M^\dagger\|, \quad \text{where } (M, \zeta) = \Psi(q, \zeta).$$

8.2. *Outline of proof of Theorem 3.6.* Let $\rho_{\mathcal{H}_d}$ denote the density of the Gaussian $N(\bar{q}, \sigma^2 \mathbf{I})$ on \mathcal{H}_d , where $\bar{q} \in \mathcal{H}_d$ and $\sigma > 0$. For fixed $\zeta \in \mathbb{S}^n$ we decompose the mean \bar{q} as

$$\bar{q} = \bar{k}_\zeta + \bar{g}_\zeta + \bar{h}_\zeta \in C_\zeta \oplus L_\zeta \oplus R_\zeta$$

according to (8.1). If we denote by ρ_{C_ζ} , ρ_{L_ζ} , and ρ_{R_ζ} the densities of the Gaussian distributions in the spaces C_ζ , L_ζ , and R_ζ with covariance matrices $\sigma^2\mathbf{I}$ and means $\bar{k}_\zeta, \bar{M}_\zeta$, and \bar{h}_ζ , respectively, then the density $\rho_{\mathcal{H}_d}$ factors as

$$(8.5) \quad \rho_{\mathcal{H}_d}(k + g + h) = \rho_{C_\zeta}(k) \cdot \rho_{L_\zeta}(g) \cdot \rho_{R_\zeta}(h).$$

The Gaussian density ρ_{L_ζ} on L_ζ induces a Gaussian density ρ_{W_ζ} on the fiber W_ζ with the covariance matrix $\sigma^2\mathbf{I}$ via the isometrical linear map (8.2), so $\rho_{W_\zeta}(M) = \rho_{L_\zeta}(g_{M,\zeta})$.

We derive now from the given Gaussian distribution $\rho_{\mathcal{H}_d}$ on \mathcal{H}_d a probability distribution on V as follows (naturally extending ρ_{st} introduced in §3.2). Think of choosing (q, ζ) at random from V by first choosing $q \in \mathcal{H}_d$ from $N(\bar{q}, \sigma^2\mathbf{I})$, then choosing one of its \mathcal{D} zeros $[\zeta] \in \mathbb{P}^n$ at random from the uniform distribution on $\{1, \dots, \mathcal{D}\}$, and finally choosing a representative ζ in the unit circle $[\zeta] \cap \mathbb{S}^n$ uniformly at random. (An explicit expression of the corresponding probability density ρ_V on V is given in (8.23).)

The plan to show Theorem 3.6 is as follows. The forthcoming Lemma 8.8 tells us that

$$(8.6) \quad \mathbb{E}_{\mathcal{H}_d} \left(\frac{\mu_2^2(q)}{\|q\|^2} \right) = \mathbb{E}_V \left(\frac{\mu_{\text{norm}}^2(q, \zeta)}{\|q\|^2} \right),$$

where $\mathbb{E}_{\mathcal{H}_d}$ and \mathbb{E}_V refer to the expectations with respect to the distribution $N(\bar{q}, \sigma^2\mathbf{I})$ on \mathcal{H}_d and the probability density ρ_V on V , respectively. Moreover, by equation (8.4),

$$\mathbb{E}_V \left(\frac{\mu_{\text{norm}}^2(q, \zeta)}{\|q\|^2} \right) = \mathbb{E}_{\mathcal{M}} (\|M^\dagger\|^2),$$

where $\mathbb{E}_{\mathcal{M}}$ denotes the expectation with respect to the pushforward density $\rho_{\mathcal{M}}$ of ρ_V with respect to the map $p_1 \circ \Psi: V \rightarrow \mathcal{M}$ (for more on pushforwards, see § 8.3).

Of course, we need to better understand the density $\rho_{\mathcal{M}}$. Let $M \in \mathcal{M}$ be of rank n and $\zeta \in \mathbb{S}^n$ with $M\zeta = 0$. The following formula

$$(8.7) \quad \rho_{\mathcal{M}}(M) = \rho_{C_\zeta}(0) \cdot \frac{1}{2\pi} \int_{\lambda \in S^1} \rho_{W_{\lambda\zeta}}(M) dS^1$$

can be heuristically explained as follows. We decompose a random $q \in \mathcal{H}_d$ according to the decomposition $\mathcal{H}_d = C_\zeta \oplus L_\zeta \oplus R_\zeta$ as $q = k + g + h$. Choose $\lambda \in \mathbb{C}$ with $|\lambda| = 1$ uniformly at random in the unit circle. Then we have $\Psi(q, \lambda\zeta) = (M, \lambda\zeta)$ if and only if $k = 0$ and g is mapped to M under the isometry in (8.2). The probability density for the event $k = 0$ equals $\rho_{C_\zeta}(0)$. The second event, conditioned on λ , has the probability density $\rho_{W_{\lambda\zeta}}(M)$.

By general principles (cf. §8.3) we have

$$(8.8) \quad \mathbb{E}_{\mathcal{M}} (\|M^\dagger\|^2) = \mathbb{E}_{\zeta \sim \rho_{\mathbb{S}^n}} \left(\mathbb{E}_{M \sim \rho_{W_\zeta}} (\|M^\dagger\|^2) \right),$$

where $\rho_{\mathbb{S}^n}$ is the pushforward density of ρ_V with respect to $p_2 \circ \Psi: V \rightarrow \mathbb{S}^n$ and $\tilde{\rho}_{W_\zeta}$ denotes a “conditional density” on the fiber W_ζ . This conditional density turns out to be of the form

$$(8.9) \quad \tilde{\rho}_{W_\zeta}(M) = c_\zeta^{-1} \cdot \det(MM^*) \rho_{W_\zeta}(M)$$

(c_ζ denoting a normalization factor). In the case $\zeta = (1, 0, \dots, 0)$ we can identify W_ζ with $\mathbb{C}^{n \times n}$ and $\tilde{\rho}_{W_\zeta}$ takes the form (7.1) studied in Section 7. Proposition 7.1 and unitary invariance imply that for all $\zeta \in \mathbb{S}^n$,

$$(8.10) \quad \mathbb{E}_{M \sim \tilde{\rho}_{W_\zeta}} (\|M^\dagger\|^2) \leq \frac{e(n+1)}{2\sigma^2}.$$

This implies by (8.8) that

$$\mathbb{E}_{\mathcal{M}} (\|M^\dagger\|^2) \leq \frac{e(n+1)}{2\sigma^2}$$

and completes the outline of the proof of Theorem 3.6.

The formal proof of the stated facts (8.7)–(8.9) is quite involved and will be given in the remainder of this section.

8.3. *Coarea formula.* We begin by recalling the coarea formula that tells us how probability distributions on Riemannian manifolds transform.

Suppose that X, Y are Riemannian manifolds of dimensions m, n , respectively such that $m \geq n$. Let $\varphi: X \rightarrow Y$ be differentiable. By definition, the derivative $D\varphi(x): T_x X \rightarrow T_{\varphi(x)} Y$ at a regular point $x \in X$ is surjective. Hence the restriction of $D\varphi(x)$ to the orthogonal complement of its kernel yields a linear isomorphism. The absolute value of its determinant is called the *normal Jacobian* of φ at x and denoted $\text{NJ}\varphi(x)$. We set $\text{NJ}\varphi(x) := 0$ if x is not a regular point. We note that the fiber $F_y := \varphi^{-1}(y)$ is a Riemannian submanifold of X of dimension $m - n$ if y is a regular value of φ . Sard’s lemma states that almost all $y \in Y$ are regular values.

The following result is the coarea formula, sometimes also called Fubini’s theorem for Riemannian manifolds. A proof can be found e.g., in [16, Appendix].

PROPOSITION 8.3. *Suppose that X and Y are Riemannian manifolds of dimensions m and n , respectively, and let $\varphi: X \rightarrow Y$ be a surjective differentiable map. Put $F_y = \varphi^{-1}(y)$. Then, for any function $\chi: X \rightarrow \mathbb{R}$ that is integrable with respect to the volume measure of X , we have that*

$$\int_X \chi dX = \int_{y \in Y} \left(\int_{F_y} \frac{\chi}{\text{NJ}\varphi} dF_y \right) dY.$$

Now suppose that we are in the situation described in the statement of Proposition 8.3, and we have a probability measure on X with density ρ_X . For

a regular value $y \in Y$ we set

$$(8.11) \quad \rho_Y(y) = \int_{F_y} \frac{\rho_X}{\text{NJ}\varphi} dF_y.$$

The coarea formula implies that for all measurable sets $B \subseteq Y$, we have

$$\int_{\varphi^{-1}(B)} \rho_X dX = \int_B \rho_Y dY.$$

Hence ρ_Y is a probability density on Y . We call it the *pushforward* of ρ_X with respect to φ .

For a regular value $y \in Y$ and $x \in F_y$ we define

$$(8.12) \quad \rho_{F_y}(x) = \frac{\rho_X(x)}{\rho_Y(y)\text{NJ}\varphi(x)}.$$

Clearly, this defines a probability density on F_y . The coarea formula implies that for all measurable functions $\chi: X \rightarrow \mathbb{R}$,

$$\int_X \chi \rho_X dX = \int_{y \in Y} \left(\int_{F_y} \chi \rho_{F_y} dF_y \right) \rho_Y(y) dY,$$

provided the left-hand integral exists. Therefore, we can interpret ρ_{F_y} as the *density of the conditional distribution* of x on the fiber F_y and briefly express the formula above in probabilistic terms as

$$(8.13) \quad \mathbb{E}_{x \sim \rho_X} (\chi(x)) = \mathbb{E}_{y \sim \rho_Y} \left(\mathbb{E}_{x \sim \rho_{F_y}} (\chi(x)) \right).$$

To put these formulas at use in our context, we must compute the normal Jacobians of some maps.

8.4. Normal Jacobians. We start with a general comment. Note that the \mathbb{R} -linear map $\mathbb{C} \rightarrow \mathbb{C}, z \mapsto \lambda z$ with $\lambda \in \mathbb{C}$ has determinant $|\lambda|^2$. More generally, let φ be an endomorphism of a finite dimensional complex vector space. Then $|\det \varphi|^2$ equals the determinant of φ , seen as a \mathbb{R} -linear map.

We describe now the normal Jacobian of the projection $p_1: W \rightarrow \mathcal{M}$ following [23].

LEMMA 8.4. *We have $\text{NJ}p_1(M, \zeta) = \prod_{i=1}^n (1 + \sigma_i^{-2})^{-1}$, where $\sigma_1, \dots, \sigma_n$ are the singular values of M .*

Proof. First note that $T_\zeta \mathbb{S}^n = \{ \dot{\zeta} \in \mathbb{C}^{n+1} \mid \text{Re} \langle \zeta, \dot{\zeta} \rangle = 0 \}$. The tangent space $T_{(M, \zeta)} W$ consists of the $(\dot{M}, \dot{\zeta}) \in \mathcal{M} \times T_\zeta \mathbb{S}^n$ such that $\dot{M}\zeta + M\dot{\zeta} = 0$.

By unitary invariance we may assume that $\zeta = (1, 0, \dots, 0)$. Then the first column of M vanishes, and we denote by $A = [m_{ij}] \in \mathbb{C}^{n \times n}$ the remaining part of M . Without loss of generality we may assume that A is invertible. Further, let $\dot{u} \in \mathbb{C}^n$ denote the first column of \dot{M} and $\dot{A} \in \mathbb{C}^{n \times n}$ its remaining part. We

may thus identify $T_{(M,\zeta)}W$ with the product $E \times \mathbb{C}^{n \times n}$ via $(\dot{M}, \dot{\zeta}) \mapsto ((\dot{u}, \dot{\zeta}), \dot{A})$, where E denotes the subspace

$$E := \left\{ (\dot{u}, \dot{\zeta}) \in \mathbb{C}^n \times \mathbb{C}^{n+1} \mid \dot{u}_i + \sum_{j=1}^n m_{ij} \dot{\zeta}_j = 0, 1 \leq i \leq n, \dot{\zeta}_0 \in i\mathbb{R} \right\}.$$

We also note that $E \simeq \text{graph}(-A) \times i\mathbb{R}$. The derivative of p_1 is described by the following commutative diagram:

$$\begin{array}{ccc} T_{(M,\zeta)}W & \xrightarrow{\simeq} & (\text{graph}(-A) \times i\mathbb{R}) \times \mathbb{C}^{n \times n} \\ Dp_1(M,\zeta) \downarrow & & \downarrow \text{pr} \times \text{id} \\ \mathcal{M} & \xrightarrow{\simeq} & \mathbb{C}^n \times \mathbb{C}^{n \times n}, \end{array}$$

where $\text{pr}(\dot{u}, \dot{\zeta}) = \dot{u}$. Using the singular value decomposition we may assume that $A = \text{diag}(\sigma_1, \dots, \sigma_n)$. Then the pseudoinverse of the projection pr is given by the \mathbb{R} -linear map

$$\varphi: \mathbb{C}^n \rightarrow \text{graph}(-A), \dot{u} \mapsto (\dot{u}, -\sigma_1^{-1} \dot{u}_1, \dots, -\sigma_n^{-1} \dot{u}_n).$$

It is easy to see that $\det \varphi = \prod_{i=1}^n (1 + \sigma_i^{-2})$. To complete the proof we note that $1/\text{NJ}p_1(M, \zeta) = \det \varphi$. □

We have already seen that the condition number $\mu_{\text{norm}}(q, \zeta)$ can be described in terms of the map Ψ introduced in (8.3). As a stepping stone towards the analysis of the normal Jacobian of Ψ we introduce now the related bundle map

$$\Phi: V \rightarrow W, (q, \zeta) \mapsto (Dq(\zeta), \zeta),$$

whose normal Jacobian turns out to be constant. (This crucial observation is due to Beltrán and Pardo in [6].)

PROPOSITION 8.5. *We have $\text{NJ}\Phi(q, \zeta) = \mathcal{D}^n$ for all $(q, \zeta) \in V$.*

Proof. By unitary invariance we may assume without loss of generality that $\zeta = (1, 0, \dots, 0)$. If we write $N = (n_{ij}) = Dq(\zeta) \in \mathcal{M}$, then we must have $n_{i0} = 0$ since $N\zeta = 0$. Moreover, according to the orthogonal decomposition (8.1) and Lemma 8.1, we have, for $1 \leq i \leq n$,

$$q_i = X_0^{d_i-1} \sum_{j=1}^n n_{ij} X_j + h_i$$

for some $h = (h_1, \dots, h_n) \in R_\zeta$. We further express $\dot{q}_i \in T_q \mathcal{H}_d = \mathcal{H}_d$ as

$$\dot{q}_i = \dot{u}_i X_0^{d_i} + \sqrt{d_i} X_0^{d_i-1} \sum_{j=1}^n \dot{a}_{ij} X_j + \dot{h}_i$$

in terms of the coordinates $\dot{u} = (\dot{u}_i) \in \mathbb{C}^n$, $\dot{A} = (\dot{a}_{ij}) \in \mathbb{C}^{n \times n}$, and $\dot{h} = (\dot{h}_i) \in R_\zeta$. The reason to put the factor $\sqrt{d_i}$ here is that

$$(8.14) \quad \|\dot{q}\|^2 = \sum_i |\dot{u}_i|^2 + \sum_{ij} |\dot{a}_{ij}|^2 + \sum_i \|\dot{h}_i\|^2$$

by the definition of the Bombieri-Weyl inner product.

The tangent space $T_{(q,\zeta)}V$ consists of the $(\dot{q}, \dot{\zeta}) \in \mathcal{H}_d \times T_\zeta \mathbb{S}^n$ such that $\dot{q}(\zeta) + N\dot{\zeta} = 0$; see [8, §10.3, Prop. 1]. This condition can be expressed in coordinates as

$$(8.15) \quad \dot{u}_i + \sum_{j=1}^n n_{ij} \dot{\zeta}_j = 0, \quad i = 1, \dots, n.$$

By (8.14) the inner product on $T_{(q,\zeta)}V$ is given by the standard inner product in the chosen coordinates $\dot{u}_i, \dot{a}_{ij}, \dot{\zeta}_j$ if $\dot{h}_i = 0$. Thinking of the description of $T_{(N,\zeta)}W$ given in the proof of Lemma 8.4, we may therefore isometrically identify $T_{(q,\zeta)}V$ with the product $T_{(N,\zeta)}W \times R_\zeta$ via $(\dot{q}, \dot{\zeta}) \mapsto ((\dot{u}, \dot{A}, \dot{\zeta}), \dot{h})$. The derivative of π_1 is then described by the commutative diagram

$$(8.16) \quad \begin{array}{ccc} T_{(q,\zeta)}V & \xrightarrow{\cong} & T_{(N,\zeta)}W \times R_\zeta \\ D\pi_1(q,\zeta) \downarrow & & \downarrow Dp_1(N,\zeta) \times \text{id} \\ \mathcal{H}_d & \xrightarrow{\cong} & \mathcal{M} \times R_\zeta. \end{array}$$

We shall next calculate the derivative of Φ . For this, we will use the shorthand $\partial_k q$ for the partial derivative $\partial_{X_k} q$, etc. A short calculation yields, for $j > 0$,

$$(8.17) \quad \partial_0 \dot{q}_i(\zeta) = d_i \dot{u}_i, \quad \partial_j \dot{q}_i(\zeta) = \sqrt{d_i} \dot{a}_{ij}, \quad \partial_{0j}^2 q_i(\zeta) = (d_i - 1) n_{ij}.$$

Similarly, we obtain $\partial_0 q_i(\zeta) = 0$ and $\partial_j q_i(\zeta) = n_{ij}$ for $j > 0$.

The derivative of $D\Phi(q, \zeta): T_{(q,\zeta)}V \rightarrow T_{(N,\zeta)}W$ is determined by

$$D\Phi(q, \zeta)(\dot{q}, \dot{\zeta}) = (\dot{N}, \dot{\zeta}), \quad \text{where } \dot{N} = D\dot{q}(\zeta) + D^2q(\zeta)(\dot{\zeta}, \cdot).$$

Introducing the coordinates $\dot{N} = (\dot{n}_{ij})$ this can be written as

$$(8.18) \quad \dot{n}_{ij} = \partial_j \dot{q}_i(\zeta) + \sum_{k=1}^n \partial_{jk}^2 q_i(\zeta) \dot{\zeta}_k.$$

For $j > 0$, this gives, using (8.17),

$$(8.19) \quad \dot{n}_{ij} = \sqrt{d_i} \dot{a}_{ij} + \sum_{k=1}^n \partial_{jk}^2 q_i(\zeta) \dot{\zeta}_k.$$

For $j = 0$, we obtain from (8.18), using (8.17) and (8.15),

$$(8.20) \quad \dot{n}_{i0} = \partial_0 \dot{q}_i(\zeta) + \sum_{k=1}^n \partial_{0k}^2 q_i(\zeta) \dot{\zeta}_k = d_i \dot{u}_i + (d_i - 1) \sum_{k=1}^n n_{ik} \dot{\zeta}_k = \dot{u}_i.$$

Note the crucial cancellation taking place here!

From (8.19) and (8.20) we see that the kernel K of $D\Phi(q, \zeta)$ is determined by the conditions $\dot{\zeta} = 0, \dot{u} = 0, \dot{A} = 0$. Hence, recalling $T_{(q, \zeta)}V \simeq T_{(N, \zeta)}W \times R_\zeta$, we have $K \simeq 0 \times R_\zeta$ and $K^\perp \simeq T_{(N, \zeta)}W \times 0$. Moreover, as in the proof of Lemma 8.4 (but replacing M by N) we write

$$E := \left\{ (\dot{u}, \dot{\zeta}) \in \mathbb{C}^n \times \mathbb{C}^{n+1} \mid \dot{u}_i + \sum_{j=1}^n n_{ij} \dot{\zeta}_j = 0, 1 \leq i \leq n, \dot{\zeta}_0 \in i\mathbb{R} \right\}$$

and identify $T_{(N, \zeta)}W$ with $E \times \mathbb{C}^{n \times n}$. Using this identification of spaces, (8.19) and (8.20) imply that $D\Phi(q, \zeta)_{K^\perp}$ has the following structure:

$$\begin{aligned} D\Phi(q, \zeta)_{K^\perp} : E \times \mathbb{C}^{n \times n} &\rightarrow E \times \mathbb{C}^{n \times n}, \\ ((\dot{u}, \dot{\zeta}), \dot{A}) &\mapsto ((\dot{u}, \dot{\zeta}), \lambda(\dot{A}) + \rho(\dot{\zeta})), \end{aligned}$$

where the linear map $\lambda: \mathbb{C}^{n \times n} \rightarrow \mathbb{C}^{n \times n}, \dot{A} \mapsto (\sqrt{d_i} \dot{a}_{ij})$, multiplies the i th row of \dot{A} with $\sqrt{d_i}$ and $\rho: \mathbb{C}^{n+1} \rightarrow \mathbb{C}^{n \times n}$ is given by $\rho(\dot{\zeta})_{ij} = \sum_{k=1}^n \partial_{j k}^2 \dot{q}_i(\zeta) \dot{\zeta}_k$.

By definition we have $\text{NJ}\Phi(q, \zeta) = |\det D\Phi(q, \zeta)_{K^\perp}|$. The triangular form of $D\Phi(q, \zeta)_{K^\perp}$ shown above implies that $|\det D\Phi(q, \zeta)_{K^\perp}| = \det \lambda$. Finally, using the diagonal form of λ , we obtain $\det \lambda = \prod_{i=1}^n \sqrt{d_i}^2 = \mathcal{D}^n$, which completes the proof. \square

Remark 8.6. An inspection of the proof of Proposition 8.5 reveals that the second order derivatives occurring in $D\Phi$ do not have any impact on the normal Jacobian $\text{NJ}\Phi$. Its value \mathcal{D}^n occurs as a result of the chosen Bombieri-Weyl inner product on \mathcal{H}_d . With respect to the naive inner product on \mathcal{H}_d (where the monomials form an orthonormal basis), the normal Jacobian of Φ at (q, ζ) would be equal to one at $\zeta = (1, 0, \dots, 0)$. However unitary invariance would not hold and the normal Jacobian would take different values elsewhere.

Before proceeding we note the following consequence of equation (8.16):

$$(8.21) \quad \text{NJ}\pi_1(q, \zeta) = \text{NJ}p_1(N, \zeta), \quad \text{where } N = Dq(\zeta).$$

The normal Jacobian of the map $\Psi: V \rightarrow W$ is not constant and takes a more complicated form in terms of the normal Jacobians of the projection $p_1: W \rightarrow \mathcal{M}$. For obtaining an expression for $\text{NJ}\Psi$ we need the following lemma.

LEMMA 8.7. *The scaling map $\gamma: W \rightarrow W, (N, \zeta) \mapsto (M, \zeta)$ with $M = \Delta^{-1}N$ of rank n satisfies*

$$\det D\gamma(N, \zeta) = \frac{1}{\mathcal{D}^{n+1}} \cdot \frac{\text{NJ}p_1(N, \zeta)}{\text{NJ}p_1(M, \zeta)}.$$

Proof. If $W_{\mathbb{P}}$ denotes the solution variety in $\mathcal{M} \times \mathbb{P}^n$ analogous to W , then we have $T_{(M, \zeta)}W = T_{(M, \zeta)}W_{\mathbb{P}} \oplus \mathbb{R}i\zeta$. Let $p'_1: W_{\mathbb{P}} \rightarrow \mathcal{M}$ denote the projection.

The derivative $D\gamma_{\mathbb{P}}(N, \zeta)$ of the corresponding scaling map $\gamma_{\mathbb{P}}: W_{\mathbb{P}} \rightarrow W_{\mathbb{P}}$ is determined by the commutative diagram

$$\begin{array}{ccc} T_{(N, \zeta)} W_{\mathbb{P}} & \xrightarrow{D\gamma_{\mathbb{P}}(N, \zeta)} & T_{(M, \zeta)} W_{\mathbb{P}} \\ Dp'_1(N, \zeta) \downarrow & & \downarrow Dp'_1(M, \zeta) \\ \mathcal{M} & \xrightarrow{\text{sc}} & \mathcal{M}, \end{array}$$

where the vertical arrows are linear isomorphisms. The assertion follows by observing that $\text{NJ}p_1(N, \zeta) = \det Dp'_1(N, \zeta)$, $\text{NJ}\gamma(N, \zeta) = \det D\gamma_{\mathbb{P}}(N, \zeta)$, and using that the \mathbb{R} -linear map $\text{sc}: \mathcal{M} \rightarrow \mathcal{M}$, $N \mapsto M = \Delta^{-1}N$ has the determinant $1/\mathcal{D}^{n+1}$. □

Proposition 8.5 combined with Lemma 8.7 immediately gives

$$(8.22) \quad \text{NJ}\Psi(q, \zeta) = \frac{1}{\mathcal{D}} \cdot \frac{\text{NJ}p_1(N, \zeta)}{\text{NJ}p_1(M, \zeta)}$$

for $N = Dq(\zeta)$, $M = \Delta^{-1}N$.

8.5. *Induced probability distributions.* By Bézout’s theorem, the fiber $V(q)$ of the projection $\pi_1: V \rightarrow \mathcal{H}_{\mathbf{d}}$ at $q \in \mathcal{H}_{\mathbf{d}}$ is a disjoint union of $\mathcal{D} = d_1 \cdots d_n$ unit circles and therefore has the volume $2\pi\mathcal{D}$, provided q does not lie in the discriminant variety.

Recall that $\rho_{\mathcal{H}_{\mathbf{d}}}$ denotes the density of the Gaussian distribution $N(\bar{q}, \sigma^2\mathbf{I})$ for fixed $\bar{q} \in \mathcal{H}_{\mathbf{d}}$ and $\sigma > 0$ and $\mathbb{E}_{\mathcal{H}_{\mathbf{d}}}$ stands for expectation taken with respect to that density. We associate with $\rho_{\mathcal{H}_{\mathbf{d}}}$ the function $\rho_V: V \rightarrow \mathbb{R}$ defined by

$$(8.23) \quad \rho_V(q, \zeta) := \frac{1}{2\pi\mathcal{D}} \rho_{\mathcal{H}_{\mathbf{d}}}(q) \text{NJ}\pi_1(q, \zeta).$$

The next result shows that ρ_V is the probability density function of the distribution on V we described in §8.2.

- LEMMA 8.8. (1) *The function ρ_V is a probability density on V .*
 (2) *The expectation of a function $\varphi: V \rightarrow \mathbb{R}$ with respect to ρ_V can be expressed as $\mathbb{E}_V(\varphi) = \mathbb{E}_{\mathcal{H}_{\mathbf{d}}}(\varphi_{\text{av}})$, where*

$$\varphi_{\text{av}}(q) := \frac{1}{2\pi\mathcal{D}} \int_{V(q)} \varphi \, dV(q).$$

- (3) *The pushforward of ρ_V with respect to $\pi_1: V \rightarrow \mathcal{H}_{\mathbf{d}}$ equals $\rho_{\mathcal{H}_{\mathbf{d}}}$.*
 (4) *For $q \notin \Sigma$, the conditional density on the fiber $V(q)$ is the density of the uniform distribution on $V(q)$.*
 (5) *The probability density ρ_{st} on $V_{\mathbb{P}}$ introduced in Section 3.2 is obtained from the density ρ_V in the case $\bar{q} = 0$, $\sigma = 1$ as the pushforward under the canonical map $V \rightarrow V_{\mathbb{P}}$, $(f, \zeta) \mapsto (f, [\zeta])$. Explicitly, we have*

$$\rho_{\text{st}}(q, [\zeta]) = \frac{1}{\mathcal{D}} \frac{1}{(2\pi)^N} e^{-\frac{1}{2}\|q\|^2} \text{NJ}\pi_1(q, \zeta).$$

Proof. The coarea formula (Proposition 8.3) applied to $\pi_1 : V \rightarrow \mathcal{H}_d$ implies

$$\begin{aligned} \int_V \varphi \rho_V dV &= \int_{q \in \mathcal{H}_d} \left(\int_{\zeta \in V(q)} \varphi(q, \zeta) \frac{\rho_V(q, \zeta)}{\text{NJ}\pi_1(q, \zeta)} dV(q) \right) d\mathcal{H}_d \\ &= \int_{q \in \mathcal{H}_d} \varphi_{\text{av}}(q) \rho_{\mathcal{H}_d}(q) d\mathcal{H}_d. \end{aligned}$$

Taking $\varphi = 1$ reveals that ρ_V is a density, proving the first assertion. The above formula also shows the second assertion.

By equation (8.11) the pushforward density ρ of ρ_V with respect to π_1 satisfies

$$\rho(q) = \int_{\zeta \in V(q)} \frac{\rho_V(q, \zeta)}{\text{NJ}\pi_1(q, \zeta)} dV(q) = \rho_{\mathcal{H}_d}(q),$$

as $\int dV(q) = 2\pi\mathcal{D}$. This shows the third assertion. By (8.12) the conditional density satisfies

$$\rho_{V(q)}(q) = \frac{\rho_V(q, \zeta)}{\rho_{\mathcal{H}_d}(q) \text{NJ}\pi_1(q, \zeta)} = \frac{1}{2\pi\mathcal{D}},$$

which shows the fourth assertion. The fifth assertion is trivial. □

We can now determine the various probability distributions induced by ρ_V .

PROPOSITION 8.9. *We have*

$$\frac{\rho_V}{\text{NJ}\Psi}(g_{M,\zeta} + h, \zeta) = \rho_W(M, \zeta) \cdot \rho_{R_\zeta}(h),$$

where the pushforward density ρ_W of ρ_V with respect to $\Psi : V \rightarrow W$ satisfies

$$\rho_W(M, \zeta) = \frac{1}{2\pi} \rho_{C_\zeta}(0) \cdot \rho_{W_\zeta}(M) \cdot \text{NJ}p_1(M, \zeta).$$

Proof. Using the factorization of Gaussians (8.5) and equation (8.21), the density ρ_V can be written as

$$\rho_V(g_{M,\zeta} + h, \zeta) = \frac{1}{2\pi\mathcal{D}} \rho_{C_\zeta}(0) \rho_{W_\zeta}(M) \rho_{R_\zeta}(h) \text{NJ}p_1(N, \zeta),$$

where $N = \Delta M$. It follows from (8.22) that

$$(8.24) \quad \frac{\rho_V}{\text{NJ}\Psi}(g_{M,\zeta} + h, \zeta) = \frac{1}{2\pi} \rho_{C_\zeta}(0) \rho_{W_\zeta}(M) \rho_{R_\zeta}(h) \text{NJ}p_1(M, \zeta).$$

This implies, using (8.11) for $\Psi : V \rightarrow W$ and the isometry $\Psi^{-1}(M, \zeta) \simeq R_\zeta$ for the fiber at ζ , that

$$\begin{aligned} \rho_W(M, \zeta) &= \int_{h \in R_\zeta} \frac{\rho_V}{\text{NJ}\Psi}(g_{M,\zeta} + h, \zeta) dR_\zeta \\ &= \frac{1}{2\pi} \rho_{C_\zeta}(0) \cdot \rho_{W_\zeta}(M) \cdot \text{NJ}p_1(M, \zeta) \int_{h \in R_\zeta} \rho_{R_\zeta}(h) dR_\zeta \\ &= \frac{1}{2\pi} \rho_{C_\zeta}(0) \cdot \rho_{W_\zeta}(M) \cdot \text{NJ}p_1(M, \zeta) \end{aligned}$$

as claimed. Replacing in (8.24) we therefore obtain

$$\frac{\rho_V}{\text{NJ}\Psi}(g_{M,\zeta} + h, \zeta) = \rho_W(M, \zeta) \rho_{R_\zeta}(h). \quad \square$$

The claimed formula (8.7) for the pushforward density $\rho_{\mathcal{M}}$ of ρ_W with respect to $p_1: W \rightarrow \mathcal{M}$ immediately follows from Proposition 8.9 by integrating $\frac{\rho_W}{\text{NJ}p_1}$ over the fibers of p_1 .

LEMMA 8.10. *Let c_ζ denote the expectation of $\det(MM^*)$ with respect to ρ_{W_ζ} . We have*

$$\frac{\rho_W}{\text{NJ}p_2}(M, \zeta) = \rho_{\mathbb{S}^n}(\zeta) \cdot \tilde{\rho}_{W_\zeta}(M),$$

where $\rho_{\mathbb{S}^n}(\zeta) = \frac{c_\zeta}{2\pi} \rho_{C_\zeta}(0)$ is the pushforward density of ρ_W with respect to $p_2: W \rightarrow \mathbb{S}^n$, and where the conditional density $\tilde{\rho}_{W_\zeta}$ on the fiber W_ζ of p_2 is given by

$$\tilde{\rho}_{W_\zeta}(M) = c_\zeta^{-1} \cdot \det(MM^*) \rho_{W_\zeta}(M).$$

Proof. In [23] (see also [8, §13.2, Lemmas 2–3]) it is shown that

$$(8.25) \quad \frac{\text{NJ}p_1}{\text{NJ}p_2}(M, \zeta) = \det(MM^*).$$

Combining this with Proposition 8.9 we get

$$\frac{\rho_W}{\text{NJ}p_2}(M, \zeta) = \frac{1}{2\pi} \rho_{C_\zeta}(0) \cdot \rho_{W_\zeta}(M) \cdot \det(MM^*).$$

Integrating over W_ζ we get $\rho_{\mathbb{S}^n}(\zeta) = \frac{1}{2\pi} \rho_{C_\zeta}(0) \cdot c_\zeta$, and finally (cf. (8.12))

$$\tilde{\rho}_{W_\zeta}(M) = \frac{\rho_W(M, \zeta)}{\rho_{\mathbb{S}^n}(\zeta) \text{NJ}p_2(M, \zeta)} = c_\zeta^{-1} \cdot \rho_{W_\zeta}(M) \cdot \det(MM^*)$$

as claimed. □

This lemma shows that the conditional density $\tilde{\rho}_{W_\zeta}$ has the form stated in (8.9) and therefore completes the proof of Theorem 3.6.

8.6. *Expected number of real zeros.* As a further illustration of the interplay of Gaussians with the coarea formula in our setting, we give a simplified proof of one of the main results of [23]. This subsection is not needed for understanding the remainder of the paper.

Our developments so far took place over the complex numbers \mathbb{C} , but much of what has been said carries over the situation over \mathbb{R} . However, we note that algorithm ALH would not work over \mathbb{R} since the lifting of the segment $E_{f,g}$ will likely contain a multiple zero (over \mathbb{C} this happens with probability zero since the real codimension of the discriminant variety equals two).

Let $\mathcal{H}_{\mathbf{d},\mathbb{R}}$ denote the space of real polynomial systems in $\mathcal{H}_{\mathbf{d}}$ endowed with the Bombieri-Weyl inner product. The standard Gaussian distribution on $\mathcal{H}_{\mathbf{d},\mathbb{R}}$ is well defined, and we denote its density with $\rho_{\mathcal{H}_{\mathbf{d},\mathbb{R}}}$.

COROLLARY 8.11. *The average number of zeros of a standard Gaussian random $f \in \mathcal{H}_{\mathbf{d},\mathbb{R}}$ in the real projective space $\mathbb{P}^n(\mathbb{R})$ equals $\sqrt{\mathcal{D}}$.*

Proof. Let $\chi(q)$ denote the number of real zeros in $\mathbb{P}^n(\mathbb{R})$ of $q \in \mathcal{H}_{\mathbf{d},\mathbb{R}}$. Thus the number of real zeros in the sphere $S^n = S(\mathbb{R}^{n+1})$ equals $2\chi(q)$. The real solution variety $V_{\mathbb{R}} \subseteq \mathcal{H}_{\mathbf{d},\mathbb{R}} \times S^n$ is defined in the obvious way and so is $W_{\mathbb{R}} \subseteq \mathcal{M}_{\mathbb{R}} \times S^n$, where $\mathcal{M}_{\mathbb{R}} = \mathbb{R}^{n \times (n+1)}$.

The same proof as for Proposition 8.5 shows that the normal Jacobian of the map $\Phi_{\mathbb{R}}: V_{\mathbb{R}} \rightarrow W_{\mathbb{R}}, (q, \zeta) \mapsto (Dq(\zeta), \zeta)$ has the constant value $\mathcal{D}^{n/2}$ (the 2 in the exponent due to the considerations opening §8.4).

Applying the coarea formula to the projection $\pi_1: V_{\mathbb{R}} \rightarrow \mathcal{H}_{\mathbf{d},\mathbb{R}}$ yields

$$\begin{aligned} \int_{\mathcal{H}_{\mathbf{d},\mathbb{R}}} \chi \rho_{\mathcal{H}_{\mathbf{d},\mathbb{R}}} d\mathcal{H}_{\mathbf{d},\mathbb{R}} &= \int_{q \in \mathcal{H}_{\mathbf{d},\mathbb{R}}} \rho_{\mathcal{H}_{\mathbf{d},\mathbb{R}}}(q) \frac{1}{2} \int_{\pi_1^{-1}(q)} d\pi_1^{-1}(q) d\mathcal{H}_{\mathbf{d},\mathbb{R}} \\ &= \int_{V_{\mathbb{R}}} \frac{1}{2} \rho_{\mathcal{H}_{\mathbf{d},\mathbb{R}}} \text{NJ}\pi_1 dV_{\mathbb{R}}. \end{aligned}$$

We can factor the standard Gaussian $\rho_{\mathcal{H}_{\mathbf{d}}}$ into standard Gaussian densities $\rho_{C_{\zeta}}$ and $\rho_{L_{\zeta}}$ on C_{ζ} and L_{ζ} , respectively, as was done in Section 8.5 over \mathbb{C} (denoting them by the same symbol will not cause any confusion). We also have an isometry $W_{\zeta} \rightarrow L_{\zeta}$ as in (8.2) and $\rho_{L_{\zeta}}$ induces the standard Gaussian density $\rho_{W_{\zeta}}$ on W_{ζ} . The fiber of $\Phi_{\mathbb{R}}: V_{\mathbb{R}} \rightarrow W_{\mathbb{R}}, (q, \zeta) \mapsto (N, \zeta)$ over (N, ζ) has the form $\Phi_{\mathbb{R}}^{-1}(N, \zeta) = \{(g_{M,\zeta} + h, \zeta) \mid h \in R_{\zeta}\}$, where $M = \Delta^{-1}N$; cf. Lemma 8.2. We therefore have $\rho_{\mathcal{H}_{\mathbf{d},\mathbb{R}}}(g_{M,\zeta} + h) = \rho_{C_{\zeta}}(0) \rho_{W_{\zeta}}(M) \rho_{R_{\zeta}}(h)$.

The coarea formula applied to $\Phi_{\mathbb{R}}: V_{\mathbb{R}} \rightarrow W_{\mathbb{R}}$, using equation (8.21), yields

$$\begin{aligned} &\int_{V_{\mathbb{R}}} \frac{1}{2} \rho_{\mathcal{H}_{\mathbf{d},\mathbb{R}}} \text{NJ}\pi_1 dV_{\mathbb{R}} \\ &= \frac{1}{2 \text{NJ}\Phi_{\mathbb{R}}} \int_{(N,\zeta) \in W_{\mathbb{R}}} \rho_{C_{\zeta}}(0) \rho_{W_{\zeta}}(M) \text{NJ}p_1(N, \zeta) \int_{h \in R_{\zeta}} \rho_{R_{\zeta}}(h) dR_{\zeta} dW_{\mathbb{R}} \\ &= \frac{1}{2 \text{NJ}\Phi_{\mathbb{R}}} \int_{(N,\zeta) \in W_{\mathbb{R}}} \rho_{C_{\zeta}}(0) \rho_{W_{\zeta}}(M) \text{NJ}p_1(N, \zeta) dW_{\mathbb{R}}. \end{aligned}$$

Applying the coarea formula to the projection $p_1: W_{\mathbb{R}} \rightarrow \mathcal{M}_{\mathbb{R}}$, we can simplify the above to

$$\begin{aligned} &\frac{1}{\text{NJ}\Phi_{\mathbb{R}}} \int_{N \in \mathcal{M}_{\mathbb{R}}} \rho_{C_{\zeta}}(0) \rho_{W_{\zeta}}(M) \frac{1}{2} \int_{\zeta \in p_1^{-1}(N)} dp_1^{-1}(N) d\mathcal{M}_{\mathbb{R}} \\ &= \frac{1}{\text{NJ}\Phi_{\mathbb{R}}} \int_{N \in \mathcal{M}_{\mathbb{R}}} \rho_{C_{\zeta}}(0) \rho_{W_{\zeta}}(M) d\mathcal{M}_{\mathbb{R}} \\ &= \frac{\mathcal{D}^{\frac{n+1}{2}}}{\text{NJ}\Phi_{\mathbb{R}}} \int_{M \in \mathcal{M}_{\mathbb{R}}} \rho_{C_{\zeta}}(0) \rho_{W_{\zeta}}(M) d\mathcal{M}_{\mathbb{R}}, \end{aligned}$$

where the last equality is due to the change of variables $\mathcal{M}_{\mathbb{R}} \rightarrow \mathcal{M}_{\mathbb{R}}, N \mapsto M$ that has the Jacobian determinant $\mathcal{D}^{-\frac{n+1}{2}}$. Now we note that

$$\rho_{C_{\zeta}}(0) \cdot \rho_{W_{\zeta}}(M) = (2\pi)^{-n/2} (2\pi)^{-n^2/2} \exp\left(-\frac{1}{2}\|M\|_F^2\right)$$

is the density of the standard Gaussian distribution on $\mathcal{M}_{\mathbb{R}} \simeq \mathbb{R}^{n \times (n+1)}$, so that the last integral (over $M \in \mathcal{M}_{\mathbb{R}}$) equals one. Altogether, we obtain, using $\text{NJ}\Phi_{\mathbb{R}} = \mathcal{D}^{n/2}$,

$$\int_{\mathcal{H}_{\mathbf{d},\mathbb{R}}} \chi \rho_{\mathcal{H}_{\mathbf{d},\mathbb{R}}} d\mathcal{H}_{\mathbf{d},\mathbb{R}} = \frac{\mathcal{D}^{\frac{n+1}{2}}}{\text{NJ}\Phi_{\mathbb{R}}} = \sqrt{\mathcal{D}}. \quad \square$$

9. Effective sampling in the solution variety

We turn now to the question of effective sampling in the solution variety endowed with the measure ρ_{st} introduced in Section 3.2. The goal is to provide the proof of Proposition 3.3.

PROPOSITION 9.1. *In the setting of Section 8.5 suppose $\bar{q} = 0, \sigma = 1$. Then the pushforward density $\rho_{\mathcal{M}}$ of ρ_W with respect to $p_1: W \rightarrow \mathcal{M}$ equals the standard Gaussian distribution in \mathcal{M} . The conditional distributions on the fibers of p_1 are uniform distributions on unit circles. Finally, the conditional distribution on the fibers of $\Psi: V \rightarrow W$ is induced from the standard Gaussian in R_{ζ} via the isometry (8.2).*

Proof. Since $\rho_{\mathcal{H}_{\mathbf{d}}}$ is standard Gaussian, the induced distributions on C_{ζ} , L_{ζ} , and R_{ζ} are standard Gaussian as well. Hence $\rho_{W_{\zeta}}$ equals the standard Gaussian distribution on the fiber W_{ζ} . Moreover, $\rho_{C_{\zeta}}(0) = (\sqrt{2\pi})^{-2n}$. Equation (8.7) implies that

$$\rho_{\mathcal{M}}(M) = \rho_{C_{\zeta}}(0) \cdot \rho_{W_{\zeta}}(M) = (2\pi)^{-n} (2\pi)^{-n^2} \exp\left(-\frac{1}{2}\|M\|_F^2\right),$$

which equals the density of the standard Gaussian distribution on \mathcal{M} .

Lemma 8.10 combined with (8.25) gives

$$\frac{\rho_W}{\text{NJ}p_1}(M, \zeta) = \frac{1}{2\pi} \rho_{C_{\zeta}}(0) \cdot \rho_{W_{\zeta}}(M) = \frac{1}{2\pi} \rho_{\mathcal{M}}(M).$$

Hence the conditional distributions on the fibers of p_1 are uniform. (Note that this is not true in the case of nonstandard Gaussians.) The assertion on the conditional distributions on the fibers of Ψ follows from Proposition 8.9. \square

Proof of Proposition 3.3. Proposition 9.1 (with Lemma 8.8) shows that the following procedure generates the distribution ρ_{st} :

- (1) choose $M \in \mathcal{M}$ from the standard Gaussian distribution (almost surely M has rank n),
- (2) compute the unique $[\zeta] \in \mathbb{P}^n$ such that $M\zeta = 0$,
- (3) choose a representative ζ uniformly at random in $[\zeta] \cap \mathbb{S}^n$,
- (4) compute $g_{M,\zeta}$, cf. (8.2),
- (5) choose $h \in R_{\zeta}$ from the standard Gaussian distribution,
- (6) compute $q = g_{M,\zeta} + h$ and return $(q, [\zeta])$.

An elegant way of choosing h in step 5 is to draw $f \in \mathcal{H}_d$ from $N(0, I)$ and then to compute the image h of f under the orthogonal projection $\mathcal{H}_\zeta \rightarrow R_\zeta$. Since the orthogonal projection of a standard Gaussian is a standard Gaussian, this amounts to draw h from a standard Gaussian in R_ζ . For computing the projection h we note that the orthogonal decomposition $f = k + g_{M,\zeta} + h$ with $k \in C_\zeta$, $M = [m_{ij}] \in \mathcal{M}$, and $h \in R_\zeta$ is obtained as

$$\begin{aligned} k_i &= f_i(\zeta)\langle X, \zeta \rangle^{d_i}, \\ m_{ij} &= d_i^{-1/2}(\partial_{X_j} f_i(\zeta) - d_i f_i(\zeta)\bar{\zeta}_j), \\ h &= f - k - g_{M,\zeta}. \end{aligned}$$

(Recall $Dg_{M,\zeta}(\zeta) = \Delta M$ and note $\frac{\partial}{\partial X_j} \langle X, \zeta \rangle^{d_i}(\zeta) = d_i \bar{\zeta}_j$.)

It is easy to check that $\mathcal{O}(N)$ samples from the standard Gaussian distribution on \mathbb{R} are sufficient for implementing this procedure. As for the operation count: step (4) turns out to be the most expensive one and can be done, e.g., as follows. Suppose that all the coefficients of $\langle X, \zeta \rangle^{k-1}$ have already been computed. Then each coefficient of $\langle X, \zeta \rangle^k = (X_0 \bar{\zeta}_0 + \dots + X_n \bar{\zeta}_n)\langle X, \zeta \rangle^{k-1}$ can be obtained by $\mathcal{O}(n)$ arithmetic operations, hence all the coefficients of $\langle X, \zeta \rangle^k$ are obtained with $\mathcal{O}(n \binom{n+k}{n})$ operations. It follows that $\langle X, \zeta \rangle^{d_i}$ can be computed with $\mathcal{O}(d_i n N_i)$ operations, hence $\mathcal{O}(DnN)$ operations suffice for the computation of $g_{M,\zeta}$. It is clear that this is also an upper bound on the cost of computing (g, ζ) . \square

10. Homotopies with a fixed extremity

We provide now the proof of the remaining results stated in Section 3. The next two cases we wish to analyze (the condition-based analysis of LV and a solution for Smale’s 17th problem with moderate degrees) share the feature that one endpoint of the homotopy segment is fixed, not randomized. This sharing actually allows one to derive both corresponding results (Theorems 3.7 and 3.8, respectively) as a consequence of the following statement.

THEOREM 10.1. *For $g \in S(\mathcal{H}_d) \setminus \Sigma$, we have*

$$\mathbb{E}_{f \in S(\mathcal{H}_d)} \left(d_S(f, g) \int_0^1 \mu_2^2(q_\tau) d\tau \right) \leq 818 D^{3/2} N(n+1) \mu_{\max}^2(g) + 0.01.$$

The idea to prove Theorem 10.1 is simple. For small values of τ , the system q_τ is close to g and therefore, the value of $\mu_2^2(q_\tau)$ can be bounded by a small multiple of $\mu_{\max}^2(g)$. For the remaining values of τ , the corresponding $t = t(\tau)$ is bounded away from 0 and therefore so is the variance σ_t^2 in the distribution $N(\bar{q}_t, \sigma_t^2 I)$ for q_t . This allows one to control the denominator in the right-hand side of Theorem 3.6 when using this result. Here are the precise details.

In the following fix $g \in S(\mathcal{H}_d) \setminus \Sigma$. First note that we may again replace the uniform distribution of f on $S(\mathcal{H}_d)$ by the truncated Gaussian $N_A(0, I)$. As before we chose $A := \sqrt{2N}$. We therefore need to bound the quantity

$$Q_g := \mathbb{E}_{f \sim N_A(0, I)} \left(d_{\mathbb{S}}(f, g) \int_0^1 \mu_2^2(q_\tau) d\tau \right).$$

To simplify notation, we set as before $\varepsilon = 0.13$, $C = 0.025$, $\lambda = 6.67 \cdot 10^{-3}$, and define

$$\delta_0 := \frac{\lambda}{D^{3/2} \mu_{\max}^2(g)}, \quad t_A := \frac{1}{1 + A + 1.00001 \frac{A}{\delta_0}}.$$

PROPOSITION 10.2. *We have*

$$Q_g \leq (1 + \varepsilon)^2 \delta_0 \mu_{\max}^2(g) + \frac{A}{P_{A,1}} \int_{t_A}^1 \mathbb{E}_{q_t \sim N(\bar{q}_t, t^2 I)} \left(\frac{\mu_2^2(q_t)}{\|q_t\|^2} \right) dt,$$

where $\bar{q}_t = (1 - t)g$.

Proof. Let $\zeta^{(1)}, \dots, \zeta^{(D)}$ be the zeros of g and denote by $(q_\tau, \zeta_\tau^{(j)})_{\tau \in [0,1]}$ the lifting of $E_{f,g}$ in V corresponding to the initial pair $(g, \zeta^{(j)})$ and final system $f \in \mathcal{H}_d \setminus \Sigma$.

Equation (4.1) for $i = 0$ in the proof of Theorem 3.1 shows the following: for all j and all $\tau \leq \frac{\lambda}{d_{\mathbb{S}}(f,g) D^{3/2} \mu_{\text{norm}}^2(g, \zeta^{(j)})}$, we have

$$\mu_{\text{norm}}(q_\tau, \zeta_\tau^{(j)}) \leq (1 + \varepsilon) \mu_{\text{norm}}(g, \zeta^{(j)}) \leq (1 + \varepsilon) \mu_{\max}(g).$$

In particular, this inequality holds for all j and all $\tau \leq \frac{\delta_0}{d_{\mathbb{S}}(f,g)}$ and hence, for all such τ , we have

$$(10.1) \quad \mu_2(q_\tau) \leq (1 + \varepsilon) \mu_{\max}(g).$$

Splitting the integral in Q_g at $\tau_0(f) := \min \left\{ 1, \frac{\delta_0}{d_{\mathbb{S}}(f,g)} \right\}$ we obtain

$$Q_g = \mathbb{E}_{f \sim N_A(0, I)} \left(d_{\mathbb{S}}(f, g) \int_0^{\tau_0(f)} \mu_2^2(q_\tau) d\tau \right) + \mathbb{E}_{f \sim N_A(0, I)} \left(d_{\mathbb{S}}(f, g) \int_{\tau_0(f)}^1 \mu_2^2(q_\tau) d\tau \right).$$

Using (10.1) we bound the first term in the right-hand side as follows:

$$\mathbb{E}_{f \sim N_A(0, I)} \left(d_{\mathbb{S}}(f, g) \int_0^{\tau_0(f)} \mu_2^2(q_\tau) d\tau \right) \leq (1 + \varepsilon)^2 \delta_0 \mu_{\max}(g)^2.$$

To bound the second term, we without loss of generality assume that $\tau_0(f) \leq 1$. We apply Proposition 5.2 to obtain, for a fixed f ,

$$d_{\mathbb{S}}(f, g) \int_{\tau_0(f)}^1 \mu_2^2(q_\tau) d\tau \leq \int_{t_0(f)}^1 \|f\| \frac{\mu_2^2(q_t)}{\|q_t\|^2} dt,$$

where $t_0(f)$ is given by

$$t_0(f) = \frac{1}{1 + \|f\|(\sin \alpha \cot \delta_0 - \cos \alpha)}, \quad \alpha := d_{\mathbb{S}}(f, g).$$

Now note that $\|f\| \leq A$ since we draw f from $N_A(0, I)$. This will allow us to bound $t_0(f)$ from below by a quantity independent of f . For $\|f\| \leq A$, we have

$$0 \leq \sin \alpha \cot \delta_0 - \cos \alpha \leq \frac{1}{\sin \delta_0} - \cos \alpha \leq \frac{1}{\sin \delta_0} + 1,$$

and, moreover, $\sin \delta_0 \geq 0.99999 \delta_0$ since $\delta_0 \leq 2^{-3/2} \lambda \leq 0.00236$. We can therefore bound $t_0(f)$ as

$$t_0(f) \geq \frac{1}{1 + A + \frac{A}{\sin(\delta_0)}} \geq \frac{1}{1 + A + 1.00001 \frac{A}{\delta_0}} = t_A.$$

We can now bound the second term in Q_g as follows:

$$\begin{aligned} \mathbb{E}_{f \sim N_A(0, I)} \left(d_{\mathbb{S}}(f, g) \int_{\tau_0(f)}^1 \mu_2^2(q_\tau) d\tau \right) &\leq \mathbb{E}_{f \sim N_A(0, I)} \left(A \int_{t_A}^1 \frac{\mu_2^2(q_t)}{\|q_t\|^2} dt \right) \\ &= A \int_{t_A}^1 \mathbb{E}_{f \sim N_A(0, I)} \left(\frac{\mu_2^2(q_t)}{\|q_t\|^2} \right) dt \leq \frac{A}{P_{A,1}} \int_{t_A}^1 \mathbb{E}_{f \sim N(0, I)} \left(\frac{\mu_2^2(q_t)}{\|q_t\|^2} \right) dt. \end{aligned}$$

To conclude, note that, for fixed t and when f is distributed following $N(0, I)$, the variable $q_t = (1 - t)g + tf$ follows the Gaussian $N(\bar{q}_t, t^2I)$, where $\bar{q}_t = (1 - t)g$. \square

Proof of Theorem 10.1. By homogeneity we can replace the uniform distribution on $S(\mathcal{H}_d)$ by $N_A(0, I)$, so that we only need to estimate Q_g by the right-hand side of Proposition 10.2. In order to bound the first term there we note that

$$(1 + \varepsilon)^2 \delta_0 \mu_{\max}^2(g) = (1 + \varepsilon)^2 \lambda D^{-3/2} \leq (1 + \varepsilon)^2 \lambda \leq 0.01.$$

For bounding the second term we apply Theorem 3.6 to deduce that

$$\begin{aligned} \int_{t_A}^1 \mathbb{E}_{q_t \sim N(\bar{q}_t, t^2I)} \left(\frac{\mu_2^2(q_t)}{\|q_t\|^2} \right) dt &\leq \int_{t_A}^1 \frac{e(n+1)}{2t^2} dt = \frac{e(n+1)}{2} \left(\frac{1}{t_A} - 1 \right) \\ &= \frac{e(n+1)A}{2} \left(1 + \frac{1.00001}{\delta_0} \right). \end{aligned}$$

Replacing this bound in Proposition 10.2 we obtain

$$\begin{aligned} Q_g &\leq \frac{eA^2(n+1)}{2P_{A,1}} \left(1 + \frac{1.00001}{\lambda} D^{3/2} \mu_{\max}^2(g) \right) + 0.01 \\ &\leq 2eN(n+1)D^{3/2} \mu_{\max}^2(g) \left(\frac{1}{D^{3/2}} + \frac{1.00001}{\lambda} \right) + 0.01 \\ &\leq 818 N(n+1)D^{3/2} \mu_{\max}^2(g) + 0.01, \end{aligned}$$

where we used $D \geq 2$ for the last inequality. \square

10.1. *Condition-based analysis of LV (proof).*

Proof of Theorem 3.7. The result follows immediately by combining Proposition 5.1 with Theorem 10.1, with the roles of f and g swapped. \square

10.2. *The complexity of a deterministic homotopy continuation.* We next prove Theorem 3.8, beginning with some general considerations. The unitary group $\mathcal{U}(n+1)$ naturally acts on \mathbb{P}^n as well as on $\mathcal{H}_{\mathbf{d}}$ via $(\nu, f) \mapsto f \circ \nu^{-1}$. The following lemma results from the unitary invariance of our setting. The proof is immediate.

LEMMA 10.3. *Let $g \in \mathcal{H}_{\mathbf{d}}$, $\zeta \in \mathbb{P}^n$ be a zero of g , and $\nu \in \mathcal{U}(n+1)$. Then $\mu_{\text{norm}}(g, \zeta) = \mu_{\text{norm}}(g \circ \nu^{-1}, \nu\zeta)$. Moreover, for $f \in \mathcal{H}_{\mathbf{d}}$, we have $K(f, g, \zeta) = K(f \circ \nu^{-1}, g \circ \nu^{-1}, \nu\zeta)$. \square*

Recall $\bar{U}_i = \frac{1}{\sqrt{2n}}(X_0^{d_i} - X_i^{d_i})$ and denote by $z_{(i)}$ a d_i th primitive root of unity. The \mathcal{D} zeros of $\bar{U} = (\bar{U}_1, \dots, \bar{U}_n)$ are the points $\mathbf{z}_j = (1 : z_{(1)}^{j_1} : \dots : z_{(n)}^{j_n}) \in \mathbb{P}^n$ for all the possible tuples $j = (j_1, \dots, j_n)$ with $j_i \in \{0, \dots, d_i - 1\}$. Clearly, each \mathbf{z}_j can be obtained from $\mathbf{z}_1 := (1 : 1 : \dots : 1)$ by a unitary transformation ν_j , which leaves \bar{U} invariant; that is,

$$\nu_j \mathbf{z}_1 = \mathbf{z}_j, \quad \bar{U} \circ \nu_j^{-1} = \bar{U}.$$

Hence Lemma 10.3 implies $\mu_{\text{norm}}(\bar{U}, \mathbf{z}_j) = \mu_{\text{norm}}(\bar{U}, \mathbf{z}_1)$ for all j . In particular, $\mu_{\text{max}}(\bar{U}) = \mu_{\text{norm}}(\bar{U}, \mathbf{z}_1)$.

PROPOSITION 10.4. $K_{\bar{U}}(f) = K(f, \bar{U}, \mathbf{z}_1)$ satisfies

$$\mathbb{E}_{f \in S(\mathcal{H}_{\mathbf{d}})} K_{\bar{U}}(f) = \mathbb{E}_{f \in S(\mathcal{H}_{\mathbf{d}})} \frac{1}{\mathcal{D}} \sum_{j=1}^{\mathcal{D}} K(f, \bar{U}, \mathbf{z}_j).$$

Proof. Lemma 10.3 implies, for all j ,

$$K(f, \bar{U}, \mathbf{z}_1) = K(f \circ \nu_j^{-1}, \bar{U} \circ \nu_j^{-1}, \nu_j \mathbf{z}_1) = K(f \circ \nu_j^{-1}, \bar{U}, \mathbf{z}_j).$$

It follows that

$$K_{\bar{U}}(f) = K(f, \bar{U}, \mathbf{z}_1) = \frac{1}{\mathcal{D}} \sum_{j=1}^{\mathcal{D}} K(f \circ \nu_j^{-1}, \bar{U}, \mathbf{z}_j).$$

The assertion follows now since, for all measurable functions $\varphi: S(\mathcal{H}_{\mathbf{d}}) \rightarrow \mathbb{R}$ and all $\nu \in \mathcal{U}(n+1)$, we have

$$\mathbb{E}_{f \in S(\mathcal{H}_{\mathbf{d}})} \varphi(f) = \mathbb{E}_{f \in S(\mathcal{H}_{\mathbf{d}})} \varphi(f \circ \nu),$$

due to the isotropy of the uniform measure on $S(\mathcal{H}_{\mathbf{d}})$, \square

LEMMA 10.5. *We have*

$$\mu_{\max}^2(\bar{U}) \leq 2n \max_i \frac{1}{d_i} (n+1)^{d_i-1} \leq 2(n+1)^D.$$

Proof. Recall $\mu_{\max}(\bar{U}) = \mu_{\text{norm}}(\bar{U}, \mathbf{z}_1)$, so it suffices to bound $\mu_{\text{norm}}(\bar{U}, \mathbf{z}_1)$. Consider $M := \text{diag}(d_i^{-\frac{1}{2}} \|\mathbf{z}_1\|^{1-d_i}) D\bar{U}(\mathbf{z}_1) \in \mathbb{C}^{n \times (n+1)}$. By definition we have (cf. §2.3)

$$\mu_{\text{norm}}(\bar{U}, \mathbf{z}_1) = \|\bar{U}\| \|M^\dagger\| = \|M^\dagger\| = \frac{1}{\sigma_{\min}(M)},$$

where $\sigma_{\min}(M)$ denotes the smallest singular value of M . It can be characterized as a constrained minimization problem as follows:

$$\sigma_{\min}^2(M) = \min_u \|Mu\|^2 \quad \text{subject to } u \in (\ker M)^\perp, \|u\|^2 = 1.$$

In our situation, $\ker M = \mathbb{C}(1, \dots, 1)$ and $D\bar{U}(\mathbf{z}_1)$ is given by the following matrix, shown here for $n = 3$:

$$D\bar{U}(\mathbf{z}_1) = \frac{1}{\sqrt{2n}} \begin{bmatrix} d_1 & -d_1 & 0 & 0 \\ d_2 & 0 & -d_2 & 0 \\ d_3 & 0 & 0 & -d_3 \end{bmatrix}.$$

Hence for $u = (u_0, \dots, u_n) \in \mathbb{C}^{n+1}$,

$$\|Mu\|^2 = \frac{1}{2n} \sum_{i=1}^n \frac{d_i}{(n+1)^{d_i-1}} |u_i - u_0|^2 \geq \frac{1}{2n} \min_i \frac{d_i}{(n+1)^{d_i-1}} \cdot \sum_{i=1}^n |u_i - u_0|^2.$$

A straightforward calculation shows that

$$\sum_{i=1}^n |u_i - u_0|^2 \geq 1 \quad \text{if} \quad \sum_{i=0}^n u_i = 0, \quad \sum_{i=0}^n |u_i|^2 = 1.$$

The assertion follows by combining these observations. □

Proof of Theorem 3.8. Equation (5.1) in the proof of Proposition 5.1 implies for $g = \bar{U}$ that

$$\frac{1}{D} \sum_{i=1}^D K(f, \bar{U}, \mathbf{z}_i) \leq 245 D^{3/2} d_{\mathbb{S}}(f, \bar{U}) \int_0^1 \mu_2^2(q_\tau) d\tau.$$

Using Proposition 10.4 we get

$$\mathbb{E}_{f \in \mathcal{S}(\mathcal{H}_d)} K_{\bar{U}}(f) \leq 245 D^{3/2} \mathbb{E}_{f \in \mathcal{S}(\mathcal{H}_d)} \left(d_{\mathbb{S}}(f, \bar{U}) \int_0^1 \mu_2^2(q_\tau) d\tau \right).$$

Applying Theorem 10.1 with $g = \bar{U}$ we obtain

$$\mathbb{E}_{f \in \mathcal{S}(\mathcal{H}_d)} K_{\bar{U}}(f) \leq 245 D^{3/2} (818 D^{3/2} N(n+1) \mu_{\max}^2(\bar{U}) + 0.01).$$

We now plug in the bound $\mu_{\max}(\bar{U})^2 \leq 2(n+1)^D$ of Lemma 10.5 to obtain

$$\mathbb{E}_{f \in S(\mathcal{H}_d)} K_{\bar{U}}(f) \leq 400820 D^3 N(n+1)^{D+1} + 2.45 D^{3/2}.$$

This is bounded from above by $400821 D^3 N(n+1)^{D+1}$, which completes the proof. \square

11. A near solution to Smale’s 17th problem

We finally proceed with the proof of Theorem 3.9. The algorithm we will exhibit uses different routines for $D \leq n$ and $D > n$. Our exposition reflects this structure.

11.1. *The case $D \leq n$.* Theorem 3.8 bounds the number of iterations of Algorithm MD as

$$\mathbb{E}_{f \in S(\mathcal{H}_d)} K_{\bar{U}}(f) = \mathcal{O}(D^3 N n^{D+1}).$$

For comparing the order of magnitude of this upper bound to the input size $N = \sum_{i=1}^n \binom{n+d_i}{n}$ we need the following technical lemma (which will be useful for the case $D > n$ as well).

LEMMA 11.1. (1) *For $D \leq n$, $n \geq 4$, we have*

$$n^D \leq \binom{n+D}{D}^{\ln n}.$$

(2) *For $D^2 \geq n \geq 1$, we have*

$$\ln n \leq 2 \ln \ln \binom{n+D}{n} + 4.$$

(3) *For $0 < c < 1$, there exists K such that for all n, D ,*

$$D \leq n^{1-c} \implies n^D \leq \binom{n+D}{n}^K.$$

(4) *For $D \leq n$, we have*

$$n^D \leq N^{2 \ln \ln N + \mathcal{O}(1)}.$$

(5) *For $n \leq D$, we have*

$$D^n \leq N^{2 \ln \ln N + \mathcal{O}(1)}.$$

Proof. Stirling’s formula states $n! = \sqrt{2\pi n} n^{n+\frac{1}{2}} e^{-n} e^{\frac{\Theta_n}{12n}}$ with $0 < \Theta_n < 1$. Let $H(x) = x \ln \frac{1}{x} + (1-x) \ln \frac{1}{1-x}$ denote the binary entropy function, defined for $0 < x < 1$. By a straightforward calculation we get from Stirling’s formula

the following asymptotics for the binomial coefficient: for any $0 < m < n$, we have

$$(11.1) \quad \ln \binom{n}{m} = nH\left(\frac{m}{n}\right) + \frac{1}{2} \ln \frac{n}{m(n-m)} - 1 + \varepsilon_{n,m},$$

where $-0.1 < \varepsilon_{n,m} < 0.2$. This formula holds as well for the extension of binomial coefficients on which m is not necessarily integer.

(1) The first claim is equivalent to $e^D \leq \binom{n+D}{D}$. The latter is easily checked for $D \in \{1, 2, 3\}$ and $n \geq 4$. So assume $n \geq D \geq 4$. By monotonicity it suffices to show that $e^D \leq \binom{2D}{D}$ for $D \geq 4$. Equation (11.1) implies

$$\ln \binom{2D}{D} > 2D \ln 2 + \frac{1}{2} \ln \frac{2}{D} - 1.1,$$

and the right-hand side is easily checked to be at least D for $D \geq 4$.

(2) Put $m := \sqrt{n}$. If $D \geq m$, then $\binom{n+D}{n} \geq \binom{n+\lceil m \rceil}{n}$, so it is enough to show that $\ln n \leq 2 \ln \ln \binom{n+\lceil m \rceil}{n} + 4$. Equation (11.1) implies

$$\ln \binom{n + \lceil m \rceil}{n} \geq \ln \binom{n + m}{n} \geq (n + m)H\left(\frac{m}{n + m}\right) + \frac{1}{2} \ln \frac{1}{m} - 1.1.$$

The entropy function can be bounded as

$$H\left(\frac{m}{n + m}\right) \geq \frac{m}{n + m} \ln \left(1 + \frac{n}{m}\right) \geq \frac{m}{n + m} \ln m.$$

It follows that

$$\ln \binom{n + \lceil m \rceil}{n} \geq \frac{1}{2} \sqrt{n} \ln n - \frac{1}{4} \ln n - 1.1 \geq \frac{1}{4} \sqrt{n} \ln n,$$

where the right-hand inequality holds for $n \geq 10$. Hence, for $n \geq 10$,

$$\ln \ln \binom{n + \lceil m \rceil}{n} \geq \frac{1}{2} \ln n + \ln \ln n - \ln 4 \geq \frac{1}{2} \ln n - 2.$$

This shows the second claim for $n \geq 10$. The cases $n \leq 9$ are easily directly checked.

(3) Writing $D = n\delta$ we obtain from equation (11.1)

$$\ln \binom{n + D}{n} = (n + D)H\left(\frac{\delta}{1 + \delta}\right) - \frac{1}{2} \ln D + \mathcal{O}(1).$$

Estimating the entropy function yields

$$H\left(\frac{\delta}{1 + \delta}\right) \geq \frac{\delta}{1 + \delta} \ln \left(1 + \frac{1}{\delta}\right) \geq \frac{\delta}{2} \ln \frac{1}{\delta} = \frac{\delta\varepsilon}{2} \ln n,$$

where ε is defined by $\delta = n^{-\varepsilon}$. By assumption, $\varepsilon \geq c$. From the last two lines we get

$$\frac{1}{D \ln n} \ln \binom{n + D}{n} \geq \frac{c}{2} - \frac{1 - c}{2D} + \mathcal{O}\left(\frac{1}{\ln n}\right).$$

In the case $c \leq \frac{3}{4}$ we have $D \geq n^{1/4}$ and we bound the above by

$$\frac{c}{2} - \frac{1}{2n^{1/4}} + \mathcal{O}\left(\frac{1}{\ln n}\right),$$

which is greater than $c/4$ for sufficiently large n . In the case $c \geq \frac{3}{4}$ we bound as follows:

$$\frac{1}{D \ln n} \ln \binom{n+D}{n} \geq \frac{c}{2} - \frac{1-c}{2} + \mathcal{O}\left(\frac{1}{\ln n}\right) = c - \frac{1}{2} + \mathcal{O}\left(\frac{1}{\ln n}\right) \geq \frac{1}{5}$$

for sufficiently large n .

We have shown that for $0 < c < 1$, there exists n_c such that for $n \geq n_c$, $D \leq n^{1-c}$, we have

$$n^D \leq \binom{n+D}{n}^{K_c},$$

where $K_c := \max\{4/c, 5\}$. By increasing K_c we can achieve that the above inequality holds for all n, D , with $D \leq n^{1-c}$.

(4) Clearly, $N \geq \binom{n+D}{n}$. If $D \leq \sqrt{n}$ then, by part (3), there exists K such that

$$n^D \leq \binom{n+D}{n}^K \leq N^K.$$

Otherwise $D \in [\sqrt{n}, n]$ and the desired inequality is an immediate consequence of parts (1) and (2).

(5) Use $\binom{n+D}{n} = \binom{n+D}{D}$ and swap the roles of n and D in part (4) above. \square

Theorem 3.8 combined with Lemma 11.1(4) implies that

$$(11.2) \quad \mathbb{E}_f K_{\overline{U}}(f) = N^{2 \ln \ln N + \mathcal{O}(1)} \quad \text{if } D \leq n.$$

Note that this bound is nearly polynomial in N . Moreover, if $D \leq n^{1-c}$ for some fixed $0 < c < 1$, then Lemma 11.1(3) implies

$$(11.3) \quad \mathbb{E}_f K_{\overline{U}}(f) = N^{\mathcal{O}(1)}.$$

In this case, the expected running time is polynomially bounded in the input size N .

11.2. *The case $D > n$.* The homotopy continuation algorithm MD is not efficient for large degrees — the main problem being that we do not know how to deterministically compute a starting system g with small $\mu_{\max}(g)$. However, it turns out that an algorithm due to Jim Renegar [18], based on the factorization of the u -resultant, computes approximate zeros and is fast for large degrees.

Before giving the specification of Renegar’s algorithm, we need to fix some notation. We identify $\mathbb{P}_0^n := \{(x_0 : \cdots : x_n) \in \mathbb{P}^n \mid x_0 \neq 0\}$ with \mathbb{C}^n via the bijection $(x_0 : \cdots : x_n) \mapsto \underline{x} := (x_1/x_0, \dots, x_n/x_0)$. If $x \in \mathbb{P}_0^n$, then we denote by $\|x\|_{\text{aff}}$ the Euclidean norm of \underline{x} , i.e.,

$$\|x\|_{\text{aff}} := \|\underline{x}\| = \left(\sum_{i=1}^n \left| \frac{x_i}{x_0} \right|^2 \right)^{\frac{1}{2}},$$

and we put $\|x\|_{\text{aff}} = \infty$ if $x \in \mathbb{P}^n \setminus \mathbb{P}_0^n$. Furthermore, for $x, y \in \mathbb{P}_0^n$, we shall write $d_{\text{aff}}(x, y) := \|\underline{x} - \underline{y}\|$ and we set $d_{\text{aff}}(x, y) := \infty$ otherwise. An elementary argument shows that $d_{\mathbb{P}}(x, y) \leq d_{\text{aff}}(x, y)$ for $x, y \in \mathbb{P}_0^n$.

By a δ -approximation of a zero $\zeta \in \mathbb{P}_0^n$ of $f \in \mathcal{H}_{\mathbf{d}}$ we understand an $x \in \mathbb{P}_0^n$ such that $d_{\text{aff}}(x, \zeta) \leq \delta$. The following result relates δ -approximations to the approximate zeros in the sense of Definition 2.1.

PROPOSITION 11.2. *Let x be a δ -approximation of a zero ζ of f . Recall $C = 0.025$. If $D^{3/2}\mu_{\text{norm}}(f, x)\delta \leq C$, then x is an approximate zero of f .*

Proof. We have $d_{\mathbb{P}}(x, \zeta) \leq d_{\text{aff}}(x, \zeta) \leq \delta$. Suppose that $D^{3/2}\mu_{\text{norm}}(f, x)\delta \leq C$. Then, by Proposition 4.1 with $g = f$, we have

$$\mu_{\text{norm}}(f, \zeta) \leq (1 + \varepsilon)\mu_{\text{norm}}(f, x)$$

with $\varepsilon = 0.13$. Hence

$$D^{3/2}\mu_{\text{norm}}(f, \zeta)d_{\mathbb{P}}(x, \zeta) \leq (1 + \varepsilon)D^{3/2}\mu_{\text{norm}}(f, x)\delta \leq (1 + \varepsilon)C.$$

We have $(1 + \varepsilon)C \leq u_0 = 3 - \sqrt{7}$. Now use Theorem 2.2. □

Consider now $R \geq \delta > 0$. Renegar’s Algorithm $\text{Ren}(R, \delta)$ from [18] takes as input $f \in \mathcal{H}_{\mathbf{d}}$, decides whether its zero set $V(f) \subseteq \mathbb{P}^n$ is finite, and if so, computes δ -approximations x to at least all zeros ζ of f satisfying $\|\zeta\|_{\text{aff}} \leq R$. (The algorithm even finds the multiplicities of those zeros ζ ; see [18] for the precise statement.)

Renegar’s Algorithm can be formulated in the BSS-model over \mathbb{R} . Its running time on input f (the number of arithmetic operations and inequality tests) is bounded by

$$(11.4) \quad \mathcal{O}\left(nD^4(\log D)\left(\log \log \frac{R}{\delta}\right) + n^2D^4\binom{1 + \sum_i d_i}{n}^4\right).$$

To find an approximate zero of f we may use $\text{Ren}(R, \delta)$ together with Proposition 11.2 and iterate with $R = 4^k$ and $\delta = 2^{-k}$ for $k = 1, 2, \dots$ until we are successful. More precisely, we consider the following algorithm:

Algorithm **ItRen**

input $f \in \mathcal{H}_d$

for $k = 1, 2, \dots$ **do**

run $\text{Re}(4^k, 2^{-k})$ **on input** f

for all δ -approximations x **found**

if $D^{3/2}\mu_{\text{norm}}(f, x)\delta \leq C$ **stop and RETURN** x

Let $\Sigma_0 := \Sigma \cup \{f \in \mathcal{H}_d \mid V(f) \cap \mathbb{P}_0^n = \emptyset\}$. It is obvious that **ItRen** stops on inputs $f \notin \Sigma_0$. In particular, **ItRen** stops almost surely.

The next result bounds the probability **Probfail** that the main loop of **ItRen**, with parameters R and δ , fails to output an approximate zero for a standard Gaussian input $f \in \mathcal{H}_d$ (and given R, δ). We postpone its proof to Section 11.3.

LEMMA 11.3. *We have $\text{Probfail} = \mathcal{O}(n^3 N^2 D^6 \mathcal{D} \delta^4 + nR^{-2})$.*

Let $T(f)$ denote the running time of algorithm **ItRen** on input f .

PROPOSITION 11.4. *We have for standard Gaussian $f \in \mathcal{H}_d$*

$$\mathbb{E}_f T(f) = (nN\mathcal{D})^{\mathcal{O}(1)}.$$

Proof. The probability that **ItRen** stops in the $(k+1)$ th loop is bounded above by the probability p_k that $\text{Re}(4^k, 2^{-k})$ fails to produce an approximate zero. Lemma 11.3 tells us that

$$p_k = \mathcal{O}(n^3 N^2 D^6 \mathcal{D} 16^{-k}).$$

If A_k denotes the running time of the $(k+1)$ th loop, then we conclude

$$\mathbb{E}_f T(f) \leq \sum_{k=0}^{\infty} A_k p_k.$$

According to (11.4), A_k is bounded by

$$\mathcal{O}\left(n\mathcal{D}^4(\log \mathcal{D})(\log k) + n^2\mathcal{D}^4\left(\frac{1 + \sum_i d_i}{n}\right)^4 + (N + n^3)\mathcal{D}\right),$$

where the last term accounts for the cost of the tests. The assertion now follows by distributing the products $A_k p_k$ and using that the series $\sum_{k \geq 1} 16^{-k}$, and $\sum_{k \geq 1} 16^{-k} \log k$ have finite sums. \square

Proof of Theorem 3.9. We use Algorithm **MD** if $D \leq n$ and Algorithm **ItRen** if $D > n$. We have already shown (see (11.2), (11.3)) that the assertion holds if $D \leq n$. For the case $D > n$ we use Proposition 11.4 together with the inequality $\mathcal{D}^{\mathcal{O}(1)} \leq D^{\mathcal{O}(n)} \leq N^{\mathcal{O}(\log \log N)}$ which follows from Lemma 11.1(5). Moreover, in the case $D \geq n^{1+\varepsilon}$, Lemma 11.1(3) implies $\mathcal{D} \leq D^n \leq N^{\mathcal{O}(1)}$. \square

11.3. *Proof of Lemma 11.3.* Let \mathcal{E} denote the set of $f \in \mathcal{H}_d$ such that there is an x on the output list of $\text{Ren}(R, \delta)$ on input f that satisfies $C < D^{3/2}\mu_{\text{norm}}(f, x)\delta$. Then

$$\text{Probfail} \leq \text{Prob}_{f \in \mathcal{H}_d} \left\{ \min_{\zeta \in V(f)} \|\zeta\|_{\text{aff}} \geq R \right\} + \text{Prob } \mathcal{E}.$$

Lemma 11.3 follows immediately from the following two results.

LEMMA 11.5. *For $R > 0$ and standard Gaussian $f \in \mathcal{H}_d$, we have*

$$\text{Prob}_{f \in \mathcal{H}_d} \left\{ \min_{\zeta \in V(f)} \|\zeta\|_{\text{aff}} \geq R \right\} \leq \frac{n}{R^2}.$$

Proof. Choose $f \in \mathcal{H}_d$ standard Gaussian and pick one of the \mathcal{D} zeros $\zeta_f^{(1)}, \dots, \zeta_f^{(\mathcal{D})}$ of f uniformly at random, call it ζ . Then the resulting distribution of (f, ζ) in $V_{\mathbb{P}}$ has the density ρ_{st} . Lemma 8.8 implies that ζ is uniformly distributed in \mathbb{P}^n . Therefore,

$$\text{Prob}_{f \in \mathcal{H}_d} \left\{ \min_i \|\zeta_f^{(i)}\|_{\text{aff}} \geq R \right\} \leq \text{Prob}_{\zeta \in \mathbb{P}^n} \left\{ \|\zeta\|_{\text{aff}} \geq R \right\}.$$

To estimate the right-hand side probability we observe that

$$\|\zeta\|_{\text{aff}} \geq R \iff d_{\mathbb{P}}(\zeta, \mathbb{P}^{n-1}) \leq \frac{\pi}{2} - \theta,$$

where θ is defined by $R = \tan \theta$ and $\mathbb{P}^{n-1} := \{x \in \mathbb{P}^n \mid x_0 = 0\}$. Therefore,

$$\text{Prob}_{\zeta \in \mathbb{P}^n} \left\{ \|\zeta\|_{\text{aff}} \geq R \right\} = \frac{\text{vol} \left\{ x \in \mathbb{P}^n \mid d_{\mathbb{P}}(x, \mathbb{P}^{n-1}) \leq \frac{\pi}{2} - \theta \right\}}{\text{vol}(\mathbb{P}^n)}.$$

Due to [11, Lemma 2.1] and using $\text{vol}(\mathbb{P}^n) = \pi^n/n!$, this can be bounded by

$$\frac{\text{vol}(\mathbb{P}^{n-1})\text{vol}(\mathbb{P}^1)}{\text{vol}(\mathbb{P}^n)} \sin^2 \left(\frac{\pi}{2} - \theta \right) = n \cos^2 \theta = \frac{n}{1 + R^2} \leq \frac{n}{R^2}. \quad \square$$

LEMMA 11.6. *We have $\text{Prob } \mathcal{E} = \mathcal{O}(n^3 N^2 D^6 \mathcal{D} \delta^4)$.*

Proof. Assume that $f \in \mathcal{E}$. Then, there exist $\zeta, x \in \mathbb{P}_0^n$ such that $f(\zeta) = 0$, $\|\zeta\|_{\text{aff}} \leq R$, $d_{\text{aff}}(x, \zeta) \leq \delta$, Ren returns x , and $D^{3/2}\mu_{\text{norm}}(f, x)\delta > C$.

We proceed by cases. Suppose first that $\delta \leq \frac{C}{D^{3/2}\mu_{\text{norm}}(f, \zeta)}$. Then, by Proposition 4.1,

$$(1 + \varepsilon)^{-1}C < (1 + \varepsilon)^{-1}D^{3/2}\mu_{\text{norm}}(f, x)\delta \leq D^{3/2}\mu_{\text{norm}}(f, \zeta)\delta;$$

hence

$$\mu_{\text{max}}(f) \geq \mu_{\text{norm}}(f, \zeta) \geq (1 + \varepsilon)^{-1}CD^{-3/2}\delta^{-1}.$$

If, on the other hand, $\delta > \frac{C}{D^{3/2}\mu_{\text{norm}}(f, \zeta)}$, then we have

$$\mu_{\text{max}}(f) \geq \mu_{\text{norm}}(f, \zeta) \geq CD^{-3/2}\delta^{-1}.$$

Therefore, for any $f \in \mathcal{E}$,

$$\mu_{\text{max}}(f) \geq (1 + \varepsilon)^{-1}CD^{-3/2}\delta^{-1}.$$

Theorem C of [23] states that $\text{Prob}_f\{\mu_{\max}(f) \geq \rho^{-1}\} = \mathcal{O}(n^3 N^2 \mathcal{D} \rho^4)$ for all $\rho > 0$. Therefore, as claimed, we get

$$\text{Prob } \mathcal{E} \leq \text{Prob}_{f \in \mathcal{H}_a} \left\{ \mu_{\max}(f) \geq (1 + \varepsilon)^{-1} C D^{-3/2} \delta^{-1} \right\} = \mathcal{O}(n^3 N^2 \mathcal{D} \delta^4). \quad \square$$

Note added in proof. Since the posting of this manuscript on September 2009, at [arXiv:0909.2114](https://arxiv.org/abs/0909.2114), a number of references have been added to the literature. The nonconstructive character of the main result in [21] — the bound in (1.3) — had also been noticed by Carlos Beltrán. In a recent paper (A continuation method to solve polynomial systems, and its complexity, *Num. Math.* **117** (2011), 89–113), Beltrán proves a very general constructive version of this result. Our Theorem 3.1 can be seen as a particular case (with a correspondingly shorter proof) of Beltrán’s paper main result. We understand that yet another constructive version for the bound in (1.3) is the subject of a paper in preparation by J.-P. Dedieu, G. Malajovich, and M. Shub.

Also, Beltrán and Pardo have recently rewritten their paper [6] (Fast linear homotopy to find approximate zeros of polynomial systems, *Found. Comput. Math.* **11**, (2011), 95–129). This revised version, which increases the length of the manuscript by a factor of about three, adds considerable detail to a number of issues only briefly sketched in [6]. In particular, the effective sampling from the solution variety is now given a full description (which is slightly different to the one we give in §9).

References

- [1] M. AGRAWAL, N. KAYAL, and N. SAXENA, PRIMES is in P, *Ann. of Math.* **160** (2004), 781–793. MR 2123939. Zbl 1071.11070. <http://dx.doi.org/10.4007/annals.2004.160.781>.
- [2] D. AMELUNXEN and P. BÜRGISSER, Robust smoothed analysis of a condition number of linear programming, *Math. Program., Ser. A.* <http://dx.doi.org/10.1007/s10107-010-0346-x>.
- [3] W. BAUR and V. STRASSEN, The complexity of partial derivatives, *Theoret. Comput. Sci.* **22** (1983), 317–330. MR 0693063. Zbl 0498.68028. [http://dx.doi.org/10.1016/0304-3975\(83\)90110-X](http://dx.doi.org/10.1016/0304-3975(83)90110-X).
- [4] C. BELTRÁN and L. M. PARDO, On Smale’s 17th problem: a probabilistic positive solution, *Found. Comput. Math.* **8** (2008), 1–43. MR 2403529. Zbl 1153.65048. <http://dx.doi.org/10.1007/s10208-005-0211-0>.
- [5] ———, Smale’s 17th problem: average polynomial time to compute affine and projective solutions, *J. Amer. Math. Soc.* **22** (2009), 363–385. MR 2476778. Zbl 1206.90173. <http://dx.doi.org/10.1090/S0894-0347-08-00630-9>.
- [6] ———, Fast linear homotopy to find approximate zeros of polynomial systems, *Found. Comput. Math.* **11** (2011), 95–129. MR 2754191. <http://dx.doi.org/10.1007/s10208-010-9078-9>.
- [7] C. BELTRÁN and M. SHUB, Complexity of Bezout’s theorem. VII. Distance estimates in the condition metric, *Found. Comput. Math.* **9** (2009), 179–195. MR 2496559. <http://dx.doi.org/10.1007/s10208-007-9018-5>.

- [8] L. BLUM, F. CUCKER, M. SHUB, and S. SMALE, *Complexity and Real Computation*, Springer-Verlag, New York, 1998. MR 1479636. Zbl 0872.68036.
- [9] L. BLUM, M. SHUB, and S. SMALE, On a theory of computation and complexity over the real numbers: NP-completeness, recursive functions and universal machines, *Bull. Amer. Math. Soc.* **21** (1989), 1–46. MR 0974426. Zbl 0681.03020. <http://dx.doi.org/10.1090/S0273-0979-1989-15750-9>.
- [10] P. BÜRGISSER and F. CUCKER, Smoothed analysis of Moore-Penrose inversion, *SIAM J. Matrix Anal. Appl.* **31** (2010), 2769–2783. MR 2740632. <http://dx.doi.org/10.1137/100782954>.
- [11] P. BÜRGISSER, F. CUCKER, and M. LOTZ, Smoothed analysis of complex conic condition numbers, *J. Math. Pures Appl.* **86** (2006), 293–309. MR 2257845. Zbl 1113.65044. <http://dx.doi.org/10.1016/j.matpur.2006.06.001>.
- [12] ———, The probability that a slightly perturbed numerical analysis problem is difficult, *Math. Comp.* **77** (2008), 1559–1583. MR 2398780. Zbl 1195.65018. <http://dx.doi.org/10.1090/S0025-5718-08-02060-7>.
- [13] K. P. CHOI, On the medians of gamma distributions and an equation of Ramanujan, *Proc. Amer. Math. Soc.* **121** (1994), 245–251. MR 1195477. Zbl 0803.62007. <http://dx.doi.org/10.2307/2160389>.
- [14] F. CUCKER, H. DIAO, and Y. WEI, Smoothed analysis of some condition numbers, *Numer. Linear Algebra Appl.* **13** (2006), 71–84. MR 2194973. Zbl 1198.65084. <http://dx.doi.org/10.1002/nla.464>.
- [15] F. CUCKER, R. HAUSER, and M. LOTZ, Adversarial smoothed analysis, *J. Complexity* **26** (2010), 255–262. MR 2657364. Zbl 05763691. <http://dx.doi.org/10.1016/j.jco.2009.12.005>.
- [16] R. HOWARD, The kinematic formula in Riemannian homogeneous spaces, *Mem. Amer. Math. Soc.* **106** (1993), vi+69. MR 1169230. Zbl 0810.53057.
- [17] M. O. RABIN, Probabilistic algorithm for testing primality, *J. Number Theory* **12** (1980), 128–138. MR 0566880. Zbl 0426.10006. [http://dx.doi.org/10.1016/0022-314X\(80\)90084-0](http://dx.doi.org/10.1016/0022-314X(80)90084-0).
- [18] J. RENEGAR, On the worst-case arithmetic complexity of approximating zeros of systems of polynomials, *SIAM J. Comput.* **18** (1989), 350–370. MR 0986672. Zbl 0676.65045. <http://dx.doi.org/10.1137/0218024>.
- [19] A. SANKAR, D. A. SPIELMAN, and S.-H. TENG, Smoothed analysis of the condition numbers and growth factors of matrices, *SIAM J. Matrix Anal. Appl.* **28** (2006), 446–476. MR 2255338. Zbl 1179.65033. <http://dx.doi.org/10.1137/S0895479803436202>.
- [20] M. SHUB, Some remarks on Bezout’s theorem and complexity theory, in *From Topology to Computation: Proceedings of the Smalefest* (Berkeley, CA, 1990), Springer-Verlag, New York, 1993, pp. 443–455. MR 1246139. Zbl 0811.13021.
- [21] ———, Complexity of Bezout’s theorem. VI. Geodesics in the condition (number) metric, *Found. Comput. Math.* **9** (2009), 171–178. MR 2496558. <http://dx.doi.org/10.1007/s10208-007-9017-6>.

- [22] M. SHUB and S. SMALE, Complexity of Bézout's theorem. I. Geometric aspects, *J. Amer. Math. Soc.* **6** (1993), 459–501. MR 1175980. Zbl 0821.65035. <http://dx.doi.org/10.2307/2152805>.
- [23] ———, Complexity of Bezout's theorem. II. Volumes and probabilities, in *Computational Algebraic Geometry* (Nice, 1992), *Progr. Math.* **109**, Birkhäuser, Boston, MA, 1993, pp. 267–285. MR 1230872. Zbl 0851.65031. <http://dx.doi.org/10.1006/jcom.1993.1002>.
- [24] ———, Complexity of Bezout's theorem. III. Condition number and packing, *J. Complexity* **9** (1993), 4–14. MR 1213484. Zbl 0846.65018. <http://dx.doi.org/10.1006/jcom.1993.1002>.
- [25] ———, Complexity of Bezout's theorem. V. Polynomial time, *Theoret. Comput. Sci.* **133** (1994), 141–164. MR 1294430. Zbl 0846.65022. [http://dx.doi.org/10.1016/0304-3975\(94\)90122-8](http://dx.doi.org/10.1016/0304-3975(94)90122-8).
- [26] ———, Complexity of Bezout's theorem. IV. Probability of success; extensions, *SIAM J. Numer. Anal.* **33** (1996), 128–148. MR 1377247. Zbl 0843.65035. <http://dx.doi.org/10.1137/0733008>.
- [27] S. SMALE, Newton's method estimates from data at one point, in *The Merging of Disciplines: New Directions in Pure, Applied, and Computational Mathematics* (Laramie, Wyo., 1985), Springer-Verlag, New York, 1986, pp. 185–196. MR 0870648. Zbl 0613.65058.
- [28] ———, Complexity theory and numerical analysis, in *Acta Num., Acta Numer.* **6**, Cambridge Univ. Press, Cambridge, 1997, pp. 523–551. MR 1489262. Zbl 0883.65125.
- [29] ———, Mathematical problems for the next century, in *Mathematics: Frontiers and Perspectives*, Amer. Math. Soc., Providence, RI, 2000, pp. 271–294. MR 1754783. Zbl 1031.00005.
- [30] R. SOLOVAY and V. STRASSEN, A fast Monte-Carlo test for primality, *SIAM J. Comput.* **6** (1977), 84–85. MR 0429721. Zbl 0345.10002. <http://dx.doi.org/10.1137/0206006>.
- [31] D. A. SPIELMAN and S.-H. TENG, Smoothed analysis of algorithms, in *Proceedings of the International Congress of Mathematicians, Vol. I* (Beijing, 2002), Higher Ed. Press, Beijing, 2002, pp. 597–606. MR 1989210. Zbl 1056.65148.
- [32] ———, Smoothed analysis of algorithms: why the simplex algorithm usually takes polynomial time, *J. ACM* **51** (2004), 385–463. MR 2145860. Zbl 0563.03027. <http://dx.doi.org/10.1145/990308.990310>.
- [33] M. WSCHEBOR, Smoothed analysis of $\kappa(A)$, *J. Complexity* **20** (2004), 97–107. MR 2031560. Zbl 1065.15029. <http://dx.doi.org/10.1016/j.jco.2003.09.003>.

(Received: September 17, 2009)

(Revised: October 4, 2010)

UNIVERSITY OF PADERBORN, PADERBORN, GERMANY
E-mail: pbuerg@upb.de

CITY UNIVERSITY OF HONG KONG, KOWLOON TONG, HONG KONG
E-mail: macucker@cityu.edu.hk