

Optimal asymptotic bounds for spherical designs

By ANDRIY BONDARENKO, DANYLO RADCHENKO, and MARYNA VIAZOVSKA

Abstract

In this paper we prove the conjecture of Korevaar and Meyers: for each $N \geq c_d t^d$, there exists a spherical t -design in the sphere S^d consisting of N points, where c_d is a constant depending only on d .

1. Introduction

Let S^d be the unit sphere in \mathbb{R}^{d+1} with the Lebesgue measure μ_d normalized by $\mu_d(S^d) = 1$.

A set of points $x_1, \dots, x_N \in S^d$ is called a *spherical t -design* if

$$\int_{S^d} P(x) d\mu_d(x) = \frac{1}{N} \sum_{i=1}^N P(x_i)$$

for all polynomials in $d+1$ variables, of total degree at most t . The concept of a spherical design was introduced by Delsarte, Goethals, and Seidel [12]. For each $t, d \in \mathbb{N}$, denote by $N(d, t)$ the minimal number of points in a spherical t -design in S^d . The following lower bound,

$$(1) \quad N(d, t) \geq \begin{cases} \binom{d+k}{d} + \binom{d+k-1}{d} & \text{if } t = 2k, \\ 2 \binom{d+k}{d} & \text{if } t = 2k+1, \end{cases}$$

is proved in [12].

Spherical t -designs attaining this bound are called *tight*. The vertices of a regular $t+1$ -gon form a tight spherical t -design in the circle, so $N(1, t) = t+1$. Exactly eight tight spherical designs are known for $d \geq 2$ and $t \geq 4$. All such configurations of points are highly symmetrical, and optimal from many

different points of view (see Cohn, Kumar [10] and Conway, Sloane [11]). Unfortunately, tight designs rarely exist. In particular, Bannai and Damerell [2], [3] have shown that tight spherical designs with $d \geq 2$ and $t \geq 4$ may exist only for $t = 4, 5, 7$, or 11 . Moreover, the only tight 11-design is formed by minimal vectors of the Leech lattice in dimension 24. The bound (1) has been improved by Delsarte's linear programming method for most pairs (d, t) ; see [22].

On the other hand, Seymour and Zaslavsky [20] have proved that spherical t -designs exist for all $d, t \in \mathbb{N}$. However, this proof is nonconstructive and gives no idea of how big $N(d, t)$ is. So, a natural question is to ask how $N(d, t)$ differs from bound (1). Generally, to find the exact value of $N(d, t)$ even for small d and t is a surprisingly hard problem. For example, everybody believes that 24 minimal vectors of the D_4 root lattice form a 5-design with minimal number of points in S^3 , although it is only proved that $22 \leq N(3, 5) \leq 24$; see [6]. Further, Cohn, Conway, Elkies, and Kumar [9] conjectured that every spherical 5-design consisting of 24 points in S^3 is in a certain 3-parametric family. Recently, Musin [17] has solved a long standing problem related to this conjecture. Namely, he proved that the kissing number in dimension 4 is 24.

In this paper we focus on asymptotic upper bounds on $N(d, t)$ for fixed $d \geq 2$ and $t \rightarrow \infty$. Let us give a brief history of this question. First, Wagner [21] and Bajnok [1] proved that $N(d, t) \leq C_d t^{C_d d^4}$ and $N(d, t) \leq C_d t^{C_d d^3}$, respectively. Then, Korevaar and Meyers [14] have improved these inequalities by showing that $N(d, t) \leq C_d t^{(d^2+d)/2}$. They have also conjectured that

$$N(d, t) \leq C_d t^d.$$

Note that (1) implies $N(d, t) \geq c_d t^d$. Here and in what follows we denote by C_d and c_d sufficiently large and sufficiently small positive constants depending only on d , respectively.

The conjecture of Korevaar and Meyers attracted the interest of many mathematicians. For instance, Kuijlaars and Saff [19] emphasized the importance of this conjecture for $d = 2$ and revealed its relation to minimal energy problems. Mhaskar, Narcowich, and Ward [16] have constructed positive quadrature formulas in S^d with $C_d t^d$ points having *almost* equal weights. Very recently, Chen, Frommer, Lang, Sloan, and Womersley [7], [8] gave a computer-assisted proof that spherical t -designs with $(t + 1)^2$ points exist in S^2 for $t \leq 100$.

For $d = 2$, there is an even stronger conjecture by Hardin and Sloane [13] saying that $N(2, t) \leq \frac{1}{2}t^2 + o(t^2)$ as $t \rightarrow \infty$. Numerical evidence supporting the conjecture was also given.

In [4], we have suggested a nonconstructive approach for obtaining asymptotic bounds for $N(d, t)$ based on the application of the Brouwer fixed point theorem. This led to the following result:

For each $N \geq C_d t^{\frac{2d(d+1)}{d+2}}$, there exists a spherical t -design in S^d consisting of N points.

Instead of the Brouwer fixed point theorem, in this paper we use the following result from the Brouwer degree theory [18, Ths. 1.2.6 and 1.2.9].

THEOREM A. *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a continuous mapping and Ω an open bounded subset, with boundary $\partial\Omega$, such that $0 \in \Omega \subset \mathbb{R}^n$. If $\langle x, f(x) \rangle > 0$ for all $x \in \partial\Omega$, then there exists $x \in \Omega$ satisfying $f(x) = 0$.*

We employ this theorem to prove the conjecture of Korevaar and Meyers.

THEOREM 1. *For each $N \geq C_d t^d$, there exists a spherical t -design in S^d consisting of N points.*

Note that [Theorem 1](#) is slightly stronger than the original conjecture because it guarantees the existence of spherical t -designs for *each* N greater than $C_d t^d$.

This paper is organized as follows. In [Section 2](#) we explain the main idea of the proof. Then in [Section 3](#) we present some auxiliary results. Finally, we prove [Theorem 1](#) in [Section 4](#).

2. Preliminaries and the main idea

Let \mathcal{P}_t be the Hilbert space of polynomials P on S^d of degree at most t such that

$$\int_{S^d} P(x) d\mu_d(x) = 0,$$

equipped with the usual inner product

$$\langle P, Q \rangle = \int_{S^d} P(x)Q(x) d\mu_d(x).$$

By the Riesz representation theorem, for each point $x \in S^d$, there exists a unique polynomial $G_x \in \mathcal{P}_t$ such that

$$\langle G_x, Q \rangle = Q(x) \text{ for all } Q \in \mathcal{P}_t.$$

Then a set of points $x_1, \dots, x_N \in S^d$ forms a spherical t -design if and only if

$$(2) \quad G_{x_1} + \dots + G_{x_N} = 0.$$

The gradient of a differentiable function $f : \mathbb{R}^{d+1} \rightarrow \mathbb{R}$ is denoted by

$$\frac{\partial f}{\partial x} := \left(\frac{\partial f}{\partial \xi_1}, \dots, \frac{\partial f}{\partial \xi_{d+1}} \right), \quad x = (\xi_1, \dots, \xi_{d+1}).$$

For a polynomial $Q \in \mathcal{P}_t$, we define the *spherical gradient*

$$(3) \quad \nabla Q(x) := \frac{\partial}{\partial x} \left(Q \left(\frac{x}{|x|} \right) \right),$$

where $|\cdot|$ is the Euclidean norm in \mathbb{R}^{d+1} .

We apply [Theorem A](#) to the open subset Ω of a vector space \mathcal{P}_t :

$$(4) \quad \Omega := \left\{ P \in \mathcal{P}_t \mid \int_{S^d} |\nabla P(x)| d\mu_d(x) < 1 \right\}.$$

Now we observe that the existence of a continuous mapping $F : \mathcal{P}_t \rightarrow (S^d)^N$, such that for all $P \in \partial\Omega$

$$(5) \quad \sum_{i=1}^N P(x_i(P)) > 0, \text{ where } F(P) = (x_1(P), \dots, x_N(P)),$$

readily implies the existence of a spherical t -design in S^d consisting of N points. Indeed, consider a mapping $L : (S^d)^N \rightarrow \mathcal{P}_t$ defined by

$$(x_1, \dots, x_N) \xrightarrow{L} G_{x_1} + \dots + G_{x_N},$$

and the following composition mapping $f = L \circ F : \mathcal{P}_t \rightarrow \mathcal{P}_t$. Clearly

$$\langle P, f(P) \rangle = \sum_{i=1}^N P(x_i(P))$$

for each $P \in \mathcal{P}_t$. Thus, applying [Theorem A](#) to the mapping f , the vector space \mathcal{P}_t , and the subset Ω defined by (4), we obtain that $f(Q) = 0$ for some $Q \in \mathcal{P}_t$. Hence, by (2), the components of $F(Q) = (x_1(Q), \dots, x_N(Q))$ form a spherical t -design in S^d consisting of N points.

The most naive approach to construct such F is to start with a certain well-distributed collection of points x_i ($i = 1, \dots, N$), put $F(0) := (x_1, \dots, x_N)$, and then move each point along the spherical gradient vector field of P . Note that this is the most greedy way to increase each $P(x_i(P))$ and make $\sum_{i=1}^N P(x_i(P))$ positive for each $P \in \partial\Omega$. Following this approach we will give an explicit construction of F in [Section 4](#), which will immediately imply the proof of [Theorem 1](#).

3. Auxiliary results

To construct the corresponding mapping F for each $N \geq C_d t^d$, we extensively use the following notion of an area-regular partition.

Let $\mathcal{R} = \{R_1, \dots, R_N\}$ be a finite collection of closed sets $R_i \subset S^d$ such that $\cup_{i=1}^N R_i = S^d$ and $\mu_d(R_i \cap R_j) = 0$ for all $1 \leq i < j \leq N$. The partition \mathcal{R} is called area-regular if $\mu_d(R_i) = 1/N$, $i = 1, \dots, N$. The partition norm for \mathcal{R} is defined by

$$\|\mathcal{R}\| := \max_{R \in \mathcal{R}} \text{diam } R,$$

where $\text{diam } R$ stands for the maximum geodesic distance between two points in R . We need the following fact on area-regular partitions (see Bourgain, Lindenstrauss [5] and Kuijlaars, Saff [15]).

THEOREM B. *For each $N \in \mathbb{N}$, there exists an area-regular partition $\mathcal{R} = \{R_1, \dots, R_N\}$ with $\|\mathcal{R}\| \leq B_d N^{-1/d}$ for some constant B_d large enough.*

We will also use a result that is an easy corollary of Theorem 3.1 in [16].

THEOREM C. *There exists a constant r_d such that for each area-regular partition $\mathcal{R} = \{R_1, \dots, R_N\}$ with $\|\mathcal{R}\| < \frac{r_d}{m}$, each collection of points $x_i \in R_i$ ($i = 1, \dots, N$), and each polynomial P of total degree m , the inequality*

$$(6) \quad \frac{1}{2} \int_{S^d} |P(x)| d\mu_d(x) \leq \frac{1}{N} \sum_{i=1}^N |P(x_i)| \leq \frac{3}{2} \int_{S^d} |P(x)| d\mu_d(x)$$

holds.

Theorem 3.1 in [16] was stated for slightly different definition of an area-regular partition. Namely, it was additionally assumed that each R_i is a spherical region. However the proof clearly works for our more general definition as well; see [16, §3.3].

COROLLARY 1. *For each area-regular partition $\mathcal{R} = \{R_1, \dots, R_N\}$ with $\|\mathcal{R}\| < \frac{r_d}{m+1}$, each collection of points $x_i \in R_i$ ($i = 1, \dots, N$), and each polynomial P of total degree m ,*

$$(7) \quad \frac{1}{3\sqrt{d}} \int_{S^d} |\nabla P(x)| d\mu_d(x) \leq \frac{1}{N} \sum_{i=1}^N |\nabla P(x_i)| \leq 3\sqrt{d} \int_{S^d} |\nabla P(x)| d\mu_d(x).$$

Proof. For a point $x = (\xi_1, \dots, \xi_{d+1}) \in S^d$, we get by (3) that

$$|\nabla P(x)| = \sqrt{P_1^2(x) + \dots + P_{d+1}^2(x)},$$

where

$$P_j(x) := \frac{\partial P}{\partial \xi_j}(x) - \sum_{k=1}^{d+1} \xi_j \xi_k \frac{\partial P}{\partial \xi_k}(x)$$

are polynomials of total degree at most $m+1$. Thus, using a simple inequality

$$\frac{1}{\sqrt{d+1}} \sum_{k=1}^{d+1} |P_k(x_i)| \leq \sqrt{\sum_{k=1}^{d+1} P_k^2(x_i)} \leq \sum_{k=1}^{d+1} |P_k(x_i)|$$

and then applying (6) to polynomials P_k , we obtain the statement of the corollary. \square

4. Proof of Theorem 1

In this section we construct the map F introduced in Section 2 and thereby finish the proof of Theorem 1.

For $d, t \in \mathbb{N}$, take $C_d > (54dB_d/r_d)^d$, where B_d is as in Theorem B and r_d is as in Theorem C, and fix $N \geq C_d t^d$. Now we are in a position to give an exact

construction of the mapping $F : \mathcal{P}_t \rightarrow (S^d)^N$, which satisfies [condition \(5\)](#). Take an area-regular partition $\mathcal{R} = \{R_1, \dots, R_N\}$ with

$$(8) \quad \|\mathcal{R}\| \leq B_d N^{-1/d} < \frac{r_d}{54dt}$$

as provided by [Theorem B](#), and choose an arbitrary $x_i \in R_i$ for each $i = 1, \dots, N$. Put $\varepsilon = \frac{1}{6\sqrt{d}}$, and consider the function

$$h_\varepsilon(u) := \begin{cases} u & \text{if } u > \varepsilon, \\ \varepsilon & \text{otherwise.} \end{cases}$$

Take a mapping $U : \mathcal{P}_t \times S^d \rightarrow \mathbb{R}^{d+1}$ such that

$$U(P, y) = \frac{\nabla P(y)}{h_\varepsilon(|\nabla P(y)|)}.$$

For each $i = 1, \dots, N$, let $y_i : \mathcal{P}_t \times [0, \infty) \rightarrow S^d$ be the map satisfying the differential equation

$$(9) \quad \frac{d}{ds} y_i(P, s) = U(P, y_i(P, s))$$

with the initial condition

$$y_i(P, 0) = x_i$$

for each $P \in \mathcal{P}_t$. Note that each mapping y_i has its values in S^d by definition of spherical gradient [\(3\)](#). Since the mapping $U(P, y)$ is Lipschitz continuous in both P and y , each y_i is well defined and continuous in both P and s , where the metric on \mathcal{P}_t is given by the inner product. Finally, put

$$(10) \quad F(P) = (x_1(P), \dots, x_N(P)) := \left(y_1\left(P, \frac{r_d}{3t}\right), \dots, y_N\left(P, \frac{r_d}{3t}\right) \right).$$

By definition, the mapping F is continuous on \mathcal{P}_t . So, as explained in [Section 2](#), to finish the proof of [Theorem 1](#) it suffices to prove

LEMMA 1. *Let $F : \mathcal{P}_t \rightarrow (S^d)^N$ be the mapping defined by [\(10\)](#). Then for each $P \in \partial\Omega$,*

$$\frac{1}{N} \sum_{i=1}^N P(x_i(P)) > 0,$$

where Ω is given by [\(4\)](#).

Proof. Fix $P \in \partial\Omega$; that is,

$$\int_{S^d} |\nabla P(x)| d\mu_d(x) = 1.$$

For the sake of simplicity, we write $y_i(s)$ in place of $y_i(P, s)$. By the Newton-Leibniz formula, we have

$$(11) \quad \begin{aligned} \frac{1}{N} \sum_{i=1}^N P(x_i(P)) &= \frac{1}{N} \sum_{i=1}^N P(y_i(r_d/3t)) \\ &= \frac{1}{N} \sum_{i=1}^N P(x_i) + \int_0^{r_d/3t} \frac{d}{ds} \left[\frac{1}{N} \sum_{i=1}^N P(y_i(s)) \right] ds. \end{aligned}$$

Now to prove [Lemma 1](#), we first estimate the value

$$\left| \frac{1}{N} \sum_{i=1}^N P(x_i) \right|$$

from above and then estimate the value

$$\frac{d}{ds} \left[\frac{1}{N} \sum_{i=1}^N P(y_i(s)) \right]$$

from below for each $s \in [0, r_d/3t]$. We have

$$\begin{aligned} \left| \frac{1}{N} \sum_{i=1}^N P(x_i) \right| &= \left| \sum_{i=1}^N \int_{R_i} P(x_i) - P(x) d\mu_d(x) \right| \leq \sum_{i=1}^N \int_{R_i} |P(x_i) - P(x)| d\mu_d(x) \\ &\leq \frac{\|\mathcal{R}\|}{N} \sum_{i=1}^N \max_{z \in S^d: \text{dist}(z, x_i) \leq \|\mathcal{R}\|} |\nabla P(z)|, \end{aligned}$$

where $\text{dist}(z, x_i)$ denotes the geodesic distance between z and x_i . Hence, for $z_i \in S^d$ such that $\text{dist}(z_i, x_i) \leq \|\mathcal{R}\|$ and

$$|\nabla P(z_i)| = \max_{z \in S^d: \text{dist}(z, x_i) \leq \|\mathcal{R}\|} |\nabla P(z)|,$$

we obtain

$$\left| \frac{1}{N} \sum_{i=1}^N P(x_i) \right| \leq \frac{\|\mathcal{R}\|}{N} \sum_{i=1}^N |\nabla P(z_i)|.$$

Consider another area-regular partition $\mathcal{R}' = \{R'_1, \dots, R'_N\}$ defined by $R'_i = R_i \cup \{z_i\}$. Clearly $\|\mathcal{R}'\| \leq 2\|\mathcal{R}\|$ and so, by [\(8\)](#), we get $\|\mathcal{R}'\| < r_d/(27dt)$. Applying [inequality \(7\)](#) to the partition \mathcal{R}' and the collection of points z_i , we obtain that

$$(12) \quad \left| \frac{1}{N} \sum_{i=1}^N P(x_i) \right| \leq 3\sqrt{d}\|\mathcal{R}\| \int_{S^d} |\nabla P(x)| d\mu_d(x) < \frac{r_d}{18\sqrt{d}t}$$

for any $P \in \partial\Omega$. On the other hand, the differential equation (9) implies

$$\begin{aligned}
 (13) \quad \frac{d}{ds} \left[\frac{1}{N} \sum_{i=1}^N P(y_i(s)) \right] &= \frac{1}{N} \sum_{i=1}^N \frac{|\nabla P(y_i(s))|^2}{h_\varepsilon(|\nabla P(y_i(s))|)} \\
 &\geq \frac{1}{N} \sum_{i: |\nabla P(y_i(s))| \geq \varepsilon} |\nabla P(y_i(s))| \\
 &\geq \frac{1}{N} \sum_{i=1}^N |\nabla P(y_i(s))| - \varepsilon.
 \end{aligned}$$

Since

$$\left| \frac{\nabla P(y)}{h_\varepsilon(|\nabla P(y)|)} \right| \leq 1$$

for each $y \in S^d$, it follows again from (9) that $\left| \frac{dy_i(s)}{ds} \right| \leq 1$. Hence we arrive at

$$\text{dist}(x_i, y_i(s)) \leq s.$$

Now for each $s \in [0, r_d/3t]$, we consider the area-regular partition $\mathcal{R}'' = \{R''_1, \dots, R''_N\}$ given by $R''_i = R_i \cup \{y_i(s)\}$. By (8), we have

$$\|\mathcal{R}''\| < \frac{r_d}{54dt} + \frac{r_d}{3t},$$

so we can apply (7) to the partition \mathcal{R}'' and the collection of points $y_i(s)$. This and inequality (13) yield

$$\begin{aligned}
 (14) \quad \frac{d}{ds} \left[\frac{1}{N} \sum_{i=1}^N P(y_i(s)) \right] &\geq \frac{1}{N} \sum_{i=1}^N |\nabla P(y_i(s))| - \frac{1}{6\sqrt{d}} \\
 &\geq \frac{1}{3\sqrt{d}} \int_{S^d} |\nabla P(x)| d\mu_d(x) - \frac{1}{6\sqrt{d}} = \frac{1}{6\sqrt{d}}
 \end{aligned}$$

for each $P \in \partial\Omega$ and $s \in [0, r_d/3t]$. Finally, equation (11) and inequalities (12) and (14) imply

$$(15) \quad \frac{1}{N} \sum_{i=1}^N P(x_i(P)) > \frac{1}{6\sqrt{d}} \frac{r_d}{3t} - \frac{r_d}{18\sqrt{d}t} = 0.$$

Lemma 1 is proved. □

Acknowledgements. We thank Joaquim Bruna, Jacob Korevaar, Pieter Moree, Michael Ontiveros, Joaquim Ortega-Cerdà, Andriy Prymak, Edward Saff, Igor Shevchuk, and Sergey Tikhonov for several fruitful discussions. We also thank the referee for valuable suggestions.

References

- [1] B. BAJNOK, Construction of spherical t -designs, *Geom. Dedicata* **43** (1992), 167–179. MR 1180648. Zbl 0765.05032. <http://dx.doi.org/10.1007/BF00147866>.
- [2] E. BANNAI and R. M. DAMERELL, Tight spherical designs. I, *J. Math. Soc. Japan* **31** (1979), 199–207. MR 0519045. Zbl 0403.05022. <http://dx.doi.org/10.2969/jmsj/031110199>.
- [3] E. BANNAI and R. M. DAMERELL, Tight spherical designs. II, *J. London Math. Soc.* **21** (1980), 13–30. MR 0576179. Zbl 0436.05018. <http://dx.doi.org/10.1112/jlms/s2-21.1.13>.
- [4] A. V. BONDARENKO and M. S. VIAZOVSKA, Spherical designs via Brouwer fixed point theorem, *SIAM J. Discrete Math.* **24** (2010), 207–217. MR 2600661. Zbl 1229.05057. <http://dx.doi.org/10.1137/080738313>.
- [5] J. BOURGAIN and J. LINDENSTRAUSS, Distribution of points on spheres and approximation by zonotopes, *Israel J. Math.* **64** (1988), 25–31. MR 0981745. Zbl 0667.52001. <http://dx.doi.org/10.1007/BF02767366>.
- [6] P. BOYVALENKOV, D. DANEV, and S. NIKOVA, Nonexistence of certain spherical designs of odd strengths and cardinalities, *Discrete Comput. Geom.* **21** (1999), 143–156. MR 1661307. Zbl 0921.05016. <http://dx.doi.org/10.1007/PL00009406>.
- [7] X. CHEN, A. FROMMER, and B. LANG, Computational existence proofs for spherical t -designs, *Numer. Math.* **117** (2011), 289–305. MR 2754852. Zbl 1208.65032. <http://dx.doi.org/10.1007/s00211-010-0332-5>.
- [8] X. CHEN and R. S. WOMERSLEY, Existence of solutions to systems of under-determined equations and spherical designs, *SIAM J. Numer. Anal.* **44** (2006), 2326–2341. MR 2272596. Zbl 1129.65035. <http://dx.doi.org/10.1137/050626636>.
- [9] H. COHN, J. H. CONWAY, N. D. ELKIES, and A. KUMAR, The D_4 root system is not universally optimal, *Experiment. Math.* **16** (2007), 313–320. MR 2367321. Zbl 1137.05020. <http://dx.doi.org/10.1080/10586458.2007.10129008>.
- [10] H. COHN and A. KUMAR, Universally optimal distribution of points on spheres, *J. Amer. Math. Soc.* **20** (2007), 99–148. MR 2257398. Zbl 1198.52009. <http://dx.doi.org/10.1090/S0894-0347-06-00546-7>.
- [11] J. H. CONWAY and N. J. A. SLOANE, *Sphere Packings, Lattices and Groups*, third ed., *Grundle Math. Wissen.* **290**, Springer-Verlag, New York, 1999. MR 1662447. Zbl 0915.52003.
- [12] P. DELSARTE, J. M. GOETHALS, and J. J. SEIDEL, Spherical codes and designs, *Geometriae Dedicata* **6** (1977), 363–388. MR 0485471. Zbl 0376.05015. <http://dx.doi.org/10.1007/BF03187604>.
- [13] R. H. HARDIN and N. J. A. SLOANE, McLaren’s improved snub cube and other new spherical designs in three dimensions, *Discrete Comput. Geom.* **15** (1996), 429–441. MR 1384885. Zbl 0858.05024. <http://dx.doi.org/10.1007/BF02711518>.
- [14] J. KOREVAAR and J. L. H. MEYERS, Spherical Faraday cage for the case of equal point charges and Chebyshev-type quadrature on the sphere, *Integral Transform. Spec. Funct.* **1** (1993), 105–117. MR 1421438. Zbl 0823.41026. <http://dx.doi.org/10.1080/10652469308819013>.

- [15] A. B. J. KUIJLAARS and E. B. SAFF, Asymptotics for minimal discrete energy on the sphere, *Trans. Amer. Math. Soc.* **350** (1998), 523–538. MR 1458327. Zbl 0896.52019. <http://dx.doi.org/10.1090/S0002-9947-98-02119-9>.
- [16] H. N. MHASKAR, F. J. NARCOWICH, and J. D. WARD, Spherical Marcinkiewicz-Zygmund inequalities and positive quadrature, *Math. Comp.* **70** (2001), 1113–1130. MR 1710640. Zbl 0980.76070. <http://dx.doi.org/10.1090/S0025-5718-00-01240-0>.
- [17] O. R. MUSIN, The kissing number in four dimensions, *Ann. of Math.* **168** (2008), 1–32. MR 2415397. Zbl 1169.52008. <http://dx.doi.org/10.4007/annals.2008.168.1>.
- [18] D. O'REGAN, Y. J. CHO, and Y.-Q. CHEN, *Topological Degree Theory and Applications*, Ser. Math. Anal. Appl. **10**, Chapman & Hall/CRC, Boca Raton, FL, 2006. MR 2223854. Zbl 1095.47001. <http://dx.doi.org/10.1201/9781420011487>.
- [19] E. B. SAFF and A. B. J. KUIJLAARS, Distributing many points on a sphere, *Math. Intelligencer* **19** (1997), 5–11. MR 1439152. Zbl 0901.11028. <http://dx.doi.org/10.1007/BF03024331>.
- [20] P. D. SEYMOUR and T. ZASLAVSKY, Averaging sets: a generalization of mean values and spherical designs, *Adv. in Math.* **52** (1984), 213–240. MR 0744857. Zbl 0596.05012. [http://dx.doi.org/10.1016/0001-8708\(84\)90022-7](http://dx.doi.org/10.1016/0001-8708(84)90022-7).
- [21] G. WAGNER, On averaging sets, *Monatsh. Math.* **111** (1991), 69–78. MR 1089385. Zbl 0721.65011. <http://dx.doi.org/10.1007/BF01299278>.
- [22] V. A. YUDIN, Lower bounds for spherical designs, *Izv. Ross. Akad. Nauk Ser. Mat.* **61** (1997), 213–223. MR 1478566. Zbl 0890.05015. <http://dx.doi.org/10.1070/im1997v061n03ABEH000132>.

(Received: March 10, 2011)

CENTRE DE RECERCA MATEMÀTICA, BELLATERRA (BARCELONA), SPAIN and
 NATIONAL TARAS SHEVCHENKO UNIVERSITY, KYIV, UKRAINE and
 NORWEGIAN UNIVERSITY OF SCIENCE AND TECHNOLOGY, TRONDHEIM, NORWAY
E-mail: bonda@univ.kiev.ua

MAX PLANCK INSTITUTE FOR MATHEMATICS, BONN, GERMANY and
 NATIONAL TARAS SHEVCHENKO UNIVERSITY, KYIV, UKRAINE
E-mail: danrad@mpim-bonn.mpg.de

UNIVERSITY OF COLOGNE, COLOGNE, GERMANY and
 MAX PLANCK INSTITUTE FOR MATHEMATICS, BONN, GERMANY
E-mail: mviazovs@math.uni-koeln.de