

Euler systems for Rankin–Selberg convolutions of modular forms

By ANTONIO LEI, DAVID LOEFFLER, and SARAH LIVIA ZERBES

Dedicated to Kazuya Kato

Abstract

We construct a Euler system in the cohomology of the tensor product of the Galois representations attached to two modular forms, using elements in the higher Chow groups of products of modular curves. We use these elements to prove a finiteness theorem for the strict Selmer group of the Galois representation when the associated p -adic Rankin–Selberg L -function is nonvanishing at $s = 1$.

Contents

1. Outline	653
2. Generalized Beilinson–Flach elements	655
3. Norm relations for generalized Beilinson–Flach elements	673
4. Relation to complex L -values	704
5. Relation to p -adic L -values	713
6. Families of cohomology classes	722
7. Bounding strict Selmer groups	743
8. Conjectures on higher-rank Euler systems	756
Appendix A. Ancillary results	761
References	767

1. Outline

In [BDR12], Bertolini, Darmon and Rotger studied certain canonical global cohomology classes (the “Beilinson–Flach elements,” obtained from the constructions of [Bei84] and [Fla92]) in the cohomology of the tensor products of the p -adic Galois representations of pairs of weight 2 modular forms. They related their image under the Bloch–Kato logarithm maps to the values of p -adic

The authors’ research is supported by the following grants: CRM-ISM Postdoctoral Fellowship (Lei); Royal Society University Research Fellowship (Loeffler); EPSRC First Grant EP/J018716/1 (Zerbes).

© 2014 Department of Mathematics, Princeton University.

Rankin–Selberg L -functions. These Beilinson–Flach elements are constructed as the image of elements of the higher Chow group of a product of modular curves.

In this paper, we construct a form of Euler system — a compatible system of cohomology classes over cyclotomic fields — of which the Beilinson–Flach elements are the bottom layer. We first define elements of higher Chow groups of the product of two (affine) modular curves over a cyclotomic field,

$$c\Xi_{m,N,j} \in \mathrm{CH}^2(Y_1(N)^2 \otimes \mathbb{Q}(\mu_m), 1)$$

for integers $m \geq 1$, $N \geq 5$, and $j \in \mathbb{Z}/m\mathbb{Z}$ (cf. Definition 2.7.3). These are obtained by considering the images of various maps from higher level modular curves to the surface $Y_1(N)^2$, together with modular units (Siegel units) on these curves. For $m = 1$, our elements reduce to those considered in [BDR12], and as in op. cit., we show that after tensoring with \mathbb{Q} we can construct preimages of our elements in the higher Chow group of the self-product of the projective modular curve $X_1(N)$; however, in this paper (as in [Kat04]) we shall take the affine versions as the principal objects of study.

We show two forms of compatibility relation for our generalized Beilinson–Flach elements: firstly, relating $c\Xi_{m,N,j}$ to the pushforward of $c\Xi_{m,Np,j}$ where p is prime (Theorem 3.1.2); secondly, relating $c\Xi_{m,N,j}$ to the pushforward (or Galois norm) of $c\Xi_{mp,N,j}$ where p is prime and either $p \mid N$ (Theorem 3.3.2) or $p \nmid mN$ (Theorem 3.4.1).

We next turn to the relation between our elements and L -values. Theorem 4.3.7 shows, following an argument due to Beilinson, that the images of the elements $c\Xi_{m,N,j}$ under the Beilinson regulator map into complex de Rham cohomology are related to the derivatives at $s = 1$ of Rankin–Selberg L -functions of weight 2 modular forms. Theorem 5.6.4 is a p -adic analogue of this result, generalizing a theorem of Bertolini–Darmon–Rotger [BDR12]; it gives a formula for the image of our element for $m = 1$ under the p -adic syntomic regulator, for a prime $p \nmid N$, in terms of Hida’s p -adic Rankin–Selberg L -functions.

Next we consider the images of our elements in étale cohomology. Applying Huber’s “continuous étale realization” functor and the Hochschild–Serre exact sequence, and projecting into the isotypical component corresponding to a pair of eigenforms (f, g) of level N , allows us to construct Galois cohomology classes

$$c\mathbf{Z}_m^{(f,g,N)} \in H^1(\mathbb{Q}(\mu_m), V_f^* \otimes V_g^*)$$

from the elements $c\Xi_{m,N,j}$; see Definition 6.4.4. Using the second norm relation in the p -adic cyclotomic tower, we can modify these to construct elements of Iwasawa cohomology groups of pairs of modular forms (under a strong “ordinarity” hypothesis; this is Theorem 6.8.6) or of the tensor product of Iwasawa cohomology groups with the algebra of distributions (under a weaker “small

slope” hypothesis; see Theorem 6.8.4). These elements satisfy compatibility relations of Euler-system type when additional primes are added to m .

Using the first norm relation, we also obtain variation in Hida families. More specifically, if one of the two forms (say g) is ordinary, we may deform our cohomology classes analytically as g varies over a Hida family; cf. Theorem 6.9.5. (In the special case $m = 1$, such results have been independently obtained by Bertolini–Darmon–Rotger.) When f and g are both ordinary, we obtain three-variable families that also incorporate cyclotomic twists; cf. Theorem 6.9.8.

As an application of these constructions, we prove (under some technical hypotheses) a finiteness theorem for the strict Selmer group of a product of modular forms (Theorem 7.4.2) when the associated p -adic Rankin–Selberg L -function is nonvanishing at $s = 1$, and (under very slightly stronger hypotheses) we give an explicit bound for the order of the strict Selmer group in terms of the p -adic L -value (Theorem 7.5.1).

Remark. It is a pleasure to acknowledge the very deep debt this article owes to the magisterial work of Kato [Kat04]. We have adopted many aspects of the strategy and methods of Kato’s work, reused a number of his results, and even in many cases adopted his notation. It is a pleasure to dedicate this work to Professor Kato as a humble gift on the occasion of his 60th birthday.

Acknowledgements. Part of this work was done while the second and the third author were visiting Montréal and Bielefeld in Spring 2012; they would like to thank Henri Darmon and Thomas Zink for their hospitality. We would also like to thank Massimo Bertolini, Francois Brunault, Francesc Castella, John Coates, Henri Darmon, Mathias Flach, Kazuya Kato, Masato Kurihara, Andreas Langer, Alan Lauder, Jan Nekovar, Ken Ribet, Victor Rotger, Karl Rubin, Tony Scholl, Xin Wan and Andrew Wiles for helpful comments. Finally, we are also grateful to the anonymous referee for a number of valuable suggestions and corrections, to Samit Dasgupta for pointing out several errors in an earlier draft and to Alan Lauder for computing the 17-adic L -value appearing in the example of Section 7.6.

2. Generalized Beilinson–Flach elements

In this section we shall construct elements of motivic cohomology groups of products of modular curves, using the explicit description of the motivic cohomology given by the Gersten complex.

2.1. *Modular curves.* We begin by fixing notation in regard to modular curves. We follow the conventions of [Kat04] very closely; see, in particular, Sections 1, 2, and 5 of op. cit.

Definition 2.1.1. For $N \geq 3$, let $Y(N)$ denote the smooth affine curve over \mathbb{Q} that represents the functor on the category of \mathbb{Q} -schemes

$$S \mapsto \left\{ \begin{array}{l} \text{isomorphism classes of triples } (E, e_1, e_2), \\ E \text{ an elliptic curve over } S, \\ e_1, e_2 \text{ sections of } E/S \text{ generating } E[N] \end{array} \right\}.$$

The variety $Y(N)$ is equipped with a left action of $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$: the element $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ maps (E, e_1, e_2) to (E, e'_1, e'_2) where

$$\begin{pmatrix} e'_1 \\ e'_2 \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} e_1 \\ e_2 \end{pmatrix}.$$

In particular, this action factors through $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})/\langle \pm 1 \rangle$.

There is an obvious surjective morphism $Y(N) \rightarrow \mu_N^\circ$, where μ_N° is the scheme of primitive N -th roots of unity, which sends (E, e_1, e_2) to $\langle e_1, e_2 \rangle_{E[N]}$, where $\langle -, - \rangle_{E[N]}$ denotes the Weil pairing on $E[N]$. Because the Weil pairing is nondegenerate and alternating, the induced action of $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$ on μ_N° is given by $\sigma \cdot \zeta = \zeta^{\det \sigma}$; and the fibre of $Y(N)(\mathbb{C})$ over the point $e^{2\pi i/N} \in \mu_N^\circ(\mathbb{C})$ is canonically and $\mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$ -equivariantly identified with $\Gamma(N) \backslash \mathcal{H}$, where \mathcal{H} is the upper half-plane and $\Gamma(N)$ the principal congruence subgroup of level N in $\mathrm{SL}_2(\mathbb{Z})$, via the map

$$\tau \mapsto (\mathbb{C}/(\mathbb{Z} + \mathbb{Z}\tau), \tau/N, 1/N).$$

We shall mainly be working with certain quotients of the curves $Y(N)$, which we now define.

Definition 2.1.2. Let $M, N \geq 1$, and choose $L \geq 3$ divisible by both M and N . We let $Y(M, N)$ be the quotient of $Y(L)$ by the action of the group

$$\left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}_2(\mathbb{Z}/L\mathbb{Z}) : \begin{array}{l} a = 1, b = 0 \pmod{M}, \\ c = 0, d = 1 \pmod{N} \end{array} \right\}.$$

(This is easily seen to be independent of the choice of L .)

If $M + N \geq 5$, this curve represents This curve $Y(M, N)$ represents the functor of triples (E, e_1, e_2) where e_1 has order M , e_2 has order N , and e_1, e_2 generate a subgroup of E of order MN .

Definition 2.1.3. We write $Y_1(N)$ for the smooth affine curve over \mathbb{Q} representing the functor

$$S \mapsto \left\{ \begin{array}{l} \text{isomorphism classes of pairs } (E, P), \\ E \text{ an elliptic curve over } S, \\ P \text{ a section of } E/S \text{ of exact order } N \end{array} \right\}.$$

Remark 2.1.4. Note that the cusp ∞ , which corresponds to the generalized elliptic curve $(\mathbb{G}_m/q^{\mathbb{Z}}, \zeta_N)$, is not defined over $\mathbb{Q}[[q]]$ but rather over $\mathbb{Q}(\mu_N)[[q]]$, so the q -expansions of elements of $\mathcal{O}(Y_1(N))$ do not necessarily lie in $\mathbb{Q}((q))$ but rather in $\mathbb{Q}(\mu_N)((q))$. See, e.g., [DI95, §9.3] for further discussion.

It is clear that $Y_1(N) = Y(1, N)$. More generally, we may use the following proposition to identify $Y_1(N) \times \mu_m^\circ$, for $m, N \geq 1$, with a quotient of a principal modular curve:

PROPOSITION 2.1.5. *If $N \geq 3, m \geq 1$, and $L \geq 3$ is divisible by both N and m , then the map*

$$\begin{aligned} Y(L) &\longrightarrow Y_1(N) \otimes \mu_m^\circ \\ (E, e_1, e_2) &\longmapsto \left[\left(E, \frac{L}{N}e_2 \right), \left\langle \frac{L}{m}e_1, \frac{L}{m}e_2 \right\rangle_{E[m]} \right] \end{aligned}$$

identifies $Y_1(N) \times \mu_m^\circ$ with the quotient of $Y(L)$ by the subgroup of $\mathrm{GL}_2(\mathbb{Z}/L\mathbb{Z})$ given by

$$\left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : \begin{array}{l} c = 0, d = 1 \pmod{N}, \\ ad - bc = 1 \pmod{m} \end{array} \right\}.$$

We shall be most interested in the curves $Y(m, mN)$ for $m \geq 1, N \geq 1$. Note that $Y(m, mN)$ maps naturally to μ_m° , with geometrically connected fibres. It has a left action of the group

$$\left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : c = 0 \pmod{N} \right\},$$

compatible with the determinant action on μ_m° : if $x = (E, e_1, e_2)$ is a point of $Y(m, mN)$, so e_1 has order m and e_2 has order mN , and $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}_2(\mathbb{Z}/mN\mathbb{Z})$ with $N|c$, then

$$g \cdot x = (E, ae_1 + bNe_2, c/Ne_1 + de_2).$$

We shall introduce some notation for maps between these curves.

Definition 2.1.6. Let $m, N \geq 1$.

(1) We write t_m for the morphism $Y(m, mN) \rightarrow Y_1(N) \times \mu_m^\circ$ given by

$$(E, e_1, e_2) \mapsto \left[(E/\langle e_1 \rangle, [me_2]), \langle e_1, Ne_2 \rangle_{E[m]} \right].$$

(2) For $a \geq 1$, we write τ_a for the morphism $Y(am, amN) \rightarrow Y(m, mN)$ given by

$$(E, e_1, e_2) \mapsto (E/C, [e_1], [ae_2]),$$

where C is the cyclic subgroup of order a generated by me_1 , and $[e_1], [ae_2]$ denote the images of e_1 and ae_2 on E/C .

PROPOSITION 2.1.7. *Let m, N, a as above.*

(1) *We have a commutative diagram*

$$\begin{array}{ccc} Y(am, amN) & \xrightarrow{\tau_a} & Y(m, mN) \\ \downarrow t_{am} & & \downarrow t_m \\ Y_1(N) \times \mu_{am}^\circ & \longrightarrow & Y_1(N) \times \mu_m^\circ \end{array}$$

where the bottom horizontal arrow is the identity map on $Y_1(N)$ and the map $\mu_{am}^\circ \rightarrow \mu_m^\circ$ given by $\zeta \mapsto \zeta^a$.

- (2) For $b \in (\mathbb{Z}/mN\mathbb{Z})^\times$, the map t_m intertwines the action of $\begin{pmatrix} b & 0 \\ 0 & 1 \end{pmatrix}$ with the automorphism $\sigma_b : \zeta \mapsto \zeta^b$ of μ_m° and $\begin{pmatrix} b^{-1} & 0 \\ 0 & b \end{pmatrix}$ with the diamond operator $\langle b \rangle$ on $Y_1(N)$.

Proof. The first statement is immediate from the definition of the maps and properties of the Weil pairing, and the second is an easy verification (cf. [Kat04, 5.7.1]). □

Remark 2.1.8. The use of the maps τ_a is forced on us by the nature of our construction of zeta elements. It would be much more satisfying to use the natural degeneracy maps $Y(am, amN) \rightarrow Y(m, mN)$ given by $\tau'_a : (E, e_1, e_2) \mapsto (E, ae_1, ae_2)$, but we do not know how to construct elements compatible under these maps; see Section 2.9 below.

Notation. For compatibility with [Kat04], we shall use the alternative notation $Y_1(N) \otimes_{\mathbb{Q}} \mathbb{Q}(\mu_m)$ for $Y_1(N) \times \mu_m^\circ$.

We shall also have to deal with products of two modular curves.

Definition 2.1.9. We shall write $Y(N)^2$ (slightly abusively) for the fibre product $Y(N) \times_{\mu_N^\circ} Y(N)$. This is a subvariety of $Y(N) \times_{\text{Spec}(\mathbb{Q})} Y(N)$ preserved by the subgroup

$$\{(\sigma, \tau) \in \text{GL}_2(\mathbb{Z}/N\mathbb{Z})^2 : \det(\sigma) = \det(\tau)\}.$$

Similarly, we shall write $Y(m, mN)^2$ for $Y(m, mN) \times_{\mu_m^\circ} Y(m, mN)$, which is acted upon by the group

$$G = \left\{ (\sigma, \tau) \in \text{GL}_2(\mathbb{Z}/mN\mathbb{Z})^2 : \det(\sigma) = \det(\tau) \pmod{m}, \sigma, \tau = \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N} \right\}.$$

Evidently, the image of $Y(m, mN) \times_{\mu_m^\circ} Y(m, mN)$ under $t_m \times t_m$ lands in

$$(Y_1(N) \times \mu_m^\circ) \times_{\mu_m^\circ} (Y_1(N) \times \mu_m^\circ) = Y_1(N)^2 \times \mu_m^\circ,$$

so we may consider $t_m \times t_m$ as a morphism $Y(m, mN)^2 \rightarrow Y_1(N)^2 \times \mu_m^\circ$, which intertwines the action of $\left(\begin{pmatrix} b & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} b & 0 \\ 0 & 1 \end{pmatrix}\right) \in G$ with σ_b .

We shall also have to consider, occasionally, some more general classes of modular curves. Here we shall only consider models over $\overline{\mathbb{Q}}$.

Definition 2.1.10. If $\Gamma \subseteq \text{SL}_2(\mathbb{Z})$ is a congruence subgroup, then we shall write $Y(\Gamma)$ for the variety

$$(\Gamma \backslash Y(L)) \otimes_{\mathbb{Q}(\mu_L)} \overline{\mathbb{Q}},$$

where L is any integer ≥ 3 such that $\Gamma \supseteq \Gamma(L)$; this variety is independent of the choice of L .

Remark 2.1.11. If $\alpha \in \mathrm{GL}_2^+(\mathbb{Q})$, then the isomorphism of Riemann surfaces $Y(\Gamma)(\mathbb{C}) \cong Y(\alpha\Gamma\alpha^{-1})(\mathbb{C})$ mapping $\tau \in \mathcal{H}$ to $\alpha\tau$ extends to an algebraic isomorphism defined over $\overline{\mathbb{Q}}$; similarly for degeneracy maps $Y(\Gamma) \rightarrow Y(\Gamma')$ for $\Gamma \subseteq \Gamma'$.

The above constructions with affine modular curves also have projective analogues where the cusps are taken into account; we shall write $X(-)$ for the compactified version of $Y(-)$ in line with the standard notation.

2.2. *Siegel units.*

Definition 2.2.1. For $(\alpha, \beta) \in (\mathbb{Q}/\mathbb{Z})^2 - \{(0, 0)\}$ of order dividing N , and $c > 1$ coprime to $6N$, let ${}_c g_{\alpha, \beta} \in \mathcal{O}(Y(N))^\times$ denote Kato’s Siegel unit, as defined in [Kat04, §1.4].

We identify ${}_c g_{\alpha, \beta}$ with a holomorphic function on the upper half-plane, via the identification of the fibre of $Y(N)(\mathbb{C})$ over $e^{2\pi i/N} \in \mu_N^\circ(\mathbb{C})$ with $\Gamma(N) \backslash \mathcal{H}$ given in the previous section. (Note that ${}_c g_{\alpha, \beta}$ is defined over \mathbb{Q} as a function on $Y(N)$, but in order to interpret it as a holomorphic function on \mathcal{H} we must make a choice of N -th root of unity, and the q -expansion coefficients of ${}_c g_{\alpha, \beta}$ are in $\mathbb{Q}(\mu_N)$.)

Recall that there is an element $g_{\alpha, \beta} \in \mathcal{O}(Y(N))^\times \otimes \mathbb{Q}$ such that ${}_c g_{\alpha, \beta} = c^2 g_{\alpha, \beta} - g_{c\alpha, c\beta}$.

PROPOSITION 2.2.2 (Distribution relations). *Let $m \geq 1$, and let c be a nonzero integer coprime to $6m$ and the orders of α, β . Then the following relations hold:*

$$(1a) \quad {}_c g_{\alpha, \beta}(mz) = \prod_{\beta'} {}_c g_{\alpha, \beta'}(z),$$

where the product is over $\beta' \in \mathbb{Q}/\mathbb{Z}$ such that $m\beta' = \beta$;

$$(1b) \quad {}_c g_{\alpha, \beta}(z/m) = \prod_{\alpha'} {}_c g_{\alpha', \beta}(z),$$

where the product is over $\alpha' \in \mathbb{Q}/\mathbb{Z}$ such that $m\alpha' = \alpha$; and

$$(1c) \quad {}_c g_{\alpha, \beta}(z) = \prod_{\alpha', \beta'} {}_c g_{\alpha', \beta'}(z),$$

where the product is over pairs $(\alpha', \beta') \in (\mathbb{Q}/\mathbb{Z})^2$ such that $(m\alpha', m\beta') = (\alpha, \beta)$.

Proof. Formula (1a) is Lemma 2.12 of [Kat04]. Formula (1b) can be proved similarly, or it can be deduced directly from (1a) using the action of $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. Formula (1c), which is Lemma 1.7(2) of op. cit., is immediate by combining (1) and (2). □

Remark 2.2.3. The three formulae above admit the following common generalization: let M be a 2×2 integer matrix with positive determinant D . Then we have

$${}_c g_{\alpha,\beta}(M \cdot z) = \prod_{\alpha',\beta'} {}_c g_{\alpha',\beta'},$$

where the product is over all (α', β') such that $(\alpha', \beta')M' = (\alpha, \beta)$, where $M' = (\det M)M^{-1}$ is the adjugate matrix of M . Cases (1), (2) and (3) correspond to taking $M = \begin{pmatrix} m & 0 \\ 0 & 1 \end{pmatrix}$, $\begin{pmatrix} 1 & 0 \\ 0 & m \end{pmatrix}$ and $\begin{pmatrix} m & 0 \\ 0 & m \end{pmatrix}$ respectively. The case where M is invertible is (part of) Lemma 1.7(1) of op. cit.

We are most interested in the units ${}_c g_{0,1/N}$, which descend to units on $Y_1(N)$. These have the following compatibility property:

THEOREM 2.2.4 (Kato). *If $M, N, N' \geq 1$ are integers with $\text{prime}(N') = \text{prime}(N)$, and α is the natural projection $Y(M, N') \rightarrow Y(M, N)$ (which induces a norm map $\alpha_* : \mathcal{O}(Y(M, N'))^\times \rightarrow \mathcal{O}(Y(M, N))^\times$), then we have*

$$\alpha_*({}_c g_{0,1/N'}) = {}_c g_{0,1/N}.$$

If $N' = N\ell$, where ℓ is prime and $\ell \nmid MN$, then we have

$$\alpha_*({}_c g_{0,1/N'}) = {}_c g_{0,1/N} \cdot ({}_c g_{0, \ell^{-1}/N})^{-1},$$

where “ ℓ^{-1} ” signifies the inverse of ℓ modulo N .

Proof. These statements are proved in [Kat04, §§2.11, 2.13] (in the course of proving the norm-compatibility relation for Kato’s elements of K_2 , Propositions 2.3 and 2.4 of op. cit.). We reproduce the proofs briefly here.

Firstly, let us suppose $\text{prime}(mN') = \text{prime}(mN)$. Let $a = N'/N$. Since $\text{prime}(mN') = \text{prime}(mN)$, for each $(x, y) \in (\mathbb{Z}/a\mathbb{Z})^2$ we may choose an element $s_{xy} \in \text{GL}_2(\mathbb{Z}/mN'\mathbb{Z})$ of the form

$$\begin{pmatrix} 1 & 0 \\ mNx & 1 + mNy \end{pmatrix}.$$

These elements s_{xy} are coset representatives for the quotient of the two subgroups of $\text{GL}_2(\mathbb{Z}/mN'\mathbb{Z})$ corresponding to $Y(m, mN)$ and $Y(m, mN')$, so we have

$$\alpha_*({}_c g_{0,1/N'}) = \prod_{x,y} s_{xy}^* ({}_c g_{0,1/N'}).$$

For any $M \geq 1$, any $u \in \text{GL}_2(\mathbb{Z}/M\mathbb{Z})$ and any $\alpha, \beta \in (\frac{1}{M}\mathbb{Z}/\mathbb{Z})$, we have

$$u^* ({}_c g_{\alpha,\beta}) = {}_c g_{\alpha',\beta'},$$

where

$$(\alpha', \beta') = (\alpha, \beta) \cdot u;$$

applying this to the formula above we deduce that

$$\alpha_* (c\mathcal{G}_{0,1/mN'}) = \prod_{x,y \in \mathbb{Z}/a\mathbb{Z}} (c\mathcal{G}_{x/a,1/mN'+y/a}).$$

The latter expression is equal to the product of $c\mathcal{G}_{\gamma,\delta}$ over all pairs (γ, δ) such that $(a\gamma, a\delta) = (0, 1/mN)$; so using the distribution property of equation (1a), the product is $c\mathcal{G}_{0,1/mN}$ as required.

In the second case, where $N' = \ell N$ for $\ell \nmid MN$, we pass via the intermediate modular curves $Y(M, N(\ell))$ and $Y(M(\ell), N)$ described in [Kat04, §2.8]. Let $\varphi_\ell : Y(M, N(\ell)) \rightarrow Y(M(\ell), N)$ be the map defined in op. cit., corresponding to $z \mapsto \ell z$ on \mathcal{H} . We factor the projection α as $\alpha_1 \circ \alpha_2$, where α_1 and α_2 are the natural maps

$$Y(M, N\ell) \xrightarrow{\alpha_2} Y(M, N(\ell)) \xrightarrow{\alpha_1} Y(M, N).$$

By [Kat04, Step 2 of §2.13 and (2.13.2)], we have

$$\begin{aligned} (\alpha_2)_* (c\mathcal{G}_{0,1/N\ell}) &= \varphi_\ell^* (c\mathcal{G}_{0,1/N}) \cdot (c\mathcal{G}_{0,“\ell-1”/N})^{-1}, \\ (\alpha_1)_* \varphi_\ell^* (c\mathcal{G}_{0,1/N}) &= c\mathcal{G}_{0,1/N} \cdot (c\mathcal{G}_{0,“\ell-1”/N})^\ell, \\ (\alpha_1)_* (c\mathcal{G}_{0,“\ell-1”/N}) &= (c\mathcal{G}_{0,“\ell-1”/N})^{\ell+1} \end{aligned}$$

(the last formula owing to the fact that the degree of α_1 is $\ell + 1$). Hence, on combining these three equations, we obtain

$$\begin{aligned} \alpha_* (c\mathcal{G}_{0,1/N\ell}) &= (\alpha_1)_* (\alpha_2)_* (c\mathcal{G}_{0,1/N\ell}) \\ &= (\alpha_1)_* \left[\varphi_\ell^* (c\mathcal{G}_{0,1/N}) \cdot (c\mathcal{G}_{0,“\ell-1”/N})^{-1} \right] \\ &= \left(c\mathcal{G}_{0,1/N} \cdot (c\mathcal{G}_{0,“\ell-1”/N})^\ell \right) \cdot \left((c\mathcal{G}_{0,“\ell-1”/N})^{-1} \right)^{\ell+1} \\ &= c\mathcal{G}_{0,1/N} \cdot (c\mathcal{G}_{0,“\ell-1”/N})^{-1}. \quad \square \end{aligned}$$

Remark 2.2.5. In the above proposition we excluded from consideration the case when $N' = N\ell$ where $\ell \mid M$ (but $\ell \nmid N$). This case can also be treated using Kato’s methods, or deduced directly from Step 1 of §2.13 of op. cit. by applying the element $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, and one finds that in this case we have

$$\alpha_* (c\mathcal{G}_{0,1/N\ell}) = c\mathcal{G}_{0,1/N} \cdot (\varphi_\ell^{-1})^* (c\mathcal{G}_{0,“\ell-1”/N}).$$

However, we shall solely be working with modular curves of the form $Y(m, mN)$, so we will not need this formula.

2.3. *Integral models of modular curves.* The following theorem is well known:

THEOREM 2.3.1 (Igusa). *There exists a smooth scheme $\mathcal{Y}(N)$ over the ring $\mathbb{Z}[\zeta_N, 1/N]$, representing the functor of Definition 2.1.1 on the category of $\mathbb{Z}[1/N]$ -schemes.*

For a sketch of the proof, see, e.g., [DR73].

PROPOSITION 2.3.2. *The Siegel units ${}_c g_{\alpha,\beta}$, for all $(\alpha, \beta) \in (\frac{1}{N}\mathbb{Z}/\mathbb{Z})^2 - \{(0, 0)\}$, are elements of $\mathcal{O}(\mathcal{Y}(N))^\times$.*

Proof. As shown in [Kat04, Prop 1.3], given an arbitrary scheme S , an elliptic curve E/S , and an integer $c > 1$ coprime to 6, there exists a canonical element ${}_c \theta_E \in \mathcal{O}(E - E[c])^\times$ whose divisor is $c^2(0) - E[c]$. As noted in [Sch98, §1.3], if the base S is integral and E has a torsion section $x : S \rightarrow E$ of order N , where $N > 1$ is coprime to c and either N is invertible on S or N has at least two prime factors, then $x^* {}_c \theta_E \in \mathcal{O}(S)^\times$. Applying this with $S = \mathcal{Y}(N)$, E the universal elliptic curve over S , and x the section $ae_1 + be_2$ where $(\alpha, \beta) = (a/N, b/N)$, we deduce that ${}_c g_{\alpha,\beta}$ extends from $Y(N)$ to a unit on the integral model $\mathcal{Y}(N)$. \square

Remark 2.3.3. By passage to the quotient we also see that for any $b \in \mathbb{Z}/N\mathbb{Z}$, $b \neq 0$, the Siegel unit ${}_c g_{0,b/N}$ is a unit on the canonical $\mathbb{Z}[1/N]$ -model $\mathcal{Y}_1(N)$ of $Y_1(N)$.

2.4. Hecke correspondences. We now recall how elements of the Hecke algebra can be interpreted as correspondences between modular curves or, equivalently, as 1-cycles on a product of two modular curves.

LEMMA 2.4.1. *Let $\alpha \in \mathrm{GL}_2^+(\mathbb{Q})$ and Γ_1, Γ_2 be finite-index subgroups of $\mathrm{SL}_2(\mathbb{Z})$. Then there is a unique morphism of varieties over $\overline{\mathbb{Q}}$,*

$$\sigma : Y(\Gamma_1 \cap \alpha^{-1}\Gamma_2\alpha) \rightarrow Y(\Gamma_1) \times Y(\Gamma_2),$$

such that the diagram

$$\begin{array}{ccc} \mathcal{H} & \xrightarrow{1 \times \alpha} & \mathcal{H} \times \mathcal{H} \\ \downarrow & & \downarrow \\ Y(\Gamma_1 \cap \alpha^{-1}\Gamma_2\alpha)(\mathbb{C}) & \xrightarrow{\sigma} & (Y(\Gamma_1) \times Y(\Gamma_2))(\mathbb{C}) \end{array}$$

commutes (where the vertical arrows are the natural projection maps). The image of σ is an irreducible closed subvariety of $Y(\Gamma_1) \times Y(\Gamma_2)$, and the map σ is a birational equivalence onto its image.

Proof. After Definition 2.1.10 and the remarks following, the only assertion that needs checking is that σ is birational. However, by Proposition A.1.4 in the appendix (applied to the subgroups Γ_1 and $\alpha^{-1}\Gamma_2\alpha$) we know that σ is injective away from a finite set. \square

Remark 2.4.2. This proposition is well known in the special case $\Gamma = \mathrm{SL}_2(\mathbb{Z})$ and $\alpha = \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}$ for a prime p , where it shows that $Y_0(p)$ is the normalization of the subvariety of \mathbb{A}^2 cut out by the classical modular equation of level p ; see, e.g., [DR73, §VI.6].

LEMMA 2.4.3. *Let Γ, Γ' be as above, let $\alpha_1, \alpha_2 \in \mathrm{GL}_2^+(\mathbb{Q})$, and for $i = 1, 2$, let C_i be the curve in $Y(\Gamma) \times Y(\Gamma')$ that is the image of points of the form $(z, \alpha_i z)$. If the double cosets $\Gamma' \alpha_1 \Gamma$ and $\Gamma' \alpha_2 \Gamma$ are distinct as subsets of $\mathrm{PGL}_2^+(\mathbb{Q})$, then $C_1 \cap C_2$ is a finite set.*

Proof. Suppose $P \in C_1 \cap C_2$. Then P admits liftings to $\mathcal{H} \times \mathcal{H}$ of the form $(z_1, \alpha_1 z_1)$ and $(z_2, \alpha_2 z_2)$; and since both of these points are preimages of P , we can find $\gamma \in \Gamma$ and $\gamma' \in \Gamma'$ such that $z_1 = \gamma z_2$ and $\alpha_1 z_1 = \gamma' \alpha_2 z_2$. Consequently, z_2 is fixed by the element

$$\gamma^{-1} \alpha_1^{-1} \gamma' \alpha_2 \in \Gamma \cdot \alpha_1^{-1} \alpha_2 \cdot (\alpha_2^{-1} \Gamma' \alpha_2).$$

By Lemma A.1.2, either $\gamma^{-1} \alpha_1^{-1} \gamma' \alpha_2$ is the identity in $\mathrm{PGL}_2^+(\mathbb{Q})$, in which case $\Gamma' \alpha_1 \Gamma$ and $\Gamma' \alpha_2 \Gamma$ have the same projective image; or z_2 lies in one of a finite set of orbits under the action of $\Gamma \cap \alpha_2^{-1} \Gamma' \alpha_2$, which implies that P lies in one of a finite set of points of C_2 , as required. \square

LEMMA 2.4.4. *Let $\Gamma_1, \Gamma_2 \subseteq \mathrm{SL}_2(\mathbb{Z})$, and let $\Gamma'_1 \subseteq \Gamma_1$ and $\Gamma'_2 \subseteq \Gamma_2$, with all four subgroups having finite index in $\mathrm{SL}_2(\mathbb{Z})$. Let $\alpha \in \mathrm{GL}_2^+(\mathbb{Q})$, and suppose $\beta_1, \dots, \beta_h \in \mathrm{GL}_2^+(\mathbb{Q})$ are such that we have*

$$\Gamma_2 \alpha \Gamma_1 = \bigsqcup_{i=1}^h \Gamma'_2 \beta_i \Gamma'_1.$$

Let C be the curve in $Y(\Gamma_1) \times Y(\Gamma_2)$ that is the image of $Y(\Gamma_1 \cap \alpha^{-1} \Gamma_2 \alpha)$ under the map σ of Lemma 2.4.1. Then the preimage of C in $Y(\Gamma'_1) \times Y(\Gamma'_2)$ is the union of h distinct curves D_1, \dots, D_h , where D_i is the image of the map

$$\begin{aligned} \sigma_i : Y(\Gamma'_1 \cap \beta_i^{-1} \Gamma'_2 \beta_i) &\longrightarrow Y(\Gamma'_1) \times Y(\Gamma'_2) \\ z &\longmapsto (z, \beta_i z). \end{aligned}$$

Moreover, if for each i we choose some $\gamma_i \in \Gamma_1$ such that $\beta_i \in \Gamma_2 \alpha \gamma_i$, then we have a commutative diagram

$$(2) \quad \begin{array}{ccc} Y(\Gamma'_1 \cap \beta_i^{-1} \Gamma'_2 \beta_i) & \xrightarrow{\sigma_i} & Y(\Gamma'_1) \times Y(\Gamma'_2) \\ \downarrow z \mapsto \gamma_i z & & \downarrow \\ Y(\Gamma_1 \cap \alpha_i^{-1} \Gamma_2 \alpha_i) & \xrightarrow{\sigma} & Y(\Gamma_1) \times Y(\Gamma_2), \end{array}$$

where the right-hand vertical arrow is the natural projection map.

Proof. The definition of γ_i implies that diagram (2) commutes, from which it is clear that D_i is a lifting of C . By Lemma 2.4.3, the D_i are distinct.

It remains only to check that the union of the D_i exhausts the preimage of C . Let $P \in C$, and let \tilde{P} be any lifting of P to $\mathcal{H} \times \mathcal{H}$. Then we have $P = (\gamma_1 z, \gamma_2 \alpha z)$ for some $\gamma_1 \in \Gamma_1$ and $\gamma_2 \in \Gamma_2$; so $P = (w, \gamma_2 \alpha \gamma_1^{-1} w)$, where $w = \gamma_1 z \in \mathcal{H}$. We have $\gamma_2 \alpha \gamma_1^{-1} \in \Gamma'_2 \beta_i \Gamma'_1$ for some $i \in \{1, \dots, h\}$ so, in particular, the image of \tilde{P} in $Y(\Gamma'_1) \times Y(\Gamma'_2)$ lies in D_i as required. \square

LEMMA 2.4.5. *Let Γ be a finite-index subgroup of $SL_2(\mathbb{Z})$, and let Γ_1, Γ_2 be finite-index subgroups of Γ such that $\Gamma_1 \Gamma_2 = \Gamma$. Then, in the diagram of modular curves*

$$\begin{array}{ccc} Y(\Gamma_1 \cap \Gamma_2) & \xrightarrow{\alpha} & Y(\Gamma_1) \\ \beta \downarrow & & \downarrow \gamma \\ Y(\Gamma_2) & \xrightarrow{\delta} & Y(\Gamma), \end{array}$$

where $\alpha, \beta, \gamma, \delta$ are the natural projection maps, the two maps $\mathcal{O}(Y(\Gamma_1))^\times \rightarrow \mathcal{O}(Y(\Gamma_2))^\times$ given by $\beta_* \circ \alpha^*$ and $\delta^* \circ \gamma_*$ coincide, and similarly the maps $\mathcal{O}(Y(\Gamma_2))^\times \rightarrow \mathcal{O}(Y(\Gamma_1))^\times$ given by $\alpha_* \circ \beta^*$ and $\gamma^* \circ \delta_*$ coincide.

Proof. Note that the hypotheses are symmetric in Γ_1 and Γ_2 , so it suffices to show that $\beta_* \circ \alpha^* = \delta^* \circ \gamma_*$. Moreover, since all of the morphisms in the diagram are surjective, the corresponding pullback morphisms are injective, so it suffices to show that

$$\beta^* \circ \beta_* \circ \alpha^* = \beta^* \circ \delta^* \circ \gamma_*.$$

Since the diagram commutes, this is equivalent to

$$(\beta^* \circ \beta_*) \circ \alpha^* = \alpha^* \circ (\gamma^* \circ \gamma_*).$$

However, the map $(\beta^* \circ \beta_*)$ is given by the product over translates by coset representatives for $(\Gamma_1 \cap \Gamma_2) \backslash \Gamma_2$, and the map $(\gamma^* \circ \gamma_*)$ is given by the product over coset representatives for $\Gamma_1 \backslash \Gamma$. However, since $\Gamma_1 \Gamma_2 = \Gamma$, the natural map

$$(\Gamma_1 \cap \Gamma_2) \backslash \Gamma_2 \rightarrow \Gamma_1 \backslash \Gamma$$

is surjective. Thus these two quotients admit a common set of coset representatives, so the two maps coincide. \square

Remark 2.4.6. One can interpret this more “categorically” as follows: our hypotheses imply that the diagram in the statement of the lemma is Cartesian (in the category of curves and dominant rational maps), so $Y(\Gamma_1 \cap \Gamma_2)$ is birational to the fibre product of $Y(\Gamma_1)$ and $Y(\Gamma_2)$ over $Y(\Gamma)$. The symmetry of pushforward and pullback is then a general property of fibre products.

2.5. *Motivic cohomology, higher Chow groups and the Gersten complex.*

We now recall the definition of the higher Chow group $\mathrm{CH}^2(X, 1)$ of a variety X and how it may be explicitly calculated using the Gersten complex. In this section k may be any field of characteristic 0. Let $\mathrm{Var}(k)$ be the category of varieties over k , by which we mean separated schemes of finite type over k . Let $\mathrm{Sm}(k)$ be the full subcategory of smooth varieties. Let $A = \mathbb{Q}$ or \mathbb{Z} be the coefficient ring.

Definition 2.5.1 (Voevodsky; cf. [MVW06, Def. 3.4]). Let $X \in \mathrm{Sm}(k)$, and $p, q \in \mathbb{Z}$ with $q \geq 0$. Define the *motivic cohomology* of X to be

$$H_{\mathcal{M}}^p(X, A(q)) = \mathbb{H}_{\mathrm{Zar}}^p(X, \mathbb{Z}(q) \otimes A),$$

where $\mathbb{Z}(q)$ denotes Voevodsky’s motivic complex of sheaves on X and $\mathbb{H}_{\mathrm{Zar}}^p$ denotes hypercohomology (with respect to the Zariski topology).

Remark 2.5.2.

- (1) We use a slightly different notation than Voevodsky; the notation used in op. cit. is $H^{i,j}(X, A)$. Our choice of notation follows [Hub00] and [Lev04].
- (2) Note that $H_{\mathcal{M}}^p(X, A(q))$ is zero for $p > \inf(2q, q + \dim X)$. It is *not* known to be zero for $p < 0$ since the motivic complex is not bounded below.

We shall not use the definition of motivic cohomology directly; we shall rather use the fact that these groups are isomorphic to Bloch’s higher Chow groups:

THEOREM 2.5.3. *For any $X \in \mathrm{Sm}(k)$ and any $p, q \geq 0$, there is a natural isomorphism*

$$H^p(X, \mathbb{Z}(q)) \cong \mathrm{CH}^q(X, 2q - p).$$

Here, the higher Chow groups are those defined by Bloch.

Proof. See [Voe02, Cor. 2] or [Lev04, Th. 1.2]. □

We also have an alternative description of these groups in terms of Quillen K -theory. We will actually be interested in the special case when $p = 3$ and $q = 2$. Here, we use a result of Landsburg [Lan91]. For X smooth over a field, $m \geq 0$ and $0 \leq p \leq m$, he constructs a map

$$\Psi_{m,p} : \mathrm{CH}^m(X, m - p) \longrightarrow H^p(X, \mathcal{K}_m),$$

where \mathcal{K}_m is the sheafification of $U \mapsto K_m(U)$ on X . Here, K_m denotes the m -th Quillen K -group.

THEOREM 2.5.4. *The map $\Psi_{m,p}$ is an isomorphism for $p = m - 1$.*

Proof. See [Lan91, Th. 2.5]. □

Remark 2.5.5. For $p < m - 1$, the map $\Psi_{m,p}$ may not be an isomorphism in general. As pointed out to us by Landsburg in a discussion on <http://mathoverflow.net/>, if $X = \text{Spec}(k)$, then $\text{CH}^m(X, m)$ is the Milnor K -group $K_m^M(k)$ (by a theorem of Nesterenko–Suslin) and the map $\Psi_{m,0} : K_m^M(k) \rightarrow H^0(X, \mathcal{K}_m) = K_m(k)$ is the natural map from Milnor to Quillen K -theory, which is not generally an isomorphism for $m > 2$.

Finally, we address the question of how to explicitly describe elements of these groups.

PROPOSITION 2.5.6. *Suppose that X is a smooth variety of finite type over a field k . Then there is a resolution of the sheaf \mathcal{K}_m*

$$0 \longrightarrow \mathcal{K}_m \longrightarrow \coprod_{x \in X^0} (i_x)_* K_m(k(x)) \longrightarrow \coprod_{x \in X^1} (i_x)_* K_{m-1}(k(x)) \longrightarrow \cdots .$$

Proof. See [Qui73]. □

COROLLARY 2.5.7. *The group $H^1(X, \mathcal{K}_2)$ is the first homology group of the “Gersten complex”*

$$(3) \quad \text{Gerst}_2(X) : \coprod_{x \in X^0} K_2(k(x)) \xrightarrow{d^0} \coprod_{x \in X^1} k(x)^\times \xrightarrow{d^1} \coprod_{x \in X^2} \mathbb{Z},$$

where d^0 is the tame symbol map and d^1 maps a function to its divisor (cf. [Fla92, §2]).

Combining the above results, we get the following statement:

PROPOSITION 2.5.8. *Assume that X is a smooth variety of finite type over a field k . Then we have isomorphisms*

$$H^1(\text{Gerst}_2(X)) \cong H^1(X, \mathcal{K}_2) \cong \text{CH}^2(X, 1) \cong H_{\mathcal{M}}^3(X, \mathbb{Z}(2)).$$

We shall use these to identify $\text{CH}^2(X, 1)$ with $H^1(\text{Gerst}_2(X))$; it is the latter group in which we shall actually construct elements.

Notation. We shall write $Z^2(X, 1)$ to denote the kernel of the boundary map d^1 in the Gersten complex $\text{Gerst}_2(X)$, so

$$Z^2(X, 1) = \left\{ \sum_i (C_i, \phi_i) : C_i \in X^1, \phi_i \in k(C_i)^\times, \sum_i \text{div}(\phi_i) = 0 \right\}.$$

This is a slight abuse of notation since in Bloch’s theory of higher Chow groups, $Z^2(X, 1)$ is used to denote something slightly different (a certain subgroup of the codimension 2 cycles on $X \times \mathbb{A}^1$); but we shall not use Bloch’s construction directly in this paper, so this abuse should cause no confusion.

Remark 2.5.9. We shall, in fact, construct a “Euler system” in the groups $Z^2(X, 1)$ as X varies over a family of modular surfaces; that is, our compatibility properties will hold at the level of cycles, rather than just after quotienting out by the image of tame symbols. The groups $Z^2(X, 1)$ are much easier to work with, as they have good descent properties: for a finite surjective map $X \rightarrow Y$, the pullback $Z^2(Y, 1) \rightarrow Z^2(X, 1)$ is injective.

This is, in a sense, analogous to the fact that in the construction of [Kat04] the compatibility properties of the Euler system in K_2 of modular curves are proved at the level of $K_1 \otimes K_1$, before quotienting by elements of the form $x \otimes (1 - x)$.

2.6. *Zeta elements on $Y(m, mN)$.* We begin by defining elements of the group $Z^2(Y(m, mN)^2, 1)$, which we shall call *zeta elements*.

Definition 2.6.1. For $m, N \geq 1$, the curve $\mathcal{C}_{m,N,j} \subseteq Y(m, mN)^2$ is defined as the subvariety

$$\left(u, v : v = \begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix} u \right).$$

For $c > 1$ coprime to $6mN$, we define

$${}_c\mathcal{Z}_{m,N,j} = (\mathcal{C}_{m,N,j}, \phi) \in Z^2(Y(m, mN)^2, 1),$$

where $\phi \in \mathcal{O}(C)^\times$ is the pullback of ${}_c g_{0,1/mN}$ along either of the projections $\mathcal{C}_{m,N,j} \rightarrow Y(m, mN)$.

The first properties of these elements are the following:

PROPOSITION 2.6.2. *The elements ${}_c\mathcal{Z}_{m,N,j}$ have the following properties:*

- (1) *We have $\rho^* {}_c\mathcal{Z}_{m,N,j} = {}_c\mathcal{Z}_{m,N,-j}$, where ρ is the involution of $Y(m, mN)^2$ that interchanges the factors.*
- (2) *For $c, d > 1$ coprime to $6mN$, the element*

$$\left[d^2 - \left(\begin{pmatrix} d & 0 \\ 0 & d \end{pmatrix}, \begin{pmatrix} d & 0 \\ 0 & d \end{pmatrix} \right)^* \right] \cdot {}_c\mathcal{Z}_{m,N,j}$$

is symmetric in c and d . In particular, there exists a unique element

$$\mathcal{Z}_{m,N,j} \in Z^2(Y(m, mN)^2, 1) \otimes \mathbb{Q}$$

such that ${}_c\mathcal{Z}_{m,N,j} = [c^2 - \left(\begin{pmatrix} c & 0 \\ 0 & c \end{pmatrix}, \begin{pmatrix} c & 0 \\ 0 & c \end{pmatrix} \right)^] \mathcal{Z}_{m,N,j}$ for any c .*

- (3) *We have*

$$\left(\begin{pmatrix} b & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} b & 0 \\ 0 & 1 \end{pmatrix} \right)^* {}_c\mathcal{Z}_{m,N,j} = {}_c\mathcal{Z}_{m,N,b^{-1}j}$$

for any $b \in (\mathbb{Z}/mN\mathbb{Z})^\times$.

Proof. Part (1) is obvious, and part (2) follows immediately from the fact that the Siegel units ${}_c g_{\alpha,\beta}$ satisfy

$$(d^2 {}_c g_{\alpha,\beta} - {}_c g_{d\alpha,d\beta}) = (c^2 d g_{\alpha,\beta} - d g_{c\alpha,c\beta})$$

for any $\alpha, \beta \in \frac{1}{N}\mathbb{Z}/\mathbb{Z} - \{(0, 0)\}$ and any $c, d > 1$ coprime to $6mN$ (cf. [Kat04, Prop. 1.3(2)]). We may then define $\mathcal{Z}_{m,N,j} = (c^2 - 1)^{-1} {}_c \mathcal{Z}_{m,N,j}$ for any $c > 1$ congruent to 1 modulo mN .

Property (3) follows from the identity $\begin{pmatrix} b & 0 \\ 0 & 1 \end{pmatrix}^{-1} \begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & b^{-1}j \\ 0 & 1 \end{pmatrix} \begin{pmatrix} b & 0 \\ 0 & 1 \end{pmatrix}^{-1}$. □

2.7. Generalized Beilinson–Flach elements. The Beilinson–Flach elements of [BDR12] are elements of $\text{CH}^2(Y_1(N)^2, 1)$ defined as (Δ, ϕ) , where Δ is the diagonal and ϕ is a suitable modular unit on Δ . Our generalization of this is motivated by the observation that one can recover the twists of a modular form by Dirichlet characters modulo m from the “shifted” forms $f(x + a/m)$ for $a \in (\mathbb{Z}/m\mathbb{Z})^\times$; this is also the idea underlying the construction of the p -adic L -function of a single modular form using modular symbols.

LEMMA 2.7.1. *Let $m, N \geq 1$ with $m^2N \geq 5$, and let $j \in \mathbb{Z}$. Then there is a unique morphism of algebraic varieties over \mathbb{C} ,*

$$\kappa_j : Y_1(m^2N) \otimes \mathbb{C} \rightarrow Y_1(N) \otimes \mathbb{C},$$

such that the diagram of morphisms of complex-analytic manifolds

$$\begin{array}{ccc} \mathcal{H} & \xrightarrow{z \mapsto z + j/m} & \mathcal{H} \\ \downarrow & & \downarrow \\ Y_1(m^2N)(\mathbb{C}) & \xrightarrow{\kappa_j} & Y_1(N)(\mathbb{C}) \end{array}$$

commutes. The morphism κ_j is defined over $\mathbb{Q}(\mu_m)$ and depends only on the residue class of $j \pmod m$.

Proof. The existence of such a map at the level of quotients of \mathcal{H} follows immediately from the inclusion of matrix groups

$$\begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix} \Gamma_1(m^2N) \begin{pmatrix} 1 & -j \\ 0 & 1 \end{pmatrix} \subseteq \Gamma_1(N).$$

However, in order to descend to an algebraic morphism over $\mathbb{Q}(\mu_m)$ we use the canonical models above.

We first consider the map $Y(m^2N) \rightarrow Y(m, mN)$ which maps (E, e_1, e_2) to $(E/\langle me_2 \rangle, [mNe_1], [e_2])$. This factors through the quotient by the subgroup

$\begin{pmatrix} u & * \\ 0 & 1 \end{pmatrix} : u = 1 \pmod{m\mathbb{Z}}$, which we have identified with $Y_1(m^2N) \otimes \mathbb{Q}(\mu_m)$. This map is compatible with $z \mapsto mz$ on \mathcal{H} . We now consider the composition

$$Y_1(m^2N) \times \mu_m \longrightarrow Y(m, mN) \xrightarrow{\begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix}} Y(m, mN) \xrightarrow{t_m} Y_1(N) \times \mu_m,$$

where t_m is as in Definition 2.1.6. All three morphisms are maps of $\mathbb{Q}(\mu_m)$ -varieties (i.e., they commute with the projection maps to μ_m); and on the fibre over $\zeta_m \in \mu_m(\mathbb{C})$ they correspond to $z \mapsto mz$, $z \mapsto z + j$, and $z \mapsto z/m$, so the composition corresponds to $z \mapsto z + j/m$. \square

Definition 2.7.2. For m, N, j as above, let $\iota_{m,N,j}$ be the map

$$(1, \kappa_j) : Y_1(m^2N) \times \mu_m \rightarrow Y_1(N)^2 \times \mu_m,$$

and let $C_{m,N,j}$ be the irreducible curve in $Y_1(N)^2$ that is the image of $\iota_{m,N,j}$.

We now use these curves $C_{m,N,j}$ to define a class in $\text{CH}^2(Y_1(N)^2 \times \mu_m, 1)$, using the presentation of the latter group given by the Gersten complex.

Definition 2.7.3. Let $N \geq 5$, $m \geq 1$, $j \in \mathbb{Z}/m\mathbb{Z}$ as above. Let $c \geq 1$ be coprime to $6mN$, and let $\alpha \in \mathbb{Z}/m^2N\mathbb{Z}$. We define the *generalized Beilinson–Flach element*

$${}_c\Xi_{m,N,j,\alpha} \in \text{CH}^2(Y_1(N)^2 \otimes \mathbb{Q}(\mu_m), 1)$$

as the class of the pair

$$(C_{m,N,j}, (\iota_{m,N,j})^*(c\mathfrak{g}_{0,\alpha/m^2N})) \in Z^2(Y_1(N)^2 \times \mu_m, 1).$$

When $\alpha = 1$, we drop it from the notation and write simply ${}_c\Xi_{m,N,j}$.

The following proposition shows that these zeta elements are simply the “ Y_1 -versions” of those defined in the previous section:

PROPOSITION 2.7.4. *The generalized Beilinson–Flach element ${}_c\Xi_{m,N,j,\alpha}$ is the pushforward of the element*

$$\left(\begin{pmatrix} \alpha & 0 \\ 0 & \alpha \end{pmatrix}, \begin{pmatrix} \alpha & 0 \\ 0 & \alpha \end{pmatrix} \right)^* {}_c\mathcal{Z}_{m,N,j} \in Z^2(Y(m, mN)^2, 1)$$

along the map $t_m \times t_m : Y(m, mN)^2 \rightarrow Y_1(N)^2 \times \mu_m$ introduced in Section 2.1.

Proof. It is clear from the construction of the map $\kappa_{m,N,j}$ that $C_{m,N,j}$ is the image of $\mathcal{C}_{m,N,j}$ under $t_m \times t_m$. So it suffices to show that the pushforward of $c\mathfrak{g}_{0,1/m^2N}$ from $Y_1(m^2N) \otimes \mathbb{Q}(\mu_m)$ to $Y(m, mN)$ along the map constructed above is $c\mathfrak{g}_{0,1/mN}$.

Let U be the subgroup of $\text{GL}_2(\mathbb{Z}/m^2N\mathbb{Z})$ consisting of elements $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ that satisfy $c = 0, d = 1 \pmod{m^2N}$ and $a = 1 \pmod{m}$ (and b arbitrary). This is clearly contained in the subgroup U' of elements satisfying $a = 1 \pmod{m}$, $c = 0 \pmod{m^2N}$ and $d = 1 \pmod{mN}$, and a set of coset representatives for

U/U' is given by the matrices

$$\left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 + mNt \end{pmatrix} : 0 \leq t < m \right\}.$$

Hence the pushforward of $c\mathcal{G}_{0,1/m^2N}$ from $U \backslash Y(m^2N)$ to $U' \backslash Y(m^2N)$ is given by

$$\prod_{0 \leq t < m} \begin{pmatrix} 1 & 0 \\ 0 & 1 + mNt \end{pmatrix}^* c\mathcal{G}_{0,1/m^2N} = \prod_{0 \leq t < m} c\mathcal{G}_{0,1/m^2N+t/m}.$$

By Proposition 2.2.2(2), this is equal to $\varphi_m^*(c\mathcal{G}_{0,1/mN})$. However, conjugation by $\begin{pmatrix} m & 0 \\ 0 & 1 \end{pmatrix}$ sends U' to the subgroup

$$U'' = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : \begin{array}{l} a = 1, b = 0 \pmod{m}, \\ c = 0, d = 1 \pmod{mN} \end{array} \right\},$$

and we have $U'' \backslash Y(m^2N) = Y(m, mN)$. □

We now record some properties of the generalized Beilinson–Flach elements.

PROPOSITION 2.7.5. *The elements above have the following properties:*

- (1) *The element $c\Xi_{m,N,j,\alpha}$ only depends on the congruence class of α modulo mN (not m^2N).*
- (2) *The involution of $Y_1(N)^2 \otimes \mathbb{Q}(\mu_m)$ given by switching the two factors interchanges $c\Xi_{m,N,j}$ and $c\Xi_{m,N,-j}$.*
- (3) *For $q \in (\mathbb{Z}/m\mathbb{Z})^\times$, we have $\sigma_q^*(c\Xi_{m,N,j,\alpha}) = c\Xi_{m,N,q^{-1}j,\alpha}$, where $\sigma_q \in \text{Gal}(\mathbb{Q}(\mu_m)/\mathbb{Q})$ is the arithmetic Frobenius at q .*
- (4) *For any $r \in (\mathbb{Z}/mN\mathbb{Z})^\times$, we have*

$$c\Xi_{m,N,j,r\alpha} = \langle d \times d \rangle^* c\Xi_{m,N,k,\alpha},$$

where $k = r^{-2}j \in \mathbb{Z}/m\mathbb{Z}$, d is the image of r in $(\mathbb{Z}/N\mathbb{Z})^\times$, and $\langle d \times d \rangle$ denotes the action on $Y_1(N)^2 \otimes \mathbb{Q}(\mu_m)$ of the element

$$\begin{pmatrix} d^{-1} & 0 \\ 0 & d \end{pmatrix} \times \begin{pmatrix} d^{-1} & 0 \\ 0 & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z}/N\mathbb{Z})^2.$$

- (5) *For c, d coprime to $6mN$, the expression*

$$d^2 c\Xi_{m,N,j,\alpha} - c\Xi_{m,N,j,d\alpha}$$

is symmetric in c and d . In particular, there exist well-defined elements $\Xi_{m,N,j,\alpha} \in \text{CH}^2(Y_1(N)^2 \otimes \mathbb{Q}(\mu_m), 1) \otimes \mathbb{Q}$ such that we have

$$c\Xi_{m,N,j,\alpha} = c^2\Xi_{m,N,j,\alpha} - \Xi_{m,N,j,c\alpha} = (c^2 - \langle c \times c \rangle^* \sigma_c^2) \Xi_{m,N,j,\alpha}.$$

Proof. After Proposition 2.7.4, parts (1) and (2) are immediate. The remaining statements follow from Proposition 2.6.2, together with the fact that t_m intertwines the action of $\begin{pmatrix} d & 0 \\ 0 & 1 \end{pmatrix}$ on $Y(m, mN)$ with the arithmetic Frobenius σ_d on $Y_1(N) \times \mu_m$ (Proposition 2.1.7(2)). \square

2.8. *Cuspidal components.* In the preceding sections we have constructed elements of the higher Chow groups of affine surfaces. In order to be able to apply results on regulator maps, it is convenient to have elements of Chow groups of projective surfaces instead. We shall show that this can be achieved, but not in a canonical way, and only at the cost of tensoring with \mathbb{Q} .

THEOREM 2.8.1. *Let N, m, j be as in Definition 2.7.3. Then the element $\Xi_{m,N,j}$ of $\text{CH}^2(Y_1(N)^2 \otimes \mathbb{Q}(\mu_m), 1) \otimes \mathbb{Q}$ is in the image of the pullback map*

$$\text{CH}^2(X_1(N)^2 \otimes \mathbb{Q}(\mu_m), 1) \otimes \mathbb{Q} \rightarrow \text{CH}^2(Y_1(N)^2 \otimes \mathbb{Q}(\mu_m), 1) \otimes \mathbb{Q}$$

induced by the open embedding $Y_1(N) \hookrightarrow X_1(N)$.

We will actually prove a slightly more precise statement; see Proposition 2.8.5 below.

Recall that we constructed $c\Xi_{m,N,j}$ as the class in $\text{CH}^2(Y_1(N)^2 \otimes \mathbb{Q}(\mu_m), 1)$ of an explicit element of $Z^2(Y_1(N)^2 \otimes \mathbb{Q}(\mu_m), 1)$, which we shall temporarily denote by $c\mathfrak{Y}_{m,N,j}$. It is clear that we may also regard $c\mathfrak{Y}_{m,N,j}$ as an element of $\text{Gerst}_2^1(X_1(N)^2 \otimes \mathbb{Q}(\mu_m))$, whose divisor is not necessarily trivial, but is supported on the cuspidal locus.

We will need a preparatory lemma. Let K be any number field.

Definition 2.8.2. We shall call an element of $\text{Gerst}_2^1(X_1(N)^2 \otimes K)$ *negligible* if it is supported on a finite union of curves of the form $\{c\} \times X_1(N)$ or $X_1(N) \times \{d\}$ for points $c, d \in X_1(N) \setminus Y_1(N)$.

Remark 2.8.3. Here by “point” we mean a 0-dimensional point of the K -scheme $X_1(N) \setminus Y_1(N)$, i.e., a $\text{Gal}(\bar{K}/K)$ -orbit of points in the naive sense. Note that this is slightly more restrictive than the definition of “negligible” in [BDR12].

Before proving the theorem, we will need the following preparatory lemma:

LEMMA 2.8.4. *Let K be any number field, and let u, v, x, y be cuspidal points of $X_1(N) \otimes K$. Then there exists a negligible element in $\text{Gerst}_2^1(X_1(N)^2 \otimes K) \otimes \mathbb{Q}$ with divisor $(u, v) - (x, y)$.*

Proof. By the Manin–Drinfeld theorem [Dri73], there exist elements $f, g \in \mathcal{O}(Y_1(N) \otimes K)^\times \otimes \mathbb{Q}$ whose divisors are $v - y$ and $u - x$, respectively. The the element

$$(\{u\} \times X_1(N), f) + (X_1(N) \times \{y\}, g)$$

has the required property. \square

We can now prove the following proposition:

PROPOSITION 2.8.5. *Let N, m, j, c be as in Definition 2.7.3. Then there exist an integer $r \geq 1$ and a negligible element Θ such that*

$$R \cdot {}_c\mathfrak{Y}_{m,N,j} + \Theta \in Z^1(\text{Gerst}_2(X_1(N)^2 \otimes \mathbb{Q}(\mu_m))).$$

Proof. Recall that $\iota_{m,N,j}$ is the map

$$\iota_{m,N,j} = (1, \kappa_j) : Y_1(m^2N) \rightarrow Y_1(N)^2,$$

where κ_j is induced from the map $\mathcal{H} \rightarrow \mathcal{H}$ given by $z \mapsto z + \frac{j}{m}$. It follows that, if we regard ${}_c\mathfrak{Y}_{m,N,j}$ as an element of $Z^2(X_1(N)^2 \otimes \mathbb{Q}(\mu_m), 1)$, then $\text{Div}({}_c\mathfrak{Y}_{m,N,j})$ is a linear combination of divisors of the form $(c_1, c_1 + \frac{j}{m}) - (c_2, c_2 + \frac{j}{m})$. But Lemma 2.8.4 implies that there exists a negligible element $\Theta \in \text{Gerst}_2^1(X_1(N) \otimes \mathbb{Q}(\mu_m))$ such that $\text{Div}(\Theta) = \text{Div}({}_c\mathfrak{Y}_{m,N,j})$. Then the element

$${}_c\mathfrak{X}_{m,N,j} := {}_c\mathfrak{Y}_{m,N,j} - \Theta \in Z^2(X_1(N)^2 \otimes \mathbb{Q}(\mu_m), 1) \otimes \mathbb{Q}$$

has the required properties. □

This clearly implies Theorem 2.8.1.

Remark 2.8.6.

- (1) Note that the negligible element Θ is not uniquely determined. However, as we will see below, this will not matter for the evaluation of the element via the Beilinson or the syntomic regulator.
- (2) Since $X_1(N)^2$ and $Y_1(N)^2$ have the same rational function field, any element of $\text{CH}^2(X_1(N)^2 \otimes \mathbb{Q}(\mu_m), 1) \otimes \mathbb{Q}$ lifting ${}_c\Xi_{m,N,j}$ is necessarily the class of an element of $Z^2(X_1(N)^2 \otimes \mathbb{Q}(\mu_m), 1) \otimes \mathbb{Q}$ differing from ${}_c\mathfrak{Y}_{m,N,j}$ by a negligible element.
- (3) Since there are only finitely many cusps on $X_1(N)$, the constant R may be chosen to be independent of c, m and j , although it may of course depend on N .

2.9. *Zeta elements versus generalized Beilinson–Flach elements.* At the referee’s request, we shall briefly clarify the relations between the two classes of elements we have introduced (the zeta elements ${}_c\mathcal{Z}_{m,N,j}$ and the generalized Beilinson–Flach elements ${}_c\Xi_{m,N,j}$) and how they would relate to a hypothetical “optimal” construction. Recall that the element ${}_c\mathcal{Z}_{m,N,j}$ lies in the group $\text{CH}^2(Y(m, mN)^2, 1)$, and the element ${}_c\Xi_{m,N,j} \in \text{CH}^2(Y_1(N)^2 \otimes \mathbb{Q}(\mu_m), 1)$ is the pushforward of ${}_c\mathcal{Z}_{m,N,j}$ via the morphism $t_m \times t_m$ of Section 2.1. It is the elements ${}_c\Xi_{m,N,j}$ that will be used in Sections 4–7 of this paper in order to bound Selmer groups.

One reason for introducing the elements ${}_c\mathcal{Z}_{m,N,j}$ is that they are somewhat easier to work with than the ${}_c\Xi_{m,N,j}$. In the next section we shall prove

norm-compatibility relations for the ${}_c\mathcal{Z}_{m,N,j}$ and deduce norm relations for the ${}_c\Xi_{m,N,j}$ as a consequence; given the somewhat opaque map $\iota_{m,N,j}$ entering into the definition of the elements ${}_c\Xi_{m,N,j}$, it seems unlikely that these norm relations could be proved without the introduction of some auxilliary higher-level modular curve.

A second reason to consider the elements ${}_c\mathcal{Z}_{m,N,j}$ is the following optimistic idea: Let us fix a prime p , and a level N coprime to p , and let us consider the curves $Y(p^r, Np^r)$ for $r \geq 0$, and their self-products $Y(p^r, Np^r)^2$. These form a tower of surfaces with Galois group $\mathrm{GL}_2(\mathbb{Z}_p) \times_{\det} \mathrm{GL}_2(\mathbb{Z}_p)$. Let us imagine that we could construct a norm-compatible family of elements in the higher Chow groups of this tower, analogous to the compatible family of elements in K_2 of the $\mathrm{GL}_2(\mathbb{Z}_p)$ -tower of modular curves constructed by Kato in [Kat04]. Then one could potentially perform a “non-abelian twisting” operation analogous to equation (8.4.3) of op. cit. in order to obtain classes in the cohomology groups attached to pairs of modular forms of arbitrary weights $k, \ell \geq 2$.

The elements ${}_c\mathcal{Z}_{p^r,N,j}$ represent our best attempt to realize this dream. They do indeed live on the surfaces $Y(p^r, Np^r)^2$; but the norm-compatibility relation they satisfy (Theorem 3.3.1) involves the “twisted” degeneracy map $\tau_p : Y(p^{r+1}, Np^{r+1}) \rightarrow Y(p^r, Np^r)$ of Definition 2.1.6, given by $z \mapsto z/p$ on the upper half-plane \mathcal{H} , rather than the natural one corresponding to the identity map on \mathcal{H} . The norm-compatibility relation also involves a Hecke operator at p , which does not appear in the setting of [Kat04]. Consequently, our methods will only allow us to construct cohomology classes for Rankin–Selberg convolutions of higher weight forms under additional ordinarity assumptions, when we can use Hida’s theory of p -adic families in order to pass from weight 2 to general weights.

3. Norm relations for generalized Beilinson–Flach elements

3.1. *The first norm relation: varying N .* We now consider the relation between the zeta elements at different levels N (for fixed m and j).

THEOREM 3.1.1 (First norm relation). *Let α be the natural projection $Y(m, mN') \rightarrow Y(m, mN)$, where N and N' are positive integers such that $N \mid N'$.*

- (1) *If $\mathrm{prime}(N') \subseteq \mathrm{prime}(mN)$, then the pushforward map*
 $(\alpha \times \alpha)_* : \mathrm{CH}^2(Y(m, mN')^2, 1) \rightarrow \mathrm{CH}^2(Y(m, mN)^2, 1)$

maps ${}_c\mathcal{Z}_{m,N',j}$ to ${}_c\mathcal{Z}_{m,N,j}$.

- (2) *If $N' = N\ell$, where $\ell \nmid mN$ is prime, then*

$$(\alpha \times \alpha)_*({}_c\mathcal{Z}_{m,N\ell,j}) = \left[1 - \left(\begin{pmatrix} \ell^{-1} & 0 \\ 0 & \ell^{-1} \end{pmatrix}, \begin{pmatrix} \ell^{-1} & 0 \\ 0 & \ell^{-1} \end{pmatrix} \right)^* \right] {}_c\mathcal{Z}_{m,N,j},$$

where $\begin{pmatrix} \ell^{-1} & 0 \\ 0 & \ell^{-1} \end{pmatrix}$ is considered as an element of $\mathrm{GL}_2(\mathbb{Z}/mN\mathbb{Z})$.

Proof. It is clear that the map α commutes with the action of $\begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix}$, so we have $(\alpha \times \alpha)(\mathcal{C}_{m,N',j}) = \mathcal{C}_{m,N,j}$; more precisely, we have a commutative diagram

$$\begin{array}{ccc} Y(m, mN') & \xrightarrow{\left(1, \begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix}\right)} & \mathcal{C}_{m,N',j} \\ \alpha \downarrow & & \downarrow \alpha \times \alpha \\ Y(m, mN) & \xrightarrow{\left(1, \begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix}\right)} & \mathcal{C}_{m,N,j}. \end{array}$$

From Theorem 2.2.4, we know that if $\text{prime}(mN') = \text{prime}(mN)$, then $\alpha_* (c\mathcal{G}_{0,1/mN'}) = c\mathcal{G}_{0,1/mN}$, so part (1) of the theorem follows. For part (2), we deduce from the second part of Theorem 2.2.4 that

$$(\alpha \times \alpha)_* (c\mathcal{Z}_{m,N',j}) = c\mathcal{Z}_{m,N,j} - c\mathcal{Z}_{m,N,j}^{\ell^{-1}},$$

where we write $c\mathcal{Z}_{m,N,j,a}$ for the element formed with $c\mathcal{G}_{0,a/mN}$ in place of $c\mathcal{G}_{0,1/mN}$. However, we have

$$c\mathcal{G}_{0,\ell^{-1}/mN} = \begin{pmatrix} \ell^{-1} & 0 \\ 0 & \ell^{-1} \end{pmatrix}^* c\mathcal{G}_{0,1/mN}$$

as elements of $\mathcal{O}(Y(m, mN))^\times$, and the action of $\begin{pmatrix} \ell^{-1} & 0 \\ 0 & \ell^{-1} \end{pmatrix}$ evidently commutes with that of $\begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix}$. □

We now deduce a compatibility relation for zeta elements on $Y_1(N)^2 \otimes \mathbb{Q}(\mu_m)$.

THEOREM 3.1.2 (First norm relation on $Y_1(N)$). *Let α be the natural projection $Y_1(N') \rightarrow Y_1(N)$, where N, N' are positive integers such that $N \mid N'$.*

If $\text{prime}(mN') = \text{prime}(mN)$, then we have

$$(\alpha \times \alpha)_* (c\Xi_{m,N',j}) = c\Xi_{m,N,j}.$$

If $N' = \ell N$ where $\ell \nmid mN$, then we have

$$(\alpha \times \alpha)_* (c\Xi_{m,N',j}) = \left[1 - (\langle \ell^{-1} \rangle, \langle \ell^{-1} \rangle)^* \sigma_\ell^{-2}\right] c\Xi_{m,N,j},$$

where σ_ℓ denotes the arithmetic Frobenius at ℓ .

Proof. This follows immediately from Theorem 3.1.1, since the map $\pi_{m,N} : Y(m, mN) \rightarrow Y_1(N) \times \mu_m$ intertwines $\begin{pmatrix} \ell^{-1} & 0 \\ 0 & \ell \end{pmatrix}$ with the diamond operator $\langle \ell \rangle$, and $\begin{pmatrix} \ell & 0 \\ 0 & 1 \end{pmatrix}$ with the Frobenius σ_ℓ . □

3.2. Hecke operators. We define Hecke operators, following [Kat04, §§2.9, 4.9]. Let ℓ be prime, and let $M, N \geq 1$. (We allow $\ell \mid M$ or $\ell \mid N$.) We define a

correspondence on $Y(M, N)$ as follows. We have a diagram of modular curves

$$\begin{array}{ccc}
 & Y(M(\ell), N) & \\
 & \downarrow \pi_1 \quad \searrow \pi_2 & \\
 Y(M, N) & & Y(M, N),
 \end{array}$$

where π_1 is the natural degeneracy map, corresponding to the identity on \mathcal{H} , and π_2 is the “twisted” degeneracy map, corresponding to $z \mapsto z/\ell$ on \mathcal{H} . (In the notation introduced in the proof of Theorem 2.2.4 above, π_1 was denoted pr_1 , and π_2 is the composite of $\varphi_\ell^{-1} : Y(M(\ell), N) \rightarrow Y(M, N(\ell))$ with the natural projection $Y(M, N(\ell)) \rightarrow Y(M, N)$.)

We denote the correspondence $(\pi_2)_*(\pi_1)^*$ by T'_ℓ if $\ell \nmid MN$ and by U'_ℓ if $\ell \mid MN$. We denote the operator $(\pi_1)_*(\pi_2)^*$ by T_ℓ (resp. U_ℓ); these latter operators T_ℓ, U_ℓ are the familiar Hecke operators of the transcendental theory, but it is the T'_ℓ, U'_ℓ that will concern us most here.

3.3. *The second norm relation for $\ell \mid N$.* Our goal in this section is to prove the following theorem:

THEOREM 3.3.1 (Second norm relation, $\ell \mid N$ case). *Let $m \geq 1, N \geq 5$, and let ℓ be a prime dividing N . Let τ_ℓ denote the degeneracy map*

$$Y(m\ell, m\ell N) \rightarrow Y(m, mN)$$

of Definition 2.1.6, compatible with $z \mapsto z/\ell$ on \mathcal{H} . Then for any $j \in (\mathbb{Z}/\ell m\mathbb{Z})^\times$, and $c > 1$ coprime to $6\ell mN$, we have

$$(\tau_\ell \times \tau_\ell)_*(c\mathcal{Z}_{\ell m, N, j}) = \begin{cases} (U'_\ell \times U'_\ell)(c\mathcal{Z}_{m, N, j}) & \text{if } \ell \mid m, \\ (U'_\ell \times U'_\ell - \Delta_\ell^*)(c\mathcal{Z}_{m, N, j}) & \text{if } \ell \nmid m, \end{cases}$$

where Δ_ℓ denotes the action of any element of $\text{GL}_2(\mathbb{Z}/mN\mathbb{Z})^2$ of the form $((\begin{smallmatrix} x & 0 \\ 0 & 1 \end{smallmatrix}), (\begin{smallmatrix} x & 0 \\ 0 & 1 \end{smallmatrix}))$ with $x = \ell \pmod m$.

We shall prove Theorem 3.3.1 below. First, we note that it implies the following property of the generalized Beilinson–Flach elements $c\Xi_{m, N, j}$ on $Y_1(N)$:

THEOREM 3.3.2 (Second norm relation on $Y_1(N)$, $\ell \mid N$ case). *Let $m \geq 1, N \geq 5, \ell$ a prime dividing $N, j \in (\mathbb{Z}/\ell m\mathbb{Z})^\times$, and $c \in (\mathbb{Z}/\ell mN\mathbb{Z})^\times$. Then we have*

$$\text{norm}_m^{\ell m}(c\Xi_{\ell m, N, j}) = \begin{cases} (U'_\ell \times U'_\ell)(c\Xi_{m, N, j}) & \text{if } \ell \mid m, \\ (U'_\ell \times U'_\ell - \sigma_\ell)(c\Xi_{m, N, j}) & \text{if } \ell \nmid m, \end{cases}$$

where $\text{norm}_m^{\ell m}$ denotes the Galois norm map, and σ_ℓ , for $\ell \nmid m$, denotes the arithmetic Frobenius at ℓ in $\text{Gal}(\mathbb{Q}(\mu_m)/\mathbb{Q})$.

Proof of Theorem 3.3.2 (assuming Theorem 3.3.1). Let

$$t_m \times t_m : Y(m, mN)^2 \rightarrow Y_1(N)^2 \otimes \mathbb{Q}(\mu_m)$$

be the map of Section 2.1. This map commutes with the actions of (U'_ℓ, U'_ℓ) on both sides and intertwines the action of Δ_ℓ with the arithmetic Frobenius σ_ℓ . Since ${}_c\Xi_{m,N,j} = (t_m \times t_m)_*({}_c\mathcal{Z}_{m,N,j})$ by Proposition 2.7.4, Theorem 3.3.2 follows from Theorem 3.3.1. \square

Proof of Theorem 3.3.1. Since we are assuming $\ell \mid N$, let us write $N' = N/\ell$. We have the following commutative diagram of modular curves:

$$\begin{array}{ccccc} Y(\ell m, \ell mN) & \xrightarrow{\alpha} & Y(\ell m, mN) & \xrightarrow{\text{pr}} & Y(m(\ell), mN) \\ & \searrow \tau_\ell & & \swarrow \pi_2 & \downarrow \pi_1 \\ & & Y(m, mN) & & Y(m, mN). \end{array}$$

Here α is the natural projection $Y(\ell m, \ell mN) \rightarrow Y(\ell m, \ell mN') = Y(\ell m, mN)$, and pr is the natural projection map. Consequently, we have a commutative diagram of surfaces

$$\begin{array}{ccccc} Y(\ell m, \ell mN)^2 & \xrightarrow{\alpha \times \alpha} & Y(\ell m, mN)^2 & \xrightarrow{\text{pr} \times \text{pr}} & Y(m(\ell), mN)^2 \\ & \searrow \tau_\ell \times \tau_\ell & & \swarrow \pi_2 \times \pi_2 & \downarrow \pi_1 \times \pi_1 \\ & & Y(m, mN)^2 & & Y(m, mN)^2. \end{array}$$

Applying Theorem 3.1.1, we see that $(\alpha \times \alpha)_*({}_c\mathcal{Z}_{\ell m,N,j}) = {}_c\mathcal{Z}_{\ell m,N',j}$. Since $Y(m(\ell), mN)^2$ is the quotient of $Y(\ell m, mN)^2$ by the subgroup

$$\left\{ \left(\begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix} \right) : x \in \mathbb{Z}/\ell m\mathbb{Z}, x = 1 \pmod m \right\},$$

we have

$$(\text{pr} \times \text{pr})^*(\text{pr} \times \text{pr})_*({}_c\mathcal{Z}_{\ell m,N',j}) = \sum_{\substack{x \in \mathbb{Z}/\ell m\mathbb{Z} \\ x=1 \pmod m}} {}_c\mathcal{Z}_{\ell m,N',xj}.$$

Let us now compute $(\text{pr} \times \text{pr})^*(\pi_1 \times \pi_1)_*({}_c\mathcal{Z}_{m,N,j})$. Since $Y(m, mN)^2$ is the quotient of $Y(\ell m, mN)^2$ by the group

$$\left\{ \left(\begin{pmatrix} x & y \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} x & z \\ 0 & 1 \end{pmatrix} \right) : \begin{array}{l} x, y, z \in \mathbb{Z}/\ell m\mathbb{Z}, x = 1 \pmod m, \\ y, z = 0 \pmod m \end{array} \right\},$$

we see that the preimage of $\mathcal{C}_{m,N,k}$ is the union of the curves $\mathcal{C}_{\ell m,N',k}$, for $k \in \mathbb{Z}/\ell m\mathbb{Z}$ congruent to j modulo m , each of which is isomorphic to $Y(\ell m, mN)$.

By counting degrees, they must be distinct. The modular units $c\mathcal{g}_{0,1/mN}$ and $c\mathcal{g}_{0,1/\ell mN'}$ coincide, and thus we have

$$(\mathrm{pr} \times \mathrm{pr})^*(\pi_1 \times \pi_1)^* c\mathcal{Z}_{m,N,j} = \sum_{\substack{k \in \mathbb{Z}/\ell m\mathbb{Z} \\ k \equiv j \pmod{m}}} c\mathcal{Z}_{\ell m,N',k}.$$

By hypothesis, j is invertible modulo ℓm . Thus if $\ell \mid m$, the sets $\{xj : x = 1 \pmod{m}\}$ and $\{k : k \equiv j \pmod{m}\}$ coincide, and since $(\mathrm{pr} \times \mathrm{pr})^*$ is clearly injective, we conclude that

$$(\mathrm{pr} \times \mathrm{pr})_* (c\mathcal{Z}_{\ell m,N',j}) = (\pi_1 \times \pi_1)^* c\mathcal{Z}_{m,N,j}.$$

Applying $(\pi_2 \times \pi_2)_*$ gives the result in this case.

If $\ell \nmid m$, there is exactly one lifting j_0 of j to $\mathbb{Z}/\ell m\mathbb{Z}$ that is not a unit. The matrix $\begin{pmatrix} 1 & j_0 \\ 0 & 1 \end{pmatrix}$ normalizes the subgroup of $\mathrm{GL}_2(\mathbb{Z}/mN\mathbb{Z})$ corresponding to $Y(m(\ell), mN)$ and thus defines a curve \mathcal{A} in $Y(m(\ell), mN)^2$, which is isomorphic to $Y(m(\ell), mN)$, consisting of points (u, v) with $v = \begin{pmatrix} 1 & j_0 \\ 0 & 1 \end{pmatrix} u$; and we have

$$(\pi_1 \times \pi_1)^* c\mathcal{Z}_{m,N,j} = (\mathrm{pr} \times \mathrm{pr})_* (c\mathcal{Z}_{\ell m,N',j}) + (\mathcal{A}, c\mathcal{g}_{0,1/mN}).$$

The image of \mathcal{A} under π_2 is $\mathcal{C}_{m,N,\ell^{-1}j_0}$; moreover, we have a diagram

$$\begin{array}{ccc} Y(m(\ell), mN) & \xrightarrow{\cong} & \mathcal{A} \\ \pi_2 \downarrow & & \pi_2 \times \pi_2 \downarrow \\ Y(m, mN) & \xrightarrow{\cong} & \mathcal{C}_{m,N,\ell^{-1}j_0}. \end{array}$$

We claim that $(\pi_2)_*(c\mathcal{g}_{0,1/mN}) = c\mathcal{g}_{0,1/mN}$. However, $(\pi_2)_*c\mathcal{g}_{0,1/mN}$ is the pushforward of $\varphi_\ell^*(c\mathcal{g}_{0,1/mN}) \in \mathcal{O}(Y(m, mN(\ell)))^\times$ along the natural projection $\mathcal{O}(Y(m, mN(\ell)))^\times \rightarrow \mathcal{O}(Y(m, mN))^\times$, and the distribution relation of equation (1b) shows that the pushforward of $\varphi_\ell^*(c\mathcal{g}_{0,1/mN})$ is $c\mathcal{g}_{0,1/mN}$, as required. Hence,

$$(\pi_2 \times \pi_2)_*(\mathcal{A}, c\mathcal{g}_{0,1/mN}) = (\mathcal{C}_{m,N,\ell^{-1}j_0}, c\mathcal{g}_{0,1/mN}) = c\mathcal{Z}_{m,N,\ell^{-1}j_0} = \Delta_\ell^*(c\mathcal{Z}_{m,N,j}),$$

as required. □

3.4. *The second norm relation for $p \nmid mN$.* In this section, we shall assume that $N \geq 5$, $m \geq 1$, $j \in \mathbb{Z}/m\mathbb{Z}$, and p is a prime such that $p \nmid mN$. Our aim is to prove the following theorem:

THEOREM 3.4.1. *We have*

$$\sum_{\substack{k \in \mathbb{Z}/m\mathbb{Z} \\ k \equiv j \pmod m \\ p \nmid k}} c\Xi_{mp,N,k} = \left(-\sigma_p + (T'_p, T'_p) + [(p+1)(\langle p^{-1} \rangle, \langle p^{-1} \rangle) - (\langle p^{-1} \rangle, T_p'^2) - (T_p'^2, \langle p^{-1} \rangle)] \sigma_p^{-1} + (\langle p^{-1} \rangle T'_p, \langle p^{-1} \rangle T'_p) \sigma_p^{-2} - p(\langle p^{-2} \rangle, \langle p^{-2} \rangle) \sigma_p^{-3} \right) c\Xi_{m,N,j}.$$

Remark 3.4.2. One can formulate a version of this theorem for the zeta elements $c\mathcal{Z}_{m,N,j}$, from which Theorem 3.4.1 would follow in the same way as Theorem 3.3.2 follows from Theorem 3.3.1. The argument given below can easily be extended to prove this slightly stronger result; however, we shall not pursue this here, as the above statement suffices for our applications.

We begin the proof of Theorem 3.4.1 by rewriting the $T_p'^2$ terms using a related Hecke operator S'_p .

PROPOSITION 3.4.3. *As elements of the Hecke algebra of $\Gamma_1(N)$, we have $T_p'^2 = S'_p + (p+1)\langle p^{-1} \rangle R_p$, where S'_p is the double coset of $\begin{pmatrix} p^2 & 0 \\ 0 & 1 \end{pmatrix}$ and R_p is the double coset of $\begin{pmatrix} p & 0 \\ 0 & p \end{pmatrix}$.*

Proof. This is a simple computation from the definition of multiplication in the Hecke algebra. □

Since R_p acts trivially on everything in sight, the formula of Theorem 3.4.1 can be written as

$$(4) \quad \sum_{\substack{k \in (\mathbb{Z}/m\mathbb{Z})^\times \\ k \equiv j \pmod m}} c\Xi_{mp,N,k} = \left(((T'_p, T'_p) - \sigma_p - p(\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-1}) (1 + (\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-2}) - [(\langle p^{-1} \rangle, S'_p) + (S'_p, \langle p^{-1} \rangle)] \sigma_p^{-1} \right) c\Xi_{m,N,j}.$$

3.4.1. *Evaluation of $(T'_p, T'_p)c\Xi_{m,N,j}$.* First we shall make a careful study of the operator (T'_p, T'_p) .

PROPOSITION 3.4.4. *If $G = \mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z})$ and B is the lower-triangular Borel subgroup, then $B \backslash G/B$ has exactly two elements, B and its complement (the “big Bruhat cell”).*

Proof. Well known. □

COROLLARY 3.4.5. *Let Γ be any congruence subgroup of $\mathrm{SL}_2(\mathbb{Z})$ of level prime to p , and let $\alpha \in \mathrm{SL}_2(\mathbb{Q})$ be integral at p . Then the double coset $\Gamma\alpha\Gamma$ is the union of exactly two double cosets of $\Gamma' = \Gamma \cap \Gamma^0(p)$, corresponding to those elements whose reductions modulo p land in the two double cosets of B in $\mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z})$.*

Proof. This is a consequence of strong approximation for $SL_2(\mathbb{Z})$. Since Γ has level prime to p , it surjects onto $SL_2(\mathbb{Z}/p\mathbb{Z})$. Hence we may assume (by left or right multiplying α by an appropriate element of Γ) that the reduction of α modulo p is the identity.

We first note that $\Gamma \cap \alpha^{-1}\Gamma\alpha$ is also a congruence subgroup of level prime to p , so (by the strong approximation theorem) we see that $\Gamma \backslash \Gamma'$ admits a set of coset representatives lying in $\Gamma \cap \alpha^{-1}\Gamma\alpha$ and thus $\Gamma\alpha\Gamma = \Gamma\alpha\Gamma'$.


Now let $x = \gamma\alpha\gamma' \in \Gamma\alpha\Gamma'$. We consider the reduction \bar{x} of x modulo p . If this lies in B , then (since $\bar{\alpha}$ and $\bar{\gamma}'$ are in B) we must have $\bar{x} \in B$ and hence $\gamma \in \Gamma'$; thus $x \in \Gamma'\alpha\Gamma'$.

On the other hand, let μ be any element of Γ that is not in $\Gamma^0(p)$. If $\bar{\gamma} \notin B$, then $\bar{\gamma} \in B\bar{\mu}B$; so there is some $\sigma \in \Gamma \cap \alpha^{-1}\Gamma\alpha \cap \Gamma^0(p)$ such that $\bar{\gamma} \in B\bar{\mu}\bar{\sigma}$. So $\bar{\gamma}\bar{\sigma}^{-1}\bar{\mu}^{-1} \in B$ and thus $\gamma \in \Gamma'\mu\sigma$. Hence $x \in \Gamma'\mu\sigma\alpha\Gamma'$; but $\alpha^{-1}\sigma\alpha \in \Gamma'$ (since by hypothesis $\alpha = 1 \pmod p$ and thus conjugation by α fixes $\Gamma^0(p)$) and thus $x \in \Gamma'\mu\alpha\Gamma'$. \square

COROLLARY 3.4.6. *For $k \in \mathbb{Z}/mp\mathbb{Z}$, let $D_{m,N,k}$ denote the curve in $Y(\Gamma_1(N) \cap \Gamma^0(p))^2$ consisting of points of the form $(z, z + \frac{k}{m})$. Then the preimage $\pi_1^{-1}(C_{m,N,j}) \subseteq Y(\Gamma_1(N) \cap \Gamma^0(p))^2$ consists of exactly two components: one is the curve $D_{m,N,k}$ where k is the unique lifting of j to $\mathbb{Z}/mp\mathbb{Z}$ that is zero modulo p , and the other is the curve $D_{m,N,k}$ where k is any lifting of j to $\mathbb{Z}/mp\mathbb{Z}$ that is a unit modulo p (the resulting curve being independent of the choice of lifting).*

Proof. The preimage of $C_{m,N,j}$ in $\mathcal{H} \times \mathcal{H}$ is exactly the set of (u, v) such that $v = \gamma u$ for some γ in the double coset $\Gamma_1(N) \begin{pmatrix} 1 & j/m \\ 0 & 1 \end{pmatrix} \Gamma_1(N)$. The above proposition describes the decomposition of this set into double cosets of $\Gamma_1(N) \cap \Gamma^0(p)$, hence the result. \square

For any k lifting j (unit or nonunit), we may erect the following diagram of modular curves:



Here U_k is the preimage in $SL_2(\mathbb{Z})$ of the subgroup

$$B \cap \begin{pmatrix} 1 & k \\ 0 & m \end{pmatrix}^{-1} B \begin{pmatrix} 1 & k \\ 0 & m \end{pmatrix} = B \cap \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}^{-1} B \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}.$$

(The equality follows from the fact that conjugation by $\begin{pmatrix} 1 & 0 \\ 0 & m \end{pmatrix}$ fixes B .) This subgroup is just B if $k \in p\mathbb{Z}$; otherwise it is the subgroup

$$\left\{ \begin{pmatrix} u^{-1} & 0 \\ k^{-1}(u - u^{-1}) & u \end{pmatrix} : u \in (\mathbb{Z}/p\mathbb{Z})^\times \right\},$$

which is a maximal torus in $SL_2(\mathbb{Z}/p\mathbb{Z})$. The square at the bottom left of the diagram is Cartesian. The maps α , ρ_k and π_1 are the natural projection maps, and the remaining maps are defined by

$$\begin{aligned} \iota'_{m,N,j} &: z &\mapsto & \left(\frac{z}{m}, \frac{z+j}{m} \right) \\ \iota'_{mp,N,k} &: z &\mapsto & \left(\frac{z}{mp}, \frac{z+k}{mp} \right) \\ \pi_2 &: (u, v) &\mapsto & \left(\frac{u}{p}, \frac{v}{p} \right) \\ \lambda_k &: z &\mapsto & \left(\frac{z}{m}, \frac{z+k}{m} \right). \end{aligned}$$

Definition 3.4.7. Let a and b be the unique elements of $\mathbb{Z}/pm\mathbb{Z}$ congruent to j modulo m and such that $a = 0 \pmod p$ and $b = 1 \pmod p$.

An application of Lemma 2.4.5 shows that we have

COROLLARY 3.4.8. *For any $\alpha \in \mathbb{Z}/mN\mathbb{Z}$, we have*

$$\begin{aligned} (T'_p, T'_p)(c\Xi_{m,N,j,\alpha}) &= (C_{mp,N,a}, (\pi_2 \circ \lambda_a) * c g_{0,\alpha/mN}) \\ &\quad + (C_{mp,N,b}, (\pi_2 \circ \lambda_b) * c g_{0,\alpha/mN}). \end{aligned}$$

(It is convenient to allow $\alpha \neq 1$ here, for reasons that will become clear below.)

We first consider the term for a . Here we have $U_a = \Gamma^0(p)$, so $\Gamma_1(mN) \cap \Gamma^0(m) \cap U_a = \Gamma_1(mN) \cap \Gamma^0(mp)$. Since $p \mid a$, we see that $\pi_2 \circ \lambda_a$ can also be expressed as a composition

$$(5) \quad \begin{array}{c} Y(\Gamma_1(mN) \cap \Gamma^0(mp)) \\ \downarrow z \mapsto z/p \\ Y(\Gamma_1(mN) \cap \Gamma^0(m)) \\ \downarrow z \mapsto \left(\frac{z}{m}, \frac{z + \text{“}p^{-1}\text{”}j}{m} \right) \\ C_{mp,N,a} = C_{m,N,\text{“}p^{-1}\text{”}j}, \end{array}$$

where “ p^{-1} ” is the inverse of p in $\mathbb{Z}/m\mathbb{Z}$.

PROPOSITION 3.4.9. *The pushforward of $c\mathcal{G}_{0,\alpha/mN}$ to $Y(\Gamma_1(mN) \cap \Gamma^0(m))$ along the first map in (5) is $c\mathcal{G}_{0,\alpha/mN} \cdot (c\mathcal{G}_{0,“p^{-1}”\alpha/mN})^p$.*

Proof. See [Kat04, 2.13.2]. □

The second map in (5) is just $l'_{m,N,“p^{-1}”j}$, so we deduce that

$$\begin{aligned} & (C_{mp,N,a}, (\pi_2 \circ \lambda_a) * c\mathcal{G}_{0,\alpha/mN}) \\ &= \left(C_{m,N,“p^{-1}”a}, (l'_{m,N,“p^{-1}”j}) * (c\mathcal{G}_{0,\alpha/mN} \cdot (c\mathcal{G}_{0,“p^{-1}”\alpha/mN})^p) \right) \\ &= c\Xi_{m,N,“p^{-1}”j,\alpha} + p c\Xi_{m,N,“p^{-1}”j,“p^{-1}”\alpha} \\ &= (\sigma_p + p(\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-1}) c\Xi_{m,N,j,\alpha}. \end{aligned}$$

COROLLARY 3.4.10. *For any $\alpha \in (\mathbb{Z}/mN\mathbb{Z})^\times$, we have*

$$\begin{aligned} & ((T'_p, T'_p) - \sigma_p - p(\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-1}) c\Xi_{m,N,j,\alpha} \\ &= \left(C_{mp,N,b}, (\pi_2 \circ \lambda_b) * (c\mathcal{G}_{0,\alpha/mN}) \right). \end{aligned}$$

In particular,

$$\begin{aligned} & ((T'_p, T'_p) - \sigma_p - p(\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-1}) (1 + (\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-2}) c\Xi_{m,N,j} \\ &= \left(C_{mp,N,b}, (\pi_2 \circ \lambda_b) * (c\mathcal{G}_{0,1/mN} \cdot c\mathcal{G}_{0,“p^{-1}”/mN}) \right). \end{aligned}$$

Proof. The first formula is immediate from (5) and the evaluation of the $C_{mp,N,a}$ term above. The second formula follows by summing the first formula for $\alpha = 1$ and for $\alpha = p^{-1}$. □

3.4.2. *Evaluation of the norm term.* We want to compare the right-hand side of the formula in Corollary 3.4.10 with the sum of the $c\Xi_{mp,N,k}$ for all unit liftings k of j . To do this, we shall use the fact that all the terms $c\Xi_{mp,N,k}$ may be written as the pushforwards of modular units on the same modular curve $C_{mp,N,b}$. More precisely, if k and ℓ are liftings of j to $\mathbb{Z}/mp\mathbb{Z}$ that are both units modulo p , we have a diagram

$$\begin{array}{ccc} \mathcal{H} & \xrightarrow{\alpha_{k\ell}} & \mathcal{H} \\ \downarrow & & \downarrow \\ Y(\Gamma_1(mN) \cap \Gamma^0(m) \cap U_k) & \xrightarrow{\cong} & Y(\Gamma_1(mN) \cap \Gamma^0(m) \cap U_\ell) \\ \downarrow \lambda_k & & \downarrow \lambda_\ell \\ D_{m,N,\ell} & \xlongequal{\quad\quad\quad} & D_{m,N,k} \end{array}$$

where α_{kl} is any matrix of the form $\begin{pmatrix} 1 & 0 \\ v & 1 \end{pmatrix}$ with $v \in mN\mathbb{Z}$ congruent to $\frac{1}{k} - \frac{1}{\ell}$ modulo p .

Consequently, we can write $c\Xi_{mp,N,k} = (C_{mp,N,b}, (\pi_2 \circ \lambda_b)_* f_k)$, where

$$f_k = \alpha_{bk}^*(\rho_k)_* (c\mathcal{G}_{0,1/mpN}).$$

We may regard $\mathcal{O}(Y(\Gamma_1(mpN) \cap \Gamma^0(mp)))^\times$ as a $\mathrm{SL}_2(\mathbb{F}_p)$ -module in the obvious way, since $\Gamma_1(mpN) \cap \Gamma^0(mp)$ is the kernel of the surjective reduction map $\Gamma_1(mN) \cap \Gamma^0(m) \rightarrow \mathrm{SL}_2(\mathbb{F}_p)$. With this convention, we have

$$(\rho_k)_* (c\mathcal{G}_{0,1/mpN}) = \prod_{u \in \mathbb{F}_p^\times} \begin{pmatrix} u^{-1} & 0 \\ k^{-1}(u - u^{-1}) & u \end{pmatrix}^* c\mathcal{G}_{0,1/mpN}$$

and thus

$$\begin{aligned} f_k &= \prod_{u \in \mathbb{F}_p^\times} \left[\begin{pmatrix} u^{-1} & 0 \\ k^{-1}(u - u^{-1}) & u \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 - k^{-1} & 1 \end{pmatrix} \right]^* c\mathcal{G}_{0,1/mpN} \\ &= \prod_{u \in \mathbb{F}_p^\times} \begin{pmatrix} u^{-1} & 0 \\ u - k^{-1}u^{-1} & u \end{pmatrix}^* c\mathcal{G}_{0,1/mpN}. \end{aligned}$$

Let K be the set of possible values of k , i.e., the set of elements of $\mathbb{Z}/mp\mathbb{Z}$ congruent to j modulo m and not divisible by p . Then as k varies over K , for each fixed u , the expression $u - k^{-1}u^{-1}$ takes every value in \mathbb{F}_p exactly once with the exception of u , since $k^{-1}u^{-1}$ takes every value except 0. So

$$\prod_{k \in K} f_k = \prod_{\substack{u, v \in \mathbb{F}_p \\ u \neq 0 \\ v \neq u}} c\mathcal{G}_{v_0/mpN, u_1/mpN}.$$

Here by x_1 and x_0 for $x \in \mathbb{F}_p$ we mean any element of $\mathbb{Z}/mpN\mathbb{Z}$ congruent to $x \pmod p$ and to 1 (resp. 0) modulo mN .

We find that

$$(6a) \quad \prod_{u, v \in \mathbb{F}_p} c\mathcal{G}_{v_0/mpN, u_1/mpN} = c\mathcal{G}_{0,1/mN},$$

$$(6b) \quad \prod_{v \in \mathbb{F}_p} c\mathcal{G}_{v_0/mpN, 0_1/mpN} = \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}^* c\mathcal{G}_{0,\beta/mN},$$

$$\begin{aligned} (6c) \quad \prod_{u \in \mathbb{F}_p} c\mathcal{G}_{u_0/mpN, u_1/mpN} &= \begin{pmatrix} 1 & 0 \\ 1_0 & 1 \end{pmatrix}^* \prod_{u \in \mathbb{F}_p} c\mathcal{G}_{0, u_1/mpN} \\ &= \begin{pmatrix} 1 & 0 \\ 1_0 & 1 \end{pmatrix}^* \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}^* c\mathcal{G}_{0,1/mN}, \end{aligned}$$

$$(6d) \quad c\mathcal{G}_{0_0/mpN, 0_1/mpN} = c\mathcal{G}_{0,\beta/mN},$$

where β is the inverse of p in $\mathbb{Z}/mN\mathbb{Z}$.

Combining the above, we have

$$\begin{aligned}
 (7) \quad \sum_{\substack{k \in \mathbb{Z}/m\mathbb{Z} \\ k \equiv j \pmod m \\ p \nmid k}} c\Xi_{mp,N,k} &= \left(C_{mp,N,b}, (\pi_2 \circ \lambda_b)_* (c\mathfrak{g}_{0,1/mN} \cdot c\mathfrak{g}_{0,\beta/mN}) \right) \\
 &\quad - \left(C_{mp,N,b}, (\pi_2 \circ \lambda_b)_* \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}^* c\mathfrak{g}_{0,\beta/mN} \right) \\
 &\quad - \left(C_{mp,N,b}, (\pi_2 \circ \lambda_b)_* \begin{pmatrix} 1 & 0 \\ 1_0 & 1 \end{pmatrix}^* \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}^* c\mathfrak{g}_{0,1/mN} \right).
 \end{aligned}$$

Combining the first term on the right-hand side of (7) with Corollary 3.4.8, we see that Theorem 3.4.1 is equivalent to

PROPOSITION 3.4.11. *We have*

$$\begin{aligned}
 & \left([(\langle p^{-1} \rangle, S'_p) + (S'_p, \langle p^{-1} \rangle)] \sigma_p^{-1} \right) c\Xi_{m,N,j} \\
 &= \left(C_{mp,N,b}, (\pi_2 \circ \lambda_b)_* \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}^* c\mathfrak{g}_{0,\beta/mN} \right) \\
 &+ \left(C_{mp,N,b}, (\pi_2 \circ \lambda_b)_* \begin{pmatrix} 1 & 0 \\ 1_0 & 1 \end{pmatrix}^* \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}^* c\mathfrak{g}_{0,1/mN} \right).
 \end{aligned}$$

3.4.3. *The first term in Proposition 3.4.11.* We now calculate how S'_p acts on $c\Xi_{m,N,j}$. We may describe the correspondence S'_p in terms of the subgroup $\Gamma_0^0(p) = \Gamma_0(p) \cap \Gamma^0(p)$; we have $S'_p = (\pi'_2)_* (\pi'_1)^*$, where π'_1 and π'_2 are the two maps from $Y_1(\Gamma_1(N) \cap \Gamma_0^0(p))$ to $Y_1(N)$ given by $z \mapsto pz$ and $z \mapsto z/p$.

An application of the strong approximation theorem shows (as usual) that the preimage $(\pi'_1 \times 1)^{-1} C_{m,N,j} \subseteq Y_1(\Gamma_1(N) \cap \Gamma_0^0(p)) \times Y_1(N)$ is the single curve $F_{m,N,j}$ given by the set of points of the form $(\frac{z}{p}, z + \frac{j}{m})$.

Applying Lemma 2.4.5 once more, we have a Cartesian square of curves (up to birational equivalence)

$$\begin{array}{ccc}
 & Y(\Gamma_1(mN) \cap \Gamma^0(m) \cap \Gamma_0^0(p)) & \\
 \swarrow pz \leftarrow z & & \searrow z \mapsto \left(\frac{z}{m}, \frac{pz+j}{m} \right) \\
 Y(\Gamma_1(mN) \cap \Gamma^0(m)) & & F_{m,N,j} \\
 \searrow z \mapsto \left(\frac{z}{m}, \frac{z+j}{m} \right) & & \swarrow (pu, v) \leftarrow (u, v) \\
 & C_{m,N,j} &
 \end{array}$$

The functoriality of pushforward maps gives the following:

PROPOSITION 3.4.12. *We have*

$$(S'_p, 1)_{c\Xi_{m,N,j,\alpha}} = \left((\pi'_2 \times 1)F_{m,N,j}, \phi_* \left(\begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}^* c\mathcal{G}_{0,\alpha/mN} \right) \right),$$

where ϕ is the map

$$Y(\Gamma_1(mN) \cap \Gamma^0(m) \cap \Gamma^0_0(p)) \rightarrow (\pi'_2 \times 1)F_{m,N,j} \subset Y_1(N)^2$$

$$z \mapsto \left(\frac{z}{mp}, \frac{pz+j}{m} \right).$$

We first identify the curve $(\pi'_2 \times 1)F_{m,N,j}$.

PROPOSITION 3.4.13. *We have $(\pi'_2 \times 1)F_{m,N,j} = (1 \times \langle p \rangle)^{-1}C_{mp,N,k}$, for any integer k congruent to $p^{-1}j$ modulo m and not divisible by p .*

More precisely, if $k = 1 \pmod p$, and γ'' is a suitable element of $\mathrm{SL}_2(\mathbb{Z})$ that we shall construct below, then there is a commutative diagram

$$\begin{array}{ccc} Y(\Gamma_1(mN) \cap \Gamma^0(m) \cap \Gamma^0_0(p)) & \xrightarrow{z \mapsto \gamma''z} & Y(\Gamma_1(mN) \cap \Gamma^0(m) \cap U_k) \\ \downarrow z \mapsto \left(\frac{z}{mp}, \frac{pz+j}{m} \right) & & \downarrow z \mapsto \left(\frac{z}{mp}, \frac{z+k}{mp} \right) \\ Y_1(N)^2 & \xrightarrow{(1 \times \langle p \rangle)} & Y_1(N)^2, \end{array}$$

where U_k is the level p congruence subgroup from the previous section.

Proof. We note the following matrix identity, which is easy to verify (although tedious to find): for any elements p, x, y of a field F , we have

$$\begin{pmatrix} \frac{y}{x} & 0 \\ \frac{1}{x} - \frac{p}{y} & \frac{x}{y} \end{pmatrix} \begin{pmatrix} 1 & \frac{x}{p} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ \frac{p^2}{y} - \frac{p}{x} & 1 \end{pmatrix} = \begin{pmatrix} p & \frac{y}{p} \\ 0 & \frac{1}{p} \end{pmatrix}.$$

In particular, taking $F = \mathbb{Q}_p$ and $x, y \in \mathbb{Z}_p^\times$, we see that the double cosets of $\mathrm{SL}_2(\mathbb{Z}_p)$ in $\mathrm{SL}_2(\mathbb{Q}_p)$ generated by $\begin{pmatrix} 1 & x/p \\ 0 & 1 \end{pmatrix}$ and $\begin{pmatrix} p & y/p \\ 0 & 1/p \end{pmatrix}$ are equal to each other and independent of x and y .

Since both $\begin{pmatrix} p & y/p \\ 0 & 1/p \end{pmatrix}$ and its inverse have entries in $\frac{1}{p}\mathbb{Z}_p$, it follows that

$$\begin{pmatrix} 1 & x/p \\ 0 & 1 \end{pmatrix} \gamma \in \mathrm{SL}_2(\mathbb{Z}_p) \begin{pmatrix} p & y/p \\ 0 & 1/p \end{pmatrix}$$

for any $\gamma \in \mathrm{SL}_2(\mathbb{Z}_p)$ congruent to $\begin{pmatrix} 1 & 0 \\ -p/x & 1 \end{pmatrix}$ modulo p^2 . (In fact, one can check that it suffices for the matrix to lie in $\begin{pmatrix} 1+p\mathbb{Z}_p & \mathbb{Z}_p \\ -p/x+p^2\mathbb{Z}_p & 1+p\mathbb{Z}_p \end{pmatrix}$.)

If x, y are in $\mathbb{Z}_p^\times \cap \mathbb{Q}$, and we choose γ to be in $\mathrm{SL}_2(\mathbb{Z})$ and congruent to $\begin{pmatrix} 1 & 0 \\ -p/x & 1 \end{pmatrix}$ modulo p^2 , then the matrix

$$\gamma' = \begin{pmatrix} 1 & x/p \\ 0 & 1 \end{pmatrix} \gamma \begin{pmatrix} p & y/p \\ 0 & 1/p \end{pmatrix}^{-1}$$

will be in $\mathrm{SL}_2(\mathbb{Q})$ and will be p -adically integral. If we choose γ to be ℓ -adically close to the identity for some prime $\ell \neq p$, then γ' will be ℓ -adically close to $\begin{pmatrix} 1 & x/p \\ 0 & 1 \end{pmatrix} \gamma \begin{pmatrix} p & y/p \\ 0 & 1/p \end{pmatrix}^{-1} = \begin{pmatrix} 1/p & (px-y)/p \\ 0 & p \end{pmatrix}$.

So if $x, y \in \mathbb{Q}$ are units at p and satisfy $y = px \pmod{1}$, we may choose $\gamma, \gamma' \in \mathrm{SL}_2(\mathbb{Z})$ such that

- $\gamma \in \Gamma_1(m^2N)$,
- $\gamma = \begin{pmatrix} 1 & 0 \\ -p/x & 1 \end{pmatrix} \pmod{p^2}$,
- $\gamma' = \begin{pmatrix} * & * \\ 0 & 1/p \end{pmatrix} \pmod{N}$,
- the identity $\begin{pmatrix} 1 & x/p \\ 0 & 1 \end{pmatrix} \gamma = \gamma' \begin{pmatrix} p & y/p \\ 0 & p \end{pmatrix}$ holds.

We now take $y = j/m$, and $x = k/m$ for any k congruent to $p^{-1}j$ modulo m and invertible modulo p . Then we obtain a commutative diagram

$$\begin{array}{ccc} \mathcal{H} & \xrightarrow{z \mapsto \gamma z} & \mathcal{H} \\ \downarrow z \mapsto \left(z, p^2z + \frac{j}{m}\right) & & \downarrow z \mapsto \left(z, z + \frac{k}{mp}\right) \\ Y_1(N)^2 & \xrightarrow{(1 \times \langle p \rangle)} & Y_1(N)^2 \end{array}$$

or, equivalently,

$$\begin{array}{ccc} \mathcal{H} & \xrightarrow{z \mapsto \gamma'' z} & \mathcal{H} \\ \downarrow z \mapsto \left(\frac{z}{mp}, \frac{pz+j}{m}\right) & & \downarrow z \mapsto \left(\frac{z}{mp}, \frac{z+k}{mp}\right) \\ Y_1(N)^2 & \xrightarrow{(1 \times \langle p \rangle)} & Y_1(N)^2, \end{array}$$

where $\gamma'' = \begin{pmatrix} mp & 0 \\ 0 & 1 \end{pmatrix} \gamma \begin{pmatrix} mp & 0 \\ 0 & 1 \end{pmatrix}^{-1}$. Note that γ'' is in $\Gamma_1(mN) \cap \Gamma^0(mp)$ and is congruent modulo p to

$$\begin{pmatrix} 1 & 0 \\ -1/mx & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ -1/k & 1 \end{pmatrix}.$$

In the preceding diagram, the left vertical map factors through $Y(\Gamma_1(mN) \cap \Gamma^0(m) \cap \Gamma_0^0(p))$, and γ' conjugates this onto $\Gamma_1(mN) \cap \Gamma^0(m) \cap U_k$; so we finally obtain the diagram stated in the proposition. \square

COROLLARY 3.4.14. *In the notation of the preceding subsection, we have*

$$(S'_p, \langle p^{-1} \rangle) \cdot \sigma_p^{-1} \cdot {}_c\Xi_{m,N,j} = \left(C_{mp,N,b}, (\pi_2 \circ \lambda_b)_* \begin{pmatrix} 1 & 0 \\ 1_0 & 1 \end{pmatrix}^* \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}^* {}_c g_{0,1/mN} \right).$$

Proof. This follows from the previous proposition (and its proof) since the right-hand vertical map in the diagram of the proposition is the same as λ_k above and $(\gamma'')^{-1}$ represents the coset $\begin{pmatrix} 1 & 0 \\ 1_0 & 1 \end{pmatrix}$. □

3.4.4. *The second term in Proposition 3.4.11.* Now we are left to analyse the operator $(1, S'_p)$. To simplify the analysis we shall also consider the operator S_p given by $(\pi'_1)_*(\pi'_2)^*$ (rather than $S'_p = (\pi'_2)_*(\pi'_1)^*$); this is the operator associated to the double coset $\begin{pmatrix} 1 & 0 \\ 0 & p^2 \end{pmatrix}$ and is related to S'_p by the formula

$$S'_p = \langle p^{-2} \rangle^* S_p.$$

Again, we find that the preimage $(\pi'_2)^{-1} C_{m,N,j}$ in $Y_1(N) \times Y(\Gamma_1(N) \cap \Gamma_0^0(p))$ is a single irreducible curve $F_{m,N,j}$ given by points of the form $(z, p(z + j/m))$.

PROPOSITION 3.4.15. *We have*

$$(1, S_p) ({}_c\Xi_{m,N,j,\alpha}) = \left((1 \times \pi'_1)(F_{m,N,j}), \phi_* \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}^* {}_c g_{0,\alpha/mN} \right),$$

where the morphism ϕ is defined by

$$\begin{aligned} Y(\Gamma_1(mN) \cap \Gamma^0(m) \cap \Gamma_0^0(p)) &\rightarrow Y_1(N)^2 \\ z &\mapsto \left(\frac{z}{mp}, \frac{pz+p^2j}{m} \right). \end{aligned}$$

Proof. Closely analogous to the previous case. □

We also have a matrix identity

$$\begin{pmatrix} 1 & \frac{x}{p} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -\frac{p}{x} & 1 \end{pmatrix} = \begin{pmatrix} 0 & x \\ -\frac{1}{x} & p + \frac{p^2y}{x} \end{pmatrix} \begin{pmatrix} p & py \\ 0 & \frac{1}{p} \end{pmatrix}$$

from which we may deduce that if x, y are rational numbers that are units at p and such that $x = py \pmod{1}$, there exist $\gamma, \gamma' \in \text{SL}_2(\mathbb{Z})$ such that

- $\gamma \in \Gamma_1(m^2N)$,
- $\gamma = \begin{pmatrix} 1 & 0 \\ -\frac{p}{x} & 1 \end{pmatrix} \pmod{p^2}$,
- the identity

$$\begin{pmatrix} 1 & \frac{x}{p} \\ 0 & 1 \end{pmatrix} \gamma = \gamma' \begin{pmatrix} p & py \\ 0 & \frac{1}{p} \end{pmatrix}$$

holds,

- γ' is congruent to $\begin{pmatrix} * & * \\ 0 & p \end{pmatrix}$ modulo N .

Thus the diagram

$$\begin{array}{ccc}
 \mathcal{H} & \xrightarrow{z \mapsto \gamma z} & \mathcal{H} \\
 \downarrow z \mapsto (z, p^2(z+y)) & & \downarrow z \mapsto (z, z+x/p) \\
 Y_1(N)^2 & \xrightarrow{(1 \times \langle p \rangle)} & Y_1(N)^2
 \end{array}$$

commutes. We take $y = j/m$, and $x = k/m$ where k is congruent to pj modulo m and not divisible by p . Letting $\gamma'' = \begin{pmatrix} mp & 0 \\ 0 & 1 \end{pmatrix} \gamma \begin{pmatrix} mp & 0 \\ 0 & 1 \end{pmatrix}^{-1}$ as before, we have the diagram

$$\begin{array}{ccc}
 \mathcal{H} & \xrightarrow{z \mapsto \gamma'' z} & \mathcal{H} \\
 \downarrow z \mapsto \left(\frac{z}{mp}, \frac{pz+p^2j}{m} \right) & & \downarrow z \mapsto \left(\frac{z}{mp}, \frac{z+k}{mp} \right) \\
 Y_1(N)^2 & \xrightarrow{(1 \times \langle p \rangle)} & Y_1(N)^2.
 \end{array}$$

Again, this shows that $(1 \times \pi'_1)(F_{m,N,j}) = (1 \times \langle p \rangle)^{-1} C_{mp,N,k}$. If we choose k to be 1 modulo p , then the right vertical map factors through $\Gamma_1(mN) \cap \Gamma^0(m) \cap U_k$, and the isomorphism between the two is given by γ'' , which is in $\Gamma_1(mN) \cap \Gamma^0(p)$ and thus acts trivially on $\begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}^* c g_{0,\alpha/mN}$. Thus we have

$$(1, S_p) \sigma_p \cdot c \Xi_{m,N,j,\alpha} = (1 \times \langle p \rangle)^* \left(C_{mp,N,b}, (\pi_2 \circ \lambda_b)_* \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}^* c g_{0,\alpha/mN} \right).$$

Taking $\alpha = \beta$ (the inverse of p modulo mN) and using the formula $c \Xi_{m,N,j,t} = (\langle t \rangle, \langle t \rangle) \sigma_t^2 c \Xi_{m,N,j}$, we see that

$$\begin{aligned}
 & \left(C_{mp,N,b}, (\pi_2 \circ \lambda_b)_* \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}^* c g_{0,\beta/mN} \right) \\
 &= (1, \langle p^{-1} \rangle) (1, S_p) c \Xi_{m,N,\beta j, \beta} \\
 &= (1, \langle p^{-1} \rangle) (1, \langle p \rangle^2 S'_p) \sigma_p c \Xi_{m,N,j, \beta} \\
 &= (1, \langle p^{-1} \rangle) (1, \langle p \rangle^2 S'_p) \sigma_p (\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-2} c \Xi_{m,N,j} \\
 &= (\langle p^{-1} \rangle, S'_p) \sigma_p^{-1} c \Xi_{m,N,j}
 \end{aligned}$$

as required, completing the proof of Proposition 3.4.11 and hence of Theorem 3.4.1.

3.5. *The second norm relation: higher powers of p .* We shall also need to know how to calculate $\text{norm}_m^{p^k m} c \Xi_{p^k m, N, j}$ for $k = 2, 3$. This is less central to our theory than the $k = 1$ case, but it will be needed in order to compare the elements we construct for N coprime to p with their “ p -stabilized” versions.

THEOREM 3.5.1. *For $p \nmid mN$, we have*

$$\begin{aligned} \text{norm}_{mp^2}^{mp^2} (c\Xi_{mp^2,N,j}) &= (T'_p, T'_p)_{c\Xi_{mp,N,j}} \\ &+ \left(p(\langle p^{-1} \rangle, \langle p^{-1} \rangle) - (\langle p \rangle^{-1}, (T'_p)^2) - ((T'_p)^2, \langle p \rangle^{-1}) \right) \\ &+ 2(\langle p^{-1} \rangle T'_p, \langle p^{-1} \rangle T'_p) \sigma_p^{-1} - p(\langle p^{-2} \rangle, \langle p^{-2} \rangle) \sigma_p^{-2} \Big)_{c\Xi_{m,N,j}} \end{aligned}$$

and

$$\begin{aligned} \text{norm}_{mp^2}^{mp^3} (c\Xi_{mp^3,N,j}) &= (T'_p, T'_p)_{c\Xi_{mp^2,N,j}} \\ &+ \left(p(\langle p^{-1} \rangle, \langle p^{-1} \rangle) - (\langle p \rangle^{-1}, (T'_p)^2) - ((T'_p)^2, \langle p \rangle^{-1}) \right)_{c\Xi_{mp,N,j}} \\ &+ \left((\langle p^{-1} \rangle, \langle p^{-1} \rangle) \left(2(T'_p, T'_p) - ((\langle p \rangle^{-1}, (T'_p)^2) + ((T'_p)^2, \langle p \rangle^{-1})) \sigma_p^{-1} \right) \right)_{c\Xi_{m,N,j}}. \end{aligned}$$

Recall the operator S'_p that appeared above, satisfying $S'_p = (T'_p)^2 - (p+1)\langle p^{-1} \rangle$. In terms of these operators, the formulae we wish to prove are

$$\begin{aligned} \text{norm}_{mp^2}^{mp^2} (c\Xi_{mp^2,N,j}) &= (T'_p, T'_p)_{c\Xi_{mp,N,j}} \\ &+ \left(- (p+2)(\langle p^{-1} \rangle, \langle p^{-1} \rangle) - (\langle p \rangle^{-1}, S'_p) - (S'_p, \langle p \rangle^{-1}) \right) \\ &+ (\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-1} \left(2(T'_p, T'_p) - p(\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-1} \right) \Big)_{c\Xi_{m,N,j}} \end{aligned}$$

and

$$\begin{aligned} \text{norm}_{mp^2}^{mp^3} (c\Xi_{mp^3,N,j}) &= (T'_p, T'_p)_{c\Xi_{mp^2,N,j}} \\ &- \left((p+2)(\langle p^{-1} \rangle, \langle p^{-1} \rangle) + (\langle p \rangle^{-1}, S'_p) + (S'_p, \langle p \rangle^{-1}) \right)_{c\Xi_{mp,N,j}} \\ &+ \left((\langle p^{-1} \rangle, \langle p^{-1} \rangle) \left(2(T'_p, T'_p) - [(2p+2)(\langle p^{-1} \rangle, \langle p^{-1} \rangle) \right. \right. \\ &\quad \left. \left. + (\langle p \rangle^{-1}, S'_p) + (S'_p, \langle p \rangle^{-1})] \sigma_p^{-1} \right) \right)_{c\Xi_{m,N,j}}. \end{aligned}$$

A routine but unpleasant computation (in which the use of Sage [Sage] was found to be invaluable) shows that Theorem 3.5.1, together with Theorem 3.4.1, implies the following formulae for the norms to level prime to p :

THEOREM 3.5.2. *If $p \nmid N$, we have*

$$\begin{aligned} \text{(a)} \quad \text{norm}_m^{p^2m} (c\Xi_{p^2m,N,j}) &= p\sigma_p^2 \left((p-1)(1 - (\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-2}) \right. \\ &\quad \left. - ((T'_p, T'_p) \sigma_p^{-1} + (p-1)) P_p(p^{-1} \sigma_p^{-1}) \right), \end{aligned}$$

$$(b) \quad \text{norm}_m^{p^3 m} ({}_c \Xi_{p^3 m, N, j}) = p^2 \sigma_p^3 \left((p-1)(1 - (\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-2}) \right. \\ \left. - (p^{-1} \sigma_p^{-2}(T'_{p^2}, T'_{p^2}) + (p-1)p^{-1} \sigma_p^{-1}(T'_p, T'_p) + (p-1)) \underline{P}_p(p^{-1} \sigma_p^{-1}) \right).$$

Here \underline{P}_p is the operator-valued Euler factor at p given by

$$\underline{P}_p(X) = 1 - (T'_p, T'_p)X \\ + \left(p(\langle T'_p \rangle^2, \langle p^{-1} \rangle) + p(\langle p^{-1} \rangle, \langle T'_p \rangle^2) - 2p^2(\langle p^{-1} \rangle, \langle p^{-1} \rangle) \right) X^2 \\ - p^2(\langle p^{-1} \rangle T'_p, \langle p^{-1} \rangle T'_p) X^3 + p^4(\langle p^{-2} \rangle, \langle p^{-2} \rangle) X^4.$$

3.5.1. *Evaluation of the (T'_p, T'_p) term.* We begin with a double coset computation in $\text{SL}_2(\mathbb{Q}_p)$. We shall write $K = \text{SL}_2(\mathbb{Z}_p)$ and U for the lower-triangular Iwahori subgroup $\left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in K : b \in p\mathbb{Z}_p \right\}$.

PROPOSITION 3.5.3. *Let $j \geq 1$. Then the double coset*

$$K \begin{pmatrix} p^{-j} & 0 \\ 0 & p^j \end{pmatrix} K$$

decomposes as a disjoint union of exactly four double cosets of the Iwahori U , represented by the elements

$$\left\{ \begin{pmatrix} p^{-j} & 0 \\ 0 & p^j \end{pmatrix}, \begin{pmatrix} 0 & -p^{-j} \\ p^j & 0 \end{pmatrix}, \begin{pmatrix} p^j & 0 \\ 0 & p^{-j} \end{pmatrix}, \begin{pmatrix} 0 & -p^j \\ p^{-j} & 0 \end{pmatrix} \right\}.$$

Proof. As shown by Iwahori and Matsumoto [IM65, §2.2], we have a decomposition

$$\text{SL}_2(\mathbb{Q}_p) = \bigsqcup_{w \in D} UwU,$$

where D is the set of matrices of the form $\begin{pmatrix} p^j & 0 \\ 0 & p^j \end{pmatrix}$ or $\begin{pmatrix} 0 & -p^{-j} \\ p^j & 0 \end{pmatrix}$ for some $j \in \mathbb{Z}$. Comparing this with the well-known Cartan decomposition $\text{SL}_2(\mathbb{Q}_p) = \bigsqcup_{j \geq 0} K \begin{pmatrix} p^{-j} & 0 \\ 0 & p^j \end{pmatrix} K$ gives the statement above. \square

PROPOSITION 3.5.4. *For $\alpha \in \text{SL}_2(\mathbb{Q}_p)$, the index of $U \cap \alpha^{-1}U\alpha$ in U is as follows:*

- (a) $p^{|2j|}$ for $\alpha \in U \begin{pmatrix} p^j & 0 \\ 0 & p^{-j} \end{pmatrix} U, j \in \mathbb{Z}$;
- (b) $p^{|2j+1|}$ if $\alpha \in U \begin{pmatrix} 0 & -p^{-j} \\ p^j & 0 \end{pmatrix} U, j \in \mathbb{Z}$.

Proof. It is clear that the index concerned depends only on the double coset $U\alpha U$, so we may reduce immediately to considering the coset representatives in (a) and (b). In each of these cases we find that the intersection $U \cap \alpha^{-1}U\alpha$ is a subgroup of the form $\left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in K : p^r \mid b, p^s \mid c \right\}$ for some $r \geq 1, s \geq 0$; this clearly has index p^{r+s-1} in U , which gives the above formulae. \square

COROLLARY 3.5.5. *Let $j \in \mathbb{Z}$ and $m \geq 1$, neither divisible by p , and $k \geq 1$. Then the preimage in $Y(\Gamma_1(N) \cap \Gamma^0(p))^2$ of the curve $C_{mp^k, N, j} \subseteq Y_1(N)^2$ is the union of four distinct curves:*

- (1) *the curve D_1 given by points of the form*

$$\left(z, \begin{pmatrix} 1 & \frac{j}{mp^k} \\ 0 & 1 \end{pmatrix} z \right),$$

mapping to $C_{mp^k, N, j}$ with degree p^2 ;

- (2) *the curve D_2 given by points of the form*

$$\left(z, \gamma \begin{pmatrix} 1 & \frac{j}{mp^k} \\ 0 & 1 \end{pmatrix} z \right)$$

*for any $\gamma \in \Gamma_1(N)$ congruent to $\begin{pmatrix} 0 & * \\ * & * \end{pmatrix}$ modulo p , again mapping to $C_{mp^k, N, j}$ with degree p ;*

- (3) *the curve D_3 given by points of the form*

$$\left(z, \begin{pmatrix} 1 & \frac{j}{mp^k} \\ 0 & 1 \end{pmatrix} \gamma^{-1} z \right),$$

where γ is as before, mapping to $C_{mp^k, N, j}$ with degree p ;

- (4) *the curve D_4 given by points of the form*

$$\left(z, \gamma \begin{pmatrix} 1 & \frac{j}{mp^k} \\ 0 & 1 \end{pmatrix} \gamma^{-1} z \right),$$

mapping isomorphically to $C_{mp^k, N, j}$.

Proof. All of these curves are evidently in the preimage of $C_{mp^k, N, j}$. One checks that we have

$$\begin{aligned} \begin{pmatrix} 1 & \frac{j}{mp^k} \\ 0 & 1 \end{pmatrix} &\in U \begin{pmatrix} 0 & -p^{-k} \\ p^k & 0 \end{pmatrix} U, \\ \gamma \begin{pmatrix} 1 & \frac{j}{mp^k} \\ 0 & 1 \end{pmatrix} &\in U \begin{pmatrix} p^k & 0 \\ 0 & p^{-k} \end{pmatrix} U, \\ \begin{pmatrix} 1 & \frac{j}{mp^k} \\ 0 & 1 \end{pmatrix} \gamma^{-1} &\in U \begin{pmatrix} p^{-k} & 0 \\ 0 & p^k \end{pmatrix} U, \\ \gamma \begin{pmatrix} 1 & \frac{j}{mp^k} \\ 0 & 1 \end{pmatrix} \gamma^{-1} &\in U \begin{pmatrix} 0 & -p^k \\ p^{-k} & 0 \end{pmatrix} U. \end{aligned}$$

Hence the curves D_i exhaust the preimage of $C_{mp^k, N, j}$, by Proposition 3.5.3. The calculation of the degrees of the maps down follows from Proposition 3.5.4; and since the total degree is $(p + 1)^2$, they must be distinct. \square

We set

$$\alpha_1 = \begin{pmatrix} 1 & \frac{j}{mp^k} \\ 0 & 1 \end{pmatrix}, \quad \alpha_2 = \gamma \alpha_1, \quad \alpha_3 = \alpha_1 \gamma^{-1}, \quad \alpha_4 = \gamma \alpha_1 \gamma^{-1},$$

so D_i is the locus of points of the form $(z, \alpha_i z)$. Let us define

$$\Delta_i := (\pi_2(D_i), (\pi_2)_*(\pi_1)^*(l'_{mp^k, N, j})_* c\mathcal{G}_{0,1/mp^k N}) \in Z^2(Y_1(N)^2 \otimes \mathbb{Q}(\mu_{mp^k}, 1)).$$

Then we evidently have

$$(T'_p, T'_p)_c \Xi_{mp^k, N, j} = \Delta_1 + \Delta_2 + \Delta_3 + \Delta_4.$$

We shall evaluate each of these in turn, showing that D_1 is the norm of $c\Xi_{mp^{k+1}, N, j}$ and the remaining Δ_i can be calculated in terms of Hecke operators acting on $c\Xi_{mp^r, N, j}$ for $r < k$.

3.5.2. Evaluation of Δ_1 .

COROLLARY 3.5.6. *Pushforward and pullback commute in each of the following four diagrams:*

$$(8a) \quad \begin{array}{ccc} Y(\Gamma_1(mp^k N) \cap \Gamma^0(mp^{k+1}) \cap U) & \xrightarrow{z \mapsto \left(\frac{z}{mp^k}, \frac{z+j}{mp^k}\right)} & D_1 \\ \downarrow & & \downarrow \pi_1 \\ Y(\Gamma_1(mp^k N) \cap \Gamma^0(mp^k)) & \xrightarrow{l'_{mp^k, N, j}} & C_{mp^k, N, j} \end{array}$$

where U is the subgroup of $\Gamma(p^k)$ consisting of matrices whose reduction modulo p^{k+1} lies in the subgroup $\begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix}^{-1} U \begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix}$, and both vertical arrows have degree p^2 ;

$$(8b) \quad \begin{array}{ccc} Y(\Gamma_1(mp^k N) \cap \Gamma^0(mp^{k+1})) & \xrightarrow{z \mapsto \left(\frac{z}{mp^k}, \gamma \cdot \frac{z+j}{mp^k}\right)} & D_2 \\ \downarrow & & \downarrow \\ Y(\Gamma_1(mp^k N) \cap \Gamma^0(mp^k)) & \xrightarrow{l'_{mp^k, N, j}} & C_{mp^k, N, j} \end{array}$$

where both vertical arrows have degree p ;

$$(8c) \quad \begin{array}{ccc} Y(\Gamma_1(mp^k N) \cap \Gamma^0(mp^k) \cap U) & \xrightarrow{z \mapsto \left(\gamma \cdot \frac{z}{mp^k}, \frac{z+j}{mp^k}\right)} & D_3 \\ \downarrow & & \downarrow \\ Y(\Gamma_1(mp^k N) \cap \Gamma^0(mp^k)) & \xrightarrow{l'_{mp^k, N, j}} & C_{mp^k, N, j} \end{array}$$

where both vertical arrows again have degree p ; and

$$(8d) \quad \begin{array}{ccc} Y(\Gamma_1(mp^k N) \cap \Gamma^0(mp^k)) & \xrightarrow{z \mapsto \left(\gamma \cdot \frac{z}{mp^k}, \gamma \cdot \frac{z+j}{mp^k}\right)} & D_4 \\ \downarrow & & \downarrow \\ Y(\Gamma_1(mp^k N) \cap \Gamma^0(mp^k)) & \xrightarrow{\iota'_{mp^k, N, j}} & C_{mp^k, N, j} \end{array}$$

where both vertical arrows are isomorphisms.

Proof. Up to conjugation (and identifying $C_{mp^k, N, j}$ and the D_i with their normalizations) each diagram takes the form

$$\begin{array}{ccc} Y(\Gamma_1 \cap \Gamma_2) & \longrightarrow & Y(\Gamma_1) \\ \downarrow & & \downarrow \\ Y(\Gamma_2) & \longrightarrow & Y(\Gamma) \end{array}$$

for subgroups $\Gamma_1, \Gamma_2 \subseteq \Gamma$. So it suffices to check in each case that $\Gamma_1\Gamma_2 = \Gamma$, or equivalently that $[\Gamma : \Gamma_1] = [\Gamma_2 : \Gamma_1 \cap \Gamma_2]$; that is, that the degrees of the two vertical arrows in each diagram are the same. In each case this reduces to an elementary local computation at p . \square

PROPOSITION 3.5.7 (Evaluation of Δ_1). *We have*

$$\Delta_1 = \sum_{\substack{j' \in (\mathbb{Z}/mp^{k+1}\mathbb{Z})^\times \\ j' \equiv j \pmod{p^k}}} c\Xi_{mp, N, j'}.$$

Proof. This follows by exactly the same argument as in the case $k = 0$ considered above. From Corollary 3.5.6 we know that the modular unit $(\pi_1)^*(\iota'_{mp^k, N, j})^*c\mathfrak{g}_{0,1/mN}$ on D_1 is equal to the pushforward of $c\mathfrak{g}_{0,1/mN}$ along the top horizontal arrow in diagram (8a). The subgroups U for all j in a congruence class modulo p^k are conjugate, and by exactly the same argument as in Proposition 3.4.11, we deduce the result. \square

3.5.3. *Evaluation of Δ_2 and Δ_3 .* We now turn our attention to Δ_2 . Evidently $\pi_2(D_2)$ is the image of $Y(\Gamma_1(mp^k N) \cap \Gamma^0(mp^{k+1}))$ in $Y_1(N)^2$ under the map

$$z \mapsto \left(\begin{pmatrix} 1 & 0 \\ 0 & mp^{k+1} \end{pmatrix} z, \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix} \gamma \begin{pmatrix} 1 & j \\ 0 & mp^k \end{pmatrix} z \right).$$

Let us write $\gamma = \begin{pmatrix} pa & b \\ Nc & d \end{pmatrix}$, where $d \equiv 1 \pmod N$. Then we find that

$$\begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix} \gamma \begin{pmatrix} 1 & j \\ 0 & mp^k \end{pmatrix} = \begin{pmatrix} p & 0 \\ 0 & p \end{pmatrix} \begin{pmatrix} a & b \\ Nc & pd \end{pmatrix} \begin{pmatrix} 1 & j \\ 0 & mp^{k-1} \end{pmatrix}.$$

Since scalar matrices act trivially and $\begin{pmatrix} a & b \\ Nc & pd \end{pmatrix}$ acts on $Y_1(N)$ as the diamond operator $\langle p \rangle$, we see that $\pi_2(D_2)$ can be written as the image of

$$\begin{aligned} Y(\Gamma_1(mp^k N) \cap \Gamma^0(mp^{k+1})) &\longrightarrow Y_1(N)^2 \\ z &\longmapsto \left(\frac{z}{mp^{k+1}}, \langle p \rangle \cdot \frac{z+j}{mp^{k-1}} \right). \end{aligned}$$

This map factors through the natural projection

$$\lambda : Y(\Gamma_1(mp^k N) \cap \Gamma^0(mp^{k+1})) \rightarrow Y(\Gamma_1(mp^{k-1} N) \cap \Gamma^0(mp^{k+1})).$$

(Indeed, the first component obviously factors through $\Gamma_1(N) \cap \Gamma^0(mp^{k+1})$, and the second component factors through $Y(\Gamma_1(mp^{k-1} N) \cap \Gamma^0(mp^{k-1}))$ as the map $\iota'_{mp^{k-1}, N, j}$ constructed above composed with the automorphism $\langle p \rangle$.)

PROPOSITION 3.5.8. *We have*

$$\lambda_* (c\mathfrak{g}_{0,1/mp^k N}) = \begin{cases} c\mathfrak{g}_{0,1/mp^{k-1} N} & \text{if } k \geq 2, \\ c\mathfrak{g}_{0,1/mN} \cdot \left(\begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}^* c\mathfrak{g}_{0, \langle p^{-1} \rangle / mN} \right)^{-1} & \text{if } k = 1. \quad \square \end{cases}$$

We thus have

PROPOSITION 3.5.9. *We have $\Delta_2 = (C, \phi)$, where*

- C is the image of $Y(\Gamma_1(mp^{k-1} N) \cap \Gamma^0(mp^{k+1}))$ in $Y_1(N)^2$ under the map

$$\beta : z \mapsto \left(\frac{z}{mp^{k+1}}, \langle p \rangle \cdot \frac{z+j}{mp^{k-1}} \right),$$

- ϕ is the pushforward of $c\mathfrak{g}_{0,1/mp^{k-1} N}$ along this map if $k \geq 2$, and if $k = 1$, it is $c\mathfrak{g}_{0,1/mN} \cdot \left(\begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}^* c\mathfrak{g}_{0, \langle p^{-1} \rangle / mN} \right)^{-1}$ along this map if $k \geq 2$ (resp. if $k = 1$).

3.5.4. *Evaluation of Δ_4 .* The last, and easiest, term is Δ_4 .

PROPOSITION 3.5.10 (Evaluation of Δ_4). *We have*

$$\Delta_4 = p(\langle p^{-1} \rangle, \langle p^{-1} \rangle) \cdot \begin{cases} c\Xi_{mp^{k-1}, N, j} & \text{if } k \geq 2, \\ (1 - (\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-2}) c\Xi_{m, N, j} & \text{if } k = 1. \end{cases}$$

Proof. Contemplating diagram (8d) we know that Δ_4 is equal to the pushforward of $c\mathfrak{g}_{0,1/mp^{k-1} N}$ from $Y(\Gamma_1(mp^k N) \cap \Gamma^0(mp^k))$ to $Y_1(N)^2$ along the map

$$z \mapsto \left(\begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix} \gamma \begin{pmatrix} 1 & 0 \\ 0 & mp^k \end{pmatrix} z, \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix} \gamma \begin{pmatrix} 1 & j \\ 0 & mp^k \end{pmatrix} z \right).$$

Since $\begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix} \gamma = \langle p \rangle \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}$, this is simply

$$z \mapsto (\langle p \rangle, \langle p \rangle) \cdot \left(\begin{pmatrix} 1 & 0 \\ 0 & mp^{k-1} \end{pmatrix} z, \begin{pmatrix} 1 & j \\ 0 & mp^{k-1} \end{pmatrix} z \right).$$

This evidently factors as the projection

$$\lambda : Y(\Gamma_1(mp^k N) \cap \Gamma^0(mp^k)) \rightarrow Y(\Gamma_1(mp^{k-1} N) \cap \Gamma^0(mp^{k-1}))$$

composed with the map $(\langle p \rangle, \langle p \rangle) \circ \iota'_{mp^{k-1}, N, j}$.

On the other hand, the pushforward of $c\mathfrak{g}_{0,1/mp^k N}$ from $Y(\Gamma_1(mp^k N) \cap \Gamma^0(mp^k))$ to $Y(\Gamma_1(mp^k N) \cap \Gamma^0(mp^{k-1}))$ is clearly $(c\mathfrak{g}_{0,1/mp^k N})^p$, since the degree of the map is p ; and the pushforward from $Y(\Gamma_1(mp^k N) \cap \Gamma^0(mp^{k-1}))$ to $Y(\Gamma_1(mp^{k-1} N) \cap \Gamma^0(mp^{k-1}))$ maps $c\mathfrak{g}_{0,1/mp^k N}$ to $c\mathfrak{g}_{0,1/mp^k N}$ if $k \geq 2$ and to $c\mathfrak{g}_{0,1/mN} \cdot (c\mathfrak{g}_{0, \text{“}p^{-1}\text{”}/mN})^{-1}$ otherwise. \square

3.5.5. *Evaluation of $(S'_p, \langle p^{-1} \rangle)_{c\Xi_{mp^{k-1}, N, j}}$.* We now compute the image of $c\Xi_{mp^{k-1}, N, j}$ under the Hecke operator $(S'_p, \langle p \rangle)$.

PROPOSITION 3.5.11. *We have the following double coset decompositions in $\mathrm{SL}_2(\mathbb{Q}_p)$:*

$$K \begin{pmatrix} 1 & j/m \\ 0 & 1 \end{pmatrix} K = K = K \begin{pmatrix} 1 & j/m \\ 0 & 1 \end{pmatrix} U^0(p^2)$$

(where $K = \mathrm{SL}_2(\mathbb{Z}_p)$) and

$$\begin{aligned} K \begin{pmatrix} 1 & j/mp \\ 0 & 1 \end{pmatrix} K &= K \begin{pmatrix} p^{-1} & 0 \\ 0 & p \end{pmatrix} K \\ &= K \begin{pmatrix} 1 & j/mp \\ 0 & 1 \end{pmatrix} U^0(p^2) \sqcup K \begin{pmatrix} p^{-1} & 1 \\ 0 & p \end{pmatrix} U^0(p^2) \\ &\sqcup K \begin{pmatrix} p^{-1} & \xi \\ 0 & p \end{pmatrix} U^0(p^2) \sqcup K \begin{pmatrix} p^{-1} & 0 \\ 0 & p \end{pmatrix} U^0(p^2), \end{aligned}$$

where ξ is any quadratic nonresidue in \mathbb{Z}_p^\times .

Geometrically this is expressed as follows:

PROPOSITION 3.5.12. *The preimage in $Y(\Gamma_1(N) \cap \Gamma^0(p^2)) \times Y_1(N)$ of $C_{mp^{k-1}, N, j}$ is*

- if $k = 1$, the single curve E consisting of points of the form $(z, z + j/m)$, with degree $p(p + 1)$ over $C_{m, N, j}$;
- if $k = 2$, the union of four distinct curves E_1, E_2, E_3, E_4 , with degrees $(p, \frac{p-1}{2}, \frac{p-1}{2}, 1)$ respectively over $C_{mp, N, j}$, where
 - E_1 is the curve consisting of points of the form $(z, z + j/mp)$,

- the curve E_2 is the locus of points of the form $(\delta_2 z, z + j/mp)$ where δ_2 is any matrix in $\Gamma_1(N)$ of the form $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$, where $p \mid a$ and $\frac{a}{pb} - \frac{m}{j}$ is a quadratic residue mod p ;
- the curve E_3 is the locus of points of the form $(\delta_3 z, z + j/mp)$, where δ_3 is any matrix in $\Gamma_1(N)$ of the form $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ where $p \mid a$ and $\frac{a}{pb} - \frac{m}{j}$ is a quadratic nonresidue mod p ;
- the curve E_4 is the locus of points of the form $(\delta_4 z, z + j/mp)$, where δ_4 is any matrix in $\Gamma_1(N)$ of the form $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ where $p \mid a$ and $\frac{a}{pb} - \frac{m}{j} = 0 \pmod p$.

Proof. This follows from Lemma 2.4.3 and the previous proposition, noting the identity

$$\begin{pmatrix} 1 & \frac{j}{mp} \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ \frac{pm}{j} & 1 \end{pmatrix} \begin{pmatrix} \frac{1}{p} & \xi \\ 0 & p \end{pmatrix} \begin{pmatrix} p + \frac{p\xi m}{j} & \frac{j}{m} \\ -\frac{m}{j} & 0 \end{pmatrix}. \quad \square$$

PROPOSITION 3.5.13. *In the case $k = 1$, the image of E in $Y_1(N)^2$ under $(u, v) \mapsto (u/p^2, \langle p \rangle v)$ is the curve C of Proposition 3.5.15 above.*

In the case $k = 2$, the image of E_1 under this map is the curve C , the images of E_2 and E_3 are both $(\langle p \rangle, \langle p \rangle)C_{mp, N, j}$, and the image of E_4 is $(\langle p \rangle, \langle p \rangle)C_{m, N, \langle p^{-1} \rangle j}$.

Proof. The $k = 1$ case is clear, as is the assertion for E_1 in case $k = 2$. The remaining statements are a fiddly double coset computation. \square

PROPOSITION 3.5.14. *In the case $k = 1$, we have*

$$(S'_p, \langle p^{-1} \rangle) \cdot {}_c\Xi_{m, N, j} = (C, \beta_* c g_{0, 1/mN}),$$

in the notation of Proposition 3.5.15.

In the case $k = 2$, we have

$$(S'_p, \langle p^{-1} \rangle) \cdot {}_c\Xi_{pm, N, j} = \sum_{i=1}^4 \Theta_i,$$

where Θ_i is the term corresponding to the curve E_i of the previous proposition, and

$$\Theta_1 = (C, \beta_* c g_{0, 1/mpN}).$$

Proof. An argument using Lemma 2.4.5 in a familiar manner shows that the pullback of $(\iota'_{mp^{k-1}, N, j})(c g_{0, 1/mp^{k-1}})$ to E coincides with the pushforward of $c g_{0, 1/mp^{k-1}}$ along the map

$$\begin{aligned} Y(\Gamma_1(mp^{k-1}N) \cap \Gamma^0(mp^{k+1})) &\longrightarrow E \\ z &\longmapsto \left(\frac{z}{mp^{k-1}}, \frac{z+j}{mp^{k-1}} \right). \end{aligned}$$

So the pushforward of this along the map $E \rightarrow Y_1(N)^2$ given by $(u, v) \mapsto (u/p^2, \langle p \rangle v)$ is $(C, \beta_* c g_{0,1/mN})$, since the composition of these two maps is β . \square

Combining the preceding proposition with Proposition 3.5.15, we have $\Delta_2 = (S'_p, \langle p^{-1} \rangle) \cdot c \Xi_{m,N,j} - \Delta'_2$, where

$$\Delta'_2 = \begin{cases} (C, \beta_* \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}^* c g_{0, \langle p^{-1} \rangle / mN}) & \text{if } k = 1, \\ \Theta_2 + \Theta_3 + \Theta_4 & \text{if } k = 2. \end{cases}$$

We may express the $k = 1$ case equivalently as

$$\Delta'_2 = (C, \beta'_* c g_{0, \langle p^{-1} \rangle / mN})$$

where β' is the map

$$Y(\Gamma_1(mN) \cap \Gamma_0(p) \cap \Gamma^0(mp)) \rightarrow C$$

given by $z \mapsto (\frac{z}{mp}, \langle p \rangle \cdot \frac{pz+j}{m})$.

We have seen this map before: we showed above in Proposition 3.4.13 that there was a commutative diagram

$$\begin{array}{ccc} Y(\Gamma_1(mN) \cap \Gamma_0(p) \cap \Gamma^0(mp)) & \xrightarrow{\gamma''} & Y(\Gamma_1(mN) \cap \Gamma^0(m) \cap U_{j'}) \\ \downarrow z \mapsto (\frac{z}{mp}, \frac{pz+j}{m}) & & \downarrow z \mapsto (\frac{z}{mp}, \frac{z+j'}{mp}) \\ Y_1(N)^2 & \xrightarrow{1 \times \langle p \rangle} & Y_1(N)^2, \end{array}$$

where γ'' is a suitable element of $\Gamma_1(mN)$ (in fact of $\Gamma_1(mN) \cap \Gamma^0(mp)$, although we do not need this), $j' = p^{-1}j \bmod m$ is invertible modulo p , and $U_{j'}$ is the preimage in a conjugate of the diagonal torus in $SL_2(\mathbb{F}_p)$.

PROPOSITION 3.5.15 (Evaluation of Δ'_2). *For $k = 1$, we have*

$$\Delta'_2 = (\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-1} \left((T'_p, T'_p) - \sigma_p - p(\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-1} \right) c \Xi_{m,N,j},$$

and consequently

$$\Delta_2 = \left[(S'_p, \langle p^{-1} \rangle) - (\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-1} \left((T'_p, T'_p) - \sigma_p - p(\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-1} \right) \right] \cdot c \Xi_{m,N,j}.$$

We now consider Δ_3 . By applying the automorphism of $Y_1(N)^2$ that switches the two factors and running through essentially the same argument as above, we see that

PROPOSITION 3.5.16 (Evaluation of Δ_3). *For $k = 1$, we have*

$$\Delta_3 = \left[(\langle p^{-1} \rangle, S'_p) - (\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-1} \left((T'_p, T'_p) - \sigma_p - p(\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-1} \right) \right] \cdot {}_c \Xi_{m,N,j}.$$

We now have all the ingredients necessary for the proof in the case $k = 1$, which will be carried out in Section 3.5.8. However, for $k = 2$, there are a few more ingredients we will need.

3.5.6. *Study of Θ_4 .* Let us now consider the term Θ_4 that arises for $k = 2$. Recall that δ_4 was any element of $\Gamma_1(N)$ satisfying a certain congruence modulo p ; we may use strong approximation to make additional congruence assumptions modulo primes away from p , so we shall assume that $\delta_4 = \begin{pmatrix} pa & b \\ mNc & d \end{pmatrix}$ with $ja = mb \pmod p$.

For brevity, we shall write $\Gamma(M, N)$ for the group

$$\left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}) : \begin{array}{l} a = 1, b = 0 \pmod M, \\ c = 0, d = 1 \pmod N \end{array} \right\}.$$

PROPOSITION 3.5.17. *There is a commutative diagram*

$$\begin{array}{ccc} Y(\Gamma(mp, mpN)) & \xrightarrow{z \mapsto \left(\delta_4 \begin{pmatrix} 1 & 0 \\ 0 & mp \end{pmatrix} z, \begin{pmatrix} 1 & j \\ 0 & mp \end{pmatrix} z \right)} & E_4 \\ \downarrow z \mapsto \varepsilon z & & \downarrow (u, v) \mapsto \left(\frac{z}{p^2}, \langle p \rangle z \right) \\ Y(\Gamma(mp, mpN)) & \xrightarrow{(\langle p \rangle, \langle p \rangle) \iota'_{mp,N,j_0}} & Y_1(N)^2, \end{array}$$

where ε is a suitably chosen element of $\Gamma_1(N)$ and j_0 is the unique integer congruent to j modulo m and to 0 modulo p .

Proof. Firstly, we note that $\Gamma_1(N)$ normalizes $\Gamma(mp, mpN)$, so the left-hand vertical arrow is well defined. More subtly, the well-definedness of the top horizontal arrow follows from the inclusion

$$\delta_4 \begin{pmatrix} 1 & 0 \\ 0 & mp \end{pmatrix} \Gamma(mp, mpN) \begin{pmatrix} 1 & 0 \\ 0 & mp \end{pmatrix}^{-1} \delta_4^{-1} \subseteq \Gamma^0(p^2);$$

indeed $\delta_4 \begin{pmatrix} 1 & 0 \\ 0 & mp \end{pmatrix} = \begin{pmatrix} p & 0 \\ 0 & m \end{pmatrix} \delta'_4$ where $\delta'_4 = \begin{pmatrix} a & mb \\ Nc & d \end{pmatrix}$ normalizes $\Gamma(mp, mpN)$, so

$$\begin{aligned} \delta_4 \begin{pmatrix} 1 & 0 \\ 0 & mp \end{pmatrix} \Gamma(mp, mpN) \begin{pmatrix} 1 & 0 \\ 0 & mp \end{pmatrix}^{-1} \delta_4^{-1} &= \begin{pmatrix} m & 0 \\ 0 & p \end{pmatrix} \Gamma(mp, mpN) \begin{pmatrix} m & 0 \\ 0 & p \end{pmatrix}^{-1} \subseteq \Gamma_0(m^2N) \cap \Gamma^0(p^2). \end{aligned}$$

It remains to show that ε may be chosen so that the diagram commutes. We need to choose ε so that we have

$$\Gamma_1(N)\langle p \rangle \begin{pmatrix} 1 & 0 \\ 0 & mp \end{pmatrix} \varepsilon = \Gamma_1(N) \begin{pmatrix} p^{-1} & 0 \\ 0 & p \end{pmatrix} \delta_4 \begin{pmatrix} 1 & 0 \\ 0 & mp \end{pmatrix} = \Gamma_1(N) \begin{pmatrix} 1 & 0 \\ 0 & mp \end{pmatrix} \delta'_4$$

and so that

$$\Gamma_1(N) \begin{pmatrix} 1 & j_0 \\ 0 & mp \end{pmatrix} \varepsilon = \Gamma_1(N) \begin{pmatrix} 1 & j \\ 0 & mp \end{pmatrix},$$

where j_0 is the unique integer congruent to $j \pmod m$ and $0 \pmod p$.

These conditions are both satisfied if we take ε to be congruent to 1 modulo mN and to satisfy the same congruence modulo p as δ'_4 , so $\varepsilon = \begin{pmatrix} x & jx \\ * & * \end{pmatrix} \pmod p$ for some x . \square

COROLLARY 3.5.18. *We have*

$$\Theta_4 = (\langle p^{-1} \rangle, \langle p^{-1} \rangle)^* (C_{m,N, \langle p^{-1} \rangle j}, (\iota'_{mp,N,j_0})^*(\varepsilon^{-1})^* c\mathcal{G}_{0,1/mpN}).$$

Now we calculate the pushforward of $(\varepsilon^{-1})^* c\mathcal{G}_{0,1/mpN}$ from $Y(\Gamma(mp, mpN))$ to $Y(\Gamma(m, mN))$.

PROPOSITION 3.5.19. *Let $\alpha, \beta \in \mathbb{Z}$ be such that $\alpha = 0, \beta = 1 \pmod{mN}$ and $\beta \neq 0 \pmod p$. Then the pushforward of $c\mathcal{G}_{\alpha/mpN, \beta/mpN}$ from $Y(\Gamma(mp, mpN))$ to $Y(\Gamma(m, mN))$ along the map $z \mapsto z/p$ is $c\mathcal{G}_{0,1/mN} \cdot (c\mathcal{G}_{0, \langle p^{-1} \rangle /mN})^{-1}$, and hence*

$$\Theta_4 = (\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p (1 - (\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-2}) c\Xi_{m,N,j}.$$

Proof. A calculation using Theorem 2.2.4 shows that pushing forward to $Y(\Gamma_1(mpN) \cap \Gamma^0(m))$ gives $c\mathcal{G}_{\alpha/mN, \beta/mpN} = c\mathcal{G}_{0, \beta/mpN}$, and we are now in familiar territory. \square

3.5.7. Study of Θ_2 and Θ_3 . Let δ be any element of $\Gamma_1(N) \cap \Gamma_0(mN)$ whose top left entry is divisible by p , so $\delta = \begin{pmatrix} pa & b \\ mNc & d \end{pmatrix}$ with $pa = d = 1 \pmod N$. Let $\delta' = \begin{pmatrix} a & mb \\ Nc & pd \end{pmatrix}$, so $\delta' \in \langle p \rangle \Gamma_1(N)$ and we have $\delta \begin{pmatrix} 1 & 0 \\ 0 & mp \end{pmatrix} = \begin{pmatrix} p & 0 \\ 0 & m \end{pmatrix} \delta'$.

Let E_δ be the locus of points in $Y(\Gamma_1(N) \cap \Gamma^0(p^2)) \times Y_1(N)$ of the form $(\delta z, z + j/mp)$; this clearly maps to $C_{mp,N,j}$ under the natural projection to $Y_1(N)^2$. We then build the following diagram of modular curves:

$$\begin{array}{ccc} Y(\Gamma(mp, mpN)) & \xrightarrow[\cong]{\begin{pmatrix} p & 0 \\ 0 & m \end{pmatrix} \delta', \begin{pmatrix} 1 & j \\ 0 & mp \end{pmatrix}} & E_\delta \\ \text{id} \cong \downarrow & & \downarrow \text{pr} \\ Y(\Gamma(mp, mpN)) & \xrightarrow[\cong]{\begin{pmatrix} 1 & 0 \\ 0 & mp \end{pmatrix}, \begin{pmatrix} 1 & j \\ 0 & mp \end{pmatrix}} & C_{mp,N,j}. \end{array}$$

PROPOSITION 3.5.20. *Suppose that $\delta' = \begin{pmatrix} a' & b' \\ c' & pd \end{pmatrix}$, with $aj - b \not\equiv 0 \pmod p$. Then the intersection*

$$\left[(\delta')^{-1} \begin{pmatrix} 1 & 0 \\ 0 & m \end{pmatrix}^{-1} (\Gamma_1(N) \cap \Gamma_0^0(p)) \begin{pmatrix} 1 & 0 \\ 0 & m \end{pmatrix} \delta' \right. \\ \left. \cap \begin{pmatrix} 1 & j \\ 0 & mp \end{pmatrix}^{-1} \Gamma_1(N) \begin{pmatrix} 1 & j \\ 0 & mp \end{pmatrix} \right] \cap \Gamma(m, mN)$$

consists precisely of those matrices in $\Gamma(m, mN)$ that are congruent to ± 1 modulo p .

Proof. It suffices to show that

$$\Gamma_1(N) \cap \Gamma_0^0(p) \cap \begin{pmatrix} 1 & 0 \\ 0 & m \end{pmatrix} \delta' \begin{pmatrix} 1 & j \\ 0 & mp \end{pmatrix}^{-1} \Gamma_1(N) \begin{pmatrix} 1 & j \\ 0 & mp \end{pmatrix} (\delta')^{-1} \begin{pmatrix} 1 & 0 \\ 0 & m \end{pmatrix}^{-1}$$

consists of matrices that are ± 1 modulo p , since such matrices are clearly preserved under conjugation by $\begin{pmatrix} 1 & 0 \\ 0 & m \end{pmatrix} \delta'$ (which is integral at p).

Let $\gamma \in \Gamma_1(N) \cap \Gamma_0^0(p)$. Then γ is congruent modulo p to $\begin{pmatrix} x & 0 \\ 0 & x^{-1} \end{pmatrix}$ for some $x \in (\mathbb{Z}/p\mathbb{Z})^\times$. We require that

$$\begin{pmatrix} 1 & j \\ 0 & mp \end{pmatrix} (\delta')^{-1} \begin{pmatrix} 1 & 0 \\ 0 & m \end{pmatrix}^{-1} \gamma \begin{pmatrix} 1 & 0 \\ 0 & m \end{pmatrix} \delta' \begin{pmatrix} 1 & j \\ 0 & mp \end{pmatrix}^{-1} \in \mathrm{SL}_2(\mathbb{Z}_p)$$

or, equivalently, that

$$\begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix} (\delta')^{-1} \begin{pmatrix} x & 0 \\ 0 & x^{-1} \end{pmatrix} \delta' \begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix}^{-1} \cong \begin{pmatrix} * & 0 \\ * & * \end{pmatrix} \pmod p.$$

Substituting the entries of δ' , we find that the top right-hand entry of the product on the left is congruent modulo p to $(aj - b)cj(x - x^{-1})$. So if $aj - b$ is not divisible by p , then we must have $x - x^{-1} \equiv 0 \pmod p$; i.e., $x = \pm 1$, as required. \square

Remark 3.5.21. Conceptually, what is going on here is that we have calculated the intersection of *three* Borel subgroups of $\mathrm{SL}_2(\mathbb{F}_p)$ in general position relative to each other, which is simply the centre of the group.

COROLLARY 3.5.22. *The pullback to E_δ of $(\nu'_{mp,N,j})_* c\mathfrak{g}_{0,1/mpN}$ is equal to the pushforward along the top row of the above diagram of the modular unit*

$$\prod_{\gamma \in U_j/\{\pm 1\}} \gamma^* c\mathfrak{g}_{0,1/mpN} \in \mathcal{O}(Y(mp, mpN)^\times),$$

where U_j is (as above) the torus in $\mathrm{SL}_2(\mathbb{F}_p)$ whose preimage is

$$\begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}^{-1} K \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix} \cap \begin{pmatrix} 1 & j \\ 0 & p \end{pmatrix}^{-1} K \begin{pmatrix} 1 & j \\ 0 & p \end{pmatrix},$$

and we choose a lifting of each element of $U/\{\pm 1\}$ to an element of $\Gamma(m, mN)$.

Note that this depends only rather weakly on δ . We calculated U_j explicitly above: it consists of all matrices of the form $\begin{pmatrix} u^{-1} & 0 \\ j^{-1}(u-u^{-1}) & u \end{pmatrix}$ with $u \in \mathbb{F}_p^\times$.

We now consider the pushforward of this to $Y_1(N)^2$ along the map $(u, v) \mapsto (\langle p \rangle^{-1} \begin{pmatrix} p^{-1} & 0 \\ 0 & p \end{pmatrix} u, v)$, so the image of E_δ is one of the components of the image of $C_{mp,N,j}$ under the Hecke operator $(\langle p \rangle S'_p, 1)$.

PROPOSITION 3.5.23. *The image of E_δ under this map is $C_{mp,N,j}$ itself. More specifically, we may find $\varepsilon \in \Gamma(m, mN)$ such that there is a commutative diagram*

$$\begin{CD} Y(\Gamma(mp, mpN)) @>{(\langle p \rangle^{-1} \begin{pmatrix} 1 & 0 \\ 0 & mp \end{pmatrix} \delta', \begin{pmatrix} 1 & j \\ 0 & mp \end{pmatrix})}>> Y_1(N)^2 \\ @V{\varepsilon \cong}VV @VV{\text{id} \cong}V \\ Y(\Gamma(mp, mpN)) @>{\iota'_{mp,N,j}}>> Y_1(N)^2. \end{CD}$$

Proof. We must show that ε can be found in such a way that

$$\begin{pmatrix} 1 & 0 \\ 0 & mp \end{pmatrix} \delta' \in \Gamma_1(N) \langle p \rangle \begin{pmatrix} 1 & 0 \\ 0 & mp \end{pmatrix} \varepsilon$$

and

$$\begin{pmatrix} 1 & j \\ 0 & mp \end{pmatrix} \in \Gamma_1(N) \begin{pmatrix} 1 & j \\ 0 & mp \end{pmatrix} \varepsilon.$$

For any $\varepsilon \in \Gamma(m, mN)$, the matrices

$$\langle p \rangle^{-1} \begin{pmatrix} 1 & 0 \\ 0 & mp \end{pmatrix} \delta' \varepsilon^{-1} \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}^{-1}$$

and

$$\begin{pmatrix} 1 & j \\ 0 & mp \end{pmatrix} \varepsilon^{-1} \begin{pmatrix} 1 & j \\ 0 & mp \end{pmatrix}$$

are integral away from p and have bottom right entry congruent to 1 modulo N ; so we need only show that ε may be chosen such that both are integral at p . So we must show that we can find ε in the intersection

$$\varepsilon \in \begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix}^{-1} U^0(p) \begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix} \cap U^0(p) \delta'.$$

The nonemptiness of this intersection is equivalent to the equality of the double cosets

$$U^0(p) \begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix} U^0(p) \quad \text{and} \quad U^0(p) \begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix} (\delta')^{-1} U^0(p).$$

However, as we have seen before, there is only one double $U^0(p)$ coset in K other than $U^0(p)$ itself, so this equality is equivalent to $\begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix} (\delta')^{-1} \notin U^0(p)$, which is equivalent to our hypothesis $b \neq ja \pmod p$. \square

It remains to be shown that we can choose δ and ε in some reasonable fashion. Let $\xi \in \mathbb{F}_p^\times$. Then we can take $\varepsilon = \begin{pmatrix} j^{-1}(1-\xi) & -\xi \\ 1-j^{-2}(1-\xi) & j+\xi j^{-1} \end{pmatrix}$ and $\delta' = \begin{pmatrix} j^{-1}(1-\xi) & -\xi \\ \xi^{-1} & 0 \end{pmatrix}$. One sees easily that $\begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix} \varepsilon \begin{pmatrix} 1 & -j \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & j \\ 1-j^{-2}(1-\xi) & j^{-1} \end{pmatrix}$ is lower-triangular and that if we take ξ to be a quadratic residue or a nonresidue, then δ' satisfies the congruences stated above, so it suffices to take $\xi = 1$ and one nonsquare ξ .

Let us write $\Gamma(mp, mpN)^\pm$ for the subgroup of $\Gamma(m, mN)$ consisting of matrices that are congruent to ± 1 modulo p . Then we have a diagram

$$\begin{array}{ccccc}
 Y(\Gamma(mp, mpN)) & \xrightarrow{\sigma} & Y(\Gamma(mp, mpN)^\pm) & \xrightarrow[\cong]{\varepsilon} & Y(\Gamma(mp, mpN)^\pm) \\
 & & \searrow^{\mu_1} & & \downarrow^{\mu_3} \\
 & & & & \begin{matrix} \begin{pmatrix} 1 & j \\ 0 & p \end{pmatrix}, \begin{pmatrix} 1 & j \\ 0 & p \end{pmatrix} \\ Y(\Gamma(m, mN))^2 \end{matrix} \\
 & & \swarrow_{\mu_2} & & \\
 & & & & Y(\Gamma(m, mN))^2
 \end{array}$$

Here σ is the natural pushforward map.

The images of μ_1 and μ_2 are both given by the curve C of points of the form $(z, z + j/p)$ in $Y(\Gamma(m, mN))^2$, which maps to $C_{mp,N,j}$ under the map $(u, v) \mapsto (u/m, v/m)$ to $Y_1(N)^2$. We find that

$$\mu_1^*(\sigma \circ \mu_1)_* c\mathcal{G}_{0,1/mpN} = \prod_{u \in \mathbb{F}_p^\times} \begin{pmatrix} u^{-1} & 0 \\ j^{-1}(u - u^{-1}) & u \end{pmatrix}^* c\mathcal{G}_{0,1/mpN}$$

and hence

$$\begin{aligned}
 & (\mu_2)_* \mu_1^*(\sigma \circ \mu_1)_* c\mathcal{G}_{0,1/mpN} \\
 &= (\mu_3)_*(\varepsilon^{-1})^* \mu_1^*(\sigma \circ \mu_1)_* c\mathcal{G}_{0,1/mpN} \\
 &= (\mu_3)_*(\varepsilon^{-1})^* \prod_{u \in \mathbb{F}_p^\times} \begin{pmatrix} u^{-1} & 0 \\ j^{-1}(u - u^{-1}) & u \end{pmatrix}^* c\mathcal{G}_{0,1/mpN} \\
 &= \prod_{v \in \mathbb{F}_p^\times / \pm 1} \prod_{u \in \mathbb{F}_p^\times} \left[\begin{pmatrix} u^{-1} & 0 \\ j^{-1}(u - u^{-1}) & u \end{pmatrix} \varepsilon^{-1} \begin{pmatrix} v^{-1} & 0 \\ j^{-1}(v - v^{-1}) & v \end{pmatrix} \right]^* c\mathcal{G}_{0,1/mpN}.
 \end{aligned}$$

Conjugating by $\begin{pmatrix} 1 & 0 \\ j^{-1} & 1 \end{pmatrix}$ maps the torus U_j onto the diagonal torus and maps ε^{-1} onto the matrix $\begin{pmatrix} j & \xi \\ 0 & j^{-1} \end{pmatrix}$, and the above expression becomes

$$\begin{aligned} & \prod_{v \in \mathbb{F}_p^\times / \pm 1} \prod_{u \in \mathbb{F}_p^\times} \left[\begin{pmatrix} 1 & 0 \\ -j^{-1} & 1 \end{pmatrix} \begin{pmatrix} u^{-1} & 0 \\ 0 & u \end{pmatrix} \begin{pmatrix} j & \xi \\ 0 & j \end{pmatrix}^{-1} \begin{pmatrix} v^{-1} & 0 \\ 0 & v \end{pmatrix} \begin{pmatrix} 1 & 0 \\ j^{-1} & 1 \end{pmatrix} \right]^* c\mathcal{G}_{0,1/mpN} \\ &= \prod_{v \in \mathbb{F}_p^\times / \pm 1} \prod_{u \in \mathbb{F}_p^\times} \left[\begin{pmatrix} 1 & 0 \\ -j^{-1} & 1 \end{pmatrix} \begin{pmatrix} u^{-1}v^{-1}j & u^{-1}v\xi \\ 0 & uvj^{-1} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ j^{-1} & 1 \end{pmatrix} \right]^* c\mathcal{G}_{0,1/mpN}. \end{aligned}$$

We may change variables by letting $a = uv$ and $b = u^{-2}$. Then the product becomes

$$\begin{aligned} & \prod_{a \in \mathbb{F}_p^\times} \prod_{b \in \mathbb{F}_p^{\times 2} \cdot \xi} \left[\begin{pmatrix} 1 & 0 \\ -j^{-1} & 1 \end{pmatrix} \begin{pmatrix} j & b \\ 0 & j^{-1} \end{pmatrix} \begin{pmatrix} a^{-1} & 0 \\ 0 & a \end{pmatrix} \begin{pmatrix} 1 & 0 \\ j^{-1} & 1 \end{pmatrix} \right]^* c\mathcal{G}_{0,1/mpN} \\ &= (\mu_1 \circ \sigma)_* \prod_{b \in \mathbb{F}_p^{\times 2} \cdot \xi} \left[\begin{pmatrix} 1 & 0 \\ -j^{-1} & 1 \end{pmatrix} \begin{pmatrix} j & b \\ 0 & j^{-1} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ j^{-1} & 1 \end{pmatrix} \right]^* c\mathcal{G}_{0,1/mpN} \\ &= (\mu_1 \circ \sigma)_* \prod_{b \in \mathbb{F}_p^{\times 2} \cdot \xi} \left[\begin{pmatrix} j & b \\ -1 & j^{-1}(1-b) \end{pmatrix} \begin{pmatrix} 1 & 0 \\ j^{-1} & 1 \end{pmatrix} \right]^* c\mathcal{G}_{0,1/mpN} \\ &= (\mu_1 \circ \sigma)_* \begin{pmatrix} 1 & 0 \\ j^{-1} & 1 \end{pmatrix}^* \prod_{b \in \mathbb{F}_p^{\times 2} \cdot \xi} c\mathcal{G}_{(-1)_0/mpN, (j^{-1}(1-b))_1/mpN}. \end{aligned}$$

Considering Θ_2 and Θ_3 together corresponds to letting b vary over all of \mathbb{F}_p^\times . If we were to extend the product over all $b \in \mathbb{F}_p$ (residue, nonresidue, or zero), then we would get

$$(9) \quad (\mu_1 \circ \sigma)_* \begin{pmatrix} 1 & 0 \\ j^{-1} & 1 \end{pmatrix}^* \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}^* c\mathcal{G}_{\alpha/p, 1/mN},$$

where α is the image of $-1/mN$ in $(\mathbb{Z}/p\mathbb{Z})^\times$. (Thus $\alpha/p = (-1)_0/mN$.)

The term for $b = 0$ is just

$$(\mu_1 \circ \sigma)_* \begin{pmatrix} j & 0 \\ -1 + j^{-2} & j^{-1} \end{pmatrix}^* c\mathcal{G}_{0,1/mpN} = (\mu_1 \circ \sigma)_* c\mathcal{G}_{0,1/mpN},$$

since $\begin{pmatrix} j & 0 \\ -1 + j^{-2} & j^{-1} \end{pmatrix} \in U_j$. This is what we want: it is the definition of $c\Xi_{mp, N, j}$.

What can we say about the expression in (9)? Writing the pushforward in terms of coset representatives gives us

$$\begin{aligned}
 & \prod_{u \in \mathbb{F}_p^\times} \left[\begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ j^{-1} & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ j^{-1} & 1 \end{pmatrix}^{-1} \begin{pmatrix} u^{-1} & 0 \\ 0 & u \end{pmatrix} \begin{pmatrix} 1 & 0 \\ j^{-1} & 1 \end{pmatrix} \right]^* c\mathcal{G}_{\alpha/p,1/mN} \\
 &= \prod_{u \in \mathbb{F}_p^\times} \left[\begin{pmatrix} u^{-1} & 0 \\ 0 & u \end{pmatrix} \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ j^{-1} & 1 \end{pmatrix} \right]^* c\mathcal{G}_{\alpha/p,1/mN} \\
 &= \begin{pmatrix} 1 & 0 \\ j^{-1} & 1 \end{pmatrix}^* \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}^* \prod_{u \in \mathbb{F}_p^\times} c\mathcal{G}_{u/p,1/mN} \\
 &= \begin{pmatrix} 1 & 0 \\ j^{-1} & 1 \end{pmatrix}^* \left(c\mathcal{G}_{0,1/mN} \cdot \left(\begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}^* c\mathcal{G}_{0,1/mN} \right)^{-1} \right) \\
 &= c\mathcal{G}_{0,1/mN} \cdot \left(\begin{pmatrix} 1 & 0 \\ j^{-1} & 1 \end{pmatrix}^* \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}^* c\mathcal{G}_{0,1/mN} \right)^{-1}.
 \end{aligned}$$

The last line is justified by the fact that $\begin{pmatrix} 1 & 0 \\ j^{-1} & 1 \end{pmatrix}^*$ denotes the action of a matrix congruent to $\begin{pmatrix} 1 & 0 \\ j^{-1} & 1 \end{pmatrix}$ modulo p but to the identity modulo mN , and such a matrix will act trivially on $c\mathcal{G}_{0,1/mN}$.

We have seen both of these terms before: the class in $\mathrm{CH}^2(Y_1(N)^2 \otimes \mathbb{Q}(\mu_m), 1)$ defined by $(C_{mp,N,j}, \text{pushforward of } c\mathcal{G}_{0,1/mN})$ is $(\langle T'_p, T'_p \rangle - \sigma_p - p\langle p \times p \rangle^{-1} \sigma_p^{-1}) c\Xi_{m,N,j}$, by Corollary 3.4.10; and the term corresponding to $(C_{mp,N,j}, \text{pushforward of } \begin{pmatrix} 1 & 0 \\ j^{-1} & 1 \end{pmatrix}^* \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}^* c\mathcal{G}_{0,1/mN})$ is $(\langle p \rangle^{-1}, S'_p) c\Xi_{m,N,j}$, by Corollary 3.4.14.

3.5.8. *Conclusion of the proof.* We can now complete the proof of Theorem 3.5.1 for $k = 1$.

We know that

$$(T'_p, T'_p) c\Xi_{mp,N,j} = \Delta_1 + \Delta_2 + \Delta_3 + \Delta_4,$$

and we have shown that

$$\Delta_1 = \mathrm{norm}_{mp}^{mp^2} (c\Xi_{mp^2,N,j}) \quad (\text{Proposition 3.5.7});$$

$$\begin{aligned}
 \Delta_2 = & \left[(S'_p, \langle p^{-1} \rangle) - (\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-1} \left((T'_p, T'_p) - \sigma_p \right. \right. \\
 & \left. \left. - p(\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-1} \right) \right] \cdot c\Xi_{m,N,j} \quad (\text{Proposition 3.5.15}),
 \end{aligned}$$

$$\begin{aligned}
 \Delta_3 = & \left[(\langle p^{-1} \rangle, S'_p) - (\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-1} \left((T'_p, T'_p) - \sigma_p \right. \right. \\
 & \left. \left. - p(\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-1} \right) \right] \cdot c\Xi_{m,N,j} \quad (\text{Proposition 3.5.16}),
 \end{aligned}$$

and

$$\Delta_4 = p(\langle p^{-1} \rangle, \langle p^{-1} \rangle) \left(1 - (\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-2} \right) c\Xi_{m,N,j} \quad (\text{Proposition 3.5.10}).$$

Combining these statements gives the $k = 1$ case of Theorem 3.5.1.

In the case $k = 2$ we have again $(T'_p, T'_p) c\Xi_{mp^2,N,j} = \Delta_1 + \Delta_2 + \Delta_3 + \Delta_4$, where

$$\Delta_1 = \text{norm}_{mp^2}^{mp^3} \left(c\Xi_{mp^3,N,j} \right) \quad (\text{Proposition 3.5.7}),$$

$$\Delta_2 = (S'_p, \langle p^{-1} \rangle) \cdot c\Xi_{mp,N,j} - \Theta_2 - \Theta_3 - \Theta_4,$$

$$\Delta_3 = (\langle p^{-1} \rangle, S'_p) \cdot c\Xi_{mp,N,j} - \Theta'_2 - \Theta'_3 - \Theta'_4,$$

and

$$\Delta_4 = p(\langle p^{-1} \rangle, \langle p^{-1} \rangle) \cdot c\Xi_{mp,N,j} \quad (\text{Proposition 3.5.10}).$$

Moreover, we have

$$\Theta_4 = (\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p (1 - (\langle p^{-1} \rangle, \langle p^{-1} \rangle) \sigma_p^{-2}) c\Xi_{m,N,j} \quad (\text{Proposition 3.5.19})$$

and

$$\begin{aligned} \Theta_2 + \Theta_3 &= (\langle p^{-1} \rangle, \langle p^{-1} \rangle) \cdot c\Xi_{mp,N,j} \\ &\quad - ((T'_p, T'_p) - \sigma_p - p\langle p \times p \rangle^{-1} \sigma_p^{-1}) c\Xi_{m,N,j} + (\langle p \rangle^{-1}, S'_p) c\Xi_{m,N,j}. \end{aligned}$$

The obvious involution of $Y_1(N)^2 \otimes \mathbb{Q}(\mu_m)$ given by swapping the two factors maps $c\Xi_{p^k m, N, j}$ to $c\Xi_{p^k m, N, -j}$ for each j , and it interchanges Θ_i with Θ'_i for $i = 1, \dots, 4$, so we obtain formulae for these terms, which are identical with the nonprimed versions except $(\langle p^{-1} \rangle, S'_p)$ is interchanged with $(S'_p, \langle p^{-1} \rangle)$. Collecting terms gives Theorem 3.5.1 for $k = 2$.

4. Relation to complex L -values

4.1. *Definition of Rankin–Selberg L -functions.* We recall the definition of Rankin–Selberg L -functions of pairs of modular forms.

Definition 4.1.1. Let f, g be cuspidal new modular eigenforms (of possibly distinct weights k, ℓ and levels N_f, N_g), L a number field containing the coefficients of f and g , and p a prime. We define the local Euler factor

$$P_p(f, g, X) = \det \left(1 - X \text{Frob}_p^{-1} | (V_{L_\lambda}(f) \otimes V_{L_\lambda}(g))^{L_p} \right),$$

where λ is an arbitrary place of L of residue characteristic distinct from p , $V_{L_\lambda}(f)$ is the L_λ -linear representation of $G_{\mathbb{Q}}$ attached to f (and similarly for g) — see Section 6.3 below — and Frob_p denotes the arithmetic Frobenius at p .

This Euler factor may be defined in purely automorphic terms (cf. [Jac72, Th. 14.8]), but the above definition is convenient for our purposes. The following is an elementary calculation:

PROPOSITION 4.1.2. *If $p \nmid N_f N_g$, then*

$$P_p(f, g, X) = (1 - \alpha\gamma X)(1 - \alpha\delta X)(1 - \beta\gamma X)(1 - \beta\delta X),$$

where α, β are the roots of $X^2 - a_p(f)X + p^{k-1}\varepsilon_p(f)$ and similarly γ, δ are the roots of $X^2 - a_p(g)X + p^{\ell-1}\varepsilon_p(g)$. Completely explicitly, this becomes

$$\begin{aligned} P_p(f, g, X) &= 1 - a_p(f)a_p(g)X \\ &\quad + \left(p^{\ell-1}a_p(f)^2\varepsilon_p(g) + p^{k-1}\varepsilon_p(f)a_p(g)^2 - 2p^{k+\ell-2}\varepsilon_p(f)\varepsilon_p(g) \right) X^2 \\ &\quad - p^{k+\ell-2}\varepsilon_p(f)a_p(f)\varepsilon_p(g)a_p(g)X^3 + p^{2k+2\ell-4}\varepsilon_p(f)^2\varepsilon_p(g)^2X^4. \end{aligned}$$

PROPOSITION 4.1.3. *We may write*

$$P_p(f, g, X) = \prod_{i=1}^4 (1 - \lambda_i X),$$

where each λ_i is either 0, or a p -Weil number of weight $\leq (k + \ell - 2)$. In particular, all poles of the meromorphic function $P_p(f, g, p^{-s})^{-1}$ have real part at most $\frac{k+\ell-2}{2}$.

Proof. This is clear from Proposition 4.1.2 if p does not divide the levels of f and g . The remaining cases follow from an explicit computation of the possible local components of f and g , using the Galois-theoretic definition adopted above (since the Weil–Deligne representations attached to f and g must fall into a finite list of possible types). \square

We now define global Rankin–Selberg L -functions as a product of local terms in the usual way.

Definition 4.1.4. We let

$$L(f, g, s) = \prod_{p \text{ prime}} P_p(f, g, p^{-s})^{-1},$$

and for $N \geq 1$, we let

$$L_{(N)}(f, g, s) = \prod_{\substack{p \text{ prime} \\ p \nmid N}} P_p(f, g, p^{-s})^{-1}.$$

PROPOSITION 4.1.5. *Suppose $k \geq \ell$, and write $\Gamma_{\mathbb{C}}(s) = (2\pi)^{-s}\Gamma(s)$. Then the completed L -function*

$$\Lambda(f, g, s) = \Gamma_{\mathbb{C}}(s)\Gamma_{\mathbb{C}}(s - \ell + 1)L(f, g, s)$$

has analytic continuation to all $s \in \mathbb{C}$, except for a simple pole at $s = k$ if $\ell = k$ and $f = \bar{g}$, and it satisfies a functional equation of the form

$$\Lambda(f, g, k + \ell - 1 - s) = \varepsilon(s) \cdot \Lambda(\bar{f} \otimes \bar{g}, s),$$

where ε is a function of the form Ae^{Bs} for constants A, B .

Remark 4.1.6. The function $\varepsilon(s)$ is, as the notation suggests, a global ε -factor, but we shall not use this interpretation here.

In particular, if $k = \ell = 2$ and $s = 1$, the value $L(f, g, 1)$ vanishes (because $\Gamma_{\mathbb{C}}(s - 1)$ has a simple pole) and we have

$$(10) \quad L'(f, g, 1) = 2\pi\Lambda(f, g, 1).$$

4.2. Real-analytic Eisenstein series. We now express the Rankin–Selberg L -function in terms of the Petersson product with a nonholomorphic Eisenstein series, the original example of the Rankin–Selberg method.

Definition 4.2.1. Let $k \geq 0 \in \mathbb{Z}$ and $\alpha \in \mathbb{Q}/\mathbb{Z}$.

(1) For $\tau \in \mathcal{H}$, $s \in \mathbb{C}$ with $k + 2\Re(s) > 2$, we define

$$E_{\alpha}^{(k)}(\tau, s) = (-2\pi i)^{-k} \pi^{-s} \Gamma(s + k) \sum'_{(m,n) \in \mathbb{Z}^2} \frac{\Im(\tau)^s}{(m\tau + n + \alpha)^k |m\tau + n + \alpha|^{2s}},$$

where the prime denotes that the term $(m, n) = (0, 0)$ is omitted if $\alpha = 0$ (but not otherwise).

(2) For τ, s as above, define

$$F_{\alpha}^{(k)}(\tau, s) = (-2\pi i)^{-k} \Gamma(s + k) \pi^{-s} \sum''_{(m,n) \in \mathbb{Z}^2} \frac{e^{2\pi i \alpha m} \Im(\tau)^s}{(m\tau + n)^k |m\tau + n|^{2s}},$$

where the double prime denotes that the term $(m, n) = (0, 0)$ is omitted (always).

PROPOSITION 4.2.2. *The above series have the following properties:*

- (i) (*Automorphy*). If $N\alpha = 0$, then for fixed τ both $E_{\alpha}^{(k)}$ and $F_{\alpha}^{(k)}$ are preserved by the weight k action of $\Gamma_1(N)$. Moreover, the diamond operators act on α by multiplication in the obvious way.
- (ii) (*Action of Atkin–Lehner involutions*). If $N\alpha = 0$, then we have

$$F_{\alpha}^{(k)}(\tau, s) = N^{-k-s} \sum_{x \in \mathbb{Z}/N\mathbb{Z}} e^{2\pi i \alpha x} \tau^{-k} E_{x/N}^{(k)}\left(\frac{-1}{N\tau}, s\right).$$

- (iii) (*Differential operators*). The Maass–Shimura weight-raising differential operator

$$\delta_k := \frac{1}{2\pi i} \left(\frac{d}{d\tau} + \frac{k}{\tau - \bar{\tau}} \right)$$

(cf. [Shi76, eq. (2.8)]) acts on the Eisenstein series via

$$\delta_k E_{\alpha}^{(k)}(\tau, s) = E_{\alpha}^{(k+2)}(\tau, s - 1)$$

and similarly for $F_{\alpha}^{(k)}$.

(iv) (*Analytic continuation and functional equation*). For fixed k, τ, α , both functions $E_\alpha^{(k)}(\tau, s)$ and $F_\alpha^{(k)}(\tau, s)$ have meromorphic continuations to the whole s -plane, which are holomorphic everywhere if $k \neq 0$; and we have

$$E_\alpha^{(k)}(\tau, s) = F_\alpha^{(k)}(\tau, 1 - k - s).$$

(v) (*Relation to Siegel units*). We have

$$E_\alpha^{(0)}(\tau, 0) = 2 \log |g_{0,\alpha}(\tau)|,$$

where $g_{0,\alpha}$ is the Siegel unit of Section 2.2.

Proof. Parts (i)–(iii) are easy explicit computations. Part (iv) is a standard application of the Poisson summation formula, and (v) is [Kat04, formula (3.8.4)(iii)]. \square

Now let f, g be any two newforms of levels N_f, N_g dividing N , and weights k, ℓ , with $k > \ell$. Let $\check{f} \in S_k(\Gamma_1(N))[\pi_f]$ and $\check{g} \in S_\ell(\Gamma_1(N))[\pi_g]$ be forms in the oldspaces at level N attached to f and g (which we shall think of as “test vectors”). For any $\alpha \in \frac{1}{N}\mathbb{Z}/\mathbb{Z}$, set

$$\begin{aligned} \mathcal{D}(\check{f}, \check{g}, x, s) &= \int_{\Gamma_1(N)\backslash\mathcal{H}} \check{f}(-\bar{\tau}) \check{g}(\tau) E_\alpha^{(k-\ell)}(\tau, s - k + 1) \Im(\tau)^{k-2} dx dy \\ &= \langle \check{f}^*(\tau), \check{g}(\tau) \cdot E_\alpha^{(k-\ell)}(\tau, s - k + 1) \rangle_{\Gamma_1(N)}. \end{aligned}$$

The next theorem shows that the function $\mathcal{D}(\check{f}, \check{g}, 1/N, s)$ is an “approximation” to the completed L -function $\Lambda(f, g, s)$ of the previous section, differing from it only by possible bad Euler factors at primes $\ell \mid N$.

THEOREM 4.2.3 (Rankin–Selberg, Shimura). *We have*

$$\mathcal{D}(\check{f}, \check{g}, 1/N, s) = 2^{1-k} i^{k-\ell} N^{2s+2-k-\ell} \Lambda(f, g, s) C(\check{f}, \check{g}, s),$$

where

$$C(\check{f}, \check{g}, s) := \left(\prod_{p \mid N} P_p(f, g, p^{-s}) \right) \sum_{n \in S(N)} a_n(\check{f}) a_n(\check{g}) n^{-s}$$

is a polynomial in the variables p^{-s} for $p \mid N$; in particular, it is holomorphic for all $s \in \mathbb{C}$. Here $S(N)$ is the set of integers all of whose prime factors divide N .

Proof. See [Kat04, Prop. 7.1]; our $E_{1/N}^{(j)}(\tau, s)$ corresponds to

$$(-2\pi i)^{-j} \Gamma(s + j) \pi^{-s} \Im(\tau)^s E(j, \tau, 1/N, 2s)$$

in Kato’s notation, where $j = k - \ell$. To see that $C(\check{f}, \check{g}, s)$ is a polynomial, it suffices to consider the case when $\check{f} = f(az)$ and $\check{g} = g(bz)$ for integers $a \mid N/N_f, b \mid N/N_g$, in which case the result is clear. \square

In particular, for $s = 1$ and $k = \ell = 2$, using equation (10) and the above proposition gives

$$(11) \quad \mathcal{D}(\check{f}, \check{g}, 1/N, 1) = (4\pi)^{-1} L'(f, g, 1) C(\check{f}, \check{g}, 1).$$

Remark 4.2.4. If f, g have coprime levels N_f, N_g with $N_f N_g = N$, and if we take $\check{f} = f$ and $\check{g} = g$ to be the normalized newforms, then $C(\check{f}, \check{g}, s)$ is identically 1, so in this case $\mathcal{D}(\check{f}, \check{g}, 1/N, s)$ is $N^{2s} \Lambda(f, g, s)$ up to constants.

From the functional equation for the real-analytic Eisenstein series and the action of Atkin–Lehner involutions, we have

$$(12) \quad \begin{aligned} \mathcal{D}(\check{f}, \check{g}, x, k + \ell - 1 - s) &= \left\langle \check{f}^*(\tau), \check{g}(\tau) \cdot F_{x/N}^{(k-\ell)}(\tau, s - k + 1) \right\rangle_{\Gamma_1(N)} \\ &= N^{1-s} \sum_{y \in \mathbb{Z}/N\mathbb{Z}} e^{2\pi i xy} \mathcal{D}(w_N \check{f}, w_N \check{g}, y, s). \end{aligned}$$

Here $w_N \check{f}$ is the function $\tau \mapsto N^{-1} \tau^{-k} \check{f}(-1/(N\tau))$; that is, we have chosen our normalizations so that w_N is an involution in weight 2 (but not in more general weights).

4.3. *The Beilinson regulator.* For any smooth variety X over a subfield of \mathbb{C} , there is a canonical map, the *Beilinson regulator*, from $H_{\mathcal{M}}^3(X, \mathbb{Z}(2))$ into complex-analytic Deligne–Beilinson cohomology. These maps were introduced in [Beĭ84]. We shall only need these maps for $H_{\mathcal{M}}^3(X, \mathbb{Z}(2))$ where X is a projective surface, in which case the target group can be identified with de Rham cohomology.

THEOREM 4.3.1 (Beilinson; cf. [Jan88b, p. 45]). *Let X be a smooth projective surface over \mathbb{C} (or a subfield of \mathbb{C}). There is a homomorphism*

$$\text{reg}_{\mathbb{C}} : \text{CH}^2(X, 1) \rightarrow H_{\text{dR}}^2(X/\mathbb{C})/\text{Fil}^2 = (\text{Fil}^1 H_{\text{dR}}^2(X/\mathbb{C}))^{\vee},$$

which sends the class of $\sum_j (Z_j, g_j) \in Z^2(X, 1)$ to the linear functional

$$(13) \quad \omega \mapsto \frac{1}{2\pi i} \sum_j \int_{Z_j - Z_j^{\text{sing}}} \omega \log |g_j|.$$

We now show that the images of the generalized Beilinson–Flach elements $\Xi_{m,N,j}$ under $\text{reg}_{\mathbb{C}}$, paired with differentials corresponding to weight 2 modular forms f, g , are related to the derivatives of Rankin–Selberg L -functions at the point $s = 1$. More precisely, we shall apply $\text{reg}_{\mathbb{C}}$ to a lifting of $\Xi_{m,N,j}$ to $\text{CH}^2(X_1(N)^2 \otimes \mathbb{Q}(\mu_m), 1) \otimes \mathbb{Q}$; the result will turn out to be independent of the choice of lifting.

Definition 4.3.2. If $f \in S_2(\Gamma_1(N))$, we let $f^* \in S_2(\Gamma_1(N))$ be the form obtained by applying complex conjugation to the Fourier coefficients of f .

We let ω_f denote the holomorphic differential on $X_1(N)$ whose pullback to \mathcal{H} is $2\pi i f(z) dz$, and we let η_f^{ah} be the anti-holomorphic differential $\overline{\omega_{f^*}}$, whose pullback is $-2\pi i f(-\bar{z}) d\bar{z}$.

Remark 4.3.3.

- (1) The factor $2\pi i$ is convenient since $\frac{dq}{q} = 2\pi i dz$.
- (2) The map $f \rightarrow \eta_f^{\text{ah}}$ is \mathbb{C} -linear and Hecke-equivariant (whereas the more obvious map $f \mapsto \overline{\omega_f}$ has neither of these desirable properties).

THEOREM 4.3.4 (Beilinson; cf. [BDR12, Prop. 4.1]). *Let $\tilde{\Xi}_N$ be any element of $\text{CH}^2(X_1(N)^2, 1)$ lifting $\Xi_N := \Xi_{1,N,1} \in \text{CH}^2(Y_1(N)^2, 1)$, and let p_1, p_2 be the projections of $X_1(N)^2$ onto its two factors. Then for \check{f}, \check{g} as above, we have*

$$\langle \text{reg}_{\mathbb{C}}(\tilde{\Xi}_N), p_1^*(\eta_{\check{f}}^{\text{ah}}) \wedge p_2^*(\omega_{\check{g}}) \rangle = 2\pi \mathcal{D}(\check{f}, \check{g}, 1/N, 1) = \frac{1}{2} L'(f, g, 1) C(\check{f}, \check{g}, 1).$$

Proof. We have

$$\begin{aligned} \mathcal{D}'(f, g, 1/N, 1) &= \left(\int_{\Gamma_1(N) \backslash \mathcal{H}} f(-\bar{\tau}) g(\tau) E_{1/N}^{(0)}(\tau, 0) dx \wedge dy \right) \\ &= 2 \int_{\Gamma_1(N) \backslash \mathcal{H}} f(-\bar{\tau}) g(\tau) \log |g_{0,1/N}(\tau)| dx \wedge dy \\ &= 2 \int_{\Gamma_1(N) \backslash \mathcal{H}} f(-\bar{\tau}) g(\tau) \log |g_{0,1/N}(\tau)| \frac{(-2\pi i d\bar{z}) \wedge (2\pi i dz)}{8\pi^2 i} \\ &= \frac{1}{2\pi} \left(\frac{1}{2\pi i} \int_{Y_1(N)(\mathbb{C})} \log |g_{0,1/N}| \eta_f^{\text{ah}} \wedge \omega_g \right). \end{aligned}$$

We compare this with Beilinson’s formula for the regulator on $\text{CH}^2(X_1(N)^2, 1)$ (Theorem 4.3.1). We know that $\tilde{\Xi}_N$ can be written as the class of $(\Delta, g_{0,1/N})$ (where $\Delta = C_{1,N,1}$ is the diagonal in $X_1(N)^2$) plus a linear combination of elements supported on cuspidal components. It is clear that $p_1^*(\eta_f^{\text{ah}}) \wedge p_2^*(\omega_g)$ restricts to 0 on any horizontal or vertical component and to $\eta_f^{\text{ah}} \wedge \omega_g$ on Δ ; so we obtain

$$\langle \text{reg}_{\mathbb{C}}(\tilde{\Xi}_N), \eta_f^{\text{ah}} \wedge \omega_g \rangle = \frac{1}{2\pi i} \int_{Y_1(N)} \log |g_{0,1/N}| \eta_f^{\text{ah}} \wedge \omega_g = 2\pi \mathcal{D}'(f, g, 1)$$

as required. The final equality follows from equation (11). □

We are interested in a version of Theorem 4.3.4 for $m \geq 1$, incorporating twists by Dirichlet characters. This relation becomes easier to state if we introduce “equivariant” versions of some of our objects, as follows:

Definition 4.3.5. For $N \geq 5, m \geq 1$ as above and for cusp forms f, g of level N that are eigenforms for the Hecke operators away from N , we define

the following elements:

$$\mathbf{g}_m = \sum_{a \in (\mathbb{Z}/m\mathbb{Z})^\times} [a]^{-1} \otimes g\left(z + \frac{a}{m}\right) \in \mathbb{C}[(\mathbb{Z}/m\mathbb{Z})^\times] \otimes_{\mathbb{C}} S_2(\Gamma_1(m^2N), \mathbb{C})$$

and the $\mathbb{C}[(\mathbb{Z}/m\mathbb{Z})^\times]$ -valued Dirichlet series

$$L_{(mN)}(f, g, (\mathbb{Z}/m\mathbb{Z})^\times, s) = \prod_{\ell \mid mN} P_\ell(f, g, [\ell]\ell^{-s})^{-1}.$$

(There is no obvious way to define an equivariant Euler factor at the primes dividing m .)

PROPOSITION 4.3.6.

(a) *We have*

$$a_n(\mathbf{g}_m) = a_n(g)\tau(n, m),$$

where $\tau(n, m)$ is the “universal Gauss sum” $\sum_{a \in (\mathbb{Z}/m\mathbb{Z})^\times} [a]^{-1} e^{2\pi i na/m} \in \mathbb{C}[(\mathbb{Z}/m\mathbb{Z})^\times]$.

(b) *If we extend the Hecke operators on $S_2(\Gamma_1(N))$ linearly to $\mathbb{C}[(\mathbb{Z}/m\mathbb{Z})^\times] \otimes_{\mathbb{C}} S_2(\Gamma_1(m^2N))$, then we have*

$$\begin{aligned} T_n(\mathbf{g}_m) &= [n]t_g(n)\mathbf{g}_m \\ \langle n \rangle(\mathbf{g}_m) &= [n]^2\varepsilon_g(m)\mathbf{g}_m \end{aligned}$$

for all n such that $(n, mN) = 1$, where $t_g(n)$ and $\varepsilon_g(n)$ are the eigenvalues of g for the T_n and $\langle n \rangle$ operators respectively.

(We can interpret (b) above as stating that \mathbf{g}_m transforms under the Hecke operators away from mN as “ g twisted by the universal character of level m .”)

Proof. Part (a) is immediate by a q -expansion computation. For part (b), we note that the statement regarding the diamond operators can be verified directly — by essentially the same computation as Proposition 2.7.5(4) — and the statement for the T_n ’s now follows immediately from the standard formulae for the action of T_n on q -expansions, together with the easily verified fact that $\tau(nn', m) = [n]\tau(n', m)$ if $(n, m) = 1$. \square

We extend the Beilinson regulator to a homomorphism

$$\mathrm{reg}_{\mathbb{C}[(\mathbb{Z}/m\mathbb{Z})^\times]} : \mathrm{CH}^2(X_1(N) \otimes \mathbb{Q}(\mu_m), 1) \rightarrow \mathbb{C}[(\mathbb{Z}/m\mathbb{Z})^\times] \otimes_{\mathbb{C}} \left(\mathrm{Fil}^1 H_{\mathrm{dR}}^2(X/\mathbb{C})\right)^\vee$$

by mapping δ to $\sum_{a \in (\mathbb{Z}/m\mathbb{Z})^\times} [a] \otimes \mathrm{reg}_{\mathbb{C}}(\sigma_a \cdot \delta)$. (Note that this is *not* a homomorphism of modules over the group ring $\mathbb{C}[(\mathbb{Z}/m\mathbb{Z})^\times]$; the Poincaré duality pairing interchanges the natural action of $\mathbb{C}[(\mathbb{Z}/m\mathbb{Z})^\times]$ with its inverse.)

THEOREM 4.3.7. *Let \check{f}, \check{g} be as above, and let $\widetilde{\Xi}_{m,N,1}$ be any lifting of $\Xi_{m,N,1}$ to $X_1(N)^2$. Then as elements of $\mathbb{C}[(\mathbb{Z}/m\mathbb{Z})^\times]$, we have*

$$\begin{aligned} \left\langle \text{reg}_{\mathbb{C}[(\mathbb{Z}/m\mathbb{Z})^\times]}(\widetilde{\Xi}_{m,N,1}), p_1^*(\eta_{\check{f}}^{\text{ah}}) \wedge p_2^*(\omega_{\check{g}}) \right\rangle \\ = \frac{1}{2} L'_{(mN)}(f, g, (\mathbb{Z}/m\mathbb{Z})^\times, 1) A(\check{f}, \check{g}, m, 1), \end{aligned}$$

where we define

$$A(\check{f}, \check{g}, m, s) = \sum_{a \in (\mathbb{Z}/m\mathbb{Z})^\times} [a]^{-1} \sum_{n \in S(mN)} a_n(f) a_n(g) e^{2\pi i a n / m} n^{-s}.$$

Proof. As we showed in the previous section, $\widetilde{\Xi}_{m,N,j}$ may be represented as the class of an element in $Z^2(X_1(N)^2 \otimes \mathbb{Q}(\mu_m), 1) \otimes \mathbb{Q}[(\mathbb{Z}/m\mathbb{Z})^\times]$ that differs by negligible elements from

$$(C_{m,N,j}, (\iota_{m,N,j})^*(g_{0,1/m^2N})).$$

As in the case $m = 1$ considered above, these negligible elements pair to 0 with the differential $p_1^*(\eta_{\check{f}}^{\text{ah}}) \wedge p_2^*(\omega_g)$. Hence we have

$$\begin{aligned} \left\langle \text{reg}_{\mathbb{C}(\widetilde{\Xi}_{m,N})}, p_1^*(\eta_{\check{f}}^{\text{ah}}) \wedge p_2^*(\omega_g) \right\rangle \\ = \sum_{j \in (\mathbb{Z}/m\mathbb{Z})^\times} [j]^{-1} \int_{C_{m,N,j}} \log \left| (\iota_{m,N,j})^*(g_{0,1/m^2N}) \right| \cdot p_1^*(\eta_{\check{f}}^{\text{ah}}) \wedge p_2^*(\omega_g) \\ = \sum_{j \in (\mathbb{Z}/m\mathbb{Z})^\times} [j]^{-1} \int_{X_1(m^2N)} \log \left| g_{0,1/m^2N} \right| \\ \cdot (p_1 \circ \iota_{m,N,j})^*(\eta_{\check{f}}^{\text{ah}}) \wedge (p_2 \circ \iota_{m,N,j})^*(\omega_g). \end{aligned}$$

By construction, $p_1 \circ \iota_{m,N,j}$ is just the natural projection map $X_1(m^2N) \rightarrow X_1(N)$, so the pullback of $\eta_{\check{f}}^{\text{ah}}$ along this map is just $\eta_{\check{f}}^{\text{ah}}$ again (where now we consider f as a modular form of level m^2N). On the other hand, $p_2 \circ \iota_{m,N,j}$ corresponds to the map $z \mapsto z + \frac{j}{m}$ on the upper half-plane, so $(p_2 \circ \iota_{m,N,j})^*(\omega_g)$ is the differential whose pullback to \mathcal{H} is $2\pi i g(z + \frac{j}{m}) dz$, and hence we have

$$\sum_j [j]^{-1} (p_2 \circ \iota_{m,N,j})^*(\omega_g) = 2\pi i \mathbf{g}_m(z) dz$$

as elements of $\mathbb{C}[(\mathbb{Z}/m\mathbb{Z})^\times] \otimes \Omega_{\text{hol}}^1(X_1(m^2N))$. Hence, by exactly the same computation as above,

$$\left\langle \text{reg}_{\mathbb{C}(\widetilde{\Xi}_{m,N})}, p_1^*(\eta_{\check{f}}^{\text{ah}}) \wedge p_2^*(\omega_g) \right\rangle = 4\pi \int_{X_1(m^2N)} f(-\bar{\tau}) \mathbf{g}_m(\tau) \log \left| g_{0,1/m^2N} \right| dx \wedge dy.$$

As remarked above, \mathbf{g}_m is an eigenform for the Hecke operators away from mN ; so we may now apply exactly the same formal manipulations as in the proof of [Kat04, Prop. 7.1], but with group ring coefficients rather than \mathbb{C} coefficients, and the result follows in this case also. \square

4.4. *A nonvanishing result.* In this section, we shall use the results of the previous section, together with a deep theorem of Shahidi on the nonvanishing of Rankin–Selberg L -values, to show that the elements $\Xi_{m,N,j}$ are not all zero (which is in no way obvious from their construction).

THEOREM 4.4.1 (Shahidi; [Sha81, Th. 5.2]). *Let f, g be any two newforms of weight 2. Then the completed L -function $\Lambda(f, g, s)$ is holomorphic and nonvanishing on the line $\Re(s) = 2$, unless $f = g^*$, in which case it has a simple pole at $s = 2$.*

Remark 4.4.2. We have stated only a special case of Shahidi’s very general theorem, which applies to automorphic forms on $\mathrm{GL}_n \times \mathrm{GL}_m$ over an arbitrary number field. Note also that Shahidi’s normalizations are slightly different from ours (He normalizes the L -function so that the abscissa of symmetry is $s = \frac{1}{2}$, independently of the weights of f and g , while we normalize it to be at $s = \frac{k+\ell-1}{2} = \frac{3}{2}$.)

COROLLARY 4.4.3. *If Σ is a finite set of primes, the function $L_\Sigma(f, g, s)$ has a zero at $s = 1$ of order $r_1 + r_2$, where*

$$r_1 = \begin{cases} 1 & \text{if } f^* \neq g, \\ 0 & \text{if } f^* = g \end{cases}$$

and r_2 is the sum of the orders of the poles at $s = 1$ of the Euler factors $L_p(f, g, s)$ for primes $p \in \Sigma$.

Proof. Applying the functional equation for the completed L -function, which switches s with $3 - s$, we deduce from Shahidi’s result that $\Lambda(f, g, s)$ is holomorphic and nonvanishing (resp. has a simple pole) at $s = 1$ if $f \neq g^*$ (resp. if $f = g^*$).

However, the L -factor at ∞ , $L_\infty(f, g, s) = \Gamma_{\mathbb{C}}(s)\Gamma_{\mathbb{C}}(s - 1)$, has a simple pole at $s = 1$, so the order of vanishing of $L(f, g, s)$ is r_1 as defined above. Since $L_\Sigma(f, g, s)$ is $L(f, g, s)$ divided by the product of the L -factors at primes in Σ , the result clearly follows. \square

Remark 4.4.4. Note that if $f^* = g$, then the local L -factor vanishes at $s = 1$ for every prime, so r_2 will tend to be rather large in this case.

COROLLARY 4.4.5. *Let f, g be any two newforms, Σ any set of primes, and p any prime in Σ . Then for all but finitely many Dirichlet characters χ of p -power conductor, $\prod_{\ell \in \Sigma} L_\ell(f, g \otimes \chi, s)$ is holomorphic and nonzero at $s = 1$.*

Proof. If χ has sufficiently large p -power conductor, then the local L -factor of $f \otimes g \otimes \chi$ at p is identically 1; so it suffices to consider the L -factors at primes $\ell \neq p$. However, since χ has conductor prime to ℓ , $L_\ell(f \otimes g \otimes \chi, s) =$

$P_\ell(\chi(\ell)\ell^{-s})^{-1}$, so it suffices to arrange that $\chi(\ell)\ell^{-1}$ does not lie in the finite set of zeroes of the polynomial $L_\ell(f \otimes g, X)$. It is clear that this may also be achieved by ensuring that the conductor of χ is sufficiently big. \square

COROLLARY 4.4.6. *Given any two forms f, g of level N that are eigenvectors for all Hecke operators, and p any prime, there is $k \geq 0$ such that the projection of $\Xi_{mp^k, N, 1}$ to the (f, g) -isotypical quotient of $\mathrm{CH}^2(Y_1(N)^2 \otimes \mathbb{Q}(\mu_{mp^k}), 1)$ is nonzero.*

Proof. Immediate from the previous corollary and Theorem 4.3.7. \square

5. Relation to p -adic L -values

In this section we develop an analogue of the $m = 1$ case of Theorem 4.3.7 in the p -adic setting. This is essentially a variant of the main theorem of [BDR12].

5.1. *Holomorphic Eisenstein series.* We begin by constructing some holomorphic Eisenstein series that may be defined over a number field. We follow Chapter 3 of [Kat04] closely, but we work on $Y_1(N)$ rather than $Y(N)$. Our purpose is to define, for $\alpha \in \mathbb{Q}/\mathbb{Z}$, the following modular forms:

- $E_\alpha^{(k)} \in M_k(\Gamma_1(N))$, where $k \geq 1$, $k \neq 2$;
- $\widetilde{E}_\alpha^{(2)} \in M_2(\Gamma_1(N))$;
- $F_\alpha^{(k)} \in M_k(\Gamma_1(N))$, for $k \geq 1$, with $\alpha \neq 0$ if $k = 2$.

We set

$$E_\alpha^{(k)}(\tau) = E_\alpha^{(k)}(\tau, 0),$$

and similarly for $F^{(k)}$.

PROPOSITION 5.1.1. *If $k \geq 1$, $k \neq 2$, then $E_\alpha^{(k)}, F_\alpha^{(k)} \in M_k(\Gamma_1(N))$ for any $\alpha \in \frac{1}{N}\mathbb{Z}/\mathbb{Z}$.*

For $k = 2$, we have $F_\alpha^{(2)} \in M_2(\Gamma_1(N))$ for any $\alpha \neq 0$, and $\widetilde{E}_\alpha^{(2)} := E_\alpha^{(2)} - E_0^{(2)} \in M_2(\Gamma_1(N))$ (for any α). The function $F_0^{(2)} = E_0^{(2)}$ is a C^∞ function on \mathcal{H} invariant under the weight 2 action of $\Gamma_1(N)$, with slow growth at the cusps, but is not holomorphic.

Proof. See [Kat04, §3.8]; our $E_\alpha^{(k)}$ is Kato’s $E_{0,\alpha}^{(k)}$. \square

We have q -expansion formulae for both families. Let $\alpha \in \mathbb{Q}/\mathbb{Z}$. For $\Re(s) > 1$, define

$$\zeta(\alpha, s) = \sum_{\substack{n \in \mathbb{Q}, n > 0 \\ n = \alpha \bmod \mathbb{Z}}} n^{-s} \quad \text{and} \quad \zeta^*(\alpha, s) = \sum_{n=1}^{\infty} e^{2\pi i \alpha n} n^{-s}$$

as in [Kat04, §3.9]. Then both $\zeta(\alpha, s)$ and $\zeta^*(\alpha, s)$ have meromorphic continuation to all $s \in \mathbb{C}$ and satisfy

$$\zeta^*(\alpha, 1-s) = \frac{\Gamma(s)}{(2\pi)^s} \left(e^{-i\pi s/2} \zeta(-\alpha, s) + e^{i\pi s/2} \zeta(\alpha, s) \right),$$

a version of the standard functional equation for the Hurwitz zeta function.

PROPOSITION 5.1.2. *Let $k \geq 1$, $\alpha \in \mathbb{Q}/\mathbb{Z}$.*

(1) *Assume $k \neq 2$. Then we have*

$$E_\alpha^{(k)} = a_0 + \sum_{n \geq 1} \left(\sum_{d|n} d^{k-1} (e^{2\pi i \alpha d} + (-1)^k e^{-2\pi i \alpha d}) \right) q^n,$$

where

$$a_0 = \begin{cases} \zeta^*(\alpha, 1-k) & \text{if } k \geq 3, \\ \frac{1}{2} (\zeta^*(\alpha, 0) - \zeta^*(-\alpha, 0)) & \text{if } k = 1. \end{cases}$$

(2) *We have*

$$\tilde{E}_\alpha^{(2)} = a_0 + \sum_{n \geq 1} \left(\sum_{d|n} d (e^{2\pi i \alpha d} + e^{-2\pi i \alpha d} - 2) \right) q^n,$$

where $a_0 = \zeta^*(\alpha, -1) + \frac{1}{12}$.

(3) *Assume $\alpha \neq 0$ in the case $k = 2$. Then*

$$F_\alpha^{(k)} = a_0 + \sum_{n \geq 1} \left(\sum_{d|n} \binom{n}{d}^{k-1} (e^{2\pi i \alpha d} + (-1)^k e^{-2\pi i \alpha d}) \right) q^n,$$

where

$$a_0 = \begin{cases} \zeta(1-k) & \text{if } k \geq 2, \\ \frac{1}{2} (\zeta^*(\alpha, 0) - \zeta^*(-\alpha, 0)) & \text{if } k = 1. \end{cases}$$

Proof. This is [Kat04, Prop. 3.10]. Note that there is a typographical error in the statement of the proposition loc. cit.; there is an extra star in the formula for $\sum a_n n^{-s}$ in case (1), and the formula should read

$$\sum_{n \in \mathbb{Q}, n > 0} a_n n^{-s} = \zeta(\alpha, s) \zeta^*(\beta, s-k+1) + (-1)^k \zeta(-\alpha, s) \zeta^*(-\beta, s-k+1). \quad \square$$

5.2. *Nearly holomorphic modular forms.* For $k \geq 0$, we define (following, e.g., [Shi86], [Shi00]) the space of *nearly holomorphic* modular forms $M_k^{\text{nh}}(\Gamma_1(N), \mathbb{C})$. This is the space of C^∞ slowly-increasing functions on \mathcal{H} that are invariant under the weight k action of $\Gamma_1(N)$ and are annihilated by some power of the Maass–Shimura weight-lowering differential operator

$$\varepsilon_k = \frac{-1}{2\pi i} \mathfrak{S}(\tau)^2 \frac{d}{d\bar{\tau}}.$$

Any such function is in fact annihilated by $\varepsilon^{[k/2]+1}$ and can be expanded as a finite sum

$$(14) \quad f(\tau) = \sum_{j=0}^{[k/2]} f_j(\tau) (\pi\Im(\tau))^{-j},$$

where the f_j are holomorphic functions. (In particular, any nearly holomorphic form of weight 0 or 1 is in fact a holomorphic form.) For K a number field containing the N -th roots of unity,¹ we shall say that $f \in M_k^{\text{nh}}(\Gamma_1(N), K)$ is defined over K if the Fourier coefficients of the holomorphic functions f_j are in K , and we write $M_k^{\text{nh}}(\Gamma_1(N), K)$ for the space of such functions.

We let $S_k^{\text{nh}}(\Gamma_1(N), K)$ be the subspace of rapidly decreasing functions in $M_k^{\text{nh}}(\Gamma_1(N), K)$. If $k \geq 2$, then $S_k^{\text{nh}}(\Gamma_1(N), K)$ coincides with the space defined algebraically in [DR14, §2.4] using the ‘‘Hodge splitting’’ of the de Rham cohomology.

COROLLARY 5.2.1. *Let k, j be integers with $k \geq 1$ and $j \in [0, k - 1]$. Then for any $\alpha \in \frac{1}{N}\mathbb{Z}/\mathbb{Z}$, the function $\tau \mapsto E_\alpha^{(k)}(\tau, -j)$ lies in $M_k^{\text{nh}}(\Gamma_1(N), \mathbb{Q}(\mu_N))$.*

Proof. We first note that $E_\alpha^{(k)}, F_\alpha^{(k)} \in M_k^{\text{nh}}(\Gamma_1(N), K)$ for all $k \geq 1$. This is clear for $k = 1$ or $k \geq 3$ (since holomorphic forms are certainly nearly holomorphic), and for $k = 2$, it suffices to check that $E_0^{(2)} = F_0^{(2)}$ is nearly holomorphic, which is clear from the formula

$$E_0^{(2)}(\tau) = \frac{1}{4\pi\Im(\tau)} - \frac{1}{12} + 2 \sum_{n \geq 1} \left(\sum_{d|n} d \right) q^n.$$

With this in hand, we obtain the near-holomorphy of $E_\alpha^{(k)}(\tau, -j)$ for $0 \leq j \leq \frac{k-1}{2}$ by applying δ^j to $E_\alpha^{(k-2j)} = E_\alpha^{(k-2j)}(\tau, 0)$. Applying the same argument to $F_\alpha^{(k-2j)}$ shows that $F_\alpha^{(k)}(\tau, -j)$ is nearly holomorphic for j in the same range; but $F_\alpha^{(k)}(\tau, -j) = E_\alpha^{(k)}(\tau, 1 - k + j)$ by the functional equation, as required, so we obtain the result for all $j \in [0, k - 1]$. \square

We define the q -expansion of a nearly-holomorphic modular form f to be the Fourier expansion of the holomorphic \mathbb{Z} -periodic function f_0 , when we write f in the form (14). Then δ corresponds to $q \frac{d}{dq}$ on q -expansions, and the following is clear from Proposition 5.1.2 and the proof of the previous proposition:

¹This is for compatibility with our notation for classical modular forms, since in our model of $Y_1(N)$, the cusp ∞ is not rational.

PROPOSITION 5.2.2. *For any $j \in [0, k - 1]$, the q -expansion of the nearly-holomorphic form $E_\alpha^{(k)}(\tau, -j)$ is*

$$a_0 + \sum_{n \geq 1} \left(\sum_{d|n} d^{k-1-j} \left(\frac{n}{d}\right)^j (e^{2\pi i \alpha d} + (-1)^k e^{-2\pi i \alpha d}) \right) q^n,$$

where $a_0 = 0$ unless $j \in \{0, k - 1\}$.

5.3. *p -adic families of Eisenstein series.* We now use the q -expansion formulae above as motivation for defining a two-parameter family of p -adic Eisenstein series, which can be regarded as an analogue of the $E^{(k)}(-, s)$, with two continuous p -adic parameters ϕ_1, ϕ_2 replacing the discrete parameter k and the continuous real-analytic parameter s .

Definition 5.3.1. Choose some (sufficiently large) finite extension L/\mathbb{Q}_p , and let $\Lambda = \mathcal{O}_L[[\mathbb{Z}_p^\times]]$, the Iwasawa algebra of \mathbb{Z}_p^\times . Let $\Omega = \text{Spf } \Lambda$, the *weight space* classifying continuous characters $\phi : \mathbb{Z}_p^\times \rightarrow \mathbb{C}_p$, and consider the formal power series with coefficients in $\Lambda \widehat{\otimes} \Lambda$ given by

$$\mathcal{E}_\alpha(\phi_1, \phi_2) = \sum_{\substack{n \geq 1 \\ p \nmid n}} \left(\sum_{d|n} \phi_1(d) \phi_2\left(\frac{n}{d}\right) [e^{2\pi i \alpha d} + \varepsilon e^{-2\pi i \alpha d}] \right) q^n \in \mathcal{O}_L[[q]],$$

where $\varepsilon = -\phi_1(-1)\phi_2(-1)$.

Note 5.3.2. We consider $\mathbb{Z} \times \mathbb{Z}$ as a subset of $\Omega \times \Omega$ in the natural way. Then for $k \geq 1, k \neq 2$, we have $\mathcal{E}_\alpha(k - 1, 0) = (E_\alpha^{(k)})^{[p]}$ and $\mathcal{E}_\alpha(0, k - 1) = (F_\alpha^{(k)})^{[p]}$, where $(-)^{[p]}$ denotes the “ p -depletion” operator.

We now fix a newform $g \in S_\ell(\Gamma_1(N_g))$ for some ℓ and some $N_g \mid N$. (Although the weight ℓ will be fixed in our discussion, it is convenient to keep it in the notation as a parameter, since this will make our notation more consistent with [BDR12].) We will write \check{g} for any element of $S_\ell(N)[\pi_g]$.

Definition 5.3.3. For integers k, j , we define

$$\Xi(k, \ell, \alpha, j)^{\text{ord}, p} = e_{\text{ord}}[\mathcal{E}_\alpha(j - \ell, k - 1 - j) \cdot \check{g}].$$

This is an ordinary Λ -adic family of modular forms, parametrized by k and j . (We are taking ℓ to be fixed here, in order to avoid the need to make any ordinarity hypotheses on g .) For any (k, j) , $\Xi(k, \ell, \alpha, j)^{\text{ord}, p}$ is a p -adic modular form of weight k .

We now compare this with the complex-analytic theory. It is clear that $\mathcal{E}_\alpha(j - \ell, k - 1 - j)$ is the p -depletion of the nearly-overconvergent form $\tau \mapsto E^{(k-\ell)}(\tau, -k + j + 1)$.

Definition 5.3.4. For $\ell \leq j \leq k - 1$, let $\Xi(k, \ell, \alpha, j)$ denote the nearly-holomorphic modular form of weight k given by

$$\Xi(k, \ell, \alpha, j)(\tau) = E_\alpha^{(k-\ell)}(\tau, -k + j + 1) \cdot \check{g},$$

and let $\Xi(k, \ell, \alpha, j)^{\text{hol}}$ be its image under the holomorphic projector.

Notation. Let $H^1(Y_1(N), \mathcal{L}_{k-2}, \nabla)$ denote the de Rham cohomology of $Y_1(N)$ with coefficients in the $(k - 2)$ -nd symmetric power of the relative de Rham cohomology sheaf of the universal elliptic curve over $Y_1(N)$, endowed with its Gauss–Manin connection.

PROPOSITION 5.3.5. *Let k, ℓ be fixed, with $k > \ell$. Let f be a newform in $S_k(\Gamma_1(N_f))$, for some $N_f \mid N$, and let e_{f^*} be the projection to the f^* -isotypic component in the Hecke algebra acting on $S_k(\Gamma_1(N))$. Assume f is ordinary at p , and let $j \in [\ell, k - 1]$. Then we have the following relation in $H^1(Y_1(N), \mathcal{L}_{k-2}, \nabla)$:*

$$e_{f^*} \Xi(k, \ell, \alpha, j)^{\text{ord}, p} = \frac{\mathcal{E}(f, g, j)}{\mathcal{E}(f)} e_{f^*} e_{\text{ord}} \Xi(k, \ell, \alpha, j)^{\text{hol}},$$

where

$$\mathcal{E}(f) = 1 - p^{-1} \beta_p(f) \alpha_p(f)^{-1}$$

and

$$\begin{aligned} \mathcal{E}(f, g, j) &= (1 - p^{-j} \beta_p(f) \alpha_p(g))(1 - p^{-j} \beta_p(f) \beta_p(g)) \\ &\quad \times (1 - p^{j-1} \alpha_p(f)^{-1} \alpha_p(g)^{-1})(1 - p^{j-1} \alpha_p(f)^{-1} \beta_p(g)^{-1}). \end{aligned}$$

Here α_f, β_f are the roots of the Hecke polynomial of f at p , and similarly for g .

Proof. This follows from Corollary 4.13 of [DR14] with the f, g, h of the theorem taken to be $f, E_\alpha^{(k-\ell)}(-, -k + j + 1)$ and g . (Note that the special case $j \geq \frac{k+\ell-1}{2}$ is [BDR12, Prop. 2.7].) \square

5.4. Interpolation in Hida families. We now interpolate the left-hand side of Proposition 5.3.5 in Hida families.

Notation. Let f be a newform (of some level $N_f \mid N$), and let \mathbf{f} be the Hida family through f (with coefficients in some finite flat Λ -algebra Λ_f). Then we define the space $\mathbf{S}^{\text{ord}}(N; \Lambda_f)[\pi_f]$ for the Λ_f -module of families of oldforms at level N corresponding to f , which is simply the space of formal q -expansions spanned over Λ_f by $\mathbf{f}(q^d)$ for $d \mid N/N_f$. We write $\check{\mathbf{f}}$ for a generic element of $\mathbf{S}^{\text{ord}}(N; \Lambda_f)[\pi_f]$, which we shall think of as a “test vector” associated to \mathbf{f} .

We shall continue to write g for a newform in $S_\ell(N_g)$ for some $N_g \mid N$, and \check{g} for a generic element of $S_\ell(N; K)[\pi_g]$.

PROPOSITION 5.4.1. *For any $\check{\mathbf{f}} \in \mathbf{S}^{\text{ord}}(N; \Lambda_f)[\pi_f]$ and $\check{g} \in S_\ell(N; K)[\pi_g]$ as above, and any $\alpha \in \frac{1}{N}\mathbb{Z}/\mathbb{Z}$, there exists an element*

$$\mathcal{D}_p(\check{\mathbf{f}}, \check{g}, \alpha) \in \text{Frac}(\Lambda_f) \widehat{\otimes} \Lambda$$

such that for all integers k, j with $k \geq 2$, we have

$$\mathcal{D}_p(\check{\mathbf{f}}, \check{g}, \alpha)(k, j) = \frac{\langle \check{f}_k^*, \Xi(k, \ell, \alpha, j)^{\text{ord}, p} \rangle}{\mathcal{E}^*(f_k) \langle f_k, f_k \rangle},$$

where f_k and \check{f}_k are the eigenforms at level N whose ordinary p -stabilizations are the weight k specializations of \mathbf{f} and $\check{\mathbf{f}}$, and

$$\mathcal{E}^*(f_k) := 1 - \beta_p(f_k) \alpha_p(f_k)^{-1}.$$

Combining this with the previous proposition, we have

PROPOSITION 5.4.2. *For integers k, j with $\ell \leq j \leq k - 1$, we have*

$$\mathcal{D}_p(\check{\mathbf{f}}, \check{g}, \alpha)(k, j) = \frac{\mathcal{E}(f_k, g, j)}{\mathcal{E}(f_k) \cdot \mathcal{E}^*(f_k) \cdot \langle f_k, f_k \rangle} \mathcal{D}(\check{f}_k, \check{g}, \alpha, j).$$

Note 5.4.3. We know that the Atkin–Lehner operator gives an isomorphism

$$S_\ell(N; K)[\pi_g] \xrightarrow[\cong]{w_N} S_\ell(N; K)[\pi_{g^*}].$$

Less obviously, there is also an operator

$$\mathbf{S}^{\text{ord}}(N; \Lambda_f)[\pi_f] \xrightarrow[\cong]{w_N} \mathbf{S}^{\text{ord}}(N; \Lambda_f)[\pi_{f^*}]$$

interpolating the action of the Atkin–Lehner operators on the weight k specializations. To see this, it suffices to note that the inclusions $S_k(N) \hookrightarrow S_k(Np) \hookrightarrow S_k(Np^2) \hookrightarrow \dots$ commute with the action of w_N and this operator is continuous with respect to the p -adic norm (by the q -expansion principle); the resulting operator on the completion $S_k(Np^\infty)$ commutes with U_p and hence preserves $e_{\text{ord}} S_k(Np^\infty)$.

PROPOSITION 5.4.4. *For any $k, j \in \Omega_f \times \Omega$, we have*

$$\mathcal{D}_p(\check{\mathbf{f}}, \check{g}, \alpha)(k, k + \ell - 1 - j) = N^{1-j} \cdot \sum_{y \in \mathbb{Z}/N\mathbb{Z}} e^{2\pi i \alpha x / N} \mathcal{D}_p(w_N \check{\mathbf{f}}, w_N \check{g}, x/N)(k, j).$$

Proof. It suffices to check this result for all pairs of integers k, j with $k \geq j$ since these points are Zariski-dense in $\Omega_f \times \Omega$. By the classical functional equation, we find that for such k, j , we have

$$\mathcal{D}_p(\check{\mathbf{f}}, \check{g}, \alpha)(k, k + \ell - 1 - j) = A \cdot N^{1-j} \sum_{y \in \mathbb{Z}/N\mathbb{Z}} e^{2\pi i \alpha x / N} \mathcal{D}_p(w_N \check{\mathbf{f}}, w_N \check{g}, \alpha)(k, j),$$

where the quantity A is defined by

$$A = \frac{\mathcal{E}(f_k, g, k + \ell - 1 - j)}{\mathcal{E}(f_k)\mathcal{E}^*(f_k)\langle f_k, f_k \rangle} \cdot \left(\frac{\mathcal{E}(f_k^*, g^*, j)}{\mathcal{E}(f_k^*)\mathcal{E}^*(f_k^*)\langle f_k^*, f_k^* \rangle} \right)^{-1}.$$

We obviously have $\langle f_k^*, f_k^* \rangle = \langle f_k, f_k \rangle$. More subtly, we have $\alpha_p(f^*) = p^{k-1}/\beta_p(f)$ and $\beta_p(f^*) = p^{k-1}/\alpha_p(f)$; similarly, we have $\{\alpha_p(g^*), \beta_p(g^*)\} = \{p^{\ell-1}/\alpha_p(g), p^{\ell-1}/\beta_p(g)\}$. From these relations, it is clear that $\mathcal{E}(f_k^*) = \mathcal{E}(f_k)$, $\mathcal{E}^*(f_k^*) = \mathcal{E}^*(f_k)$, and $\mathcal{E}(f_k, g, k + \ell - 1 - j) = \mathcal{E}(f_k^*, g^*, j)$. So the ratio A is identically 1. \square

Notation. We write $\mathcal{D}_p(\check{f}, \check{g}, \alpha)$ for the restriction of $\mathcal{D}_p(\mathbf{f}, \mathbf{g}, \alpha)$ to $k \times \Omega \subset \Omega_f \times \Omega$, where $\check{\mathbf{f}}$ is any element of $\mathbf{S}^{\text{ord}}(N; \Lambda_f)[\pi_f]$ whose specialization in weight k is \check{f} . (This is independent of the choice of family $\check{\mathbf{f}}$.)

Note 5.4.5. If $k > \ell$, then the L -function $\mathcal{D}_p(\check{f}, \check{g}, \alpha)$ interpolates the critical values $\mathcal{D}(f, g, \alpha, s)$; but when $k = \ell$, there are no such critical values.

5.5. The syntomic regulator. Let p be prime and K be a finite extension of \mathbb{Q}_p with ring of integers \mathcal{O}_K , and let \mathcal{X} be a smooth proper scheme over \mathcal{O}_K with generic fibre X . Then there exists a map, the *syntomic regulator*,

$$r_{\text{syn}} : \text{CH}^2(\mathcal{X}, 1) \rightarrow H_{\text{dR}}^2(X/K)/\text{Fil}^2 = \left(\text{Fil}^1 H_{\text{dR}}^2(X/K) \right)^\vee,$$

with the property that the diagram

$$\begin{array}{ccc} \text{CH}^2(\mathcal{X}, 1) & \longrightarrow & \text{CH}^2(X, 1) \\ \downarrow r_{\text{syn}} & & \downarrow r_{\text{ét}} \\ H_{\text{dR}}^2(X/K)/\text{Fil}^2 & \xrightarrow{\text{exp}} & H^1(K, H_{\text{ét}}^2(\overline{X}, \mathbb{Q}_p)(2)) \end{array}$$

commutes (cf. [Bes00]). Here exp denotes the Bloch–Kato exponential map constructed in [BK90] for the crystalline G_K -representation $V = H_{\text{ét}}^2(\overline{X}, \mathbb{Q}_p)(2)$.

Remark 5.5.1. Note that since \mathcal{X} has good reduction, all eigenvalues of Frobenius on $\mathbb{D}_{\text{cris}}(V)$ are Weil numbers of weight -2 ; thus $\mathbb{D}_{\text{cris}}(V)^{\varphi=1} = 0$, implying that exp is injective. This also implies that $H_e^1(K, V) = H_f^1(K, V)$. Note that we do *not* necessarily have $H_f^1(K, V) = H_g^1(K, V)$ since $V^*(1)$ has all weights equal to 0. It is conjectured that the image of $r_{\text{ét}}$ is precisely $H_g^1(K, V)$, but this is only known in a few special cases; cf. [SS10, Fact 1.1].

5.6. Generalization of a theorem of Bertolini–Darmon–Rotger. Note that there is a map

$$\text{dlog} : \mathcal{O}(Y_1(N))^\times \otimes \mathbb{Q} \rightarrow M_2(\Gamma_1(N)),$$

which corresponds to $F(\tau) \mapsto \frac{F'(\tau)}{F(\tau)}$ as functions on \mathcal{H} , and this commutes with the Atkin–Lehner involutions.

PROPOSITION 5.6.1. *For any $\alpha \neq 0 \in \mathbb{Q}/\mathbb{Z}$, we have*

$$\mathrm{dlog} g_{0,\alpha} = -F_\alpha^{(2)}.$$

Proof. Immediate from comparing the q -expansion of $F_\alpha^{(2)}$ with that of $g_{0,\alpha}$, which is given in [Kat04, §1.9]. \square

We recall the following result, which is a slight reformulation and extension of the main theorem of [BDR12]:

THEOREM 5.6.2. *Let u_α be the modular unit on $\mathcal{Y}_1(N) \otimes \mathbb{Z}(\mu_N)$ such that*

$$\mathrm{dlog} u_\alpha = \widetilde{E}_\alpha^{(2)},$$

and let Δ_{u_α} be any element of $\mathrm{CH}^2(\mathcal{X}_1(N) \otimes \mathbb{Z}(\mu_N), 1)$ whose pullback to $\mathrm{CH}^2(\mathcal{Y}_1(N) \otimes \mathbb{Z}(\mu_N), 1)$ is the class of (Δ, u_α) , where Δ is the diagonal subvariety.

Let f, g be any two newforms of weight 2 and levels N_f, N_g dividing N , with f ordinary at p , and let \check{f}, \check{g} be test vectors attached to f, g as before. Then we have

$$\left(\mathcal{D}_p(\check{f}, \check{g}, \alpha)(2) - \mathcal{D}_p(\check{f}, \check{g}, 0)(2) \right) = \frac{\mathcal{E}(f, g, 2)}{\mathcal{E}(f) \cdot \mathcal{E}^*(f)} \left\langle r_{\mathrm{syn}}(\Delta_{u_\alpha}), \mathrm{pr}_1^*(\eta_{\check{f}}^{\mathrm{ur}}) \wedge \mathrm{pr}_2^*(\omega_{\check{g}}) \right\rangle.$$

Here, as in Proposition 4.6 of [DR14], $\eta_{\check{f}}^{\mathrm{ur}}$ denotes the unique class in $H_{\mathrm{dR}}^1(X_1(N)/\mathbb{Q}_p)$ that lies in the $\varphi = \alpha_p(f)$ eigenspace and whose image in $H^1(X_1(N), \mathcal{O}_{X_1(N)})$ agrees with that of $\frac{1}{\langle \check{f}, \check{f} \rangle} \eta_{\check{f}}^{\mathrm{ah}}$, where $\eta_{\check{f}}^{\mathrm{ah}}$ is as in Definition 4.3.2 above.

Proof. By Fourier inversion on the multiplicative group $(\mathbb{Z}/N\mathbb{Z})^\times$, which acts on both sides of the claimed formula, it suffices to show that for each Dirichlet character ψ modulo N , we have

$$(15) \quad \frac{\mathcal{E}(f, g, 2)}{\mathcal{E}(f) \cdot \mathcal{E}^*(f)} \left\langle r_{\mathrm{syn}}(\Delta_{u_\psi}), \mathrm{pr}_1^*(\eta_{\check{f}}^{\mathrm{ur}}) \wedge \mathrm{pr}_2^*(\omega_{\check{g}}) \right\rangle \\ = \begin{cases} \sum_{d \in (\mathbb{Z}/N\mathbb{Z})^\times} \psi(d)^{-1} \mathcal{D}_p(\check{f}, \check{g}, d\alpha)(2) & \text{if } \psi \neq 1, \\ \sum_{d \in (\mathbb{Z}/N\mathbb{Z})^\times} \left(\mathcal{D}_p(\check{f}, \check{g}, d\alpha)(2) - \mathcal{D}_p(\check{f}, \check{g}, 0)(2) \right) & \text{if } \psi = 1, \end{cases}$$

where $u_\psi = \sum_d \psi(d)^{-1} \otimes u_{d\alpha} \in \mathbb{Z}(\chi) \otimes_{\mathbb{Z}} \mathcal{O}(Y_1(N)^\times)$. However, it is clear that both sides of equation (15) are zero unless $\psi = \chi := \chi_f^{-1} \chi_g^{-1}$, so we may assume $\psi = \chi$.

If $\chi \neq 1$ and α has exact order N , then we can assume without loss of generality that $\alpha = 1/N$, and we are in the case studied in [BDR12]. In the remaining cases, the argument goes through essentially identically. \square

Remark 5.6.3. If in fact χ is primitive modulo N , then both sides are zero unless α has exact order N , so we may reduce to precisely the case covered by [BDR12].

We can now deduce our main theorem of this section.

THEOREM 5.6.4. *Let $f, g, \check{f}, \check{g}$ be as above. Then we have*

$$\mathcal{D}_p(\check{f}, \check{g}, 1/N)(1) = -\frac{\mathcal{E}(f, g, 1)}{\mathcal{E}(f) \cdot \mathcal{E}^*(f)} \left\langle r_{\text{syn}}(\Xi_{1,N,0}), \text{pr}_1^*(\eta_{\check{f}}^{\text{ur}}) \wedge \text{pr}_2^*(\omega_{\check{g}}) \right\rangle.$$

Proof. Applying the previous theorem to $w_N \check{f}$ and $w_N \check{g}$, we have

$$\begin{aligned} (\mathcal{D}_p(w_N \check{f}, w_N \check{g}, x/N)(2) - \mathcal{D}_p(w_N \check{f}, w_N \check{g}, 0)(2)) \\ = \frac{\mathcal{E}(f, g, 1)}{\mathcal{E}(f) \cdot \mathcal{E}^*(f)} \left\langle r_{\text{syn}}(\Delta_{u_{x/N}}), \text{pr}_1^*(\eta_{w_N \check{f}}^{\text{ur}}) \wedge \text{pr}_2^*(\omega_{w_N \check{g}}) \right\rangle. \end{aligned}$$

We multiply by $e^{2\pi i x/N}$ and sum over $x \in \mathbb{Z}/N\mathbb{Z}$. The left-hand side becomes

$$\sum_{x \in \mathbb{Z}/N\mathbb{Z}} e^{2\pi i x/N} \mathcal{D}_p(w_N \check{f}, w_N \check{g}, x/N)(2) = N \mathcal{D}_p(\check{f}, \check{g}, x/N)(1)$$

by the p -adic functional equation.

Meanwhile, the right-hand side is

$$\sum_{x \in \mathbb{Z}/N\mathbb{Z}} e^{2\pi i x/N} \frac{\mathcal{E}(f, g, 1)}{\mathcal{E}(f) \cdot \mathcal{E}^*(f)} \left\langle r_{\text{syn}}(\Delta_{u_{x/N}}), \text{pr}_1^*(\eta_{w_N \check{f}}^{\text{ur}}) \wedge \text{pr}_2^*(\omega_{w_N \check{g}}) \right\rangle.$$

By the functoriality of the syntomic regulator, we have

$$\left\langle r_{\text{syn}}(\Delta_{u_{x/N}}), \text{pr}_1^*(\eta_{w_N \check{f}}^{\text{ur}}) \wedge \text{pr}_2^*(\omega_{w_N \check{g}}) \right\rangle = \left\langle r_{\text{syn}}(\Delta_{(w_N^* u_{x/N})}), \text{pr}_1^*(\eta_{\check{f}}^{\text{ur}}) \wedge \text{pr}_2^*(\omega_{\check{g}}) \right\rangle.$$

As elements of $\mathbb{Q}(\mu_N) \otimes_{\mathbb{Z}} \mathcal{O}(Y_1(N))^\times$, we have

$$\sum_{x \in \mathbb{Z}/N\mathbb{Z}} e^{2\pi i x/N} \otimes w_N^*(u_{x/N}) = -N \otimes g_{0,1/N},$$

and the result follows. □

Remark 5.6.5. One could also prove this statement directly (without the extended detour via Atkin–Lehner involutions and functional equations) by generalizing some of the calculations of [BDR12] to use the weight 2 Eisenstein series $F_\chi^{(2)} = \sum_{x \in (\mathbb{Z}/N\mathbb{Z})^\times} \chi(x)^{-1} F_{x/N}^{(2)}$ in place of $E_\chi^{(2)} = \sum_{x \in (\mathbb{Z}/N\mathbb{Z})^\times} \chi(x)^{-1} E_{x/N}^{(2)}$. (Note that $F_\chi^{(2)}$ is always a holomorphic Eisenstein series if $N > 1$, while $E_\chi^{(2)}$ becomes nonholomorphic if χ is the trivial character.)

Remark 5.6.6. If we impose slightly more restrictive hypotheses, we can avoid the need for any generalization of the main theorem of [BDR12]. If χ is primitive, then it suffices to check that the main theorem of [BDR12] holds without the assumption that f, g are eigenforms for the U_ℓ with $\ell \mid N$; but this

assumption is not used anywhere in the paper, except in order to explicitly evaluate the Euler factors at $\ell \mid N$. If $N_f = N_g = N$, then we can dispense with this assumption as well.

6. Families of cohomology classes

In this section, we will construct étale cohomology classes from the generalized Beilinson–Flach elements in motivic cohomology defined above and investigate their properties.

6.1. *The étale regulator.* In [Hub00], Huber constructs a p -adic regulator map from motivic cohomology into Jannsen’s continuous étale cohomology:

PROPOSITION 6.1.1. *Assume that X is a smooth variety over a characteristic 0 field k . Then there is a regulator map*

$$(16) \quad r_{\acute{e}t} : \mathrm{CH}^2(X, 1) \longrightarrow H_{\mathrm{cont}}^3(X, \mathbb{Z}_p(2)).$$

Proof. See the second example on [Hub00, p. 772]. □

PROPOSITION 6.1.2. *If X is any smooth variety over k , then we have a Hochschild–Serre spectral sequence*

$$H^p(k, H_{\acute{e}t}^q(\bar{X}, \mathbb{Z}_p(2))) \Rightarrow H_{\mathrm{cont}}^{p+q}(X, \mathbb{Z}_p(2)),$$

where $H^*(k, -)$ denotes continuous Galois cohomology.

Proof. See [Jan88a, Rem. 3.5]. □

COROLLARY 6.1.3. *Suppose that X is a smooth affine surface over k . Then we have an edge map*

$$(17) \quad H_{\mathrm{cont}}^3(X, \mathbb{Z}_p(2)) \rightarrow H^1(k, H_{\acute{e}t}^2(\bar{X}, \mathbb{Z}_p(2))).$$

Proof. The fact that \bar{X} is defined over an algebraically closed field implies that $H_{\acute{e}t}^q(\bar{X}, \mathbb{Z}_p(2)) = 0$ for $q > 2$, as a d -dimensional affine variety over an algebraically closed field has étale cohomological dimension d [Del77, Arcata IV.6.4]. Consequently, we have $H_{\mathrm{cont}}^3(\bar{X}, \mathbb{Z}_p(2)) = 0$, and we obtain the required edge map by Proposition 6.1.2. □

COROLLARY 6.1.4. *If X is a smooth affine surface over k , the étale regulator induces a map (which we also denote by $r_{\acute{e}t}$ by abuse of notation)*

$$(18) \quad r_{\acute{e}t} : \mathrm{CH}^2(X, 1) \longrightarrow H^1(k, H_{\acute{e}t}^2(\bar{X}, \mathbb{Z}_p(2))).$$

Proof. Compose $r_{\acute{e}t}$ with the edge map (17). □

The regulator maps have the following functoriality property:

PROPOSITION 6.1.5. *The regulator maps (18) are compatible with pullback along flat morphisms of surfaces $X \rightarrow Y$ over k and pushforward along finite morphisms. In particular, they are compatible with the Galois restriction maps for arbitrary extensions k'/k and with the corestriction maps for finite ones.*

Proof. This is true essentially by construction for Huber’s regulator into continuous cohomology since it arises from a realization functor on Voevodsky’s category DM_{gm} of geometrical motives, which in turn is built up from the category (denoted by $SmCor$ in [Hub00]) whose objects are smooth varieties over k and whose morphisms are finite correspondences $X \rightrightarrows Y$. It remains only to check that the Hochschild–Serre exact sequence (17) has the required functoriality property, which is standard. \square

6.2. *The Künneth formula.* We also recall the Künneth formula for étale cohomology (cf. [Mil12, Th. 22.4]): if U and V are varieties of finite type over an algebraically closed field of characteristic 0, then we have an exact sequence

$$0 \longrightarrow \sum_{r+s=m} H_{\acute{e}t}^r(U, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} H_{\acute{e}t}^s(V, \mathbb{Z}_p) \longrightarrow H_{\acute{e}t}^m(U \times V, \mathbb{Z}_p) \\ \longrightarrow \sum_{r+s=m+1} \mathrm{Tor}_1^{\mathbb{Z}_p}(H_{\acute{e}t}^r(U, \mathbb{Z}_p), H_{\acute{e}t}^s(V, \mathbb{Z}_p)) \longrightarrow 0.$$

We are interested in the case when $m = 2$, and U and V are smooth curves. If U and V are affine, then they have étale cohomological dimension 1; so the third term vanishes, as do two of the three summands in the first term, and we have the following result:

LEMMA 6.2.1. *For affine curves U, V , the Künneth formula gives an isomorphism*

$$H_{\acute{e}t}^1(U, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} H_{\acute{e}t}^1(V, \mathbb{Z}_p) \xrightarrow{\cong} H_{\acute{e}t}^2(U \times V, \mathbb{Z}_p),$$

functorial in U and V and compatible with the Galois action.

We shall also need to consider the case when U and V are projective (and connected). In this case, we shall assume the ground field k is $\overline{\mathbb{Q}}$. By the compatibility of étale cohomology with Betti cohomology after base extension to \mathbb{C} , we find that in this case the étale cohomology is \mathbb{Z}_p in degree 0 or 2 and \mathbb{Z}_p^{2g} in degree 1, where g is the genus. Hence all the Tor terms vanish, since the cohomology groups are free \mathbb{Z}_p -modules. We conclude that $H_{\acute{e}t}^2(U \times V, \mathbb{Z}_p)$ is the direct sum of $H_{\acute{e}t}^1(U, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} H_{\acute{e}t}^1(V, \mathbb{Z}_p)$ and two other summands, which are both isomorphic (as Galois representations) to $\mathbb{Z}_p(-1)$.

6.3. *Galois representations attached to modular forms.* We recall the construction of the Galois representations attached to cuspidal modular forms of weight 2, using the cohomology of the affine modular curves $Y_1(N)$.

Notation. Let f be a cuspidal modular form of weight 2 and level N . We assume that f is a normalized eigenform for all the Hecke operators T_v (for $v \nmid N$) and U_v (for $v \mid N$). (We do *not* assume that f is new of level N .) As

usual, we write $a_v(f)$ for the v -th Fourier coefficient of f , which is its eigenvalue for T_v if $v \nmid N$ and for U_v if $v \mid N$; we also write $\varepsilon_d(f)$ for the eigenvalue of f for the $\langle d \rangle$ operator for $d \in (\mathbb{Z}/N\mathbb{Z})^\times$.

By [AS86, Prop. 4], the space $H_{\text{Betti},c}^1(Y_1(N)(\mathbb{C}), \mathbb{C})$ is isomorphic to the space of *modular symbols* of level $\Gamma_1(N)$ with coefficients in \mathbb{C} . This contains a unique 2-dimensional \mathbb{C} -linear subspace $V_{\mathbb{C}}(f)$ on which the Hecke operators T_v, U_v act as multiplication by the Fourier coefficients $a_v(f)$; and the period isomorphism relating Betti and de Rham cohomology allows us to regard f as an element of $V_{\mathbb{C}}(f)$. Moreover, if L is any finite extension of \mathbb{Q} containing the Fourier coefficients of f , $V_{\mathbb{C}}(f)$ is the base-extension of a 2-dimensional L -subspace $V_L(f) \subseteq H_{\text{Betti},c}^1(Y_1(N)(\mathbb{C}), L)$.

Let p be a prime. Invoking the comparison theorem between (compactly-supported) p -adic and Betti cohomology, we can regard $\mathbb{Q}_p \otimes_{\mathbb{Q}} V_L(f)$ as a subspace of $L \otimes_{\mathbb{Q}} H_{\text{ét},c}^1(\overline{Y_1(N)}, \mathbb{Q}_p)$. Both of these are free modules of rank 2 over $L \otimes_{\mathbb{Q}} \mathbb{Q}_p = \prod_{\mathfrak{p}|p} L_{\mathfrak{p}}$, where the product is over primes of L above p ; so we obtain for each \mathfrak{p} a 2-dimensional $L_{\mathfrak{p}}$ -linear subspace $V_{L_{\mathfrak{p}}}(f) \subseteq H_{\text{ét},c}^1(\overline{Y_1(N)}, L_{\mathfrak{p}})$.

The following proposition is well known:

PROPOSITION 6.3.1. *The Galois representation $V_{L_{\mathfrak{p}}}(f)$ is “the” irreducible $L_{\mathfrak{p}}$ -linear Galois representation attached to f . That is, for each prime $v \nmid Np$, the representation $V_{L_{\mathfrak{p}}}(f)$ is unramified at v and we have*

$$\text{trace}_{L_{\mathfrak{p}}}(\text{Frob}_v^{-1} \mid V_{L_{\mathfrak{p}}}(f)) = a_v(f),$$

where Frob_v is the arithmetic Frobenius.

We note that under Poincaré duality, the dual space $V_L(f)^*$ is identified with the maximal quotient of $H_{\text{Betti}}^1(Y_1(N)(\mathbb{C}), L)$ on which the transposes T'_v and U'_v of T_v and U_v act as multiplication by $a_v(f)$. Tensoring with \mathbb{Q}_p , and noting that Poincaré duality holds in étale cohomology with a twist by the cyclotomic character, we obtain an identification of $V_{L_{\mathfrak{p}}}(f)^*$ with a quotient of $H_{\text{ét}}^1(\overline{Y_1(N)}, L_{\mathfrak{p}})(1)$.

Definition 6.3.2. Let $\mathcal{O}_{\mathfrak{p}}$ be the ring of integers of $L_{\mathfrak{p}}$. We define $T_{\mathcal{O}_{\mathfrak{p}}}(f)^*$ as the $\mathcal{O}_{\mathfrak{p}}$ -submodule of $V_{L_{\mathfrak{p}}}(f)^*$ generated by the image of $H_{\text{ét}}^1(\overline{Y_1(N)}, \mathbb{Z}_p)(1)$, which is a $G_{\mathbb{Q}}$ -stable $\mathcal{O}_{\mathfrak{p}}$ -lattice in $V_{L_{\mathfrak{p}}}(f)^*$.

Remark 6.3.3. Note that our conventions are somewhat different from those of [Kat04, §§6.3, 8.4]: we define $V_{L_{\mathfrak{p}}}(f)$ as a subspace of compactly-supported cohomology of a modular curve, while Kato uses the same symbol to denote a quotient of the noncompactly-supported cohomology. If f is new of level N , then our $V_{\mathcal{O}_{\mathfrak{p}}}(f)^*$ coincides with the space Kato would denote by $V_{L_{\mathfrak{p}}}(\overline{f})(1)$ where \overline{f} is the complex conjugate of N , and similarly for the integral lattices. (Our $T_{\mathcal{O}_{\mathfrak{p}}}(f)^*$ is Kato’s $V_{\mathcal{O}_{\mathfrak{p}}}(\overline{f})(1)$.)

Remark 6.3.4. One can also define a lattice in $V_{L_{\mathfrak{p}}}(f)^*$ using the cohomology of the projective modular curve. The inclusion $Y_1(N) \hookrightarrow \overline{X_1(N)}$ induces a pullback map $H^1(\overline{X_1(N)}, \mathbb{Z}_p) \rightarrow H^1(\overline{Y_1(N)}, \mathbb{Z}_p)$, which is injective with cokernel isomorphic to \mathbb{Z}_p^{r-1} , where r is the number of cusps. The action of the Hecke algebra on the boundary term \mathbb{Z}_p^{r-1} is Eisenstein, so the map $H^1(\overline{X_1(N)}, \mathbb{Q}_p) \rightarrow H^1(\overline{Y_1(N)}, \mathbb{Q}_p)$ is an isomorphism on the f -isotypical component. We define $\widetilde{T}_{\mathcal{O}_{\mathfrak{p}}}(f)^*$ as the image of $H^1(\overline{X_1(N)}, \mathbb{Z}_p) \otimes \mathcal{O}_{\mathfrak{p}}$ in $V_{L_{\mathfrak{p}}}(f)^*$. Note that $\widetilde{T}_{\mathcal{O}_{\mathfrak{p}}}(f)^* \subseteq T_{\mathcal{O}_{\mathfrak{p}}}(f)^*$, and equality holds if f is not congruent modulo \mathfrak{p} to an Eisenstein series.

6.4. *Generalized Beilinson–Flach classes.* Let $N \geq 5$. Observe that $Y_1(N)^2 \otimes \mathbb{Q}(\mu_m)$ is a smooth variety over $\mathbb{Q}(\mu_m)$ for any m . By (18), for any prime p , we therefore have an étale regulator

$$r_{\text{ét}, \mathbb{Q}(\mu_m)} : \text{CH}^2(Y_1(N)^2 \otimes \mathbb{Q}(\mu_m), 1) \longrightarrow H^1(\mathbb{Q}(\mu_m), H_{\text{ét}}^2(\overline{Y_1(N)^2}, \mathbb{Z}_p(2))).$$

Definition 6.4.1. Let f, g be modular forms of level N that are normalized eigenforms for all the Hecke operators T_{ℓ} (for $\ell \nmid N$) and U_{ℓ} (for $\ell \mid N$), L a number field containing the Fourier coefficients of f and g , and \mathfrak{p} a place of L above the rational prime p .

Remark 6.4.2. In the situation of Definition 6.4.1, we can use the Künneth formula (Lemma 6.2.1) to regard

$$T_{\mathcal{O}_{\mathfrak{p}}}(f, g)^* := T_{\mathcal{O}_{\mathfrak{p}}}(f)^* \otimes_{\mathcal{O}_{\mathfrak{p}}} T_{\mathcal{O}_{\mathfrak{p}}}(g)^*$$

as a quotient of $\mathcal{O}_{\mathfrak{p}} \otimes_{\mathbb{Z}_p} H_{\text{ét}}^2(\overline{Y_1(N)^2}, \mathbb{Z}_p)(2)$.

Definition 6.4.3. Define the map

$$\kappa_{f, g, \mathbb{Q}(\mu_m)} : \text{CH}^2(Y_1(N)^2 \otimes \mathbb{Q}(\mu_m), 1) \longrightarrow H^1(\mathbb{Q}(\mu_m), T_{\mathcal{O}_{\mathfrak{p}}}(f, g)^*)$$

to be the composition of $r_{\text{ét}, \mathbb{Q}(\mu_m)}$ with the map on Galois cohomology induced by the projection

$$H_{\text{ét}}^2(\overline{Y_1(N)^2}, \mathbb{Z}_p)(2) \longrightarrow T_{\mathcal{O}_{\mathfrak{p}}}(f, g)^*.$$

Definition 6.4.4. We define the *generalized Beilinson–Flach class*

$${}_c \mathbf{z}_m^{(f, g, N)} := \kappa_{f, g, \mathbb{Q}(\mu_m)}({}_c \Xi_{m, N, 1}) \in H^1(\mathbb{Q}(\mu_m), T_{\mathcal{O}_{\mathfrak{p}}}(f, g)^*)$$

and its nonintegral version

$$\mathbf{z}_m^{(f, g, N)} := \kappa_{f, g, \mathbb{Q}(\mu_m)}(\Xi_{m, N, 1}) \in H^1(\mathbb{Q}(\mu_m), V_{\mathcal{O}_{\mathfrak{p}}}(f, g)^*).$$

The compatibility relations we have shown for the generalized Beilinson–Flach elements for varying m carry over to the cohomology classes:

COROLLARY 6.4.5. *For any integers $m \geq 1, N \geq 1$, and ℓ a prime such that $\ell \mid N$, we have*

$$\text{cores}_m^{\ell m} \left(c\mathbf{z}_{\ell m}^{(f,g,N)} \right) = \begin{cases} (\alpha_f \alpha_g) \cdot c\mathbf{z}_m^{(f,g,N)} & \text{if } \ell \mid m, \\ (\alpha_f \alpha_g - \sigma_\ell) \cdot c\mathbf{z}_m^{(f,g,N)} & \text{if } \ell \nmid m, \end{cases}$$

where α_f, α_g are the U_ℓ -eigenvalues of f and g , and in the latter case σ_ℓ is the arithmetic Frobenius element at ℓ in $\text{Gal}(\mathbb{Q}(\mu_m)/\mathbb{Q})$.

If ℓ is a prime not dividing mN , then

$$\text{cores}_m^{\ell m} \left(c\mathbf{z}_{\ell m}^{(f,g,N)} \right) = \sigma_\ell \left((\ell - 1)(1 - \varepsilon_f(\ell)\varepsilon_g(\ell)\sigma_\ell^{-2}) - \ell P_\ell(f, g, \ell^{-1}\sigma_\ell^{-1}) \right) c\mathbf{z}_m^{(f,g,N)},$$

where $P_\ell(f, g, X)$ is the local Euler factor of f and g at ℓ (cf. Proposition 4.1.2).

Proof. Immediate from Theorems 3.3.2, 3.4.1 and the compatibility of the regulator map with corestriction (Proposition 6.1.5). □

The dependence of $c\mathbf{z}_m^{(f,g,N)}$ on c is as follows:

PROPOSITION 6.4.6. *There exist classes*

$$\mathbf{z}_m^{(f,g,N)} \in H^1(\mathbb{Q}(\mu_m), V_{L_p}(f, g)^*)$$

such that the relation

$$(19) \quad c\mathbf{z}_m^{(f,g,N)} = (c^2 - \varepsilon_f(c)^{-1}\varepsilon_g(c)^{-1}[c]^2)\mathbf{z}_m^{(f,g,N)}$$

holds for any $c > 1$ coprime to $6mN$.

Proof. Immediate from Proposition 2.7.5(5). □

PROPOSITION 6.4.7. *If there exists $d \geq 1$ coprime to $6mN$ such that $d^2 - \varepsilon_f(d)^{-1}\varepsilon_g(d)^{-1}[d]^2$ is invertible in $\mathcal{O}_{\mathfrak{p}}[(\mathbb{Z}/m\mathbb{Z})^\times]$, then there exists $\mathbf{z}_m^{(f,g,N)} \in H^1(\mathbb{Q}(\mu_m), T_{\mathcal{O}_{\mathfrak{p}}}(f, g)^*)$ such that equation (19) holds in $H^1(\mathbb{Q}(\mu_m), T_{\mathcal{O}_{\mathfrak{p}}}(f, g)^*)$ (not just modulo torsion).*

In particular, this holds if the conductor of the reduction modulo \mathfrak{p} of $\varepsilon_f\varepsilon_g$ is divisible by some prime that does not divide mp .

Proof. Clear, since if such a d exists, we may define

$$\mathbf{z}_m^{(f,g,N)} := (d^2 - \varepsilon_f(d)^{-1}\varepsilon_g(d)^{-1}[d]^2)^{-1} d\mathbf{z}_m^{(f,g,N)}. \quad \square$$

6.5. *Local properties of the generalized Beilinson–Flach classes (I).* We now study the local properties of the Beilinson–Flach classes. We shall first recall some standard definitions.

Definition 6.5.1. If K is a local field and M is a topological G_K -module, we define $H_{\text{nr}}^1(K, M)$ to be the image of the inflation map

$$H^1(K^{\text{nr}}/K, M^{I_K}) \rightarrow H^1(K, M),$$

where I_K is the inertia subgroup of G_K and K^{nr} the maximal unramified extension of K .

If V is a finite-dimensional \mathbb{Q}_p -vector space and ℓ is the residue characteristic of K , we define

$$H_f^1(K, V) = \begin{cases} H_{\text{nr}}^1(K, V) & \text{if } \ell \neq p, \\ \ker(H^1(K, V) \rightarrow H^1(K, V \otimes \mathbb{B}_{\text{cris}})) & \text{if } \ell = p. \end{cases}$$

If T is a \mathbb{Z}_p -lattice in V stable under G_K , we write $H_f^1(K, T)$ for the preimage of $H_f^1(K, V)$ in $H^1(K, T)$; cf. [BK90].

PROPOSITION 6.5.2. *If T is a finite-rank free \mathbb{Z}_p -module with a continuous action of G_K that is trivial on I_K and if $\ell \neq p$, then*

$$H_f^1(K, T) = H_{\text{nr}}^1(K, T).$$

Proof. We have an inflation-restriction exact sequence

$$0 \longrightarrow H_{\text{nr}}^1(K, T) \longrightarrow H^1(K, T) \longrightarrow H^0(K^{\text{nr}}/K, H^1(I_K, T)) \longrightarrow 0$$

and a corresponding sequence for V in place of T . Suppose $x \in H_f^1(K, T)$. Then the image of x in $H^1(I_K, V)$ is zero, so the image of x in $H^1(I_K, T)$ is torsion. However, $H^1(I_K, T) = \text{Hom}(I_K, T)$ is torsion-free since T is; thus the image of x in $H^1(I_K, T)$ is zero, and hence $x \in H_{\text{nr}}^1(K, T)$. \square

Definition 6.5.3. If K is a number field and M is a topological G_K -module and v is a prime of K , we say that $x \in H^1(K, M)$ is *unramified at v* if its image in $H^1(K_v, M)$ lies in $H_{\text{nr}}^1(K, M)$. If M is a finite-rank \mathbb{Z}_p -module or \mathbb{Q}_p -vector space and v is a prime above p , we say x is *crystalline at v* if its image in $H^1(K_v, M)$ lies in $H_f^1(K_v, M)$.

PROPOSITION 6.5.4. *The generalized Beilinson–Flach class $c_{\mathcal{Z}_m}^{(f,g,N)}$ is unramified outside the primes dividing mNp . If $p \nmid mN$, it is crystalline at the primes above p .*

Proof. By the preceding proposition, it suffices to check this result after inverting p . Let us choose a prime $\ell \nmid mNp$. The compactified modular curve $X(m, mN)$ associated to $Y(m, mN)$ admits a smooth proper model $\mathcal{X}(m, mN)$ over $\mathbb{Z}[1/mN]$; hence it has such a model over \mathbb{Z}_ℓ . It is clear that the class $c_{\mathcal{Z}_{m,N,1}}$ lies in the higher Chow group $Z^2(\mathcal{Y}(m, mN), 1)$ of the integral model of $Y(m, mN)$, and we can choose the “negligible elements” of Theorem 2.8.5 in order to obtain a lifting of $c_{\mathcal{Z}_{m,N,1}}$ to $\text{CH}^2(\mathcal{X}(m, mN), 1) \otimes \mathbb{Q}$.

For proper smooth schemes \mathcal{S} over \mathbb{Z}_ℓ , with $\ell \neq p$, there is a regulator map

$$\text{CH}^2(\mathcal{S}, 1) \otimes \mathbb{Q}_p \rightarrow H_{\text{ét}}^3(\mathcal{S}, \mathbb{Q}_p(2))$$

(see, e.g., [Fla92]) compatible with the regulator map $r_{\text{ét}}$ on the generic fibre S . Moreover, the étale cohomology $H_{\text{ét}}^2(\overline{S}, \mathbb{Q}_p(2))$ is unramified as a representation of $G_{\mathbb{Q}_\ell}$, by the proper base change theorem; and the Hochschild–Serre spectral sequence maps $H_{\text{ét}}^3(\mathcal{S}, \mathbb{Q}_p(2))$ to $H^1(\mathbb{Q}_\ell^{\text{nr}}/\mathbb{Q}_\ell, H_{\text{ét}}^2(\overline{S}, \mathbb{Q}_p(2)) \subset H^1(\mathbb{Q}_\ell, H_{\text{ét}}^2(\overline{S}, \mathbb{Q}_p(2)))$, where $\overline{S} = \mathcal{S} \otimes \overline{\mathbb{Q}_\ell}$ (cf. [Fla92, Lemma 2.3]). Hence the class $c_{\mathbf{Z}_m}^{(f,g,N)}$ is unramified at the primes above ℓ , as required.

Similarly, if $p \nmid mN$, we can lift $c_{\mathcal{Z}_{m,N,1}}$ to a class in $\text{CH}^2(\mathcal{X}(m, mN), 1) \otimes \mathbb{Q}$ where $\mathcal{X}(m, mN)$ is proper and smooth over \mathbb{Z}_p . However, the regulator $r_{\text{ét}}$ for proper smooth \mathbb{Z}_p -schemes takes values in H_f^1 , as a consequence of the commutative diagram of Section 5.5 above relating $r_{\text{ét}}$ to the syntomic regulator r_{syn} ; so we are done. \square

Remark 6.5.5. We believe it is known that the regulator map $r_{\text{ét}}$ for arbitrary varieties over p -adic fields takes values in H_g^1 , as remarked in Section 5.5, which would imply that the localization at p of the Beilinson–Flach classes always lies in this subspace.

6.6. *Local properties of the generalized Beilinson–Flach classes (II).* In order to control the local properties of the generalized Beilinson–Flach classes at the “bad” primes, we shall make use of the compatibility in the p -adic cyclotomic tower, under mild additional hypotheses.

ASSUMPTION 6.6.1. *The level N is divisible by p , and the U_p -eigenvalues α_f, α_g of f and g satisfy*

$$v_p(\alpha_f \alpha_g) < 1.$$

PROPOSITION 6.6.2. *Suppose that Assumption 6.6.1 holds. Then for any $m \geq 1$ divisible by p and any prime $v \nmid p$ of $\mathbb{Q}(\mu_m)$, the cohomology class $c_{\mathbf{Z}_m}^{(f,g,N)}$ lies in $H_f^1(\mathbb{Q}(\mu_m)_v, T_{\mathcal{O}_p}(f, g)^*)$. If $v_p(\alpha_f \alpha_g) = 0$, it lies in $H_{\text{nr}}^1(\mathbb{Q}(\mu_m)_v, T_{\mathcal{O}_p}(f, g)^*)$. If $m \geq 1$ is coprime to p , and $\alpha_f \alpha_g$ is not a root of unity, then again we have $c_{\mathbf{Z}_m}^{(f,g,N)} \in H_f^1(\mathbb{Q}(\mu_m)_v, T_{\mathcal{O}_p}(f, g)^*)$.*

Proof. To lighten the notation, we write $K = \mathbb{Q}(\mu_m)_v$ and $M = T_{\mathcal{O}_p}(f, g)^*$. We assume first that $p \mid m$. We write $K_i = \mathbb{Q}(\mu_{mp^i})_v$ (after choosing one of the finitely many primes of $\mathbb{Q}(\mu_{mp^\infty})$ above v). Each K_i is contained in K^{nr} since $v \nmid p$.

For each i , there is an inflation–restriction exact sequence

$$0 \longrightarrow H^1(K^{\text{nr}}/K_i, M^{I_v}) \longrightarrow H^1(K_i, M) \longrightarrow H^0(K^{\text{nr}}/K_i, H^1(I_v, M)) \longrightarrow 0,$$

and the corestriction maps $H^1(K_{i+1}, M) \rightarrow H^1(K_i, M)$ correspond to the trace maps

$$H^0(K^{\text{nr}}/K_{i+1}, H^1(I_v, M)) \rightarrow H^0(K^{\text{nr}}/K_i, H^1(I_v, M)).$$

Since M is a finitely-generated \mathbb{Z}_p -module, $H^1(I_v, M)$ is finitely generated over \mathbb{Z}_p , by [Rub00, Prop. B.2.7(iii)]; thus the sequence of modules $M_i = H^0(K^{\text{nr}}/K_i, H^1(I_v, M))$ stabilizes at some $i_0 \gg 0$. So for $i \geq i_0$, the trace maps $M_{i+1} \rightarrow M_i$ are simply multiplication by p on $M_{i+1} = M_i = M_\infty$. Let z_i be the image of $c\mathbf{z}_{mp^i}^{(f,g,N)}$ in M_i . It then follows that for $i \geq i_0$, we have

$$(\alpha_f \alpha_g)^i z_0 = p^{i-i_0} \text{cores}_{i_0}^0(z_i).$$

If $\alpha_f \alpha_g$ is a p -adic unit, then this immediately implies that $z_0 = 0$ since it is divisible by arbitrarily high powers of p . Thus $c\mathbf{z}_m^{(f,g,N)}$ is unramified at v .

Otherwise, we can only deduce that

$$z_0 \in \left(\frac{p^{i-i_0}}{(\alpha_f \alpha_g)^i} \right) M_0 + (M_0)_{\text{tors}}$$

for all $i \gg 0$, which implies that $z_0 \in (M_0)_{\text{tors}}$ as $v_p(\alpha_f \alpha_g) < 1$. Hence the image of z_0 in $M_0 \otimes \mathbb{Q}_p$ is zero, so the image of $c\mathbf{z}_m^{(f,g,N)}$ in $H^1(\mathbb{Q}(\mu_m)_v, M \otimes \mathbb{Q}_p)$ is unramified. Thus $c\mathbf{z}_m^{(f,g,N)} \in H_f^1(\mathbb{Q}(\mu_m)_v, M)$. Now suppose $p \nmid m$. Applying the above argument with m replaced by mp and using Corollary 6.4.5, we see that $(\alpha_f \alpha_g - \sigma_p) c\mathbf{z}_m^{(f,g,N)}$ lies in H_f^1 , but since $\alpha_f \alpha_g$ is not a root of unity, the factor $(\alpha_f \alpha_g - \sigma_p)$ is invertible after extending scalars to \mathbb{Q}_p . \square

6.7. *Relation between p -stabilized and non- p -stabilized classes.* For the arguments of the previous section, we assumed throughout that $p \mid N$. If we are given forms of levels prime to p , then we can obtain forms of level divisible by p via “ p -stabilization” (choosing old eigenforms of level divisible by p with the same Hecke eigenvalues at all other primes). In this section, we shall investigate the relations between the classes obtained for the p -stabilized and non- p -stabilized forms.

Let f be a normalized eigenform of weight 2 and level N , and let p be a prime such that $p \nmid N$. Then there are two eigenforms f_α, f_β at level Np in the oldspace attached to f , whose U_p -eigenvalues are the roots α, β of the Hecke polynomial $X^2 - a_p(f)X + p\epsilon_p(f)$. (We assume, by enlarging the field if necessary, that these lie in our coefficient field L .)

Then there are projection maps

$$\begin{aligned} \text{pr}_{f_\alpha} &: H_{\text{ét}}^1(\overline{Y_1(Np)}, L_{\mathfrak{p}}) \rightarrow V_{L_{\mathfrak{p}}}(f_\alpha)^*, \\ \text{pr}_{f_\beta} &: H_{\text{ét}}^1(\overline{Y_1(Np)}, L_{\mathfrak{p}}) \rightarrow V_{L_{\mathfrak{p}}}(f_\beta)^*, \\ \text{pr}_f &: H_{\text{ét}}^1(\overline{Y_1(N)}, L_{\mathfrak{p}}) \rightarrow V_{L_{\mathfrak{p}}}(f)^* \end{aligned}$$

and a pushforward map $\pi : H_{\text{ét}}^1(\overline{Y_1(Np)}, L_{\mathfrak{p}}) \rightarrow H_{\text{ét}}^1(\overline{Y_1(N)}, L_{\mathfrak{p}})$.

PROPOSITION 6.7.1. *In the above situation, there is a nonzero, $G_{\mathbb{Q}}$ -equivariant map*

$$\pi^{(\alpha)} : V_{L_{\mathfrak{p}}}(f_\alpha)^* \rightarrow V_{L_{\mathfrak{p}}}(f)^*$$

and similarly $\pi^{(\beta)}$, with the property that

$$\pi^{(\alpha)} \circ \text{pr}_{f_\alpha} + \pi^{(\beta)} \circ \text{pr}_{f_\beta} = \text{pr}_f \circ \pi$$

as maps $H_{\text{ét}}^1(\overline{Y_1(Np)}, L_p) \rightarrow V_f^*$.

Proof. Let $H_{\text{ét}}^1(\overline{Y_1(Np)}, L_p)_{[f]}$ denote the maximal quotient of the space $H_{\text{ét}}^1(\overline{Y_1(Np)}, L_p)$ where the operators T'_v for $v \nmid Np$ and U'_v for $v \mid N$ act via $a_v(f)$. Then, by comparison with modular symbols, we see that $H_{\text{ét}}^1(\overline{Y_1(Np)}, L_p)_{[f]}$ is 4-dimensional, and the U'_p operator on this space is annihilated by the Hecke polynomial.

By [CE98, Th. 2.1], the roots α and β are distinct, so we may write $H_{\text{ét}}^1(\overline{Y_1(Np)}, L_p)_{[f]}$ as a direct sum of $G_{\mathbb{Q}}$ -stable eigenspaces, which map isomorphically onto the quotients $V_{L_p}(f_\alpha)$ and $V_{L_p}(f_\beta)$. This gives a lifting of $V_{L_p}(f_\alpha)$ to a subspace of $H_{\text{ét}}^1(\overline{Y_1(Np)}, L_p)_{[f]}$, and the map $\text{pr}_f \circ \pi$ clearly factors through $H_{\text{ét}}^1(\overline{Y_1(Np)}, L_p)_{[f]}$ as stated. \square

Now let us suppose we have two normalized weight 2 eigenforms f, g , of level N prime to p as before. Let α, β be the roots of the Hecke polynomial of f at p , and similarly γ, δ for g . By the Coleman–Edixhoven theorem cited above, we have $\alpha \neq \beta$ and $\gamma \neq \delta$.

A choice of root of each polynomial gives p -stabilized eigenforms f_α, g_γ of level Np . Then for each m , we have

- a class $\mathbf{z}_m^{(f,g,N)}$ in the cohomology of $V_{L_p}(f, g)^*$, which is a quotient of $H_{\text{ét}}^2(\overline{Y_1(N)^2}, L_p)(2)$;
- an element $\mathbf{z}_m^{(f_\alpha, g_\gamma, Np)}$ living in the cohomology of the representation $V_{L_p}(f_\alpha, g_\gamma)^*$, which is a quotient of $H_{\text{ét}}^2(\overline{Y_1(Np)^2}, L_p)(2)$.

These two representations are isomorphic as abstract Galois representations but are realized differently as quotients of étale cohomology. We can regard both as quotients of the following space:

Definition 6.7.2. Let $H_{\text{ét}}^2(\overline{Y_1(Np)^2}, L_p)_{f,g}$ denote the maximal L_p -linear quotient of $H_{\text{ét}}^2(\overline{Y_1(Np)^2}, L_p)$ on which the operators $(T'_v, 1)$ (for $v \nmid Np$) and $(U'_v, 1)$ (for $v \mid N$) act via the Fourier coefficients of f , and similarly for g .

Note 6.7.3. Using the Künneth formula and a modular symbol calculation, we see that $H_{\text{ét}}^2(\overline{Y_1(Np)^2}, L_p)_{f,g}$ has dimension 16 and can be viewed as a direct sum of four simultaneous eigenspaces for the two operators $(U'_p, 1)$ and $(1, U'_p)$, corresponding to the stabilizations (α, γ) , (α, δ) , (β, γ) and (β, δ) . Each of these is a 4-dimensional $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ -stable L_p -linear subspace.

For the remainder of this section, we shall assume the following:

ASSUMPTION 6.7.4. *We have $\alpha\gamma \neq \beta\delta$.*

Remark 6.7.5. Assumption 6.7.4 is a consequence of Assumption 6.6.1, since $v_p(\alpha\beta\gamma\delta) = v_p(p^2\varepsilon_f(p)\varepsilon_g(p)) = 2$, so if $v_p(\alpha\gamma) < 1$, then $v_p(\beta\delta) > 1$.

PROPOSITION 6.7.6. *If Assumption 6.7.4 is satisfied, then the operator*

$$J_{\alpha,\gamma} := \frac{(\mathcal{U} - \alpha\delta)(\mathcal{U} - \beta\gamma)(\mathcal{U} - \beta\delta)}{(\alpha\gamma - \alpha\delta)(\alpha\gamma - \beta\gamma)(\alpha\gamma - \beta\delta)},$$

where $\mathcal{U} = (U'_p, U'_p)$, is an idempotent in $\text{End}_{L_p} H_{\text{ét}}^2(\overline{Y_1(Np)^2}, L_p)_{f,g}$; it is equal to the identity on the (α, γ) eigenspace and zero on the other three eigenspaces.

Proof. We know that $\alpha\gamma \neq \alpha\delta$ and $\alpha\gamma \neq \beta\gamma$ by the Coleman–Edixhoven theorem, so if $\alpha\gamma \neq \beta\delta$, the $\alpha\gamma$ eigenspace for the operator \mathcal{U} coincides with the (α, γ) simultaneous eigenspace for $(U'_p, 1)$ and $(1, U'_p)$. We may thus define a projection onto this eigenspace by applying to \mathcal{U} a polynomial that is 1 at $\alpha\gamma$ and zero at the other three eigenvalues. \square

PROPOSITION 6.7.7. *There is a $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ -equivariant L_p -linear isomorphism*

$$\pi^{(\alpha,\gamma)} : V_{L_p}(f_\alpha, g_\gamma)^* \xrightarrow{\cong} V_{L_p}(f, g)^*$$

with the property that

$$\pi^{(\alpha,\gamma)} \circ \text{pr}_{(\alpha,\gamma)} = \pi \circ J_{\alpha,\gamma}$$

as maps $H_{\text{ét}}^2(\overline{Y_1(Np)^2}, L_p)_{f,g}(2) \rightarrow V_{L_p}(f, g)^*$, where

$$\pi : H_{\text{ét}}^2(\overline{Y_1(Np)^2}, L_p)_{f,g} \rightarrow V_{L_p}(f, g)^*$$

is the natural map induced by the pushforward map $Y_1(Np)^2 \rightarrow Y_1(N)^2$.

Proof. We define $\pi^{(\alpha,\gamma)}$ as $\pi \circ \iota^{(\alpha,\gamma)}$, where $\iota^{(\alpha,\gamma)}$ is the section of $\text{pr}_{\alpha,\gamma}$ identifying $V_{L_p}(f_\alpha, g_\gamma)$ with the (α, γ) -eigenspace of $H_{\text{ét}}^2(\overline{Y_1(Np)^2}, L_p)_{f,g}$. The composition $\iota^{(\alpha,\gamma)} \circ \text{pr}_{\alpha,\gamma}$ is therefore equal to the projection operator $J_{\alpha,\gamma}$ above, and the proposition follows. \square

COROLLARY 6.7.8. *For $p \nmid m$, we have*

$$\pi^{(\alpha,\gamma)} \left({}_c z_m^{(f_\alpha, g_\gamma, Np)} \right) = \frac{\alpha\gamma \left(1 - \frac{\beta\delta}{p} \sigma_p^{-1}\right) \left(1 - \frac{\alpha\delta}{p} \sigma_p^{-1}\right) \left(1 - \frac{\beta\gamma}{p} \sigma_p^{-1}\right)}{(\gamma - \delta)(\alpha - \beta)} \cdot {}_c z_m^{(f, g, N)}.$$

Proof. We shall prove this by a slightly roundabout argument, using the second norm relation “in reverse” to understand how \mathcal{U} acts on the zeta elements. Let us write the polynomial

$$\frac{(X - \alpha\delta)(X - \beta\gamma)(X - \beta\delta)}{(\alpha\gamma - \alpha\delta)(\alpha\gamma - \beta\gamma)(\alpha\gamma - \beta\delta)} \in L_p[X]$$

as $j_0 + j_1 X + j_2 X^2 + j_3 X^3$, and let ${}_c z_m^{(f, g, Np)}$ be the image of $\text{reg}_{\text{ét}}({}_c \Xi_{m, Np, j})$ in $H_{\text{ét}}^2(\overline{Y_1(Np)^2}, L_p)_{f,g}$.

Essentially, by definition, we have ${}_{c z_m}^{(f_\alpha, g_\gamma, Np)} = \text{pr}_{\alpha, \gamma} {}_{c z_m}^{(f, g, Np)}$, and hence we may apply the preceding proposition to obtain

$$\begin{aligned} \pi^{(\alpha, \gamma)} \left({}_{c z_m}^{(f_\alpha, g_\gamma, Np)} \right) &= \pi \left(J_{\alpha, \gamma} \cdot {}_{c z_m}^{(f, g, Np)} \right) \\ &= \pi \left((j_0 + j_1 \mathcal{U} + j_2 \mathcal{U}^2 + j_3 \mathcal{U}^3) {}_{c z_m}^{(f_\alpha, g_\gamma, Np)} \right). \end{aligned}$$

By the second norm relation for $p \mid N$ (Theorem 3.3.2) and induction on r we see that for $r \geq 1$, we have

$$\begin{aligned} \mathcal{U}^r \left({}_{c z_m}^{(f, g, Np)} \right) &= \text{norm}_m^{p^r m} \left({}_{c z_{p^r m}}^{(f, g, Np)} \right) \\ &\quad + \sigma_p \cdot \text{norm}_m^{p^{r-1} m} \left({}_{c z_{p^{r-1} m}}^{(f, g, Np)} \right) + \cdots + \sigma_p^r \cdot {}_{c z_m}^{(f, g, Np)}. \end{aligned}$$

On the other hand, by the first norm relation (Theorem 3.1.2) we know that for $r \geq 1$, we have

$$\pi \left(\text{norm}_m^{p^r m} \left({}_{c z_{p^r m}}^{(f, g, Np)} \right) \right) = \text{norm}_m^{p^r m} \left(\pi \left({}_{c z_{p^r m}}^{(f, g, Np)} \right) \right) = \text{norm}_m^{p^r m} {}_{c z_m}^{f, g, N}$$

while for $r = 0$, we have

$$\pi \left({}_{c z_m}^{(f, g, Np)} \right) = (1 - \varepsilon_p(f) \varepsilon_p(g) \sigma_p^{-2}) {}_{c z_m}^{(f, g, N)}.$$

Combining these statements, we have

$$\begin{aligned} \pi^{(\alpha, \gamma)} \left({}_{c z_m}^{(f_\alpha, g_\gamma, Np)} \right) &= (j_0 + j_1 \sigma_p + j_2 \sigma_p^2 + j_3 \sigma_p^3) (1 - \varepsilon_p(f) \varepsilon_p(g) \sigma_p^{-2}) {}_{c z_m}^{(f, g, N)} \\ &\quad + (j_1 + j_2 \sigma_p + j_3 \sigma_p^2) \text{norm}_m^{p m} \left({}_{c z_{p m}}^{(f, g, N)} \right) \\ &\quad + (j_2 + \sigma_p j_3) \text{norm}_m^{p^2 m} \left({}_{c z_{p^2 m}}^{(f, g, N)} \right) + j_3 \text{norm}_m^{p^3 m} \left({}_{c z_{p^3 m}}^{(f, g, N)} \right). \end{aligned}$$

The prime-to- p case of the second norm relation (Theorem 3.4.1) gives a formula for the second term, and Theorem 3.5.2 extends this to the remaining two terms. Substituting these in, the entirety of the right-hand side simplifies to a linear combination of terms, each of which is ${}_{c z_m}^{(f, g, N)}$ acted on by a polynomial in σ_p, σ_p^{-1} with coefficients given as rational functions in $\alpha, \beta, \gamma, \delta$. After a computation (which was carried out using Sage, [Sage]), one finds the polynomial simplifies to the product of Euler-type factors stated above. \square

This extremely laborious computation allows us to prove the following theorem, which will be crucial to the Iwasawa-theoretic applications of our Euler system:

COROLLARY 6.7.9. *Suppose f, g admit p -stabilizations f_α, g_γ such that $v_p(\alpha\gamma) < 1$. Suppose m is coprime to p and neither of the quantities $\alpha\delta/p, \beta\gamma/p$ is an r -th root of unity, where r is the order of p in $(\mathbb{Z}/m\mathbb{Z})^\times$.*

Then for every prime $v \nmid p$ of $\mathbb{Q}(\mu_m)$, the localization of ${}_{c \mathbf{z}_m}^{f, g, N}$ at v lies in H_f^1 .

Proof. We know from Proposition 6.6.2 that the class $c_{\mathbb{Z}_m}^{f_{\alpha}, g_{\beta}, Np}$ is in H_f^1 at all primes away from p . Since $\alpha\gamma \neq \beta\delta$ by Remark 6.7.5, the formula of the previous corollary applies.

We note that for $\lambda \in L_p$, the element $1 - \lambda\sigma_p^{-1}$ is invertible in $L_p[(\mathbb{Z}/m\mathbb{Z})^\times]$ if and only if $\lambda^r \neq 1$, where r is the order of σ_p as above. It is clear that $\frac{\beta\delta}{p}$ cannot be a root of unity of any order, as its \mathfrak{p} -adic valuation is strictly positive. By assumption neither $\alpha\delta/p$ nor $\beta\gamma/p$ is an r -th root of unity; so the quantity

$$\frac{\alpha\gamma \left(1 - \frac{\beta\delta}{p}\sigma_p^{-1}\right) \left(1 - \frac{\alpha\delta}{p}\sigma_p^{-1}\right) \left(1 - \frac{\beta\gamma}{p}\sigma_p^{-1}\right)}{(\gamma - \delta)(\alpha - \beta)}$$

is invertible in $L_p[(\mathbb{Z}/m\mathbb{Z})^\times]$. Hence $c_{\mathbb{Z}_m}^{(f, g, N)}$ is also in H_f^1 . □

Remark 6.7.10. Note that the conclusion of Corollary 6.7.9 does not explicitly mention the choice of p -stabilization (α, γ) ; we use only the fact that one exists. We conjecture that the conclusion holds much more generally.

6.8. *Iwasawa cohomology classes.*

Notation. We now let S be a finite set of places of \mathbb{Q} containing p, ∞ , and all primes whose inertia groups act nontrivially on $T_{\mathcal{O}_p}(f, g)^*$ (which can only happen for primes dividing N). Let \mathbb{Q}^S be the maximal extension of \mathbb{Q} unramified outside S .

Definition 6.8.1. For K a finite extension of \mathbb{Q} contained in \mathbb{Q}^S , $i \geq 0$, and T a topological $\mathbb{Z}_p[G_K]$ -module unramified outside S , define

$$H_S^i(K, T) = H^i(\mathbb{Q}^S/K, T).$$

If T is also a finitely-generated \mathbb{Z}_p -module and K_∞ is a p -adic Lie extension of K unramified outside S , define

$$H_{Iw, S}^i(K_\infty, T) = \varprojlim_L H_S^i(L, T),$$

where L varies over the set of finite extensions of K contained in K_∞ and the inverse limit is with respect to the corestriction maps.

Remark 6.8.2. If K_∞ contains finite extensions of K of degree divisible by arbitrarily large powers of p – for instance, if K_∞/K is Galois and its Galois group is a p -adic Lie group of positive dimension – then $H_{Iw, S}^0(K_\infty, V)$ is zero, and $H_{Iw, S}^1(K_\infty, T)$ is in fact independent of S , as long as S contains the set S_0 consisting of all primes above p or ∞ , all primes ramifying in K_∞/K and all primes at which T is ramified.

We now let f, g be eigenforms of level N , with coefficients in a field L , as in Definition 6.4.1.

PROPOSITION 6.8.3. *Suppose $m \geq 1$ and there is no Dirichlet character ψ of conductor dividing mp^∞ such that $f \sim \bar{g} \otimes \psi$, where \sim signifies that these two eigenforms have the same Hecke eigenvalues away from their levels (i.e. correspond to the same newform). Then*

$$H^0(\mathbb{Q}(\mu_{mp^\infty}), V_{L_p}(f, g)^*) = 0.$$

Proof. The space $H^0(\mathbb{Q}(\mu_{mp^\infty}), V_{L_p}(f, g)^*)$ is preserved by the residual action of the abelian group $\text{Gal}(\mathbb{Q}(\mu_{mp^\infty})/\mathbb{Q})$, so if it is nonzero, it contains a subspace on which $\text{Gal}(\mathbb{Q}(\mu_{mp^\infty})/\mathbb{Q})$ acts by some character λ (possibly after a finite extension of the field L). This gives a nonzero $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ -equivariant homomorphism

$$V_{L_p}(f) \rightarrow V_{L_p}(g)^*(\lambda) = V_{L_p}(\bar{g})(\chi\lambda),$$

where χ is the cyclotomic character. Since both sides are irreducible representations of $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$, this map must be an isomorphism; consequently, $\psi = \chi\lambda$ has finite order and $f \sim \bar{g} \otimes \psi$. \square

We can now prove the main result of this section, which shows that the elements $(\alpha_f\alpha_g)^{-i} c_{mp^i}^{(f,g,N)}$ for $i \geq 0$ can be glued together into a (possibly unbounded) Euler system.

THEOREM 6.8.4. *Let $m, N \geq 1$ with $(m, p) = 1$, and let p be a prime dividing N . Let f, g be modular forms of level N that are eigenforms for all the Hecke operators, with U_p -eigenvalues α_f and α_g such that $h := v_p(\alpha_f\alpha_g) < 1$. Suppose that $f \not\sim \bar{g} \otimes \psi$ for all Dirichlet characters ψ of conductor dividing mp^∞ . Then for any r such that $h \leq r < 1$, there is a unique element*

$$c_{\mathfrak{d}_{m,r}}^{(f,g,N)} \in \mathcal{H}_r(\Gamma) \otimes_{\Lambda(\Gamma)} H_{\text{Iw},S}^1(\mathbb{Q}(\mu_{mp^\infty}), T_{\mathcal{O}_p}(f, g)^*)$$

whose projection to $H_S^1(\mathbb{Q}(\mu_{mp^i}), V_{L_p}(f, g)^*)$ is equal to

$$(\alpha_f\alpha_g)^{-i} c_{mp^i}^{(f,g,N)}$$

if $i \geq 1$ and to

$$(1 - (\alpha_f\alpha_g)^{-1}\sigma_p) c_{\mathfrak{d}_m}^{(f,g,N)}$$

if $i = 0$.

Moreover, if ℓ is a prime not dividing mN , the corestriction map sends $c_{\mathfrak{d}_{\ell m,r}}^{(f,g,N)}$ to

$$\sigma_\ell \left((\ell - 1)(1 - \varepsilon_f(\ell)\varepsilon_g(\ell)\sigma_\ell^{-2}) - \ell P_\ell(\ell^{-1}\sigma_\ell^{-1}) \right) c_{\mathfrak{d}_{m,r}}^{(f,g,N)}.$$

Proof. The existence of $c_{\mathfrak{d}_{m,r}}^{(f,g,N)}$ satisfying the projection formula for $i \geq 1$ is immediate from Corollary 6.4.5 and Proposition A.2.10. The projection formula for $i = 0$ follows from the $i = 0$ case of Corollary 6.4.5. \square

Note 6.8.5. The elements $c_{\mathfrak{d}_{m,r}}^{(f,g,N)}$ are in fact independent of $r \in [h, 1)$, in the sense that if $v_p(\alpha_f \alpha_g) \leq r < r' < 1$, then $c_{\mathfrak{d}_{m,r'}}^{(f,g,N)}$ is the image of $c_{\mathfrak{d}_{m,r}}^{(f,g,N)}$ under the natural map

$$\begin{aligned} \mathcal{H}_r(\Gamma) \otimes_{\Lambda(\Gamma)} H_{Iw,S}^1(\mathbb{Q}(\mu_{mp^\infty}), T_{\mathcal{O}_p}(f, g)^*) \\ \longrightarrow \mathcal{H}_{r'}(\Gamma) \otimes_{\Lambda(\Gamma)} H_{Iw,S}^1(\mathbb{Q}(\mu_{mp^\infty}), T_{\mathcal{O}_p}(f, g)^*) \end{aligned}$$

induced by the inclusion $\mathcal{H}_r(\Gamma) \hookrightarrow \mathcal{H}_{r'}(\Gamma)$.

In the case when $v_p(\alpha_f \alpha_g) = 0$, we can prove a stronger result; in this case we can dispense with the assumption that f is not a twist of \bar{g} , and we even get integral coefficients.

THEOREM 6.8.6. *Assume that f and g are eigenforms of level dividing N and such that $v_p(\alpha_f \alpha_g) = 0$. Then there is a unique element $c_{\mathfrak{d}_m}^{(f,g,N)} \in H_{Iw,S}^1(\mathbb{Q}(\mu_{mp^\infty}), T_{\mathcal{O}_p}(f, g)^*)$ whose projection to $H_S^1(\mathbb{Q}(\mu_{mp^i}), T_{\mathcal{O}_p}(f, g)^*)$ is equal to*

$$\begin{cases} (\alpha_f \alpha_g)^{-i} c_{\mathfrak{z}_{mp^i}}^{(f,g,N)} & \text{if } i \geq 1, \\ (1 - (\alpha_f \alpha_g)^{-1} \sigma_p) c_{\mathfrak{z}_m}^{(f,g,N)} & \text{if } i = 0. \end{cases}$$

Proof. For $i \geq 1$, let

$$c_{\mathfrak{d}_{m,i}}^{(f,g,N)} = (\alpha_f \alpha_g)^{-i} c_{\mathfrak{z}_{mp^i}}^{(f,g,N)}.$$

As $v_p(\alpha_f \alpha_g) = 0$, we have $c_{\mathfrak{d}_{m,i}}^{(f,g,N)} \in H_S^1(\mathbb{Q}(\mu_{mp^i}), T_{\mathcal{O}_p}(f, g)^*)$. Moreover, it is clear from Corollary 6.4.5 that

$$\text{cores}_{i/i-1}(c_{\mathfrak{d}_{m,i}}^{(f,g,N)}) = c_{\mathfrak{d}_{m,i-1}}^{(f,g,N)};$$

i.e., $c_{\mathfrak{d}_m}^{(f,g,N)} = \left(c_{\mathfrak{d}_{m,i}}^{(f,g,N)} \right)_{i \geq 1}$ defines an element in $H_{Iw,S}^1(\mathbb{Q}(\mu_{mp^\infty}), T_{\mathcal{O}_p}(f, g)^*)$. The projection formula for $i = 0$ follows as before. \square

Remark 6.8.7. The difficulties arising when $v_p(\alpha_f \alpha_g) \geq 1$ are somewhat reminiscent of the “critical slope” case in the Iwasawa theory of a single modular form over the cyclotomic tower; cf. [PS11], [PS13], [LZ13]. However, in our situation the conditions we must impose are more restrictive; in particular, given forms f', g' of level prime to p , at most two of the four possible choices of p -stabilizations f, g of f', g' will be possible, and in many cases (e.g. if $a_p(f') = a_p(g') = 0$) there are no valid choices at all.

Our methods using higher Chow groups can perhaps be thought of as an “algebraic avatar” of the modular symbol computations of [AV75]. It is interesting to speculate whether the overconvergent modular symbols of [PS11] also admit such an algebraic analogue, which could conceivably be applied in the critical-slope cases.

6.8.1. *Dispensing with c .* We now investigate the extent to which the “smoothing factor” c may be removed.

Notation. If R is an integral domain, we write $Q(R)$ for its field of fractions. For integers c and m such that $(c, m) = 1$, we write $[c]$ for the image of σ_c in $\mathbb{Z}_p[\text{Gal}(\mathbb{Q}(\mu_m)/\mathbb{Q})]$.

Under the hypotheses of Theorem 6.8.6, there exists an element

$$\mathfrak{z}_m^{(f,g,N)} \in H_{\text{Iw},S}^1(\mathbb{Q}(\mu_{mp^\infty}), T_{\mathcal{O}_p}(f, g)^*) \otimes_{\Lambda(\Gamma_1)} Q(\Lambda(\Gamma_1))$$

such that

$$(20) \quad c\mathfrak{z}_m^{(f,g,N)} = (c^2 - \varepsilon_f(c)^{-1}\varepsilon_g(c)^{-1}[c]^2)\mathfrak{z}_m^{(f,g,N)}.$$

Remark 6.8.8. Note that we are identifying Γ_1 with $\text{Gal}(\mathbb{Q}(\mu_{mp^\infty})/\mathbb{Q}(\mu_{mp}))$. We may define $\mathfrak{z}_m^{(f,g,N)}$ as

$$(d^2 - \varepsilon_f(d)^{-1}\varepsilon_g(d)^{-1}[d]^2)^{-1}d\mathfrak{z}_m^{(f,g,N)}$$

for any $d > 1$ coprime to $6N$ and congruent to 1 mod mp ; then $[d]$ lies in Γ_1 , so the expression is well defined, and it is evidently independent of the choice of d .

Notation. Write $\Gamma^{(m)} = \text{Gal}(\mathbb{Q}(\mu_{mp^\infty})/\mathbb{Q}) \cong (\mathbb{Z}/m\mathbb{Z})^\times \times \Gamma$.

LEMMA 6.8.9. *Let \mathfrak{p} be a prime ideal of $\Lambda(\Gamma^{(m)})$ of height 1 that does not contain p . If the conductor of the Dirichlet character $\varepsilon_f\varepsilon_g$ does not divide mp^∞ , then there exists an integer $c > 1$ coprime to $6mpN$ such that $c^2 - \varepsilon_f(c)^{-1}\varepsilon_g(c)^{-1}[c]^2 \notin \mathfrak{p}$.*

Proof. Since \mathfrak{p} does not contain p , it corresponds to a Galois orbit of continuous characters $\Gamma^{(m)} \rightarrow \overline{\mathbb{Q}}_p$. Let $\kappa_{\mathfrak{p}}$ be a representative of this orbit. Define $\tilde{h} : \Gamma^{(m)} \rightarrow \overline{\mathbb{Q}}_p^\times$ by $\tilde{h}(x) = \kappa_{\mathfrak{p}}(x)^2/\chi(x)^2$, where $\chi : \Gamma^{(m)} \rightarrow \mathbb{Z}_p^\times$ is the p -adic cyclotomic character. We need to show that there is an integer $c \geq 1$ coprime to $6mpN$ such that $\tilde{h}([c]) \neq \varepsilon_f(c)\varepsilon_g(c)$.

However, if no such integer existed, then $\varepsilon_f\varepsilon_g$ would have to factor through the natural map $\widehat{\mathbb{Z}}^\times \rightarrow \Gamma^{(m)}$, i.e., would have to have conductor dividing mp^∞ , contrary to our hypotheses. \square

COROLLARY 6.8.10. *If $\varepsilon_f\varepsilon_g$ does not have conductor dividing mp^∞ , then*

$$\mathfrak{z}_m^{(f,g,N)} \in H_{\text{Iw},S}^1(\mathbb{Q}(\mu_{mp^\infty}), T_{\mathcal{O}_p}(f, g)^*) \otimes \mathbb{Q}.$$

Proof. Let Z_0 be the $\Lambda(\Gamma^{(m)})$ -module generated by $c\mathfrak{z}_m^{(f,g,N)}$ for all possible c , and let Z_1 be the $\Lambda(\Gamma^{(m)})$ -module generated by $\mathfrak{z}_m^{(f,g,N)}$. By (20), $Z_0 \subset Z_1$ and there exists $\mu \in \Lambda(\Gamma^{(m)})$ such that $\mu Z_1 \subset Z_0$ and $Z_0/\mu Z_1$ is p -torsion free. Hence, it is enough to show that $Z_{0,\mathfrak{p}} = Z_{1,\mathfrak{p}}$ for any prime ideal \mathfrak{p} of height 1

that does not contain p . Fix such a \mathfrak{p} . By Lemma 6.8.9, there exists c such that $c^2 - \varepsilon_f(c)^{-1}\varepsilon_g(c)^{-1}[c]^2 \notin \mathfrak{p}$, so

$$\mathfrak{z}_m^{(f,g,N)} \in Z_{0,\mathfrak{p}}$$

by (20), as required. □

Remark 6.8.11. Note that if the mod p reduction of the Dirichlet character $\varepsilon_f\varepsilon_g$ does not have conductor dividing mp^∞ , then we can even deduce that

$$\mathfrak{z}_m^{(f,g,N)} \in H_{I_w,S}^1(\mathbb{Q}(\mu_{mp^\infty}), T_{\mathcal{O}_p}(f, g)^*).$$

LEMMA 6.8.12. *If the residual representation of $T_{\mathcal{O}_p}(f, g)^*$ restricted to $G_{\mathbb{Q}_{\mu_m}}$ is irreducible, then $H_{I_w,S}^1(\mathbb{Q}(\mu_{mp^\infty}), T_{\mathcal{O}_p}(f, g)^*)$ is a free $\Lambda(\Gamma^{(m)})$ -module.*

Proof. This follows from the argument of [Kat04, §13.8], which we briefly outline here. For simplicity, let $T = T_{\mathcal{O}_p}(f, g)^*$, $\Lambda = \Lambda(\Gamma^{(m)})$ and $H^1(T) = H_{I_w,S}^1(\mathbb{Q}(\mu_{mp^\infty}), T_{\mathcal{O}_p}(f, g)^*)$. It is enough to show that if (x, y) is a maximal ideal of Λ , the two maps

$$\alpha : H^1(T) \xrightarrow{x} H^1(T) \quad \text{and} \quad \beta : H^1(T)/xH^1(T) \xrightarrow{y} H^1(T)/xH^1(T)$$

are injective.

If $x = p$, the injectivity of α follows from the fact that

$$\varprojlim_n H^0(\mathbb{Q}(\mu_{mp^n}), T/p) = 0,$$

which is a consequence of the finiteness of T/p . If x is such that $\Lambda/x\Lambda$ is p -torsion free, it is enough to show that $H^0(\mathbb{Q}(\mu_m), T \otimes \Lambda/x\Lambda) = 0$, where the action of $\sigma \in G_{\mathbb{Q}(\mu_m)}$ on Λ is given by multiplication by $\bar{\sigma}^{-1}$, where $\bar{\sigma}$ denotes the image of σ in $\Gamma^{(m)}$. But T is irreducible, so non-abelian. This implies that $H^0(\mathbb{Q}(\mu_m), T \otimes \Lambda/x\Lambda) = 0$.

To show that β is injective, it is enough to show that $H^0(\mathbb{Z}[1/mp], T \otimes \Lambda/(x, y)) = 0$. But $\Lambda/(x, y) \cong \mathcal{O}_p/\mathfrak{M}_p(r)$ for some r , so we are done by the irreducibility of T/\mathfrak{M}_p . □

COROLLARY 6.8.13. *If $\varepsilon_f\varepsilon_g$ does not have p -power conductor and the residual representation of $T_{\mathcal{O}_p}(f, g)^*$ restricted to $G_{\mathbb{Q}_{\mu_m}}$ is irreducible, then*

$$\mathfrak{z}_m^{(f,g,N)} \in H_{I_w,S}^1(\mathbb{Q}(\mu_{mp^\infty}), T_{\mathcal{O}_p}(f, g)^*).$$

Proof. This follows from the argument in [Kat04, §13.14]. Let Z_1 be the $\Lambda(\Gamma^{(m)})$ -module generated by $\mathfrak{z}_m^{(f,g,N)}$. By the proof of Corollary 6.8.10, $Z_{1,\mathfrak{p}} \subset H_{I_w,S}^1(\mathbb{Q}(\mu_{mp^\infty})_{\mathfrak{p}})$ for any prime ideal \mathfrak{p} of $\Lambda(\Gamma^{(m)})$ height 1. But $H_{I_w,S}^1(\mathbb{Q}(\mu_{mp^\infty}))$ is a free $\Lambda(\Gamma^{(m)})$ -module by Lemma 6.8.12, hence the result. □

6.9. *Variation in Hida families.* We now make use of the first norm relation (Theorem 3.1.2) to build elements in the cohomology of *towers* of modular curves, under additional ordinarity hypotheses. Let us begin by recalling some of Ohta’s results in [Oht99], [Oht00] concerning the structure of the module

$$\text{GES}_p(N, \mathbb{Z}_p) := \varprojlim_{n \geq 1} H_{\text{ét}}^1(\overline{Y_1(Np^n)}, \mathbb{Z}_p),$$

which can be roughly summarized by the statement that one can build a Hida theory for this module after replacing the usual Hecke operators with their transposes.

We note that the full Hecke algebra does not act on $\text{GES}_p(N, \mathbb{Z}_p)$, since the operator $U_p = \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}$ does not commute with the trace maps. Rather, we obtain an action of the Hecke operator U'_p , corresponding to $\begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}$. Ohta shows that one may use the operator U'_p to define an “anti-ordinary projector” $e'_{\text{ord}} = \lim_{n \rightarrow \infty} (U'_p)^{n!}$, analogous to the usual Hida ordinary projector $e_{\text{ord}} = \lim_{n \rightarrow \infty} (U_p)^{n!}$.

Ohta proves the following control theorem for the anti-ordinary part of $\text{GES}_p(N)_{\mathbb{Z}_p}$, which is naturally a module over the Iwasawa algebra of $\Gamma_1 = (1 + p\mathbb{Z}_p)^\times$ via the diamond operators:

PROPOSITION 6.9.1 ([Oht99, 1.3, 1.4]). *The module $e'_{\text{ord}} \text{GES}_p(N)_{\mathbb{Z}_p}$ is free of finite rank over $\Lambda(\Gamma_1)$, and for each $r \geq 1$, there is a $G_{\mathbb{Q}}$ -equivariant isomorphism*

$$e'_{\text{ord}} \text{GES}_p(N, \mathbb{Z}_p) / \omega_r \cong e'_{\text{ord}} H^1(\overline{Y_1(Np^r)}, \mathbb{Z}_p),$$

where ω_r is the kernel of the natural map $\Lambda(\Gamma_1) \rightarrow \mathbb{Z}_p[(\mathbb{Z}/p^r\mathbb{Z})^\times]$.

From this isomorphism, we deduce that $e'_{\text{ord}} \text{GES}_p(N, \mathbb{Z}_p)$ has an action of the Hecke algebra

$$e'_{\text{ord}} \mathcal{H}'(N, \mathbb{Z}_p) = \varprojlim_{r \geq 1} e'_{\text{ord}} \mathcal{H}'(\Gamma_1(Np^r), \mathbb{Z}_p),$$

where $\mathcal{H}'(\Gamma_1(Np^r), \mathbb{Z}_p)$ is the \mathbb{Z}_p -subalgebra of $\text{End}_{\mathbb{Q}_p} M_2(\Gamma_1(Np^r), \mathbb{Q}_p)$ generated by the Hecke operators $T'(n)$ for $n \geq 1$ and $\langle q \rangle$ for $q \in (\mathbb{Z}/N\mathbb{Z})^\times$. Here $T'(\ell)$, for ℓ prime, corresponds to the double coset $\begin{pmatrix} \ell & 0 \\ 0 & 1 \end{pmatrix}$ so, in particular, for $(m, Np) = 1$, we have $T'(m) = \langle m \rangle^{-1} T(m)$ (the adjoint of $T(m)$ with respect to the Petersson product).

The algebra $e'_{\text{ord}} \mathcal{H}'(N, \mathbb{Z}_p)$ algebra is finite and free as a $\Lambda(\Gamma_1)$ -module. (Note that it is *not* generally free as a $\Lambda(\Gamma)$ -module, although it is evidently projective.)

PROPOSITION 6.9.2 ([Oht99, 2.2]). *The Hecke algebra $e'_{\text{ord}} \mathcal{H}'(N, \mathbb{Z}_p)$ is isomorphic to the Hecke algebra $\mathcal{H}_{\text{ord}}(N, \mathbb{Z}_p) := e_{\text{ord}} \mathcal{H}(N, \mathbb{Z}_p)$ acting on the*

module $e_{\text{ord}} M_2(N, \Lambda(\Gamma_1))$ of ordinary Λ -adic modular forms (not necessarily cuspidal), via the map sending $T(n)'$ to $T(n)$ and $\langle n \rangle$ to $\langle n \rangle^{-1}$.

Definition 6.9.3. In the above situation, by a Hida family of tame level N , we mean a maximal ideal of the ring $\mathcal{H}_{\text{ord}}(N, \mathbb{Z}_p)$. For each Hida family \mathbf{g} , we define

$$T(\mathbf{g})^* = (e'_{\text{ord}} \text{GES}_p(N, \mathbb{Z}_p))_{\mathbf{g}}(1).$$

COROLLARY 6.9.4. Let g be an ordinary weight 2 Hecke eigenform of level Np^s , with coefficients in some finite extension $L_{\mathfrak{p}}/\mathbb{Q}_p$ with ring of integers $\mathcal{O}_{\mathfrak{p}}$. Then we have an isomorphism of $\mathcal{O}_{\mathfrak{p}}$ -linear Galois representations

$$\mathcal{O}_{\mathfrak{p}} \otimes_{\mathcal{H}_{\text{ord}}(N, \mathbb{Z}_p)} T(\mathbf{g})^* \cong T_{\mathcal{O}_{\mathfrak{p}}}(g)^*,$$

where $T_{\mathcal{O}_{\mathfrak{p}}}(g)^*$ is the representation defined in 6.3 above.

Proof. Clear from the definition of $T_{\mathcal{O}_{\mathfrak{p}}}(g)^*$ and the control theorem (Theorem 6.9.1). \square

THEOREM 6.9.5. Let $N \geq 1$ be prime to p . If \mathbf{g} is a Hida family of tame level N , and f is any eigenform of level Np^k for $k \geq 1$ whose U_p -eigenvalue α_f satisfies $v_p(\alpha_f) < 1$, then for each integer $m \geq 1$, there is a cohomology class

$$c_{\mathbf{z}_m}^{(f, \mathbf{g})} \in H_S^1(\mathbb{Q}(\mu_m), T_{\mathcal{O}_{\mathfrak{p}}}(f)^* \otimes_{\mathbb{Z}_p} T(\mathbf{g})^*)$$

such that for each classical weight 2 specialization g of \mathbf{g} with coefficients in L , the image of $c_{\mathbf{z}_m}^{(f, \mathbf{g})}$ in

$$H^1(\mathbb{Q}(\mu_m), T_{\mathcal{O}_{\mathfrak{p}}}(f)^* \otimes_{\mathcal{O}_{\mathfrak{p}}} T_{\mathcal{O}_{\mathfrak{p}}}(g)^*) = H^1(\mathbb{Q}(\mu_m), T_{\mathcal{O}_{\mathfrak{p}}}(f, g)^*)$$

is the generalized Beilinson–Flach element $c_{\mathbf{z}_m}^{(f, g, N')}$, where N' is the greatest common divisor of the levels of f and g .

Proof. We know that the elements $c_{\Xi_{m, Np^s, 1}}$ for $s \geq 1$ are unramified outside S and are compatible under pushforward via the natural projection maps. Hence the sequence of elements defined by pushing forward $c_{\Xi_{m, Np^s, 1}}$ to $\text{CH}^2(Y_1(Np^r) \times Y_1(Np^s) \times \mathbb{Q}(\mu_m), 1)$, for $s \geq r$, are compatible under pushforward maps in the $Y_1(Np^s)$ factor alone. Applying the étale regulator, we obtain elements of the module

$$\varprojlim_{s \geq r} H^1(\mathbb{Q}(\mu_m), H_{\text{ét}}^2(\overline{Y_1(Np^r)} \times \overline{Y_1(Np^s)}, \mathbb{Z}_p)(2)).$$

For each s , we may decompose $H_{\text{ét}}^2(\overline{Y_1(Np^r)} \times \overline{Y_1(Np^s)}, \mathbb{Z}_p)$ as the tensor product of the H^1 's of the two factors, using the Künneth formula. Projecting to the quotient $T_{\mathcal{O}_{\mathfrak{p}}}(f)$ of $H_{\text{ét}}^1(\overline{Y_1(Np^r)}, \mathbb{Z}_p)(1)$ and applying the anti-ordinary projector e'_{ord} to $H_{\text{ét}}^1(\overline{Y_1(Np^s)}, \mathbb{Z}_p)(1)$, we may argue exactly as in Proposition 6.6.2 to deduce that the elements we obtain are unramified outside Np .

Since the restricted-ramification cohomology groups $H_S^i(\mathbb{Q}(\mu_m), -)$ commute with inverse limits, we obtain an element of

$$H_S^1(\mathbb{Q}(\mu_m), T_{\mathcal{O}_p}(f)^* \otimes e'_{\text{ord}} \text{GES}_p(N)_{\mathbb{Z}_p}(1)).$$

Pushing forward along the canonical map $e'_{\text{ord}} \text{GES}_p(N)_{\mathbb{Z}_p}(1) \rightarrow T(\mathbf{g})^*$, we obtain the required elements. \square

We also obtain a corresponding result for the product of two Hida families, whose proof is essentially identical to the above:

THEOREM 6.9.6. *Let $N \geq 1$ be prime to p . If \mathbf{f}, \mathbf{g} are Hida families of tame level N , then for each integer $m \geq 1$, there is a cohomology class*

$$c_{\mathbf{z}_m}^{(\mathbf{f}, \mathbf{g})} \in H_S^1(\mathbb{Q}(\mu_m), T(\mathbf{f})^* \hat{\otimes}_{\mathbb{Z}_p} T(\mathbf{g})^*)$$

such that for classical weight 2 specializations f, g of \mathbf{f}, \mathbf{g} with coefficients in L , the image of $c_{\mathbf{z}_m}^{(\mathbf{f}, \mathbf{g})}$ in

$$H^1(\mathbb{Q}(\mu_m), T_{\mathcal{O}_p}(f)^* \otimes_{\mathcal{O}_p} T_{\mathcal{O}_p}(g)^*) = H^1(\mathbb{Q}(\mu_m), T_{\mathcal{O}_p}(f, g)^*)$$

is the generalized Beilinson–Flach element $c_{\mathbf{z}_m}^{(f, g, N')}$, where N' is the greatest common divisor of the levels of f and g .

Remark 6.9.7. We do not know if one can formulate a result analogous to Theorem 6.8.4 incorporating Hida-family variation in g since we do not know whether the results of Appendix A.2 apply for “big” Galois representations; but if f is ordinary, there are no such issues.

THEOREM 6.9.8. *In the situation of Theorem 6.9.6, for each m prime to p , there exists a cohomology class*

$$c_{\mathbf{z}_m}^{(\mathbf{f}, \mathbf{g})} \in H_{1w, S}^1(\mathbb{Q}(\mu_{mp^\infty}), T(\mathbf{f})^* \hat{\otimes}_{\mathbb{Z}_p} T(\mathbf{g})^*)$$

whose image in $H_{1w, S}^1(\mathbb{Q}(\mu_{mp^i}), T(\mathbf{f})^* \hat{\otimes}_{\mathbb{Z}_p} T(\mathbf{g})^*)$ for each $i \geq 1$ is equal to $(\alpha_f \alpha_g)^{-i} \cdot c_{\mathbf{z}_{mp^i}}^{(\mathbf{f}, \mathbf{g})}$.

6.10. Integrality of the Poincaré pairing. In this section, we prove a technical lemma that will be needed in our applications to bounding Selmer groups. We assume that $p > 3$, $N \geq 5$, and $p \nmid N$.

Recall that $X_1(N)$ admits a canonical smooth proper model over $\mathbb{Z}[1/N]$ ([DR73]) and hence over \mathbb{Z}_p . By [FM87], the integral de Rham cohomology $H^1(X_1(N), \Omega_{X_1(N)/\mathbb{Z}_p}^\bullet)$ is a filtered Dieudonné module over \mathbb{Z}_p , and

$$T(H^1(X_1(N), \Omega_{X_1(N)/\mathbb{Z}_p}^\bullet)) = H_{\text{ét}}^1(\overline{X_1(N)}, \mathbb{Z}_p),$$

where $T(-)$ is the Fontaine–Laffaille functor.

We define versions of these in the f -isotypical component by projection. As in Remark 6.3.4, we define $\widetilde{T}_{\mathcal{O}_p}(f)^*$ to be the image of $H_{\text{ét}}^1(\overline{X_1(N)}, \mathbb{Z}_p) \otimes \mathcal{O}_p$ in $V_{L_p}(f)^*$, and similarly for g . We define $\mathbb{D}_{\text{cris}}(\widetilde{T}_{\mathcal{O}_p}(f)^*)$ as the image of $H^1(X_1(N), \Omega_{X_1(N)/\mathbb{Z}_p}^\bullet) \otimes_{\mathbb{Z}_p} \mathcal{O}_p$ in $\mathbb{D}_{\text{cris}}(V_{L_p}(f)^*)$; then $\mathbb{D}_{\text{cris}}(\widetilde{T}_{\mathcal{O}_p}(f)^*)$ is a strongly divisible \mathcal{O}_p -lattice, and its image under $T(-)$ is $\widetilde{T}_{\mathcal{O}_p}(f)^*$.

Let us recall here the definition of η_f^{ur} .

Definition 6.10.1. Let $X = X_1(N)$ and

$$\eta_f^{\text{ah}} = \frac{\bar{f}^*(z)d\bar{z}}{\langle f^*, f^* \rangle_{k,N}} \in H_{\text{dR}}^1(X_{\mathbb{C}}).$$

We denote by η_f its image in $H^1(X/\mathbb{C}, \mathcal{O}_{X/\mathbb{C}})$, which lies in $H^1(X/\mathbb{Q}_p, \mathcal{O}_{X/\mathbb{Q}_p})$; then η_f^{ur} is defined to be the lift of η_f to the unit root subspace of $H_{\text{dR}}^1(X_{\mathbb{C}_p})^{f, \text{ur}}$.

Our aim is to investigate the denominator of the class η_f^{ur} relative to the sublattice

$$\mathcal{O}_L \otimes_{\mathbb{Z}_p} H^1(X_1(N), \Omega_{X_1(N)/\mathbb{Z}_p}^\bullet) \subseteq L_p \otimes_{\mathbb{Q}_p} H^1(X_1(N), \Omega_{X_1(N)/\mathbb{Q}_p}^\bullet).$$

Since the unit root lifting is obviously integral, it suffices to show that $\eta_f \in \mathcal{O}_L \otimes_{\mathbb{Z}_p} H^1(X_1(N), \mathcal{O}_{X_1(N)/\mathbb{Z}_p})$.

PROPOSITION 6.10.2. *An element of*

$$L_p \otimes H^1(X_1(N)/\mathbb{Q}_p, \mathcal{O}_{X_1(N)/\mathbb{Q}_p})$$

lies in the sublattice

$$\mathcal{O}_p \otimes H^1(X_1(N), \mathcal{O}_{X_1(N)/\mathbb{Z}_p})$$

if and only if it pairs to an element of \mathcal{O}_p with all elements of

$$\mathcal{O}_p \otimes H^0(X_1(N)/\mathbb{Z}_p, \Omega_{X_1(N)/\mathbb{Z}_p}^1).$$

Proof. Since the pairing between

$$H^1(X_1(N), \mathcal{O}_{X_1(N)/\mathbb{Z}_p}) \text{ and } H^0(X_1(N)/\mathbb{Z}_p, \Omega_{X_1(N)/\mathbb{Z}_p}^1)$$

is defined over \mathbb{Z}_p , it suffices to assume $\mathcal{O}_p = \mathbb{Z}_p$. But Serre duality shows that this pairing is perfect, i.e., identifies $H^0(X_1(N)/\mathbb{Z}_p, \Omega_{X_1(N)/\mathbb{Z}_p}^1)$ with the \mathbb{Z}_p -dual of $H^1(X_1(N)/\mathbb{Z}_p, \mathcal{O}_{X_1(N)/\mathbb{Z}_p})$. \square

LEMMA 6.10.3. *Let $\phi \in S_2(N; L_p)$. Then the element ω_ϕ of $H^0(X/L, \Omega_{X/L}^1)$ lies in $H^0(X/\mathcal{O}_L, \Omega_{X/\mathcal{O}_L}^1)$ if and only if $\phi \in S_2(N; \mathcal{O}_p)$.*

Proof. We have by definition $\omega_\phi(q) = \phi(q)dq/q$, which is defined over \mathcal{O}_L if ϕ is. \square

Definition 6.10.4. If $f \in S_2(\Gamma_1(N), L)$, let I_f denote the ideal in \mathcal{O} such that

$$\left\{ \frac{\langle f^*, \phi \rangle}{\langle f^*, f^* \rangle} : \phi \in S_2(N, \mathcal{O}) \right\} = I_f^{-1}.$$

Remark 6.10.5. Note that I_f^{-1} contains \mathcal{O} , so I_f is an integral ideal (rather than a fractional ideal). The ideal I_f essentially measures the extent to which f is congruent to other eigenforms in $S_2(N, \mathcal{O})$.

COROLLARY 6.10.6. *For any prime $\mathfrak{p} \nmid N$, we have*

$$\eta_f \in I_f^{-1} \cdot \mathcal{O}_{\mathfrak{p}} \otimes_{\mathbb{Z}_p} H^1(X_1(N), \mathcal{O}_{X_1(N)/\mathbb{Z}_p}).$$

Proof. By the construction of the class η_f , for any $\varphi \in S_2(\Gamma_1(N), \mathcal{O})$, we have

$$\langle \eta_f, \omega_{\varphi} \rangle = \frac{\langle f^*, \phi \rangle}{\langle f^*, f^* \rangle} \in I_f^{-1} \mathcal{O},$$

so the result follows by Lemma 6.10.3 and Proposition 6.10.2. \square

COROLLARY 6.10.7. *The linear functional*

$$\mathbb{D}_{\text{cris}}(V_{L_{\mathfrak{p}}}(f, g)^*) \rightarrow L_{\mathfrak{p}}$$

given by pairing with $\eta_f^{\text{ur}} \otimes \omega_g$ maps the submodule

$$\mathbb{D}_{\text{cris}}(\tilde{T}_{\mathcal{O}_{\mathfrak{p}}}(f)^*) \otimes \mathbb{D}_{\text{cris}}(\tilde{T}_{\mathcal{O}_{\mathfrak{p}}}(g)^*)$$

into $I_f^{-1} \mathcal{O}_{\mathfrak{p}}$.

PROPOSITION 6.10.8. *Let $z \in H_f^1(\mathbb{Q}_p, [\tilde{T}_{\mathcal{O}_{\mathfrak{p}}}(f) \otimes \tilde{T}_{\mathcal{O}_{\mathfrak{p}}}(g)]^*)$. Then*

$$\langle \log(z), \eta_f^{\text{ur}} \otimes \omega_g \rangle \in I_f^{-1} \cdot (1 - \alpha^{-1} \gamma^{-1})^{-1} (1 - \alpha^{-1} \delta^{-1})^{-1} \cdot \mathcal{O}_{\mathfrak{p}},$$

where α is the unit root of the Hecke polynomial of f and β, δ are the roots of the Hecke polynomial of g .

Proof. By Fontaine–Laffaille theory, for any crystalline $\mathcal{O}_{\mathfrak{p}}$ -linear $G_{\mathbb{Q}_p}$ -representation V whose Hodge filtration has length $< p$ and such that we have $\mathbb{D}_{\text{cris}}(V)^{\varphi=1} = 0$, the map

$$\log_{\mathbb{Q}_p, V} : H_f^1(\mathbb{Q}_p, V) \xrightarrow{\cong} \frac{\mathbb{D}_{\text{cris}}(V)}{\text{Fil}^0 \mathbb{D}_{\text{cris}}(V)}$$

induces an isomorphism of $\mathcal{O}_{\mathfrak{p}}$ -modules

$$\frac{H_f^1(\mathbb{Q}_p, T)}{\text{torsion}} \xrightarrow{\cong} \frac{(1 - \varphi)^{-1} D}{(1 - \varphi)^{-1} D \cap \text{Fil}^0 \mathbb{D}_{\text{cris}}(V)}$$

for any $G_{\mathbb{Q}_p}$ -stable lattice $T \subseteq V$ with corresponding strongly divisible lattice $D \subseteq \mathbb{D}_{\text{cris}}(V)$; cf. Theorem 4.1 and Lemma 4.5 of [BK90].

In our case we may take $V = W \otimes V_{L_p}(g)^*$, where W is the 1-dimensional unramified quotient of $V_{L_p}(f)^*$, since the linear functional given by pairing with $\eta_f^{\text{ur}} \otimes \omega_g$ factors through this quotient. Let us suppose that g is ordinary; using the explicit description of the strongly divisible lattices in $\mathbb{D}_{\text{cris}}(V_{L_p}(g)^*)$ given in [LZ13, §5], one checks that

$$\frac{(1 - \varphi)^{-1}D}{(1 - \varphi)^{-1}D \cap \text{Fil}^0 \mathbb{D}_{\text{cris}}(V)} \subseteq p^{-k} \cdot \frac{D}{D \cap \text{Fil}^0 \mathbb{D}_{\text{cris}}(V)},$$

where $k = v_p [(1 - \alpha^{-1}\gamma^{-1})(1 - \alpha^{-1}\delta^{-1})]$. In the nonordinary case one reasons similarly using the description of the Wach module of the (unique up to scaling) lattice in $\mathbb{D}_{\text{cris}}(V_{L_p}(g)^*)$ given in [BLZ04]. Combining this with Corollary 6.10.7 gives the result. \square

7. Bounding strict Selmer groups

Let f, g be newforms of weight 2, level N and characters χ_f and χ_g , respectively. Let L be the subfield of $\overline{\mathbb{Q}}$ generated by the coefficients of f and g . For a prime \mathfrak{p} of L , denote by $V_{L_p}(f)$ and $V_{L_p}(g)$ the L_p -representations of $G_{\mathbb{Q}}$ attached to f and g , respectively. The aim of this section is to apply Theorem 7.1.5 to the representation $V_{L_p}(f) \otimes V_{L_p}(g)$.

7.1. *The method of Euler systems.* We recall some definitions and results from [Rub00]. Let \mathcal{O} be the ring of integers of a finite extension E/\mathbb{Q}_p , and let T be a free \mathcal{O} -module of finite rank with a continuous action of $G_{\mathbb{Q}}$ that is unramified at almost all primes. Let $V = T \otimes_{\mathcal{O}} E$ and $W = V/T = T \otimes_{\mathcal{O}} E/\mathcal{O}$.

Let Σ be a finite set of primes containing p and all prime numbers at which the action of $G_{\mathbb{Q}}$ on T ramifies. Let A be a set of integers such that

- if $m \in A$, then all divisors of m are in A ;
- if $r, s \in A$, then $LCM(r, s) \in A$;
- $\ell \in A$ for all primes $\ell \notin \Sigma$.

For a prime $\ell \notin \Sigma$, define

$$p_{\ell}(X) = \det_E (1 - \text{Frob}_{\ell}^{-1} X | V^*(1)) \in \mathbb{Z}_p[X],$$

where Frob_{ℓ} is the arithmetic Frobenius at ℓ .

Definition 7.1.1 (cf. [Rub00, Def. 2.1.1]). A Euler system for (T, A, Σ) is a system of elements

$$c_m \in H^1(\mathbb{Q}(\mu_m), T) \text{ for all } m \in A$$

such that if ℓ is a prime such that $m, m\ell \in A$ and $\ell \neq \Sigma$, then the corestriction map

$$H^1(\mathbb{Q}(\mu_{m\ell}), T) \rightarrow H^1(\mathbb{Q}(\mu_m), T)$$

sends $c_{\ell m}$ to

$$\begin{cases} p_{\ell}(\sigma_{\ell}^{-1})c_m & \text{if } \ell \nmid m \text{ and } \ell \neq \Sigma, \\ c_m & \text{if } \ell \mid m \text{ or } \ell \in \Sigma. \end{cases}$$

Here, σ_{ℓ} denotes the arithmetic Frobenius of ℓ in $\text{Gal}(\mathbb{Q}(\mu_m)/\mathbb{Q})$.

Remark 7.1.2. Our notation differs slightly from that of [Rub00]. Firstly, Rubin writes T^* for the ‘‘Tate dual’’ $\text{Hom}(T, \mathbb{Z}_p(1))$, while we write this as $T^*(1)$. More significantly, Rubin considers an infinite abelian extension \mathcal{K} and a class c_F for every finite subextension F of \mathcal{K} ; in our case \mathcal{K} is the extension $\mathcal{K}_A = \mathbb{Q}(\mu_r : r \in A)$, and it suffices to specify a class for each subextension of the form $\mathbb{Q}(\mu_m)$, which is our c_m , and to fill in the remainder via corestriction.

Definition 7.1.3. For each prime ℓ , let $H_f^1(\mathbb{Q}_{\ell}, W)$ be the image of $H_f^1(\mathbb{Q}_{\ell}, V)$ in $H^1(\mathbb{Q}_{\ell}, W)$. Define

$$\mathcal{S}^{\{p\}}(\mathbb{Q}, W) = \ker \left(H^1(\mathbb{Q}, W) \longrightarrow \bigoplus_{\ell \neq p} H^1(\mathbb{Q}_{\ell}, W) / H_f^1(\mathbb{Q}_{\ell}, W) \right),$$

and define the *strict Selmer group* of W over \mathbb{Q} as

$$\mathcal{S}_{\{p\}}(\mathbb{Q}, W) = \ker \left(\mathcal{S}^{\{p\}}(\mathbb{Q}, W) \longrightarrow H^1(\mathbb{Q}_p, W) \right).$$

(Thus $\mathcal{S}_{\{p\}}$ is the Selmer group with local conditions given by the Bloch–Kato condition at primes away from p and the zero local condition at p .)

We define $\mathcal{S}^{\{p\}}(\mathbb{Q}, T)$ similarly, and also $\mathcal{S}^{\{p\}}(K, T)$ similarly, for any number field K . (We shall only need this when $K = \mathbb{Q}(\mu_m)$, see Hypothesis $\text{Hyp}(\mathcal{S}^{\{p\}}, V)$ below.)

In order to state the main theorem, we introduce the following sets of hypotheses. Note that $\text{Hyp}(\mathbb{Q}, T)$ is strictly stronger than $\text{Hyp}(\mathbb{Q}, V)$, but $\text{Hyp}(p, A)$ and $\text{Hyp}(\mathcal{S}^{\{p\}}, V)$ are independent of each other.

Hypothesis ($\text{Hyp}(\mathbb{Q}, T)$). $T \otimes \mathbf{k}$ is an irreducible $\mathbf{k}[G_{\mathbb{Q}}]$ -module, where \mathbf{k} is the residue field of \mathcal{O} , and there exists an element $\tau \in G_{\mathbb{Q}}$ that satisfies the following conditions:

- (i) τ acts trivially on μ_{p^∞} ;
- (ii) $T/(\tau - 1)T$ is free of rank 1 over \mathcal{O} .

Hypothesis ($\text{Hyp}(\mathbb{Q}, V)$). V is an irreducible $E[G_{\mathbb{Q}}]$ -module, and there exists an element $\tau \in G_{\mathbb{Q}}$ that satisfies the following conditions:

- (i) τ acts trivially on μ_{p^∞} ;
- (ii) $\dim_{\mathbb{Q}_p}(V/(\tau - 1)V) = 1$.

Hypothesis ($\text{Hyp}(p, A)$). The set A contains all powers of p .

Hypothesis ($\text{Hyp}(\mathcal{S}^{\{p\}}, V)$). The following three conditions hold:

- (i) $T^{G_{\mathbb{Q}}} = 0$;
- (ii) $c_m \in \mathcal{S}^{\{p\}}(\mathbb{Q}(\mu_m), T)$ for all $m \in A$;

(iii) there exists an element $\gamma \in G_{\mathbb{Q}}$ such that

- γ acts trivially on μ_{p^∞} ,
- $\gamma - 1$ is injective on T .

THEOREM 7.1.4. *Assume that V is not the trivial representation, and that Hypothesis $\text{Hyp}(\mathbb{Q}, V)$ and at least one of hypotheses $\text{Hyp}(p)$ and $\text{Hyp}(\mathcal{S}^{(p)}, V)$ are satisfied. If $\mathbf{c} = (c_m)_{m \in A}$ is a Euler system for (T, A, Σ) and the image of c_1 in $H^1(\mathbb{Q}, T)$ is not contained in $H^1(\mathbb{Q}, T)_{\text{tors}}$, then $\mathcal{S}_{\{p\}}(\mathbb{Q}, W^*(1))$ is finite.*

THEOREM 7.1.5. *Assume that $p > 2$ and that Hypothesis $\text{Hyp}(\mathbb{Q}, T)$ and at least one of hypotheses $\text{Hyp}(p)$ and $\text{Hyp}(\mathcal{S}^{(p)}, V)$ are satisfied. If $\mathbf{c} = (c_m)_{m \in A}$ is a Euler system for (T, A, Σ) , then*

$$\text{length}_{\mathcal{O}}(\mathcal{S}_{\{p\}}(\mathbb{Q}, W^*(1))) \leq \text{ind}_{\mathcal{O}}(\mathbf{c}) + \mathbf{n}_W + \mathbf{n}_W^*,$$

where $\text{ind}_{\mathcal{O}}(\mathbf{c})$ is the largest power of the maximal ideal by which c_1 can be divided in $H^1(\mathbb{Q}, T)/\text{torsion}$ and the quantities \mathbf{n}_W and \mathbf{n}_W^* are as defined in Theorem 2.2.2 of [Rub00].

Proofs. If $\text{Hyp}(p, A)$ holds, then Theorems 7.1.4 and 7.1.5 are Theorems 2.2.3 and 2.2.2 of [Rub00] respectively. If $\text{Hyp}(\mathcal{S}^{(p)}, V)$ holds instead, then the necessary modifications to the proofs are outlined in Section 9.1 of op. cit. \square

7.2. Verifying the hypotheses on T . The main result of this section is Proposition 7.2.18, which implies that under some mild technical assumptions there is a large supply of primes where the condition $\text{Hyp}(\mathbb{Q}, T)$ is satisfied.

7.2.1. Big image results for one modular form. We begin by some results from [Mom81] and [Rib85] regarding the image of the Galois representations attached to a modular form. Let $f = \sum_{n \geq 1} a_n q^n$ be a new eigenform of weight $k \geq 2$, level N and character ϵ , not of CM type. Let $L = \mathbb{Q}(a_n : n \geq 1)$ be its coefficient field, with ring of integers \mathcal{O}_L .

Recall that an *extra twist* of f is an element $\gamma \in \text{Gal}(L/\mathbb{Q})$ such that $\gamma(f)$ is equal to the twist of f by some Dirichlet character χ_γ . We let $\Gamma \subseteq \text{Gal}(L/\mathbb{Q})$ be the group of such γ . We let $F \subseteq L$ be the fixed field of Γ , and we let $H \subseteq \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ be the absolute Galois group of the finite abelian extension K cut out by the Dirichlet characters χ_γ .

For each prime λ of L , it is clear that the trace of the Galois representation $\rho_{L_\lambda}(f)|_H$ takes values in F_μ , where μ is the prime of F below λ .

THEOREM 7.2.1 (Momose–Ribet; see [Rib85, Th. 3.1]). *For all but finitely many λ , the image of the Galois representation $\rho_{L_\lambda}(f)|_H$ is a conjugate of the group*

$$\{g \in \text{GL}_2(\mathcal{O}_{F,\mu}) : \det(g) \in \mathbb{Z}_\ell^\times\},$$

where μ and ℓ are the primes of F and \mathbb{Q} below λ .

Remark 7.2.2. For a “generic” modular form f , there will be no extra twists if the character f is trivial, but there will always be at least one if f has nontrivial character since the complex conjugate f^* is a twist of f .

We will need the following slight strengthening:

PROPOSITION 7.2.3. *Let K' be any finite extension of K that is abelian over \mathbb{Q} , and let $H' \subseteq H$ be its absolute Galois group. Then for all but finitely many λ , the image of $\rho_{L_\lambda}(f)|_{H'}$ is a conjugate of the group $\{g \in \mathrm{GL}_2(\mathcal{O}_{F,\mu}) : \det(g) \in \mathbb{Z}_\ell^\times\}$ above.*

Proof. If λ is a prime satisfying the conclusion of the theorem, then the image of $\rho_{L_\lambda}(f)|_{H'}$ contains $\mathrm{SL}_2(\mathcal{O}_{F,\mu})$ since $\mathrm{SL}_2(\mathcal{O}_{F,\mu})$ is equal to its own commutator subgroup. But for all but finitely many primes ℓ , the field K' is linearly disjoint from $\mathbb{Q}(\mu_{\ell^\infty})$ and thus the cyclotomic character is a surjection $H' \rightarrow \mathbb{Z}_\ell$. □

7.2.2. Big image results for pairs of modular forms. We recall the following result from group theory:

PROPOSITION 7.2.4 (Goursat’s Lemma; cf. [Lan02, Exercise I.5]). *Let G_1, G_2 be groups and H a subgroup of $G = G_1 \times G_2$ such that the projections $\pi_i : H \rightarrow G_i$ are surjective. Let $N_1 = H \cap (G_1 \times \{e_2\})$ and $N_2 = H \cap (\{e_1\} \times G_2)$, which we identify with subgroups of G_1, G_2 in the obvious manner. Then the N_i are normal in G_i , and H is the graph of an isomorphism $G_1/N_1 \cong G_2/N_2$.*

COROLLARY 7.2.5. *Let \mathbb{F}, \mathbb{F}' be finite fields of the same characteristic, both of order ≥ 4 . Let H be a subgroup of $\mathrm{SL}_2(\mathbb{F}) \times \mathrm{SL}_2(\mathbb{F}')$ surjecting onto both factors. Then either H is the whole of $\mathrm{SL}_2(\mathbb{F}) \times \mathrm{SL}_2(\mathbb{F}')$, or $\mathbb{F} = \mathbb{F}'$ and H is conjugate in $\mathrm{GL}_2(\mathbb{F}) \times \mathrm{GL}_2(\mathbb{F})$ to one of the following subgroups:*

- (i) *the diagonal subgroup $\{(x, \varphi^j(x)) : x \in G\}$ for some $0 \leq j < k$, where $\mathbb{F} = \mathbb{F}_{p^k}$ and φ is the p -power Frobenius of \mathbb{F} ;*
- (ii) *the subgroup $\{(x, y) : y = \pm \varphi^j(x)\}$ for some $0 \leq j < k$.*

Proof. This follows immediately from Goursat’s lemma and a case-by-case check, given that the groups $\mathrm{PSL}_2(\mathbb{F})$ for fields \mathbb{F} of order ≥ 4 are pairwise nonisomorphic simple groups, and the automorphism groups of $\mathrm{SL}_2(\mathbb{F})$ and $\mathrm{PSL}_2(\mathbb{F})$ are both isomorphic to $\mathrm{PGL}_2(\mathbb{F}) \rtimes \langle \varphi \rangle$. □

PROPOSITION 7.2.6. *Let $\mathcal{O}, \mathcal{O}'$ be the rings of integers of any two unramified extensions of \mathbb{Q}_p , where p is a prime ≥ 5 , with residue fields \mathbb{F}, \mathbb{F}' . Then any closed subgroup $H \subseteq \mathrm{SL}_2(\mathcal{O}) \times \mathrm{SL}_2(\mathcal{O}')$ that surjects onto $\mathrm{SL}_2(\mathbb{F}) \times \mathrm{SL}_2(\mathbb{F}')$ must be the whole of $\mathrm{SL}_2(\mathcal{O}) \times \mathrm{SL}_2(\mathcal{O}')$.*

Proof. We follow the argument given for $\mathrm{SL}_2(\mathbb{Z}_p)$ by Swinnerton-Dyer in [SD73]. It suffices to show that for each $n \geq 2$, the image H_n of H in $\mathrm{SL}_2(\mathcal{O}/p^n) \times \mathrm{SL}_2(\mathcal{O}'/p^n)$ contains the subgroups $K_n \times 1$ and $1 \times K'_n$, where K_n

is the kernel of $\mathrm{SL}_2(\mathcal{O}/p^n) \rightarrow \mathrm{SL}_2(\mathcal{O}/p^{n-1})$ and similarly for K'_n . Note that for each n , the group K_n is abelian and is isomorphic (via $m \mapsto 1 + p^{n-1}m$) to the group of trace zero matrices in $M_2(\mathbb{F})$, which is generated by matrices u such that $u^2 = 0$.

We now proceed by induction on n . Let $u \in M_2(\mathbb{F})$ satisfy $u^2 = 0$. By assumption, we may then find $h \in H$ congruent to $(1 + u, 1)$ modulo p ; and, as shown in op. cit., we have $h^p = (1 + pu, 1) \bmod p^2$. Thus $(1 + pu, 1) \in H_2$, and thus $H_2 \supseteq K_2 \times 1$. Similarly, H_2 contains $1 \times K'_2$, so in fact H_2 is the whole of $\mathrm{SL}_2(\mathcal{O}/p^2) \times \mathrm{SL}_2(\mathcal{O}'/p^2)$.

Suppose $n \geq 3$ and H_{n-1} is everything. We claim H_n contains $K_n \times 1$. Again, K_n consists of matrices of the form $(1 + p^{n-1}u, 1)$, and by the induction assumption we can find $h \in H$ congruent to $(1 + p^{n-2}u, 1)$ modulo p^{n-1} . Then h^p is congruent to $(1 + p^{n-1}u, 1)$ modulo p^n , so $(1 + p^{n-1}u, 1) \in H_n$. Thus $H_n \supseteq K_n \times 1$ and similarly $H_n \supseteq 1 \times K'_n$, so we are done. \square

As a corollary, we obtain the following result:

PROPOSITION 7.2.7. *Let $\mathcal{O}, \mathcal{O}'$ be as above, with characteristic ≥ 5 , and let H be a subgroup of $\mathrm{SL}_2(\mathcal{O}) \times \mathrm{SL}_2(\mathcal{O}')$ that surjects onto both factors. Then either $H = \mathrm{SL}_2(\mathcal{O}) \times \mathrm{SL}_2(\mathcal{O}')$, or $\mathcal{O}' = \mathcal{O}$ and H is contained in the subgroup*

$$\{(x, y) \in \mathrm{SL}_2(\mathcal{O}) : x = \pm\varphi^j y \bmod p\}$$

for some j .

We shall now boost this to a statement about GL_2 . For $\mathcal{O}, \mathcal{O}'$ as before, let G denote the group

$$\{(x, y) \in \mathrm{GL}_2(\mathcal{O}) \times \mathrm{GL}_2(\mathcal{O}') : \det(x) = \det(y) \in \mathbb{Z}_p^\times\}.$$

We can regard this as a fibre product $G_1 \times_{\mathbb{Z}_p^\times} G_2$, where $G_1 = \{x \in \mathrm{GL}_2(\mathcal{O}) : \det(x) \in \mathbb{Z}_p^\times\}$ and similarly for G_2 .

PROPOSITION 7.2.8. *Let H be a subgroup of G that surjects onto G_1 and G_2 . Then either $H = G$, or we have $\mathcal{O} = \mathcal{O}'$ and H is contained in the subgroup of G given by $\{(x, y) : x = \pm\varphi^j y \bmod p\}$ for some j .*

Proof. Let $G^\circ = \mathrm{SL}_2(\mathcal{O}) \times \mathrm{SL}_2(\mathcal{O}')$, and let $H^\circ = H \cap G^\circ$. Then H° has full image in each of $\mathrm{SL}_2(\mathcal{O})$ and $\mathrm{SL}_2(\mathcal{O}')$, so either $H^\circ = G^\circ$, or $\mathcal{O}' = \mathcal{O}$ and the image of H° modulo p is contained in the subgroup $\{(x, y) : x = \pm\varphi^j y \bmod p\}$ for some j .

Suppose first that $H^\circ = G^\circ$. Then we must have $H = G$, since for each $g \in G$, there is some $h \in H$ with $\det(h) = \det(g)$, and then $h^{-1}g$ lies in G° so by assumption it must be in H .

In the remaining case, by replacing H with its image under the automorphism $\varphi^j \times 1$, we may assume without loss of generality that $j = 0$. Then any

$h \in H^\circ$ is of the form (x, y) with $x = \pm y$ modulo p . Let (x, y) be any element of H , and consider the class of $t = x^{-1}y$ in $\mathrm{PSL}_2(\mathbb{F})$; then for any $(u, v) \in H^\circ$, we have

$$[u^{-1}tu] = [u^{-1}x^{-1}yu] = [x^{-1}][(xux^{-1})^{-1}(yvy^{-1})][y][v^{-1}u] = [x^{-1}y] = [t]$$

since $(xux^{-1}, yvy^{-1}) \in H^\circ$. Thus the classes $[t]$ and $[u]$ commute in $\mathrm{PSL}_2(\mathbb{F})$. However, since H° surjects onto $\mathrm{SL}_2(\mathcal{O})$, this forces $[t]$ to be in the centre of $\mathrm{PSL}_2(\mathbb{F})$, which is trivial (since it is a simple group). Thus $x = \pm y \pmod p$ for all $(x, y) \in H$, as claimed. \square

Assume now that we have two newforms f and g , and let L be the subfield of $\overline{\mathbb{Q}}$ generated by the coefficients of f and g . For each prime \mathfrak{p} of L , we may consider the image of the Galois representation $\rho_{f,\mathfrak{p}} \times \rho_{g,\mathfrak{p}} : \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \mathrm{GL}_2(L_{\mathfrak{p}}) \times \mathrm{GL}_2(L_{\mathfrak{p}})$.

Let H be the subgroup of $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ cut out by the Dirichlet characters corresponding to the ‘‘extra twists’’ of f and g , and let K be its fixed field (an abelian extension of \mathbb{Q}). Let F, F' be the subfields of L fixed by the extra twists. By Proposition 7.2.3, we know that for all but finitely many \mathfrak{p} , the image of $\rho_{f,\mathfrak{p}}|_H$ is the group $\{x \in \mathrm{GL}_2(\mathcal{O}) : \det(x) \in \mathbb{Z}_{\mathfrak{p}}^\times\}$, where \mathcal{O} is the completion of F at the prime below \mathfrak{p} and p is the residue characteristic of \mathfrak{p} ; similarly, the image of $\rho_{g,\mathfrak{p}}|_H$ will be the group $\{x \in \mathrm{GL}_2(\mathcal{O}') : \det(x) \in \mathbb{Z}_{\mathfrak{p}}^\times\}$ where \mathcal{O}' is the completion of F' at \mathfrak{p} . Then the image of the Galois representation $\rho_{f,\mathfrak{p}} \times \rho_{g,\mathfrak{p}}$ is a subgroup of the group $G_{\mathfrak{p}} = G$ defined above that surjects onto either factor.

PROPOSITION 7.2.9. *In the above situation, either the image of H under $\rho_{f,\mathfrak{p}} \times \rho_{g,\mathfrak{p}}$ is $G_{\mathfrak{p}}$, or $\mathcal{O}' = \mathcal{O}$ and there is an element $\gamma \in \mathrm{Gal}(L/\mathbb{Q})$ and a quadratic character $\chi : H \rightarrow \{\pm 1\}$ such that the equality*

$$(21) \quad \rho_{f,\mathfrak{p}}(\sigma) = \pm \rho_{f,\mathfrak{p}}(\sigma)^\gamma \pmod{\mathfrak{p}}$$

holds for all $\sigma \in H$.

Proof. We know from above that if the image of $\rho_{f,\mathfrak{p}} \times \rho_{g,\mathfrak{p}}$ is not G , then $\mathcal{O} = \mathcal{O}'$ and $\rho_{f,\mathfrak{p}}(\sigma) = \pm \varphi^j \rho_{g,\mathfrak{p}}(\sigma) \pmod{\mathfrak{p}}$ for all $\sigma \in H$, where φ^j is the mod \mathfrak{p} Frobenius.

Now we may take γ to be any element of the decomposition group of \mathfrak{p} in $\mathrm{Gal}(L/\mathbb{Q})$ reducing to φ^j modulo \mathfrak{p} . (Of course, there will almost always be only one such element since only finitely many primes ramify in L/\mathbb{Q} .) \square

We now lift to characteristic 0. Let w be a prime of the field K ; we define $a_w(f) = \mathrm{tr} \rho_{f,\lambda}(\sigma_w^{-1})$, where σ_w is the arithmetic Frobenius at w in H and λ is some prime of L ; if w is a degree 1 prime, then this is just $a_v(f)$ where v is the rational prime below w , and for general primes w , it may be expressed as a polynomial in $a_v(f)$ and $\chi_v(f)$. In any case it is obviously independent

of the choice of auxiliary prime λ , and (since K is abelian over \mathbb{Q}) it depends only on the prime v of \mathbb{Q} below w . We define $a_w(g)$ similarly.²

Definition 7.2.10. Let us say a prime \mathfrak{p} of residue characteristic ≥ 5 is a *good prime* for the pair (f, g) if the image of H under $\rho_{f,\mathfrak{p}} \times \rho_{g,\mathfrak{p}}$ is the whole of $G_{\mathfrak{p}}$. If the image is a proper subgroup, but has full projection to either factor, we say \mathfrak{p} is a *bad prime*.

Remark 7.2.11. If \mathfrak{p} divides 2 or 3, or is such that $\rho_{f,\mathfrak{p}}$ or $\rho_{g,\mathfrak{p}}$ has small image, we consider \mathfrak{p} to be neutral, neither good nor bad. By the theorem of Momose–Ribet (Theorem 7.2.1), there are only finitely many neutral primes.

COROLLARY 7.2.12. *If there are infinitely many bad primes for (f, g) , then there is $\gamma \in \text{Gal}(L/\mathbb{Q})$ such that the equality*

$$a_w(f) = \pm \gamma(a_w(g))$$

holds for all primes w of K .

Proof. For each bad prime \mathfrak{p} , there exists a $\gamma \in \text{Gal}(L/\mathbb{Q})$ such that the congruence (21) holds and, in particular, (by taking $\sigma = \sigma_w^{-1}$) we have $a_w(f)^2 = \gamma(a_w(g)^2) \pmod{\mathfrak{p}}$ for all primes w of K .

Since $\text{Gal}(L/\mathbb{Q})$ is finite, there exists some γ such that the congruence (21) of the proposition holds for all \mathfrak{p} in an infinite set B . In this case, we have

$$a_w(f)^2 = \gamma(a_w(g)^2) \pmod{\mathfrak{p}}$$

for infinitely many \mathfrak{p} . So we must have an equality $a_w(f)^2 = \gamma(a_w(g)^2)$, since a nonzero element of a number field cannot be divisible by infinitely many primes. \square

COROLLARY 7.2.13. *If there are infinitely many bad primes for (f, g) , there exists a quadratic Groessencharacter κ of K (equivalently, a continuous quadratic character of H) such that*

$$a_w(f) = \kappa(w)a_w(g)$$

for all primes w of K .

Proof. This follows from the strong multiplicity-one theorem for SL_2/K (cf. [Ram00]): the Satake parameters of the base-change representations $\text{BC}(\pi_f)$ and $\text{BC}(\pi_g^?)$ of $\text{GL}_2(\mathbb{A}_K)$ agree up to sign at any prime w , and Ramakrishnan’s result guarantees that the sign relating the two is given by a quadratic character. \square

²Of course, we can define the quantity $a_w(f)$ intrinsically in “automorphic” terms, as (up to normalizations) it is the trace of the d -th power of the conjugacy class in $\text{GL}_2(\mathbb{C})$ that is the Satake parameter of $\pi_{f,v}$, where d is the degree of the unramified extension $[K_w : \mathbb{Q}_v]$; this makes the independence of λ self-evident.

Remark 7.2.14. Frustratingly it does not seem to be possible to show the existence of κ without such heavy automorphic machinery, even though we know that for infinitely many primes \mathfrak{p} , the sign relating ρ_f and ρ_g modulo \mathfrak{p} is given by a character.

THEOREM 7.2.15. *Suppose there are infinitely many bad primes for (f, g) . Then f is Galois-conjugate to some twist of g .*

Proof. From Ramakrishnan's theorem, we know that we have

$$\rho_{f,\lambda}|_H \cong \rho_{g,\lambda}^\gamma|_H \otimes \tau$$

for a quadratic character τ of H and any choice of prime λ . Inducing up from H to $H' = \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, we have

$$\rho_{f,\lambda} \otimes \text{Ind}_H^{H'}(1_H) = \rho_{g,\lambda} \otimes \text{Ind}_H^{H'}(\tau).$$

But the left-hand side contains $\rho_{f,\lambda}$ as a direct summand, while the right-hand side is a direct sum of representations of the form $\rho_{g,\lambda}^\gamma \otimes \mu$ where μ is an irreducible Artin representation. Hence there must be at least one μ that is 1-dimensional and such that $\rho_{f,\lambda} \cong \rho_{g,\lambda}^\gamma \otimes \mu$, in which case we must have $f = g^\gamma \otimes \mu$. \square

7.2.3. Existence of the special element. As in the previous section, let f and g be two newforms, and let L be the subfield of $\overline{\mathbb{Q}}$ generated by the coefficients of f and g . We assume that f is not Galois conjugate to a twist of g , so by Theorem 7.2.15 there are only finitely many bad primes for (f, g) . We retain the notation of the previous section.

Let \mathfrak{p} be a good prime that does not divide the levels of f and g , and p the rational prime below \mathfrak{p} .

We make the following crucial assumption:

ASSUMPTION 7.2.16. *The character $\chi = (\chi_f \chi_g)^{-1}$ is nontrivial, and its conductor is not a power of p .*

For simplification, we also make the following assumption:

ASSUMPTION 7.2.17. *We have $L_{f,\mathfrak{p}} = L_{g,\mathfrak{p}} = \mathbb{Q}_p$, so after a suitable choice of basis, we may assume that the image of $\rho_{f,\mathfrak{p}} \times \rho_{g,\mathfrak{p}}$ is contained in $\text{GL}_2(\mathbb{Z}_p) \times \text{GL}_2(\mathbb{Z}_p)$.*

We can now prove the main result of this section.

PROPOSITION 7.2.18. *There exists an element $\tau \in G_{\mathbb{Q}(\mu_{p^\infty})}$ such that $V/(\tau - 1)V$ is 1-dimensional, where $V = V_{L_p}(f, g)^*$.*

If χ is not congruent modulo \mathfrak{p} to any character of p -power conductor, then there exists τ such that $T/(\tau - 1)T$ is free of rank 1 for any $G_{\mathbb{Q}}$ -stable lattice T in V .

Proof. Choose some $\alpha \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ such that $\chi(\alpha) \neq 1$ but α is in the kernel of the p -adic cyclotomic character. Note that such α do exist, since the conductor of χ not a power of p . Consider the coset $\alpha \cdot (H \cap G_{\mathbb{Q}(\mu_{p^\infty})})$. Since \mathfrak{p} is a good prime, under $\rho_{f,\mathfrak{p}} \times \rho_{g,\mathfrak{p}}$, the coset $\alpha \cdot (H \cap G_{\mathbb{Q}(\mu_{p^\infty})})$ is mapped to

$$(\rho_{f,\mathfrak{p}}(\alpha), \rho_{g,\mathfrak{p}}(\alpha)) \cdot \text{SL}_2(\mathbb{Z}_p)^2,$$

which consists of *all* pairs (u, v) of matrices such that $\det(u) = \chi_f(\alpha)$ and $\det(v) = \chi_g(\alpha)$. In particular, it contains the pair

$$\begin{pmatrix} x & 0 \\ 0 & x^{-1}\chi_f(\alpha) \end{pmatrix} \begin{pmatrix} x^{-1} & 0 \\ 0 & x\chi_g(\alpha) \end{pmatrix}$$

for any $x \in \mathbb{Z}_p^\times$. The image of this pair under the tensor product homomorphism $\text{GL}_2 \times \text{GL}_2 \rightarrow \text{GL}_4$ is the diagonal matrix with entries

$$[1, x^{-2}\chi_f(\alpha), x^2\chi_g(\alpha), \chi_f(\alpha)\chi_g(\alpha)].$$

By choosing x appropriately, we can arrange that neither $x^{-2}\chi_f(\alpha)$ nor $x^2\chi_g(\alpha)$ is equal to 1. Thus 1 is an eigenvalue of τ on $V_{L_p}(f) \otimes V_{L_p}(g)$ with multiplicity exactly 1.

If we assume the stronger condition on χ in the statement, then we can assume that $\chi_f(\alpha)\chi_g(\alpha)$ is not 1 modulo \mathfrak{p} . By choosing x appropriately we can assume that $x^{-2}\chi_f(\alpha)$ and $x^2\chi_g(\alpha)$ are also noncongruent to 1, so it follows that $T/(\tau - 1)T$ is free of rank 1 as required (for any τ -stable \mathcal{O}_L -lattice in V and, in particular, any $G_{\mathbb{Q}}$ -stable lattice). \square

7.2.4. *The quantities \mathfrak{n}_W and \mathfrak{n}_W^* .* We recall the definitions of the quantities \mathfrak{n}_W and \mathfrak{n}_W^* in Rubin’s theory. Let T be a finite-rank free \mathcal{O} -module with a continuous action of $G_{\mathbb{Q}}$. As usual, write $V = T \otimes E$ and $W = V/T$.

Definition 7.2.19. Define Ω to be the smallest extension of \mathbb{Q} whose Galois group acts trivially on W and on μ_{p^∞} , and define

$$\begin{aligned} \mathfrak{n}_W &= \ell_{\mathcal{O}}(H^1(\Omega/\mathbb{Q}, W) \cap \mathcal{S}^{\{p\}}(K, W)), \\ \mathfrak{n}_W^* &= \ell_{\mathcal{O}}(H^1(\Omega/\mathbb{Q}, W^*(1)) \cap \mathcal{S}_{\{p\}}(K, W^*(1))). \end{aligned}$$

We now give conditions under which these quantities are zero.

PROPOSITION 7.2.20. *Suppose the centre of Ω acts on each of $T \otimes \mathbf{k}$ and $T^*(1) \otimes \mathbf{k}$ via a nontrivial character. Then $\mathfrak{n}_W = \mathfrak{n}_W^* = 0$.*

Proof. We shall show that the hypotheses imply that $H^1(\Omega/\mathbb{Q}, W) = H^1(\Omega/\mathbb{Q}, W^*(1)) = 0$. We give the argument for W ; the proof for $W^*(1)$ is similar.

Clearly we have $H^1(\Omega/\mathbb{Q}, V) = 0$, and hence $H^1(\Omega/\mathbb{Q}, W)$ is finite. So it suffices to show that $H^1(\Omega/\mathbb{Q}, W)[\varpi] = 0$, where ϖ is a uniformizer. But

we have a surjection $H^1(\Omega/\mathbb{Q}, W[\varpi]) \twoheadrightarrow H^1(\Omega/\mathbb{Q}, W)[\varpi]$, so we are reduced to showing that $H^1(\Omega/\mathbb{Q}, W[\varpi]) = H^1(\Omega/\mathbb{Q}, T \otimes \mathbf{k})$ is zero. However, this is immediate since any representation of nontrivial central character cannot have a nontrivial extension by the trivial representation. \square

7.3. *The Euler system.* As above, let f and g be newforms of weight 2, level N and characters χ_f and χ_g , respectively. Let p be a prime not dividing N . Let L be a number field containing the coefficients of f_α and g_β , and let \mathfrak{p} be a prime of L above p . Let $E = L_{\mathfrak{p}}$, and let \mathcal{O} be its ring of integers. We write $T = T_{\mathcal{O}}(f, g)^*$ and $p_\ell(X) = \det(1 - \text{Frob}_q^{-1} X | T^*(1)) = P_\ell(f, g, \ell^{-1} X) \in \mathcal{O}_L[X]$. We assume that the following conditions are satisfied:

ASSUMPTION 7.3.1.

- (i) *the character $\chi = \chi_f \chi_g$ is not trivial and, moreover, is not trivial modulo \mathfrak{p} ;*
- (ii) *there exist p -stabilizations f_α, g_γ of f and g with U_p -eigenvalues α, γ respectively such that*
 - $v_{\mathfrak{p}}(\alpha\gamma) < 1$,
 - α/γ *is not a root of unity.*

Fix $c \geq 1$ coprime to $6N$, and let A be the set of square-free integers prime to Npc . By Corollary 6.4.5, we have, for every integer $m \in A$, a cohomology class ${}_c \mathbf{z}_m^{f,g,N} \in H^1(\mathbb{Q}_m, T)$ that satisfies the following compatibility property: if $m \in A$ and ℓ is a prime coprime to $mNpc$, then the image of $\mathbf{z}_{\ell m}^{f,g,N}$ under the corestriction map $H^1(\mathbb{Q}_{\ell m}, T) \rightarrow H^1(\mathbb{Q}_m, T)$ is

$$-\sigma_\ell A_\ell(\sigma_\ell^{-1}) \mathbf{z}_m^{f,g,N},$$

where $A_\ell(X)$ is a polynomial in $\mathcal{O}_L[X]$ congruent modulo $\ell - 1$ to $p_\ell(X)$.

LEMMA 7.3.2. *There exists a system of cohomology classes*

$$\{ {}_c \tilde{\mathbf{z}}_m^{f,g,N} \in H^1(\mathbb{Q}_m, T) : m \in A \}$$

such that

$${}_c \tilde{\mathbf{z}}_1^{f,g,N} = {}_c \mathbf{z}_1^{f,g,N}$$

and if $m \in A$ and ℓ is a prime such that $m\ell \in A$, then the image of ${}_c \tilde{\mathbf{z}}_{\ell m}^{f,g,N}$ under the corestriction map $H^1(\mathbb{Q}_{\ell m}, T) \rightarrow H^1(\mathbb{Q}_m, T)$ is

$$A_\ell(\sigma_\ell^{-1}) {}_c \tilde{\mathbf{z}}_m^{f,g,N}.$$

Proof. By induction on the number of prime factors of m , we can choose (noncanonically) a system of elements $\gamma_m \in (\mathbb{Z}/m\mathbb{Z})^\times$ for every $m \in A$ such that $\gamma_{m\ell} = \ell^{-1} \gamma_m \pmod m$. Identify γ_m with an element of $\text{Gal}(\mathbb{Q}(\mu_m)/\mathbb{Q})$ via the inverse of the cyclotomic character, and define ${}_c \tilde{\mathbf{z}}_m^{f,g,N} = (-1)^{s(m)} \gamma_m \cdot {}_c \mathbf{z}_m^{f,g,N}$, where $s(m)$ is the number of prime factors of m . It is clear by construction that the elements have the required property. \square

Note 7.3.3. By Corollary 6.7.9, the classes $c_{\mathbf{z}_m}^{f,g,N}$ are in the Selmer group $\mathcal{S}^{\{p\}}(\mathbb{Q}(\mu_m), T_{\mathcal{O}_p}(f, g)^*)$. As $\mathcal{S}^{\{p\}}(\mathbb{Q}(\mu_m), T_{\mathcal{O}_p}(f, g)^*)$ is invariant under the action of $\text{Gal}(\mathbb{Q}(\mu_m)/\mathbb{Q})$, it follows that the same is true for the modified classes $c_{\tilde{\mathbf{z}}_m}^{f,g,N}$.

We now show that we can convert the classes $(c_{\mathbf{z}_m}^{f,g,N})_{m \in A}$ into a Euler system. Let T, Σ and A be as defined at the beginning of Section 7.1. Then we have the following result (cf. [Rub00, Lemma 9.6.1]):

LEMMA 7.3.4. *Suppose that for all primes $\ell \in A$ we have polynomials $r_\ell(X), s_\ell(X) \in \mathcal{O}[X]$ such that*

$$r_\ell(X) \equiv s_\ell(X) \pmod{\ell - 1},$$

and suppose we have a collection of cohomology classes $\{\tilde{c}_m \in H^1(\mathbb{Q}(\mu_m), T) : m \in A\}$ such that if $\ell \in A$ is coprime to m , then

$$\text{cores}_{\mathbb{Q}(\mu_{\ell m})/\mathbb{Q}(\mu_m)}(\tilde{c}_{\ell m}) = \begin{cases} r_\ell(\sigma_\ell^{-1})\tilde{c}_m & \text{if } \ell \nmid m, \\ \tilde{c}_m & \text{if } \ell \mid m. \end{cases}$$

Then there exists a collection of classes $\{c_m \in H^1(\mathbb{Q}(\mu_m), T) : m \in A\}$ with the following properties:

(i) *for all m ,*

$$c_m \in \mathcal{O}[(\mathbb{Z}/m\mathbb{Z})^\times] \cdot \tilde{c}_m;$$

(ii) *if ℓ is a prime such that $m, m\ell \in A$, then*

$$\text{cores}_{\mathbb{Q}(\mu_{\ell m})/\mathbb{Q}(\mu_m)}(c_{\ell m}) = \begin{cases} s_\ell(\sigma_\ell^{-1})c_m & \text{if } \ell \nmid m, \\ c_m & \text{if } \ell \mid m; \end{cases}$$

(iii) *if $m \in A$ and χ is a character of $\text{Gal}(\mathbb{Q}(\mu_m)/\mathbb{Q})$ of conductor k such that $\text{prime}(m) \subset \text{prime}(k) \cup \Sigma$, then*

$$\sum_{\gamma \in \text{Gal}(\mathbb{Q}(\mu_m)/\mathbb{Q})} \chi(\gamma)\gamma(c_m) = \sum_{\gamma \in \text{Gal}(\mathbb{Q}(\mu_m)/\mathbb{Q})} \chi(\gamma)\gamma(\tilde{c}_m).$$

Definition 7.3.5. Define $\{c_{\hat{\mathbf{z}}_m}^{f,g,N} \in H^1(\mathbb{Q}(\mu_m), T_{\mathcal{O}_p}(f, g)^*) : m \in A\}$ to be the classes obtained by applying Lemma 7.3.4 to our classes $c_{\tilde{\mathbf{z}}_m}^{f,g,N}$, where we take $r_\ell(X) = A_\ell(X)$ and $s_\ell(X) = p_\ell(X) = P_\ell(f, g, \ell^{-1}X)$ and, as above, A is the set of square-free integers coprime to Npc .

Note 7.3.6. By construction, the classes $\{c_{\hat{\mathbf{z}}_m}^{f,g,N} : m \in A\}$ are a Euler system for (T, Σ, A) in the sense of Definition 7.1.1, where Σ is the set of primes dividing Npc . Moreover, because of (i), we have $c_{\hat{\mathbf{z}}_m} \in \mathcal{S}^{\{p\}}(\mathbb{Q}(\mu_m), T)$ for all m .

7.4. Finiteness of the strict Selmer group. We now combine the above results to prove a finiteness theorem for the strict Selmer group. For the convenience of the reader, we shall recapitulate all of the assumptions we have made on f and g .

ASSUMPTION 7.4.1. *Assume that f and g are weight 2 newforms with coefficients in a number field L and that \mathfrak{p} is a prime of L above the rational prime p , with the following properties:*

- (i) *Neither f nor g is of CM type;*
- (ii) *f is not a twist of g ;*
- (iii) *The character $\varepsilon_f \varepsilon_g$ is nontrivial;*
- (iv) *$p \geq 5$;*
- (v) *p does not divide the levels of f and g ;*
- (vi) *\mathfrak{p} is totally split in the field L , so $L_{\mathfrak{p}} = \mathbb{Q}_p$;*
- (vii) *the \mathfrak{p} -adic Galois representations of f and g are surjective onto $\mathrm{GL}_2(\mathbb{Z}_{\mathfrak{p}})$;*
- (viii) *there exists some prime v such that $\chi(v) = 1$ for all inner twists χ of f or g , and $a_v(f) \not\equiv \pm a_v(g) \pmod{\mathfrak{p}}$;*
- (ix) *f is ordinary at \mathfrak{p} ;*
- (x) *there exists a root γ of the Hecke polynomial of g at p such that $v_{\mathfrak{p}}(\gamma) < 1$ and α/γ is not a root of unity, where α is the unit root of the Hecke polynomial of f .*

If we assume hypotheses (i)–(iii) (which do not depend on \mathfrak{p}), then there will be many \mathfrak{p} such that the remaining hypotheses hold.

THEOREM 7.4.2. *Suppose Assumption 7.4.1 is satisfied, and the p -adic Rankin–Selberg L -function $\mathcal{D}_{\mathfrak{p}}(f, g, 1/N)$ does not vanish at 1, where N is some integer divisible by the levels of f and g . Then*

$$\#\mathcal{S}_{\{\mathfrak{p}\}} \left(\mathbb{Q}, \frac{V_{L_{\mathfrak{p}}}(f, g)}{T_{\mathcal{O}_{\mathfrak{p}}}(f, g)}(1) \right) < \infty.$$

Proof. It suffices to show that the hypotheses of Theorem 7.1.4 are satisfied for $T = T_{\mathcal{O}_{\mathfrak{p}}}(f, g)^*$. By Proposition 7.2.18, the element τ required by Hypothesis $\mathrm{Hyp}(\mathbb{Q}, V)$ exists, and the Euler system of Definition 7.3.5 satisfies Hypothesis $\mathrm{Hyp}(\mathcal{S}^{\{p\}}, V)$ (ii). Since T is nontrivial and irreducible, $T^{G_K} = 0$, and the element γ in Hypothesis $\mathrm{Hyp}(\mathcal{S}^{\{p\}}, V)$ (iii) clearly exists.

By Theorem 5.6.4, if $\mathcal{D}_{\mathfrak{p}}(f, g, 1/N)(1) \neq 0$, the image of $\mathrm{reg}_p \Xi_{1, N, 1}$ in the (f, g) -isotypical quotient of $H_{\mathrm{dR}}^2(X_1(N)/\mathbb{Q}_p)/\mathrm{Fil}^2$ is nonzero. Hence, by the diagram of Section 5.5, the localization of the Galois cohomology class $\mathbf{z}_1^{f, g, N}$ at p is nonzero so, in particular, $c\mathbf{z}_1^{f, g, N}$ is nontorsion as an element of $H^1(\mathbb{Q}, T_{L_{\mathfrak{p}}}(f, g)^*)$ for any $c > 1$. Thus we may apply Theorem 7.1.4 to the Euler system $(c\hat{\mathbf{z}}_m^{f, g, N})_{m \in A}$ of Definition 7.3.5 to obtain the finiteness of the strict Selmer group. \square

7.5. *The order of the strict Selmer group.* Theorem 7.1.5 gives a bound for the order of the strict Selmer group, under slightly stronger hypotheses than Theorem 7.4.2.

THEOREM 7.5.1. *Suppose Assumption 7.4.1 is satisfied and, in addition, the mod \mathfrak{p} reduction of $\varepsilon_f \varepsilon_g$ is not trivial. Then we have*

$$\begin{aligned} \text{length}_{\mathbb{Z}_p} \mathcal{S}_{\{p\}} \left(\mathbb{Q}, \frac{V_{L_p}(f, g)}{T_{\mathcal{O}_p}(f, g)}(1) \right) \\ \leq v_p \left(\frac{(1 - p^{-1}\beta\alpha^{-1})}{(1 - p^{-1}\beta\gamma)(1 - p^{-1}\beta\delta)} \mathcal{D}_p(f, g, 1/N)(1) \right) + \lambda, \end{aligned}$$

where λ is the \mathfrak{p} -adic valuation of the ideal I_f of Definition 6.10.4 above.

Proof. Our condition on the mod \mathfrak{p} reduction of $\varepsilon_f \varepsilon_g$ implies that Hypothesis $\text{Hyp}(\mathbb{Q}, T)$ is satisfied (again by Proposition 7.2.18; note that the mod \mathfrak{p} reduction cannot be a nontrivial character of p -power conductor as p does not divide the levels of f and g). The condition also assures that the quantities \mathfrak{n}_W and \mathfrak{n}_W^* appearing in Theorem 7.1.5 are zero (Proposition 7.2.20).

We consider the linear functional α on $H^1(\mathbb{Q}, V_{L_p}(f, g)^*)$ given by $x \mapsto (\log_{\mathbb{Q}_p}(x), \eta_f^{\text{ur}} \otimes \omega_g)$. On the lattice $\widetilde{T}_{\mathcal{O}_p}(f, g)^*$ this takes values in

$$I_f^{-1} \cdot (1 - \alpha^{-1}\gamma^{-1})^{-1}(1 - \alpha^{-1}\delta^{-1})^{-1} \mathcal{O}_p,$$

by Corollary 6.10.7; but since the Galois representations of f and g are assumed to have big image, we have $\widetilde{T}_{\mathcal{O}_p}(f, g)^* = T_{\mathcal{O}_p}(f, g)^*$.

Theorem 5.6.4 shows that τ maps the class $\mathbf{z}_{f, g, 1}$ to

$$\frac{\mathcal{E}(f)\mathcal{E}^*(f)}{\mathcal{E}(f, g, 1)} \mathcal{D}_p(f, g, 1/N)(1).$$

Hence the index of divisibility of $\mathbf{z}_1^{f, g, N}$ is bounded above by

$$v_p \left(\frac{\mathcal{E}(f)\mathcal{E}^*(f)(1 - \alpha^{-1}\gamma^{-1})(1 - \alpha^{-1}\delta^{-1})}{\mathcal{E}(f, g, 1)} \mathcal{D}_p(f, g, 1/N)(1) \right) + \lambda.$$

We can ignore the factor $\mathcal{E}^*(f) := 1 - \beta\alpha^{-1}$, since α_f is a unit and β_f is a nonunit so $\mathcal{E}^*(f) \in \mathcal{O}_p^\times$. Substituting the definitions of $\mathcal{E}(f)$ and $\mathcal{E}(f, g, 1)$, we have

$$\frac{\mathcal{E}(f)(1 - \alpha^{-1}\gamma^{-1})(1 - \alpha^{-1}\delta^{-1})}{\mathcal{E}(f, g, 1)} = \frac{(1 - p^{-1}\beta\alpha^{-1})}{(1 - p^{-1}\beta\gamma)(1 - p^{-1}\beta\delta)}. \quad \square$$

7.6. An example. It may seem slightly unclear whether the long list of conditions in Assumption 7.4.1 may be simultaneously satisfied, so we present the following explicit example (computed using Sage [Sage]):

Let f be the unique weight 2 newform of level 11 (corresponding to the elliptic curve $E : y^2 + y = x^3 - x$). Let g be the unique newform of weight 2, level 26, and character $\chi := \left(\frac{\bullet}{13}\right)$ with $a_2(g) = i$, so the q -expansions of f and g are

$$\begin{aligned} f &= q - 2q^2 - q^3 + 2q^4 + q^5 + O(q^6), \\ g &= q + iq^2 - q^3 - q^4 - 3iq^5 + O(q^6). \end{aligned}$$

Note that χ has conductor 13, so the local component of π_g at 2 is an unramified twist of the Steinberg representation; on the other hand f is unramified principal series at 2 and Steinberg at 11. So f cannot be a twist of g , and neither f nor g is of CM type (since CM forms cannot be Steinberg at any prime). The form f has no inner twists (since it is non-CM and has coefficients in \mathbb{Q}); as for g , its Galois orbit consists of g and \bar{g} , so its only inner twist is \bar{g} .

To calculate the image of the Galois representations of f and g , we note that Sage [Sage] has a facility to compute all the exceptional primes for the Galois representation attached to an elliptic curve (i.e., those primes for which the image of the Galois representation is not $\mathrm{GL}_2(\mathbb{Z}_p)$). This speedily tells us that $\rho_{f,p}$ is surjective for all $p \neq 5$.

The form g does not correspond to an elliptic curve, but there is a Dirichlet character $\psi : (\mathbb{Z}/13\mathbb{Z})^\times \rightarrow \mathbb{Q}(i)^\times$ such that $g \otimes \psi$ corresponds to an elliptic curve E' of conductor $2 \times 13^2 = 338$ (the curve with Cremona label 338d, given by $y^2 + xy = x^3 + x^2 + 504x - 13112$), and the only exceptional primes for E' are $\{3, 5\}$. Letting $H = G_{\mathbb{Q}(\sqrt{13})}$, the kernel of the character χ , we see that for all primes $p \notin \{2, 3, 5, 13\}$, the image of H under $\rho_{g,\mathfrak{p}}$, for any prime \mathfrak{p} of $\mathbb{Q}(i)$ above p , is $\mathrm{GL}_2(\mathbb{Z}_p)$.

Moreover, the only prime such that $a_v(f) = a_v(g)$ for all v split in $\mathbb{Q}(\sqrt{13})$ is $p = 5$. We deduce that for any p congruent to 1 mod 4 and not in $\{5, 13\}$, and any prime \mathfrak{p} of $\mathbb{Q}(i)$ above p , the hypotheses (i) – (viii) are satisfied.

We check that both f and g are ordinary at the primes above 17 (it does not matter which prime we take, since $a_{17}(g) \in \mathbb{Z}$). For any choice of roots α, γ of roots of the Hecke polynomials of f and g , the minimal polynomial of α/γ over \mathbb{Q} is $x^4 + \frac{6}{17}x^3 - \frac{21}{17}x^2 + \frac{6}{17}x + 1$ so, in particular, α/γ is not a root of unity. Thus hypotheses (ix) and (x) are satisfied if \mathfrak{p} is either of the primes above 17.

Note added after submission. Since this paper was originally submitted, the 17-adic L -values appearing in this example have been computed modulo 17^3 by Alan Lauder, using the techniques described in [Lau14]. It is actually more convenient to compute $\mathcal{D}_{\mathfrak{p}}(g, f, 1/N)(1)$ rather than $\mathcal{D}_{\mathfrak{p}}(f, g, 1/N)(1)$. This quantity is indeed nonzero, so applying Theorem 7.4.2 gives finiteness of the Selmer group.

8. Conjectures on higher-rank Euler systems

We now explain how the cohomology classes constructed in the previous section may be reconciled with the general conjectural setup of cyclotomic Iwasawa theory for motivic Galois representations formulated by Perrin-Riou, and its extension to the two-variable situation as formulated by the second and third authors in [LZ12].

8.1. *Euler systems: rank 1 and higher rank.* Let us place ourselves again in the general setting of Section 7.1 above, so T is a free \mathcal{O} -module with a continuous action of $G_{\mathbb{Q}}$ unramified outside a finite set $\Sigma \ni p$, and A is a set of integers satisfying the conditions loc. cit. Suppose that all integers in A are coprime to p .

Perrin-Riou’s conjectures, as formulated in [PR98] (cf. also [Rub00, §8.5]), discuss the following class of objects:

Definition 8.1.1. A *Euler–Iwasawa system* of rank $r \geq 1$ consists of the data of, for each $m \in A$, a class

$$c_m \in \bigwedge_{\Lambda(\Gamma_m)}^r H_{\text{Iw}}^1(\mathbb{Q}(\mu_{mp^\infty}), V)$$

with the property that if ℓ is prime and $\ell, m\ell \in A$, we have

$$\text{cores}_{\mathbb{Q}(\mu_{mp^\infty})}^{\mathbb{Q}(\mu_{m\ell p^\infty})} c_{m\ell} = \begin{cases} p_\ell(\sigma_\ell^{-1})c_m & \text{if } \ell \nmid m\Sigma, \\ c_m & \text{if } \ell \mid m\Sigma. \end{cases}$$

Note that a rank 1 Euler–Iwasawa system is equivalent to the data of a Euler system for (T, A_p, Σ) in the previous sense, where $A_p = \{p^k m : m \in A, k \geq 0\}$.

As noted in [PR98, §1.2.3], a higher-rank Euler system can be used to construct rank 1 Euler systems, by pairing with appropriate “rank $r - 1$ ” elements. We make the following definition:

Definition 8.1.2. We define a *Perrin-Riou functional* to be the data of, for each squarefree m prime to S as above, an element

$$\Phi_m \in \bigwedge^{r-1} \text{Hom}_\Lambda(H_{\text{Iw},S}^1(\mathbb{Q}(\mu_{mp^\infty}), T), \Lambda)^\iota,$$

with the property that for each $\ell \nmid mS$, we have

$$\Phi_m = \Phi_{m\ell} \circ \text{res}_{\mathbb{Q}(\mu_{mp^\infty})}^{\mathbb{Q}(\mu_{m\ell p^\infty})}.$$

Lemma 1.2.3 of op. cit. shows that if $(c_p(m))$ is a Euler system of rank r , and (Φ_m) is a Perrin-Riou functional, then the elements

$$\Phi_m(c_p(m)) \in H_{\text{Iw},S}^1(\mathbb{Q}(\mu_{mp^\infty}), T)$$

define a Euler system of rank 1. (Here, as explained loc.cit., we interpret Φ_m as a map

$$\bigwedge^r H_{\text{Iw},S}^1(\mathbb{Q}(\mu_{mp^\infty}), T) \rightarrow H_{\text{Iw},S}^1(\mathbb{Q}(\mu_{mp^\infty}), T),$$

which we also denote by Φ_m .)

Appendix B of op. cit. shows that (under mild hypotheses on T) there is a plentiful supply of Perrin-Riou functionals, although there is no obvious

canonical choice. More specifically, given any m and any Φ_m , there exists a Perrin-Riou functional extending Φ_m . Hence, given as a starting point a rank r Euler system, one may construct a rank 1 Euler system (indeed many such systems) and obtain Iwasawa-theoretic results from this rank 1 system; but these rank 1 Euler systems are noncanonical and, in particular, there is no reason to expect that they should have any relation to L -values.

Remark 8.1.3. An alternative approach to bounding Selmer groups in the $r > 1$ case by directly utilizing a notion of “higher-rank Kolyvagin systems,” rather than by constructing rank 1 Euler systems, has been initiated by Mazur and Rubin (unpublished).

8.2. *Otsuki’s functionals.* We now explain a construction due to Otsuki [Ots09], who has shown how to construct *canonical* linear functionals on cohomology groups by composing the dual exponential map with an appropriate “weighted trace.” These maps do not satisfy the compatibility properties of a Perrin-Riou functional, and thus they give rise to systems of elements of group rings satisfying a modified compatibility property; we shall show that this modification is consistent with the results we have shown for our generalized Beilinson–Flach classes.

For technical reasons we shall work in the limit over the cyclotomic extension, rather than directly over $\mathbb{Q}(\mu_m)$; this avoids problems caused by zeroes of local Euler factors (cf. the discussion at the start of Section 9.1 of [Rub00]).

Choose a system of roots of unity $\zeta_m \in \overline{\mathbb{Q}}$ for all $m \geq 1$ that satisfy $\zeta_{mn}^n = \zeta_m$ for all integers m, n . Let $G_m = \text{Gal}(\mathbb{Q}(\mu_m)/\mathbb{Q})$ and $\Gamma_m = \text{Gal}(\mathbb{Q}(\mu_{mp^\infty})/\mathbb{Q})$; we identify Γ_m with $G_m \times \Gamma$ in the obvious way.

Let V be an E -linear p -adic representation of $G_{\mathbb{Q}}$, where E/\mathbb{Q}_p is a finite extension, that is crystalline at p with nonnegative Hodge–Tate weights and such that no eigenvalue of Frobenius on $\mathbb{D}_{\text{cris}}(V)$ is a root of unity. Then for all $m \geq 1$, the p -adic regulator map

$$\mathcal{L}_{\mathbb{Q}(\mu_m), V}^\Gamma : H_{\text{Iw}}^1(\mathbb{Q}(\mu_{mp^\infty}), V) \longrightarrow \mathbb{Q}(\mu_m) \otimes_{\mathbb{Q}} \mathcal{H}_E(\Gamma) \otimes_E \mathbb{D}_{\text{cris}}(V)$$

is well defined (as the sum of the local regulator maps at the primes of $\mathbb{Q}(\mu_m)$ above p).

Let $D = \mathbb{D}_{\text{cris}}(V^*) = \mathbb{D}_{\text{cris}}(V)^*$, and let $D_{mp^\infty} = \Lambda_E(\Gamma) \otimes_E D \otimes_{\mathbb{Q}} \mathbb{Q}(\mu_m)$. We regard D_{mp^∞} as a Γ_m -module, via the usual action of G_m on $\mathbb{Q}(\mu_m)$ and of Γ on $\Lambda_E(\Gamma)$. Following [Kur02] and [Ots09], we have the following definition:

Definition 8.2.1. Define a pairing

$$t_m : D_{mp^\infty} \times H_{\text{Iw}}^1(\mathbb{Q}(\mu_{mp^\infty}), V) \longrightarrow \mathcal{H}_E[\Gamma_m]$$

by

$$t_m(x, z) = \sum_{\sigma \in G_m} [\sigma] \text{trace}_{\mathbb{Q}(\mu_m)/\mathbb{Q}} \left\langle \sigma x, \mathcal{L}_{\mathbb{Q}(\mu_m), V}^\Gamma(z) \right\rangle_{\text{cris}}.$$

Here we extend $\langle, \rangle_{\text{cris}}$ to be Γ -linear in the second variable and Γ -antilinear in the first. One checks that

$$t_m(\sigma x, \tau z) = [\sigma^{-1}\tau] \cdot t_m(x, z)$$

for all $\sigma, \tau \in \Gamma_m$ (not just in Γ).

Now fix two families F_ℓ, G_ℓ of polynomials in $E[X]$, indexed by primes $\ell \notin \Sigma$, such that $F_\ell, G_\ell \in 1 + XE[X]$ for all ℓ .

Let A be the set of square-free integers prime to Σ . For each prime $\ell \notin \Sigma$ and each $m \in A$, consider the $\Lambda(\Gamma_m)$ -linear endomorphism of $\Lambda_E(\Gamma) \otimes_{\mathbb{Q}} \mathbb{Q}(\mu_m)$ given by $\hat{\sigma}_\ell(x \otimes \zeta) = \tau_\ell x \otimes \zeta^\ell$, for all roots of unity $\zeta \in \mu_m$, where τ_ℓ is the arithmetic Frobenius at ℓ in Γ . Thus $\hat{\sigma}_\ell$ is the action of the Frobenius at ℓ in Γ_m if $\ell \nmid m$ and is a possibly noninvertible endomorphism if $\ell \mid m$; and the $\hat{\sigma}_\ell$ all commute with each other.

PROPOSITION 8.2.2. *The endomorphism $F_\ell(\hat{\sigma}_\ell)$ is invertible in*

$$\text{End}_Q(Q \otimes_{\mathbb{Q}} \mathbb{Q}(\mu_m)),$$

where $Q = \text{Frac } \Lambda_E(\Gamma)$.

Proof. Clear, since the roots of the characteristic polynomial of $\hat{\sigma}_\ell$ on $\mathbb{Q}(\mu_m)$ are scalars. □

For each $m \in A$, let us define an element $x'_m \in Q \otimes_{\mathbb{Q}} \mathbb{Q}(\mu_m)$ by

$$x'_m = \left(\prod_{\ell \mid m} F_\ell(\hat{\sigma}_\ell)^{-1} G_\ell(\hat{\sigma}_\ell) \right) \cdot (1 \otimes \zeta_m).$$

PROPOSITION 8.2.3. *If $\ell \nmid m$, then we have*

$$\text{tr}_m^{m\ell}(x'_{m\ell}) = \sigma_\ell^{-1} F_\ell(\sigma_\ell)^{-1} ((\ell - 1)G_\ell(\sigma_\ell) - \ell F_\ell(\sigma_\ell)) x'_m.$$

Proof. This is a straightforward generalization of (one case of) Proposition 2.5 of [Ots09]. We define $H_\ell(X) = \frac{G_\ell(X) - F_\ell(X)}{X}$, so we have

$$F_\ell(\hat{\sigma}_\ell)^{-1} G_\ell(\hat{\sigma}_\ell) = 1 + H_\ell(\hat{\sigma}_\ell) F_\ell(\hat{\sigma}_\ell)^{-1} \hat{\sigma}_\ell$$

in $\text{End}_Q Q \otimes_{\mathbb{Q}} \mathbb{Q}(\mu_{m\ell})$. The operator $\hat{\sigma}_v$ commutes with $\text{tr}_m^{m\ell}$ whenever $v \neq \ell$. Hence

$$\begin{aligned} \text{tr}_m^{m\ell}(x'_{m\ell}) &= \text{tr}_m^{m\ell} \left(\left(\prod_{v \mid m\ell} F_v(\hat{\sigma}_v)^{-1} G_v(\hat{\sigma}_v) \right) \zeta_{m\ell} \right) \\ &= \left(\prod_{v \mid m} F_v(\hat{\sigma}_v)^{-1} G_v(\hat{\sigma}_v) \right) \text{tr}_m^{m\ell} (F_\ell(\hat{\sigma}_\ell)^{-1} G_\ell(\sigma_\ell) \zeta_{m\ell}) \end{aligned}$$

$$\begin{aligned}
 &= \left(\prod_{v|m} F_v(\hat{\sigma}_v)^{-1} G_v(\hat{\sigma}_v) \right) \mathrm{tr}_m^{m\ell} \left((1 + H_\ell(\hat{\sigma}_\ell) F_\ell(\hat{\sigma}_\ell)^{-1} \hat{\sigma}_\ell) \zeta_{m\ell} \right) \\
 &= \left(\prod_{v|m} F_v(\hat{\sigma}_v)^{-1} G_v(\hat{\sigma}_v) \right) \left[\mathrm{tr}_m^{m\ell}(\zeta_{m\ell}) + \mathrm{tr}_m^{m\ell} (H_\ell(\hat{\sigma}_\ell) F_\ell(\hat{\sigma}_\ell)^{-1} \zeta_m) \right] \\
 &= \left(\prod_{v|m} F_v(\hat{\sigma}_v)^{-1} G_v(\hat{\sigma}_v) \right) \left[(-\sigma_\ell^{-1} \zeta_m) + (\ell - 1) (H_\ell(\hat{\sigma}_\ell) F_\ell(\hat{\sigma}_\ell)^{-1} \zeta_m) \right] \\
 &= (-\sigma_\ell^{-1} + (\ell - 1) H_\ell(\sigma_\ell) F_\ell(\sigma_\ell)^{-1}) x'_m
 \end{aligned}$$

(where we have dropped the hats, since $\hat{\sigma}_\ell$ acts on $Q \otimes \mathbb{Q}(\mu_m)$ as the usual Frobenius σ_ℓ). Since $\sigma_\ell H_\ell(\sigma_\ell) = G_\ell(\sigma_\ell) - F_\ell(\sigma_\ell)$, we have

$$\begin{aligned}
 &-\sigma_\ell^{-1} + (\ell - 1) H_\ell(\sigma_\ell) F_\ell(\sigma_\ell)^{-1} \\
 &= \sigma_\ell^{-1} F_\ell(\sigma_\ell)^{-1} (-F_\ell(\sigma_\ell) + (\ell - 1) \sigma_\ell H_\ell(\sigma_\ell)) \\
 &= \sigma_\ell^{-1} F_\ell(\sigma_\ell)^{-1} (-F_\ell(\sigma_\ell) + (\ell - 1) (G_\ell(\sigma_\ell) - F_\ell(\sigma_\ell))) \\
 &= \sigma_\ell^{-1} F_\ell(\sigma_\ell)^{-1} ((\ell - 1) G_\ell(\sigma_\ell) - \ell F_\ell(\sigma_\ell)),
 \end{aligned}$$

which gives the formula stated above. □

COROLLARY 8.2.4. *If we are given, for each $m \in A$, an element*

$$z_m \in Q \otimes_{\Lambda(\Gamma)} H_{\mathrm{Iw}}^1(\mathbb{Q}(\mu_{mp^\infty}), V)$$

satisfying

$$\mathrm{cores}_m^{m\ell}(z_{m\ell}) = F_\ell(\sigma_\ell^{-1}) z_m$$

for each m and each prime $\ell \nmid m$, $\ell \notin \Sigma$, and if we define $x_m = x'_m v \in D_{mp^\infty}$ for some fixed $v \in D$, then we have the relation

$$\mathrm{pr}_m^{m\ell} t_{m\ell}(x_{m\ell}, z_{m\ell}) = \sigma_\ell \left((\ell - 1) G_\ell(\sigma_\ell^{-1}) - \ell F_\ell(\sigma_\ell^{-1}) \right) t_m(x_m, z_m).$$

By base extension we may regard $t_m(x_m, -)$ as a map $\bigwedge_{\Lambda(\Gamma_m)}^2 M_m \rightarrow M_m$, where

$$M_m = Q \otimes_{\Lambda(\Gamma)} H_{\mathrm{Iw}}^1(\mathbb{Q}(\mu_{mp^\infty}), V),$$

so it makes sense to evaluate $t_m(x_m, -)$ against a rank 2 Euler–Iwasawa system.

We now specialize to the case where $V = V_{L_\lambda}(f, g)^*$ for some weight 2 eigenforms (f, g) of levels divisible only by primes in $\Sigma - \{p\}$. We take $G_\ell(X) = 1 - \varepsilon_\ell(f)\varepsilon_\ell(g)X^2$ and $F_\ell(X) = P_\ell(\ell^{-1}X)$ as before. Choose a p -stabilization (α, γ) of f and g , and let $v = v_\alpha \otimes v_\gamma$ be the obvious φ -eigenvector in $\mathbb{D}_{\mathrm{cris}}(V^*)$ of eigenvalue $\alpha\gamma$.

PROPOSITION 8.2.5. *Let $(w_m)_{m \geq 1}$ be a Euler–Iwasawa system of rank 2 for (T, A, Σ) , for some lattice T in V , and let*

$$v_m = t_m(x_m, w_m).$$

Then we have

$$v_m \in \mathcal{H}_h(\Gamma) \otimes_{\Lambda(\Gamma)} Q \otimes_{\Lambda(\Gamma)} H^1(\mathbb{Q}(\mu_{mp^\infty}), V),$$

where $h = v_p(\alpha\gamma)$. The elements v_m satisfy the compatibility relation

$$\text{cores}_m^{m\ell} v_{m\ell} = \sigma_\ell \left((\ell - 1)G_\ell(\sigma_\ell^{-1}) - \ell F_\ell(\sigma_\ell^{-1}) \right) v_m.$$

Note that the growth condition $\mathcal{H}_h(\Gamma)$ is consistent with what we have seen for the elements $\mathfrak{z}_m^{f_\alpha, g_\gamma, Np}$ (cf. Theorem 6.8.4) and the compatibility condition between levels m and $m\ell$ is consistent with Theorem 3.4.1. This suggests the following conjecture:

CONJECTURE 8.2.6. *Then there exists a rank 2 Euler–Iwasawa system (w_m) for $(T_{\mathcal{O}_\lambda}(f, g)^*, A, \Sigma)$ with the property that for all $m \in A$, and all choices of p -stabilizations (α, γ) of (f, g) , the Iwasawa cohomology class $\mathfrak{z}_m^{f_\alpha, g_\gamma, Np}$ of Theorem 6.8.4 is given by*

$$\mathfrak{z}_m^{f_\alpha, g_\gamma, Np} = t_m(x_m, w_m)$$

in the notation above.

This gives a conceptual explanation for the (somewhat surprising) growth and compatibility properties of the generalized Beilinson–Flach elements in the context of Perrin-Riou’s theory of higher-rank Euler systems. The authors would like to express their cautious hope that similar rank 1 “shadows” of higher rank Euler systems might also exist in other contexts.

Remark 8.2.7. Note that it is implicit in this conjecture that the elements $t_m(x_m, w_m)$ have no poles (except possibly at the trivial character), so the singularities of t_m , at the characters where one of the $F_\ell(\hat{\sigma}_\ell)$ for $\ell \mid m$ fails to be invertible, must be “cancelled out” by zeroes of w_m .

Appendix A. Ancillary results

A.1. *Fixed points of double cosets.* Here we shall prove a result that is used in the proof of Theorem 3.4.1 above.

Let Γ be a discrete subgroup of $\text{PSL}_2(\mathbb{R})$. Recall that a *fundamental domain* for Γ is a closed subset D of \mathcal{H} such that

- D is equal to the closure of its interior D° ,
- $\bigcup_{\gamma \in \Gamma} \gamma D = \mathcal{H}$,
- $\gamma D^\circ \cap D^\circ = \emptyset$ for all nonidentity elements $\gamma \in \Gamma$.

We assume henceforth that Γ is a *Fuchsian group of the first kind*, i.e., that Γ admits a fundamental domain D with finite hyperbolic area. We shall say that a fundamental domain D is *polygonal* if D is the region bounded by a finite number of geodesic arcs in \mathcal{H} ; it is known that every Γ admits a polygonal fundamental domain.

LEMMA A.1.1. *Let D be a Dirichlet domain for Γ , and let $E = \alpha D$ where α lies in the commensurator $\text{Comm}(\Gamma)$. Then there are only finitely many $\gamma \in \Gamma$ such that $\alpha D \cap \gamma D \neq \emptyset$.*

Proof. It is clear that αD is a Dirichlet domain for $\alpha\Gamma\alpha^{-1}$. In particular, it is polygonal. Hence it can be decomposed as the union of a compact set M and a finite number N_i of “cusp neighbourhoods,” which are subsets bounded by two geodesics intersecting at a vertex at infinity, which is a parabolic point x_i of $\alpha\Gamma\alpha^{-1}$ on the boundary $\mathbb{P}^1(\mathbb{R})$, and an arc of a Euclidean circle tangent to the real line at x_i .

Since M is compact, it can intersect only finitely many Γ -translates of D (cf. [Kat92, Th. 3.5.1]). Moreover, since $\alpha \in \text{Comm}(D)$, the sets of parabolic points of Γ and $\alpha\Gamma\alpha^{-1}$ are the same; so for each vertex-at-infinity x of αD , we may choose some $\gamma \in \Gamma$ that maps a vertex-at-infinity y_i of D to x_i , and it is clear that N_i is contained in a finite union of translates of $\gamma\gamma'D$ where γ' lies in the stabilizer of y_i . Each of these, in turn, intersects finitely many other translates of D (since D has finitely many sides). \square

LEMMA A.1.2. *Let Γ be Fuchsian group of the first kind, and let $X \subset \text{Comm}(\Gamma)$ be a finite union of double cosets $\Gamma\alpha\Gamma$. Then the set*

$$\text{Fix}(X) = \{u \in \mathcal{H} : \gamma u = u \text{ for some } \gamma \in X, \gamma \neq 1\}$$

is a finite union of Γ -orbits in \mathcal{H} .

Proof. Since $X - \{\text{id}\}$ is preserved by conjugation by Γ , the set $\text{Fix}(X)$ is a union of orbits of Γ . So it suffices to show that $\text{Fix}(X) \cap D$ is finite, where D is a Dirichlet domain for Γ .

We claim that there are only finitely many $x \in X$ such that $xD \cap D \neq \emptyset$. From Lemma A.1.1, we know that for each $\alpha \in G$, there are finitely many $\gamma \in \Gamma$ such that $\gamma D \cap \alpha D \neq \emptyset$ and hence finitely many $x \in \Gamma\alpha$ such that $xD \cap D \neq \emptyset$. Since X is the union of finitely many left cosets $\Gamma\alpha_i$, this implies the claim.

However, each nonidentity element in the finite set $\{x \in X : xD \cap D \neq \emptyset\}$ can only have finitely many fixed points in \mathcal{H} and, in particular, in D ; so $\text{Fix}(X) \cap D$ is finite, as required. \square

LEMMA A.1.3. *Let Γ_1, Γ_2 be commensurable Fuchsian groups of the first kind. Then the set*

$$\{u \in \mathcal{H} : \exists c \in \Gamma_1, d \in \Gamma_2 \text{ such that } cd \neq 1 \text{ and } cdu = u\}$$

is a finite union of orbits under $\Gamma_1 \cap \Gamma_2$.

Proof. This follows from the previous lemma applied to $\Gamma = \Gamma_1 \cap \Gamma_2$ and $X = \Gamma_1\Gamma_2$. \square

(Note that if $\Gamma_1 = \Gamma_2$, or more generally if the group generated by Γ_1 and Γ_2 is Fuchsian, this generalizes the well-known result that Fuchsian groups of the first kind have finitely many elliptic points in Γ .)

In particular, we have the following:

PROPOSITION A.1.4. *Let Γ_1, Γ_2 be commensurable Fuchsian groups of the first kind. Then the natural map*

$$(\Gamma_1 \cap \Gamma_2) \backslash \mathcal{H} \rightarrow (\Gamma_1 \backslash \mathcal{H}) \times (\Gamma_2 \backslash \mathcal{H})$$

is injective away from a finite subset of its domain.

Proof. Let z, z' be two points of \mathcal{H} such that $z' \in \Gamma_1 z$ and $z' \in \Gamma_2 z$. Then we may write $z' = \gamma_1 z$ for some $\gamma_1 \in \Gamma_1$ and $z' = \gamma_2 z$ for some $\gamma_2 \in \Gamma_2$.

Hence $\gamma_1^{-1} \gamma_2 z = z$. So either z lies in the finite subset $\text{Fix}(\Gamma_1 \Gamma_2)$ of $(\Gamma_1 \cap \Gamma_2) \backslash \mathcal{H}$, or $\gamma_1^{-1} \gamma_2 = 1$, in which case z and z' are clearly in the same orbit under $\Gamma_1 \cap \Gamma_2$. □

A.2. *Unbounded Iwasawa cohomology.* In this section, we shall consider inverse systems of cohomology classes in \mathbb{Z}_p^\times -extensions that are not bounded (as in the usual definition of Iwasawa cohomology) but satisfy a weaker growth condition.

Let K be a finite extension of either \mathbb{Q} or \mathbb{Q}_p . If K is global, suppose that either $p \neq 2$, or K has no real places.

Notation. In order to handle the two cases in a uniform manner, we shall adopt a notation that is slightly abusive: for T a \mathbb{Z}_p -representation of $\text{Gal}(\bar{K}/K)$, the notation $H^i(K, T)$ will mean either $H^i(K, T)$ as defined above if G is local, or what we previously called $H_S^i(K, T)$ if K is global, where S is some fixed finite set of places containing all infinite places and all those dividing p . In the latter case, we will assume that S contains all primes at which T is ramified.

As before, we let $K_n = K(\mu_{p^n})$ and $K_\infty = \bigcup_n K_n$, and define Iwasawa cohomology groups $H_{\text{Iw}}^i(K_\infty, T)$ as the inverse limit of the $H^i(K_n, T)$ with respect to corestriction, with their natural module structure over $\Lambda = \Lambda_{\mathbb{Z}_p}(\Gamma)$.

PROPOSITION A.2.1 (Nekovar). *For any $j \in \{0, 1, 2\}$, we have a short exact sequence*

$$(22) \quad 0 \longrightarrow H_{\text{Iw}}^j(K_\infty, T)_{\Gamma_n} \longrightarrow H^j(K_n, T) \longrightarrow H_{\text{Iw}}^{j+1}(K_\infty, T)^{\Gamma_n} \longrightarrow 0.$$

Proof. This is Corollary 8.4.8.2 of [Nek06]. We briefly recall the proof. There are natural isomorphisms

$$\begin{aligned} H_{\text{Iw}}^i(K_\infty, T) &\cong H^i(K, \Lambda \otimes_{\mathbb{Z}_p} T), \\ H^i(K_n, T) &\cong H^i(K, \mathbb{Z}_p[\Gamma/\Gamma_n] \otimes_{\mathbb{Z}_p} T) \end{aligned}$$

(for a suitable Λ -linear action of G_K on the tensor products); see Proposition 8.3.5 of op. cit. Then the result above follows from the long exact cohomology sequence of K -cohomology attached to the short exact sequence of $\Lambda[G_K]$ -modules

$$0 \longrightarrow \Lambda \otimes_{\mathbb{Z}_p} T \xrightarrow{[\gamma_n]-1} \Lambda \otimes_{\mathbb{Z}_p} T \longrightarrow \mathbb{Z}_p[\Gamma/\Gamma_n] \otimes_{\mathbb{Z}_p} T \longrightarrow 0. \quad \square$$

PROPOSITION A.2.2 (Perrin-Riou). *There is an exact sequence*

$$0 \longrightarrow T^{G_{K_\infty}} \longrightarrow H_{\text{Iw}}^1(K_\infty, T) \longrightarrow \text{Hom}_\Lambda(H^1(K_\infty, V/T)^\vee, \Lambda)^\iota \longrightarrow (\text{finite}) \longrightarrow 0,$$

where ι signifies that the Λ -module structure is composed with the automorphism $\gamma \mapsto \gamma^{-1}$. In both cases, this exact sequence identifies $T^{G_{K_\infty}}$ with the Λ -torsion submodule of $H_{\text{Iw}}^1(K_\infty, T)$.

Proof. The local case is [PR92, Prop. 2.1.6]; note that in the local situation, Tate duality furnishes an isomorphism $H^1(K_\infty, V/T)^\vee \cong H_{\text{Iw}}^1(K_\infty, T^*(1))$ and the middle map can be interpreted as Perrin-Riou’s pairing $H_{\text{Iw}}^1(K_\infty, T) \times H_{\text{Iw}}^1(K_\infty, T^*(1)) \rightarrow \Lambda$. The global case is [PR95, Lemma 1.3.3]. \square

The main object of study in this section is the following module. Let $V = \mathbb{Q}_p \otimes_{\mathbb{Z}_p} T$.

Definition A.2.3. For K, T, V as above, and $0 \leq r < 1$, let $Y_r(K_\infty, V)$ be the space of sequences $(c_n)_{n \geq 0} \in \varprojlim_n H^1(K_n, V)$ such that there exists $\delta < \infty$ independent of n for which $p^{\lfloor rn \rfloor + \delta} c_n$ is in the image of $H^1(K_n, T)$ in $H^1(K_n, V)$.

PROPOSITION A.2.4. *For all $0 \leq r < 1$, the natural map $\lambda_r : \mathcal{H}_r(\Gamma) \otimes_{\Lambda_{\mathbb{Q}_p}(\Gamma)} H_{\text{Iw}}^1(K_\infty, V) \rightarrow \varprojlim_n H^1(K_n, V)$ has image contained in $Y_r(K_\infty, V)$.*

Proof. This is clear from the definition of $\mathcal{H}_r(\Gamma)$. \square

Remark A.2.5. The map λ_r is not necessarily injective, even for $r = 0$ (where $\mathcal{H}_r(\Gamma)$ is just $\Lambda \otimes \mathbb{Q}_p$). A counterexample is provided by the representation $T = \mathbb{Z}_p(1)$. Then the cocycle c_n given by $\sigma \mapsto \frac{\chi(\sigma)-1}{p^n}$ is well defined as an element of $H^1(K_n, T)$ (for either local or global K). The sequence (c_n) defines an element of $H_{\text{Iw}}^1(K_\infty, T)$ that is not p -torsion and thus is nonzero as an element of $H_{\text{Iw}}^1(K_\infty, V)$. But $p^n c_n$ is a coboundary for all n , so the image of c_n in $H^1(K_n, V)$ is zero for all n . Thus (c_n) lies in the kernel of the above map.

PROPOSITION A.2.6. *The kernel of λ_r is contained in $H_{\text{Iw}}^1(K_\infty, V)_{\text{tors}} \cong V^{G_{K_\infty}}$.*

Remark A.2.7. We note first that this statement does make sense, since for any $\Lambda_{\mathbb{Q}_p}(\Gamma)$ -torsion module M , tensoring with $1 \in \mathcal{H}_r(\Gamma)$ gives an isomorphism $\mathcal{H}_r(\Gamma) \otimes_{\Lambda_{\mathbb{Q}_p}(\Gamma)} M \cong M$.

Proof. Tensoring (22) with \mathbb{Q}_p , we find that the map

$$H_{\text{Iw}}^1(K_\infty, V)_{\Gamma_n} \rightarrow H^1(K_n, V)$$

is injective. Thus the kernel of λ_r consists of those elements lying in

$$\bigcap_{n \geq 0} (\gamma_n - 1) \left(\mathcal{H}_r(\Gamma) \otimes_{\Lambda_{\mathbb{Q}_p}(\Gamma)} H_{\text{Iw}}^1(K_\infty, V) \right).$$

Since $H_{\text{Iw}}^1(K_\infty, V)$ is a finitely-generated module over the subring $\Lambda_{\mathbb{Q}_p}(\Gamma_1) \subset \Lambda_{\mathbb{Q}_p}(\Gamma)$, which is a PID, we may write it as the direct sum of its torsion submodule and a complementary free submodule. Since $r < 1$, we find that

$$\bigcap_{n \geq 1} (\gamma_n - 1) \mathcal{H}_r(\Gamma) = 0,$$

and hence the kernel of λ_r is contained in the torsion part of $H_{\text{Iw}}^1(K_\infty, V)$, which is equal to $V^{G_{K_\infty}}$ by Proposition A.2.2. \square

Remark A.2.8. Although we shall not need this, it clearly follows that the kernel of λ_r is equal to $\bigcup_{n \geq 0} (\gamma_n - 1) V^{G_{K_\infty}}$, which is the unique Γ -invariant complement of $\bigcup_n H^0(K_n, V)$ in $H^0(K_\infty, V)$.

PROPOSITION A.2.9. *Let K be a p -adic field, and suppose that $V^{G_{K_\infty}} = 0$. Then the map*

$$\mathcal{H}_r(\Gamma) \otimes_{\Lambda_{\mathbb{Q}_p}(\Gamma)} H_{\text{Iw}}^1(K_\infty, V) \rightarrow Y_r(K_\infty, V)$$

is an isomorphism.

Proof. By Proposition A.2.2, our hypotheses imply that $H_{\text{Iw}}^1(K_\infty, V)$ is a torsion-free $\Lambda_{\mathbb{Q}_p}(\Gamma)$ -module; hence it is free, since $\Lambda_{\mathbb{Q}_p}(\Gamma)$ is a finite product of principal ideal domains (and the ranks of the Γ_{tors} -isotypical direct summands of $H_{\text{Iw}}^1(K_\infty, V)$ are all equal). Thus there exists a free basis x_1, \dots, x_d of $H_{\text{Iw}}^1(K_\infty, V)$, where $d = \dim_{\mathbb{Q}_p}(V)$.

For each n , the cokernel of the projection map

$$H_{\text{Iw}}^1(K, T) \rightarrow H^1(K_n, T)$$

is finite, and its order is bounded independently of n . (In fact, the cokernel of this map is isomorphic to the Γ_n -invariants of

$$H_{\text{Iw}}^2(K_\infty, T) \cong H^0(K_\infty, (V/T)^*(1))^\vee,$$

and $H^0(K_\infty, (V/T)^*(1))$ is finite, since $H^0(K_\infty, V^*(1)) = H^0(K_\infty, V)^*(1) = 0$.)

Thus the map

$$H_{\text{Iw}}^1(K_\infty, V) \rightarrow H^1(K_n, V)$$

is surjective for all n , and there is $\nu < \infty$ independent of n such that the \mathbb{Z}_p -submodule spanned by the images of x_1, \dots, x_d in $H^1(K_n, V)$ contains $p^\nu \cdot \frac{H^1(K_n, T)}{\text{torsion}}$.

Consequently, given any sequence $(c_n)_{n \geq 0} \in Y_r(K_\infty, V)$, we have for each n uniquely determined elements $b_1^{(n)}, \dots, b_d^{(n)} \in \mathbb{Q}_p[\Gamma/\Gamma_n]$ such that $\sum_{i=1}^d b_i^{(n)} x_i^{(n)} = c_n$, where $x_i^{(n)}$ is the image of x_i in $H^1(K_n, V)$. Also, for each i , the sequence $(b_i^{(n)})_{n \geq 0}$ is compatible under projection (by uniqueness), and its valuation is bounded below by $-[rn] - \delta - \nu$; so (as $r < 1$) there is a unique element $b_i \in \mathcal{H}_r(\Gamma)$ whose image at level n is $b_i^{(n)}$ for all n . Then it is clear that $c = \sum_i b_i \otimes x_i \in \mathcal{H}_r(\Gamma) \otimes_{\Lambda_{\mathbb{Q}_p}(\Gamma)} H_{\text{Iw}}^1(K_\infty, V)$ is a preimage of $(c_n)_{n \geq 0}$; by the previous proposition, it is unique. \square

In the global case we cannot prove quite such a strong result, as we do not have such good control over $H_{\text{Iw}}^2(K_\infty, T)$; the following rather more specific result (which applies to both local and global cases) will suffice for our purposes:

PROPOSITION A.2.10. *Let $r < 1$, and suppose T has the structure of a module over \mathcal{O}_E , for some finite extension E/\mathbb{Q}_p , and that $T^{G_{K_\infty}} = 0$. Let $\alpha \in \mathcal{O}_E$ such that $v_p(\alpha) \leq r$, and suppose we are given elements $x_n \in H^1(K_n, T)$ for $n \geq 0$ satisfying*

$$\text{cores}_n^{n+1}(x_{n+1}) = \alpha x_n.$$

Then there is a unique element $x \in \mathcal{H}_r(\Gamma) \otimes_{\Lambda} H_{\text{Iw}}^1(K_\infty, T)$ whose image in $H^1(K_n, V)$ is equal to $\alpha^{-n} c_n$ for all n .

Proof. We claim that the hypotheses of the theorem force each c_n to be the image of an element of $H_{\text{Iw}}^1(K_\infty, V)_{\Gamma_n}$. To prove this, we shall argue much as in the proof of Proposition 6.6.2. We begin by noting that $H_{\text{Iw}}^2(K_\infty, T)$ is a finitely-generated Λ -module and the subgroups $M_n := H_{\text{Iw}}^2(K_\infty, T)^{\Gamma_n}$ are Λ -submodules (since Γ is abelian). As Λ is a Noetherian ring, the ascending chain of submodules $(M_n)_{n \geq 0}$ must eventually stabilize; that is, there is an n_0 such that $M_n = M_{n_0}$ for all $n \geq n_0$.

The corestriction map $H^1(K_{n+1}, T) \rightarrow H^1(K_n, T)$ corresponds to the trace map $M_{n+1} \rightarrow M_n$; when $n \geq n_0$, this is simply multiplication by p on M_{n_0} . Since $v_p(\alpha) < 1$, we deduce as in Proposition 6.6.2 that for all $n \geq 0$, the image of x_n is contained in the torsion submodule of M_n . Inverting p , the torsion is killed, and the image of x_n in $H_{\text{Iw}}^2(K_\infty, V)^{\Gamma_n}$ is 0; so x_n lies in the submodule $H_{\text{Iw}}^1(K_\infty, V)_{\Gamma_n} \subseteq H^1(K_n, V)$.

Now let us choose a basis of the free $\Lambda_{\mathbb{Q}_p}(\Gamma)$ -module $H_{\text{Iw}}^1(K_\infty, V)$. In order to apply the argument of the previous proposition, we need only check that the order of the torsion subgroup of M_n is bounded independently of n ; but this is immediate from the fact that the M_n stabilize for large n (and are all finitely-generated as \mathbb{Z}_p -modules). This shows that there exists a μ such that $\alpha^{-n} c_n$ lies in $p^{-[rn] - \mu} H_{\text{Iw}}^1(K_\infty, T)$, and the argument proceeds as in Proposition A.2.9. \square

References

- [AV75] Y. AMICE and J. VÉLU, Distributions p -adiques associées aux séries de Hecke, in *Journées Arithmétiques de Bordeaux* (Conf., Univ. Bordeaux, Bordeaux, 1974), *Astérisque* **24–25**, Soc. Math. France, Paris, 1975, pp. 119–131. MR 0376534. Zbl 0332.14010.
- [AS86] A. ASH and G. STEVENS, Modular forms in characteristic l and special values of their L -functions, *Duke Math. J.* **53** (1986), 849–868. MR 0860675. Zbl 0618.10026. <http://dx.doi.org/10.1215/S0012-7094-86-05346-9>.
- [Beĭ84] A. A. BEĬLINSON, Higher regulators and values of L -functions, in *Current Problems in Mathematics, Vol. 24, Itogi Nauki i Tekhniki*, Akad. Nauk SSSR Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1984, pp. 181–238. MR 0760999.
- [BLZ04] L. BERGER, H. LI, and H. J. ZHU, Construction of some families of 2-dimensional crystalline representations, *Math. Ann.* **329** (2004), 365–377. MR 2060368. Zbl 1085.11028. <http://dx.doi.org/10.1007/s00208-004-0529-y>.
- [BDR12] M. BERTOLINI, H. DARMON, and V. ROTGER, Beilinson–Flach elements and Euler systems I: syntomic regulators and p -adic Rankin L -series, 2012, preprint. Available at <http://www-ma2.upc.edu/vrotger/docs/BDR1.pdf>.
- [Bes00] A. BESSER, Syntomic regulators and p -adic integration. I. Rigid syntomic regulators, in *Proceedings of the Conference on p -adic Aspects of the Theory of Automorphic Representations* (Jerusalem, 1998), *Israel J. Math.* **120**, 2000, pp. 291–334. MR 1809626. Zbl 1001.19003. <http://dx.doi.org/10.1007/BF02834843>.
- [BK90] S. BLOCH and K. KATO, L -functions and Tamagawa numbers of motives, in *The Grothendieck Festschrift, Vol. I, Progr. Math.* **86**, Birkhäuser, Boston, MA, 1990, pp. 333–400. MR 1086888. Zbl 0768.14001. http://dx.doi.org/10.1007/978-0-8176-4574-8_9.
- [CE98] R. F. COLEMAN and B. EDIXHOVEN, On the semi-simplicity of the U_p -operator on modular forms, *Math. Ann.* **310** (1998), 119–127. MR 1600034. Zbl 0902.11020. <http://dx.doi.org/10.1007/s002080050140>.
- [DR14] H. DARMON and V. ROTGER, Diagonal cycles and Euler systems I: A p -adic Gross–Zagier formula, *Ann. Sci. Éc. Norm. Supér.* **47** (2014), to appear.
- [Del77] P. DELIGNE, Cohomologie étale, in *Séminaire de Géométrie Algébrique du Bois-Marie SGA 4 $\frac{1}{2}$, Lecture Notes in Math.* **569**, Springer-Verlag, New York, 1977, avec la collaboration de J. F. Boutot, A. Grothendieck, L. Illusie et J. L. Verdier, pp. iv+312pp. MR 0463174. Zbl 0345.00010. <http://dx.doi.org/10.1007/BFb0091516>.
- [DR73] P. DELIGNE and M. RAPOPORT, Les schémas de modules de courbes elliptiques, in *Modular Functions of One Variable, II* (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972), *Lecture Notes in Math.* **349**, Springer-Verlag, New York, 1973, pp. 143–316. MR 0337993. Zbl 0281.14010. http://dx.doi.org/10.1007/978-3-540-37855-6_4.

- [DI95] F. DIAMOND and J. IM, Modular forms and modular curves, in *Seminar on Fermat's Last Theorem* (Toronto, ON, 1993–1994), *CMS Conf. Proc.* **17**, Amer. Math. Soc., Providence, RI, 1995, pp. 39–133. MR 1357209. Zbl 0853.11032.
- [Dri73] V. DRINFEL'D, Two theorems on modular curves, *Funct. Anal. Appl.* **7** (1973), 155–156, translation of the Russian original. MR 0318157. Zbl 0285.14006. <http://dx.doi.org/10.1007/BF01078890>.
- [Fla92] M. FLACH, A finiteness theorem for the symmetric square of an elliptic curve, *Invent. Math.* **109** (1992), 307–327. MR 1172693. Zbl 0781.14022. <http://dx.doi.org/10.1007/BF01232029>.
- [FM87] J.-M. FONTAINE and W. MESSING, p -adic periods and p -adic étale cohomology, in *Current Trends in Arithmetical Algebraic Geometry* (Arcata, Calif., 1985), *Contemp. Math.* **67**, Amer. Math. Soc., Providence, RI, 1987, pp. 179–207. MR 0902593. Zbl 0632.14016. <http://dx.doi.org/10.1090/conm/067/902593>.
- [Hub00] A. HUBER, Realization of Voevodsky's motives, *J. Algebraic Geom.* **9** (2000), 755–799. MR 1775312. Zbl 0994.14014.
- [IM65] N. IWAHORI and H. MATSUMOTO, On some Bruhat decomposition and the structure of the Hecke rings of p -adic Chevalley groups, *Inst. Hautes Études Sci. Publ. Math.* **25** (1965), 5–48. MR 0185016. Zbl 0228.20015. <http://dx.doi.org/10.1007/BF02684396>.
- [Jac72] H. JACQUET, *Automorphic Forms on $GL(2)$. Part II, Lecture Notes in Math.* **278**, Springer-Verlag, New York, 1972. MR 0562503. Zbl 0243.12005.
- [Jan88a] U. JANNSEN, Continuous étale cohomology, *Math. Ann.* **280** (1988), 207–245. MR 0929536. Zbl 0649.14011. <http://dx.doi.org/10.1007/BF01456052>.
- [Jan88b] U. JANNSEN, Deligne homology, Hodge- \mathcal{D} -conjecture, and motives, in *Beilinson's Conjectures on Special Values of L -Functions*, *Perspect. Math.* **4**, Academic Press, Boston, MA, 1988, pp. 305–372. MR 0944998. Zbl 0701.14019.
- [Kat04] K. KATO, p -adic Hodge theory and values of zeta functions of modular forms, in *Cohomologies p -Adiques et Applications Arithmétiques. III*, **295**, Soc. Math. France, Paris, 2004, pp. ix, 117–290. MR 2104361. Zbl 1142.11336. Available at http://smf4.emath.fr/en/Publications/Asterisque/2004/295/html/smf_ast_295_117-290.html.
- [Kat92] S. KATOK, *Fuchsian Groups, Chicago Lectures in Math.*, University of Chicago Press, Chicago, IL, 1992. MR 1177168. Zbl 0753.30001.
- [Kur02] M. KURIHARA, On the Tate Shafarevich groups over cyclotomic fields of an elliptic curve with supersingular reduction. I, *Invent. Math.* **149** (2002), 195–224. MR 1914621. Zbl 1033.11028. <http://dx.doi.org/10.1007/s002220100206>.

- [Lan91] S. E. LANDSBURG, Relative Chow groups, *Illinois J. Math.* **35** (1991), 618–641. MR 1115990. Zbl 0746.19007. Available at <http://projecteuclid.org/euclid.ijm/1255987675>.
- [Lan02] S. LANG, *Algebra*, third ed., *Grad. Texts in Math.* **211**, Springer-Verlag, New York, 2002. MR 1878556. Zbl 0984.00001. <http://dx.doi.org/10.1007/978-1-4613-0041-0>.
- [Lau14] A. G. B. LAUDER, Efficient computation of Rankin p -adic L -functions, in *Computations with Modular Forms: Proceedings of a Summer School and Conference, Heidelberg, August/September 2011* (G. BOECKLE and G. WIESE, eds.), Springer-Verlag, New York, 2014, pp. 181–200. http://dx.doi.org/10.1007/978-3-319-03847-6_7.
- [Lev04] M. LEVINE, K -theory and motivic cohomology for schemes, I, 2004, preprint. Available at <https://www.uni-due.de/~bm0032/publ/KthyMotI12.01.pdf>.
- [LZ12] D. LOEFFLER and S. L. ZERBES, Iwasawa theory and p -adic L -functions over \mathbf{Z}_p^2 -extensions, 2012, *Int. J. Number Theory*, to appear. arXiv 1108.5954. <http://dx.doi.org/10.1142/S1793042114500699>.
- [LZ13] D. LOEFFLER and S. L. ZERBES, Wach modules and critical slope p -adic L -functions, *J. Reine Angew. Math.* **679** (2013), 181–206. MR 3065158. Zbl 1276.11192. <http://dx.doi.org/10.1515/crelle.2012.012>.
- [MVW06] C. MAZZA, V. VOEVODSKY, and C. WEIBEL, *Lecture Notes on Motivic Cohomology*, *Clay Math. Monogr.* no. 2, Amer. Math. Soc., Providence, RI, 2006. MR 2242284. Zbl 1115.14010.
- [Mil12] J. S. MILNE, Lectures on étale cohomology, 2012. Available at <http://www.jmilne.org/math/CourseNotes/lec.html>.
- [Mom81] F. MOMOSE, On the l -adic representations attached to modular forms, *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* **28** (1981), 89–109. MR 0617867. Zbl 0482.10023.
- [Nek06] J. NEKOVÁŘ, *Selmer Complexes*, *Astérisque* **310**, Soc. Math. France, Paris, 2006. MR 2333680. Zbl 1211.11120.
- [Oht99] M. OHTA, Ordinary p -adic étale cohomology groups attached to towers of elliptic modular curves, *Compositio Math.* **115** (1999), 241–301. MR 1674001. Zbl 0967.11015. <http://dx.doi.org/10.1023/A:1000556212097>.
- [Oht00] M. OHTA, Ordinary p -adic étale cohomology groups attached to towers of elliptic modular curves. II, *Math. Ann.* **318** (2000), 557–583. MR 1800769. Zbl 0967.11016. <http://dx.doi.org/10.1007/s002080000119>.
- [Ots09] R. OTSUKI, Construction of a homomorphism concerning Euler systems for an elliptic curve, *Tokyo J. Math.* **32** (2009), 253–278. MR 2541166. Zbl 1188.11029. <http://dx.doi.org/10.3836/tjm/1249648421>.
- [PR92] B. PERRIN-RIOU, Théorie d’Iwasawa et hauteurs p -adiques, *Invent. Math.* **109** (1992), 137–185. MR 1168369. Zbl 0781.14013. <http://dx.doi.org/10.1007/BF01232022>.

- [PR95] B. PERRIN-RIOU, *Fonctions L p -Adiques des Représentations p -Adiques*, *Astérisque* **229**, Soc. Math. France, Paris, 1995. MR 1327803. Zbl 0845.11040.
- [PR98] B. PERRIN-RIOU, Systèmes d'Euler p -adiques et théorie d'Iwasawa, *Ann. Inst. Fourier (Grenoble)* **48** (1998), 1231–1307. MR 1662231. Zbl 0930.11078. <http://dx.doi.org/10.5802/aif.1655>. Available at http://www.numdam.org/item?id=AIF_1998__48_5_1231_0.
- [PS11] R. POLLACK and G. STEVENS, Overconvergent modular symbols and p -adic L -functions, *Ann. Sci. Éc. Norm. Supér.* **44** (2011), 1–42. MR 2760194. Zbl 1268.11075.
- [PS13] R. POLLACK and G. STEVENS, Critical slope p -adic L -functions, *J. Lond. Math. Soc.* **87** (2013), 428–452. MR 3046279. Zbl 06160939. <http://dx.doi.org/10.1112/jlms/jds057>.
- [Qui73] D. QUILLEN, Higher algebraic K -theory. I, in *Algebraic K-theory, I: Higher K-Theories* (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972), *Lecture Notes in Math.* **341**, Springer-Verlag, New York, 1973, pp. 85–147. MR 0338129. Zbl 0292.18004. <http://dx.doi.org/10.1007/BFb0067053>.
- [Ram00] D. RAMAKRISHNAN, Modularity of the Rankin-Selberg L -series, and multiplicity one for $SL(2)$, *Ann. of Math.* **152** (2000), 45–111. MR 1792292. Zbl 0989.11023. <http://dx.doi.org/10.2307/2661379>.
- [Rib85] K. A. RIBET, On l -adic representations attached to modular forms. II, *Glasgow Math. J.* **27** (1985), 185–194. MR 0819838. Zbl 0596.10027. <http://dx.doi.org/10.1017/S0017089500006170>.
- [Rub00] K. RUBIN, *Euler Systems*, *Ann. of Math. Stud.* **147**, Princeton Univ. Press, Princeton, NJ, 2000. MR 1749177. Zbl 0977.11001.
- [SS10] S. SAITO and K. SATO, A p -adic regulator map and finiteness results for arithmetic schemes, *Doc. Math.* (2010), 525–594. MR 2804264. Zbl 1207.14015. Available at http://www.math.uiuc.edu/documenta/vol-suslin/saito_sato.pdf.
- [Sch98] A. J. SCHOLL, An introduction to Kato's Euler systems, in *Galois Representations in Arithmetic Algebraic Geometry* (Durham, 1996), *London Math. Soc. Lecture Note Ser.* **254**, Cambridge Univ. Press, Cambridge, 1998, pp. 379–460. MR 1696501. Zbl 0952.11015. <http://dx.doi.org/10.1017/CBO9780511662010.011>.
- [Sha81] F. SHAHIDI, On certain L -functions, *Amer. J. Math.* **103** (1981), 297–355. MR 0610479. Zbl 0467.12013. <http://dx.doi.org/10.2307/2374219>.
- [Shi76] G. SHIMURA, The special values of the zeta functions associated with cusp forms, *Comm. Pure Appl. Math.* **29** (1976), 783–804. MR 0434962. Zbl 0348.10015. <http://dx.doi.org/10.1002/cpa.3160290618>.
- [Shi86] G. SHIMURA, On a class of nearly holomorphic automorphic forms, *Ann. of Math.* **123** (1986), 347–406. MR 0835767. Zbl 0593.10022. <http://dx.doi.org/10.2307/1971276>.

- [Shi00] G. SHIMURA, *Arithmeticity in the Theory of Automorphic Forms*, *Math. Surveys Monogr.* **82**, Amer. Math. Soc., Providence, RI, 2000. MR 1780262. Zbl 0967.11001. <http://dx.doi.org/10.1090/surv/082>.
- [SD73] H. P. F. SWINNERTON-DYER, On l -adic representations and congruences for coefficients of modular forms, in *Modular Functions of One Variable, III* (Proc. Internat. Summer School, Univ. Antwerp, 1972), *Lecture Notes in Math.* **350**, Springer-Verlag, New York, 1973, pp. 1–55. MR 0406931. Zbl 0267.10032. http://dx.doi.org/10.1007/978-3-540-37802-0_1.
- [Sage] THE SAGE GROUP, Sage Mathematics Software. Available at <http://www.sagemath.org/>.
- [Voe02] V. VOEVODSKY, Motivic cohomology groups are isomorphic to higher Chow groups in any characteristic, *Int. Math. Res. Not.* **2002** (2002), 351–355. MR 1883180. Zbl 1057.14026. <http://dx.doi.org/10.1155/S107379280210403X>.

(Received: February 7, 2013)

(Revised: August 21, 2013)

McGILL UNIVERSITY, MONTREAL, QC, CANADA
Current address: UNIVERSITÉ LAVAL, QUÉBEC, CANADA
E-mail: antonio.lei@mcgill.ca

MATHEMATICS INSTITUTE, UNIVERSITY OF WARWICK, COVENTRY, UK
E-mail: d.a.loeffler@warwick.ac.uk

UNIVERSITY COLLEGE LONDON, LONDON, UK
E-mail: s.zerbes@ucl.ac.uk