

# Total Betti numbers of modules of finite projective dimension

By MARK E. WALKER

## Abstract

The Buchsbaum-Eisenbud-Horrocks Conjecture predicts that the  $i^{\text{th}}$  Betti number  $\beta_i(M)$  of a nonzero module  $M$  of finite length and finite projective dimension over a local ring  $R$  of dimension  $d$  should be at least  $\binom{d}{i}$ . It would follow from the validity of this conjecture that  $\sum_i \beta_i(M) \geq 2^d$ . We prove the latter inequality holds in a large number of cases and that, when  $R$  is a complete intersection in which 2 is invertible, equality holds if and only if  $M$  is isomorphic to the quotient of  $R$  by a regular sequence of elements.

## 1. Introduction

We recall a long-standing conjecture (see [3, 1.4] and [6, Prob. 24]):

CONJECTURE (Buchsbaum-Eisenbud-Horrocks Conjecture). *Let  $R$  be a commutative Noetherian ring such that  $\text{Spec}(R)$  is connected, and let  $M$  be a nonzero, finitely generated  $R$ -module of finite projective dimension. For any finite projective resolution  $0 \rightarrow P_d \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow M \rightarrow 0$  of  $M$ , we have*

$$\text{rank}_R(P_i) \geq \binom{c}{i},$$

where  $c = \text{height}_R(\text{ann}_R(M))$ , the height of the annihilator ideal of  $M$ .

The validity of the Buchsbaum-Eisenbud-Horrocks Conjecture would imply that the “total rank” of any projective resolution of  $M$  is at least  $2^c$ . In this paper, we prove this latter inequality holds in a large number of cases:

THEOREM 1. *Assume  $R$ ,  $M$ , and  $P$  are as in the Buchsbaum-Eisenbud-Horrocks Conjecture and, in addition, that*

---

Keywords: Betti numbers, Buchsbaum-Eisenbud-Horrocks Conjecture

AMS Classification: Primary: 13D02.

This work was partially supported by grant #318705 from the Simons Foundation.

© 2017 Department of Mathematics, Princeton University.

- (1)  $R$  is locally a complete intersection and  $M$  is 2-torsion free, or
  - (2)  $R$  contains  $\mathbb{Z}/p$  as a subring for an odd prime  $p$ .
- Then  $\sum_i \text{rank}_R(P_i) \geq 2^c$ , where  $c = \text{height}_R(\text{ann}_R(M))$ .

**Theorem 2** below is the special case of **Theorem 1** in which we assume  $R$  is a local ring and  $M$  has finite length. We record it as a separate theorem since **Theorem 1** follows immediately from it and also because in the local situation we can say a bit more.

For a local ring  $R$  and a finitely generated  $R$ -module  $M$ , let  $\beta_i(M)$  be the  $i^{\text{th}}$  Betti number of  $R$ , defined to be the rank of the  $i^{\text{th}}$  free module in the minimal free resolution of  $M$ .

**THEOREM 2.** *Assume  $(R, \mathfrak{m}, k)$  is a local (Noetherian, commutative) ring of Krull dimension  $d$  and that  $M$  is a nonzero  $R$ -module of finite length and finite projective dimension. If either*

- (1)  $R$  is the quotient of a regular local ring by a regular sequence of elements and 2 is invertible in  $R$ , or
- (2)  $R$  contains  $\mathbb{Z}/p$  as a subring for an odd prime  $p$ ,

then  $\sum_i \beta_i(M) \geq 2^d$ .

Moreover, if the assumptions in (1) hold and  $\sum_i \beta_i(M) = 2^d$ , then  $M$  is isomorphic to the quotient of  $R$  by a regular sequence of  $d$  elements.

To see that **Theorem 1** follows from **Theorem 2**, with the notation of the first theorem, let  $\mathfrak{p}$  be a minimal prime containing  $\text{ann}_R(M)$  of height  $c$ . Then  $\dim(R_{\mathfrak{p}}) = c$ ,  $M_{\mathfrak{p}}$  has finite length, and  $\beta_i(M_{\mathfrak{p}}) \leq \text{rank}_R(P_i)$  for all  $i$ . Moreover, if  $M$  is 2-torsion free, then  $2 \notin \mathfrak{p}$  and hence is invertible in  $R_{\mathfrak{p}}$ .

I thank Seth Lindokken, Michael Brown, Claudia Miller, Peder Thompson and Luchezar Avramov for useful conversations about this paper.

## 2. Complete intersections of residual characteristic not 2

In this section we prove part (1) of **Theorem 2** and the assertion concerning when the equation  $\sum_i \beta_i(M) = 2^d$  holds; see **Theorem 2.4** below.

For any local ring  $(R, \mathfrak{m}, k)$ , let  $\text{Perf}^{\text{fl}}(R)$  be the category of bounded complexes of finite rank free  $R$ -modules  $F$  such that  $H_i(F)$  has finite length for all  $i$ , and define  $K_0^{\text{fl}}(R)$  to be the Grothendieck group of  $\text{Perf}^{\text{fl}}(R)$ . Recall that  $K_0^{\text{fl}}(R)$  is generated by isomorphism classes of objects of  $\text{Perf}^{\text{fl}}(R)$ , modulo relations coming from short exact sequences and quasi-isomorphisms.

Let  $\psi^2 : K_0^{\text{fl}}(R) \rightarrow K_0^{\text{fl}}(R)$  be the 2<sup>nd</sup> Adams operation, as defined by Gillet-Soulé [4]. Gillet-Soulé's definition involves the Dold-Kan correspondence between complexes and simplicial modules, but if 2 is invertible in  $R$ , then  $\psi^2$  admits a simpler description: For  $F \in \text{Perf}^{\text{fl}}(R)$ , let  $T^2(F)$  denote its second tensor power  $F \otimes_R F$  endowed with the action of the symmetric group  $\Sigma_2 = \langle \tau \rangle$

given by

$$\tau \cdot (x \otimes y) = (-1)^{|x||y|} y \otimes x.$$

Since  $\frac{1}{2} \in R$ , we have a direct sum decomposition  $T^2(F) = S^2(F) \oplus \Lambda^2(F)$ , where  $S^2(F) := \ker(\tau - \text{id})$  and  $\Lambda^2(F) := \ker(\tau + \text{id})$ . By [1, 6.14] we have

$$(2.1) \quad \psi^2[F] = [S^2(F)] - [\Lambda^2(F)] \in K_0^{\text{fl}}(R).$$

Let  $\ell_R$  denote the length of an  $R$ -module, and write  $\chi : K_0^{\text{fl}}(R) \rightarrow \mathbb{Z}$  for the Euler characteristic map:  $\chi([F]) = \sum_i (-1)^i \ell_R H_i(F)$ .

PROPOSITION 2.2 (Gillet-Soulé; see [4, 7.1]). *If  $R$  is a local complete intersection of dimension  $d$ , then  $\chi \circ \psi^2 = 2^d \cdot \chi$ .*

Definition 2.3. A local ring  $(R, \mathfrak{m}, k)$  of dimension  $d$  such that 2 is invertible in  $R$  will be called a *quasi-Roberts ring* if there we have an equality of maps  $\chi \circ \psi^2 = 2^d \cdot \chi$ .

THEOREM 2.4. *Let  $(R, \mathfrak{m}, k)$  be a local ring of dimension  $d$  such that 2 is invertible in  $R$ . If  $R$  is a quasi-Roberts ring, then for any nonzero  $R$ -module  $M$  of finite length and finite projective dimension, we have  $\sum_i \beta_i(M) \geq 2^d$ .*

Moreover, if  $\sum_i \beta_i(M) = 2^d$ , then  $M \cong R/(y_1, \dots, y_d)$  for some regular sequence of elements  $y_1, \dots, y_d \in \mathfrak{m}$ .

Proof. Let  $F$  be the minimal free resolution of  $M$ , so that  $\chi(F) = \ell_R(M)$  and  $\text{rank}_R(F_i) = \beta_i(M)$ . Using (2.1) we get

$$(2.5) \quad \begin{aligned} 2^d \cdot \ell_R(M) &= \chi(\psi^2(F)) = \sum_i (-1)^i \ell_R H_i(S^2(F)) - \sum_j (-1)^j \ell_R H_j(\Lambda^2(F)) \\ &\leq \sum_{i \text{ even}} \ell_R H_i(S^2(F)) + \sum_{i \text{ odd}} \ell_R H_i(\Lambda^2(F)). \end{aligned}$$

Since  $S^2(F)$  and  $\Lambda^2(F)$  are direct summands of  $F \otimes_R F$ ,

$$(2.6) \quad \sum_{i \text{ even}} \ell_R H_i(S^2(F)) + \sum_{i \text{ odd}} \ell_R H_i(\Lambda^2(F)) \leq \sum_i \ell_R H_i(F \otimes_R F).$$

For each  $i$ ,  $H_i(F \otimes_R F) \cong H_i(F \otimes_R M)$  is a subquotient of  $F_i \otimes_R M$  and thus

$$(2.7) \quad \ell_R H_i(F \otimes_R M) \leq \ell_R(F_i \otimes_R M) = \text{rank}(F_i) \cdot \ell_R(M) = \beta_i(M) \cdot \ell_R(M).$$

Putting the inequalities (2.5), (2.6), and (2.7) together yields

$$2^d \cdot \ell_R(M) \leq \ell_R(M) \cdot \sum_i \beta_i(M),$$

and since  $\ell_R(M) > 0$ , we conclude  $\sum_i \beta_i(M) \geq 2^d$ .

Now suppose  $\sum_i \beta_i(M) = 2^d$ . Then the inequalities (2.5), (2.6), and (2.7) must actually be equalities, which means that  $H_i(S^2(F)) = 0$  for all odd  $i$ ,  $H_j(\Lambda^2(F)) = 0$  for all even  $j$ , and  $F \otimes_R M$  has trivial differential. Since

$H_0(\Lambda^2(F.)) \cong \Lambda^2(M)$  is the classical second exterior power,  $M$  must be cyclic, i.e., of the form  $R/I$  for some ideal  $I$ . Since  $F. \otimes_R R/I$  has trivial differential,  $I/I^2 \cong \text{Tor}_1^R(R/I, R/I)$  is free as an  $R/I$ -module, and thus a result of Ferrand and Vasconcelos (see [2, 2.2.8]) gives that  $I$  is generated by a regular sequence of elements. □

### 3. Rings of odd characteristic

In this section we prove part (2) of [Theorem 2](#). The main idea is to replace the Euler characteristic  $\chi$  occurring in the proof of part (1) with the *Dutta multiplicity*.

*Definition 3.1.* Assume  $(R, \mathfrak{m}, k)$  is a complete local ring of dimension  $d$  that contains  $\mathbb{Z}/p$  as a subring for some prime  $p$  and that  $k$  is a perfect field. For  $F. \in \text{Perf}^{\text{fl}}(R)$ , define

$$\chi_{\infty}(F.) = \lim_{e \rightarrow \infty} \frac{\chi(\varphi^e F.)}{p^{de}},$$

where  $\varphi^e$  denotes extension of scalars along the  $e^{\text{th}}$  iterate of the Frobenius endomorphism of  $R$ . The limit is known to exist by, e.g., [9, 7.3.3].

*Proof of Theorem 2 part (2).* There is a faithfully flat map  $(R, \mathfrak{m}, k) \rightarrow (R', \mathfrak{m}', k')$  of local rings such that  $\mathfrak{m} \cdot R' = \mathfrak{m}'$ ,  $R'$  is complete and  $k'$  is algebraically closed; see [5, 0.10.3.1]. Letting  $M' := M \otimes_R R'$ , we have that  $M'$  is a nonzero  $R'$ -module of finite length and finite projective dimension,  $\beta_i^{R'}(M') = \beta_i^R(M)$  for all  $i$ , and  $\dim(R') = \dim(R)$ . We may therefore assume  $R$  is complete with algebraically closed residue field.

Let  $F.$  be the minimal free resolution of  $M$ . Since  $R$  is complete with perfect residue field, a result of Roberts [9, 7.3.5] gives

$$(3.2) \quad \chi_{\infty}(F.) > 0$$

and a result of Kurano-Roberts [7, 3.1] gives (using (2.1))

$$(3.3) \quad \chi_{\infty}(S^2(F.)) - \chi_{\infty}(\Lambda^2(F.)) = \chi_{\infty}(\psi^2(F.)) = 2^d \cdot \chi_{\infty}(F.).$$

For each  $e \geq 0$ , we have  $\varphi^e S^2(F.) \cong S^2(\varphi^e F.)$  and  $\varphi^e \Lambda^2(F.) \cong \Lambda^2(\varphi^e F.)$ , and thus

$$\begin{aligned} \chi_{\infty}(S^2(F.)) &= \lim_{e \rightarrow \infty} \frac{1}{p^{de}} \sum_i (-1)^i \ell_R H_i(S^2(\varphi^e F.)), \\ \chi_{\infty}(\Lambda^2(F.)) &= \lim_{e \rightarrow \infty} \frac{1}{p^{de}} \sum_i (-1)^i \ell_R H_i(\Lambda^2(\varphi^e F.)). \end{aligned}$$

As in the proof of [Theorem 2.4](#), for a fixed  $e$ , we have

$$\begin{aligned} \frac{1}{p^{de}} \sum_i (-1)^i \ell_R H_i(S^2(\varphi^e F)) - \frac{1}{p^{de}} \sum_i (-1)^i \ell_R H_i(\Lambda^2(\varphi^e F)) \\ \leq \sum_j \ell_R H_j(T^2(\varphi^e F)). \end{aligned}$$

By [\[8, 1.7\]](#), the complex  $\varphi^e(F)$  is the minimal free resolution of the finite length module  $\varphi^e(M)$  for each  $e \geq 0$ . As in the proof of [Theorem 2.4](#), for each  $i$ , we have

$$\ell_R H_i(T^2(\varphi^e F)) \leq \text{rank}(\varphi^e F_i) \cdot \ell_R(\varphi^e M) = \beta_i(M) \cdot \chi(\varphi^e F).$$

We have proven that

$$\begin{aligned} \frac{1}{p^{de}} \sum_i (-1)^i \ell_R H_i(\varphi^e S^2(F)) - \frac{1}{p^{de}} \sum_i (-1)^i \ell_R H_i(\varphi^e \Lambda^2(F)) \\ \leq \frac{1}{p^{de}} \chi(\varphi^e F) \cdot \sum_i \beta_i(M) \end{aligned}$$

holds for each  $e \geq 0$ . Taking limits and using [\(3.3\)](#) gives

$$2^d \cdot \chi_\infty(F) \leq \chi_\infty(F) \cdot \sum_i \beta_i(M).$$

Since  $\chi_\infty(F) > 0$  by [\(3.2\)](#), we conclude  $\sum_i \beta_i(M) \geq 2^d$ . □

## References

- [1] M. K. BROWN, C. MILLER, P. THOMPSON, and M. E. WALKER, Cyclic Adams operations, *J. Pure Appl. Algebra* **221** no. 7 (2017), 1589–1613. [MR 3614968](#). [Zbl 1360.19006](#). <https://doi.org/10.1016/j.jpaa.2016.12.018>.
- [2] W. BRUNS and J. HERZOG, *Cohen-Macaulay Rings*, *Cambridge Stud. Adv. Math.* **39**, Cambridge University Press, Cambridge, 1993. [MR 1251956](#). [Zbl 0788.13005](#).
- [3] D. A. BUCHSBAUM and D. EISENBUD, Algebra structures for finite free resolutions, and some structure theorems for ideals of codimension 3, *Amer. J. Math.* **99** no. 3 (1977), 447–485. [MR 0453723](#). [Zbl 0373.13006](#). <https://doi.org/10.2307/2373926>.
- [4] H. GILLET and C. SOULÉ, Intersection theory using Adams operations, *Invent. Math.* **90** no. 2 (1987), 243–277. [MR 0910201](#). [Zbl 0632.14009](#). <https://doi.org/10.1007/BF01388705>.
- [5] A. GROTHENDIECK, *Éléments de Géométrie Algébrique. I. Le Langage des Schémas*, *Inst. Hautes Études Sci. Publ. Math.* **4**, 1960. [MR 0163908](#). [Zbl 0118.36206](#). Available at [http://www.numdam.org/item?id=PMIHES\\_1960\\_4\\_5\\_0](http://www.numdam.org/item?id=PMIHES_1960_4_5_0).
- [6] R. HARTSHORNE, Algebraic vector bundles on projective spaces: a problem list, *Topology* **18** no. 2 (1979), 117–128. [MR 0544153](#). [Zbl 0417.14011](#). [https://doi.org/10.1016/0040-9383\(79\)90030-2](https://doi.org/10.1016/0040-9383(79)90030-2).

- [7] K. KURANO and P. C. ROBERTS, Adams operations, localized Chern characters, and the positivity of Dutta multiplicity in characteristic 0, *Trans. Amer. Math. Soc.* **352** no. 7 (2000), 3103–3116. MR 1707198. Zbl 0959.13004. <https://doi.org/10.1090/S0002-9947-00-02589-7>.
- [8] C. PESKINE and L. SZPIRO, Dimension projective finie et cohomologie locale. Applications à la démonstration de conjectures de M. Auslander, H. Bass et A. Grothendieck, *Inst. Hautes Études Sci. Publ. Math.* no. 42 (1973), 47–119. MR 0374130. Zbl 0268.13008. Available at [http://www.numdam.org/item?id=PMIHES\\_1973\\_42\\_47\\_0](http://www.numdam.org/item?id=PMIHES_1973_42_47_0).
- [9] P. C. ROBERTS, *Multiplicities and Chern Classes in Local Algebra*, Cambridge Tracts in Math. **133**, Cambridge Univ. Press, Cambridge, 1998. MR 1686450. Zbl 0917.13007. <https://doi.org/10.1017/CBO9780511529986>.

(Received: April 7, 2017)

(Revised: May 22, 2017)

UNIVERSITY OF NEBRASKA, LINCOLN, NE

*E-mail*: [mark.walker@unl.edu](mailto:mark.walker@unl.edu)